



# Evolution of resistance in paediatric patients with failure on antiretroviral therapy

## Citation

Orrell, C., J. Levison, A. Ciaranello, L. Bekker, D. Kuritzkes, Kenneth Alan Freedberg, and R. Wood. 2012. Evolution of resistance in paediatric patients with failure on antiretroviral therapy. *Journal of the International AIDS Society* 15(Suppl 4): 18086.

## Published Version

doi:10.7448/IAS.15.6.18086

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10586353>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

## Oral Abstract – O225

# Evolution of resistance in paediatric patients with failure on antiretroviral therapy

Orrell, C<sup>1</sup>; Levison, J<sup>2</sup>; Ciaranello, A<sup>2</sup>; Bekker, L<sup>1</sup>; Kuritzkes, D<sup>3</sup>; Freedberg, K<sup>2</sup> and Wood, R<sup>1</sup>

<sup>1</sup>University of Cape Town, Desmond Tutu HIV Centre, Cape Town, South Africa. <sup>2</sup>Massachusetts General Hospital, Division of Infectious Diseases, Boston, USA.  
<sup>3</sup>Harvard Medical School, Boston, USA.

### Introduction

HIV-1 resistance data to inform treatment sequencing are limited for children with virological failure on first- and second-line antiretroviral therapy (ART) in Sub-Saharan Africa.

### Methods

HIV-1-infected children aged  $\leq 15$  years were retrospectively identified from an ART cohort in Cape Town, South Africa (2003 to 2010). First-line ART was either non-nucleoside reverse transcriptase inhibitor (NNRTI) or lopinavir/ritonavir-based (with the exception of children  $< 6$  months old who received full-dose ritonavir as the sole protease inhibitor (PI) from 2004 to 2007). Second-line ART was the alternative regimen. Treatment outcomes, including virological failure, loss to care, death or remaining in care, were determined. Genotypic resistance testing was conducted on stored serum from children at first- or second-line virological failure (two consecutive HIV-1 RNA levels  $> 1000$  copies/ml). International AIDS Society criteria defined resistance mutations.

### Results

Of 472 children starting first-line ART, 352 (75%) remained in care, 45 (9%) were lost and 4 (1%) died on first-line treatment. Seventy-one (15%) had observed virological failure, and 37 of these children had specimens available for genotype testing. Eight children (22%) had wild-type virus, seven (19%) had thymidine analog mutations (TAMs), 24 (65%) had NNRTI resistance and two (5.4%) had multiple protease resistance (PR). Of the 78 children who received second-line ART, 54 (71%) remained in care, 6 (8%) were lost and 1 (1%) died during second-line treatment. Fifteen (20%) had observed virological failure; 13 had samples available for genotype. Three (23%) had wild-type virus, eight (62%) had TAMs, nine (69%) had NNRTI resistance, and five (38%) had multiple PI resistance all of whom had received full-dose ritonavir.

### Conclusion

Although virological failure was infrequent in children on first- and second-line ART, rates of observed resistance including multiple PR resistance after failure were high. Reasons for high rates of resistance include use of full-dose ritonavir and continued viremia. Wild-type virus was common, suggesting poor adherence or challenges in correct dosing. Genotype resistance testing in children with virological failure may optimize selection of subsequent regimens and inform recommendations for sequencing of existing ART.

Published 11 November 2012

**Copyright:** © 2012 Orrell C et al; licensee International AIDS Society. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.