



# A closer look into DNase I hypersensitivity

## Citation

He, Housheng Hansen, Clifford A Meyer, Henry Long, X Shirley Liu, and Myles Brown. 2013. A closer look into dnase i hypersensitivity. *Epigenetics & Chromatin* 6(Suppl 1): P25.

## Published Version

doi:10.1186/1756-8935-6-S1-P25

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10646782>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



**POSTER PRESENTATION**

**Open Access**

# A closer look into DNase I hypersensitivity

Housheng Hansen He<sup>1,2,3\*</sup>, Clifford A Meyer<sup>1,3</sup>, Henry Long<sup>3</sup>, X Shirley Liu<sup>1,3</sup>, Myles Brown<sup>2,3</sup>

From Epigenetics and Chromatin: Interactions and processes  
Boston, MA, USA. 11-13 March 2013

DNase I hypersensitivity (DHS) combined with next generation sequencing (DNase-seq) is an efficient way of observing, in a single experiment, the genome-wide chromatin effects associated with the binding of multiple transcription factors. Using quantitative contrasts of DHS before and after estrogen and androgen stimulation in breast and prostate cancer cell lines, we have shown that differential DHS can accurately predict hormone induced transcription factor binding. Despite its effectiveness, the DHS assay can vary significantly depending on the experimental parameters. To increase the robustness of this assay, we have systematically evaluated two major parameters, digestion level and fragment size. We found that while there is a broad range of suitable digestion level, over-digestion dramatically decreases the efficiency of detecting DHS regions. More interestingly, we found that different fragment sizes capture distinct chromatin elements, and thus represent different chromatin structures. We were able to classify different combinations of estrogen receptor coregulators that resulted in different local chromatin structures.

#### Author details

<sup>1</sup>Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute and Harvard School of Public Health, Boston, MA 02115, USA.

<sup>2</sup>Department of Medical Oncology, Dana-Farber Cancer Institute and Harvard Medical School, Boston, MA 02115, USA. <sup>3</sup>Center for Functional Cancer Epigenetics, Dana-Farber Cancer Institute, Boston, MA 02215, USA.

Published: 18 March 2013

doi:10.1186/1756-8935-6-S1-P25

Cite this article as: He et al.: A closer look into DNase I hypersensitivity. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):P25.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute and Harvard School of Public Health, Boston, MA 02115, USA  
Full list of author information is available at the end of the article