



Myocardial infarct delineation in vivo using diffusion tensor MRI and the tractographic propagation angle

Citation

Mekkaoui, Choukri, Shuning Huang, Guangping Dai, Timothy G Reese, Jeremy Ruskin, Udo Hoffmann, Marcel P Jackowski, and David E Sosnovik. 2013. Myocardial infarct delineation in vivo using diffusion tensor mri and the tractographic propagation angle. *Journal of Cardiovascular Magnetic Resonance* 15(Suppl 1): P2.

Published Version

doi:10.1186/1532-429X-15-S1-P2

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11179847>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

POSTER PRESENTATION

Open Access

Myocardial infarct delineation *in vivo* using diffusion tensor MRI and the tractographic propagation angle

Choukri Mekkaoui*, Shuning Huang, Guangping Dai, Timothy G Reese, Jeremy Ruskin, Udo Hoffmann, Marcel P Jackowski, David E Sosnovik

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Delayed gadolinium enhancement (Gd-DE) is widely used to detect scar formation following myocardial infarction (MI) [1], but cannot be performed in patients with renal impairment. Here we use the tractographic propagation angle (PA), a novel index derived from 3D diffusion tensor MRI (DTI), to detect changes in myocardial fiber architecture post-MI [2]. We compare image segmentation based on the tractographic PA to infarct delineation with Gd-DE.

Methods

Normal human (n=5) and infarcted sheep hearts (n=6) were studied *ex vivo*. Infarcted mice (n=7) were imaged *in vivo*. MI was produced in C57BL6 mice via permanent ligation of the left coronary artery. *In vivo* DTI was performed on a 9.4T scanner (Bruker) using a 3D fat-suppressed single-shot 3D spin echo EPI sequence with motion-compensated diffusion-encoding gradients in 24 directions. Other parameters were: TR/TE=2000/13.5 ms, b-value 500-700 s/mm² and isotropic resolution of 280 μm. The human and sheep hearts were imaged on a clinical 3T Siemens scanner with an isotropic resolution of 2 mm³, TR/TE=8430/96 ms, and a b-value of 2000 s/mm². The tractographic propagation angle PA was defined as the angle between two adjacent principal eigenvectors (\hat{e}_{ij} , \hat{e}_{ij+1}) relative to a given fiber (Figure 1A). PA values were computed along myofiber trajectories within the principal eigenvector field using a 4th order Runge-Kutta integration method. Gd-DE imaging was performed 10min after the injection of 0.2mmol Gd-DTPA/kg. A short axis slice

through the infarcted myocardium was acquired using a cardiac-gated inversion recovery gradient echo sequence. Infarcted regions were segmented automatically on the Gd-DE images using a threshold of 2 standard deviations above normal. A PA threshold value greater than 4 degrees was used to automatically segment infarcted myocardium. Percent infarct size was calculated with both techniques and correlated.

Results

Tractography of a normal human heart color-coded by the PA is shown in Figure 1B. PA in the normal myocardium is highly homogeneous, averaging between 2 and 4 degrees. PA in the sheep infarct is significantly elevated and allows the infarct zone to be differentiated from the rest of the myocardium (Figure 1 C-D). Both PA (Figure 2A) and Gd-DE uptake (Figure 2B) were significantly increased in the infarct zone of all the mouse hearts imaged. A PA threshold of 4 degrees robustly segmented the infarct zone (Figure 2C), and an excellent correlation ($R^2=0.94$) was seen between percent infarct size by Gd-DE and tractographic PA (Figure 2D).

Conclusions

PA detects the loss of tract coherence in infarcted myocardium and robustly delineates myocardial infarcts *in vivo*. The use of DTI, and hence the tractographic PA, does not require exogenous contrast and can be performed in all patients regardless of renal function. The technique provides a complementary and valuable adjunct to Gd-DE.

Funding

R01HL093038.

Radiology, Harvard Medical School - Massachusetts General Hospital, Boston, MA, USA




Figure 1 Tractograms color-coded by the propagation angle (PA). (A) PA is defined as the angle between two adjacent principal eigenvectors (\hat{e}_{ij} , \hat{e}_{ij+1}) relative to a given myofiber. (B) Normal human heart viewed from the base, showing a low and homogenous PA. (C) Sheep heart with a large anteroseptal infarct. (C1) A low-pass PA value of 4 degrees delineates the normal myocardium and creates a void in the infarct. (C2) Conversely, a high-pass PA value of 4 degrees robustly delineates the infarcted myocardium.




Figure 2 *In vivo* PA maps in infarcted mice. (A) PA map in a mouse with a large anterolateral infarct. (B) Delayed enhancement image at the corresponding level. It should be noted that the PA maps were acquired in mid-systole and the delayed enhancement images in mid-diastole. (C) Segmentation of the PA map using a threshold value of 4 degrees robustly segments normal from infarcted myocardium. (D) A high correlation ($R^2=0.94$) between infarct size calculated from the *In vivo* PA and infarct size measured by delayed gadolinium enhancement was obtained.

Published: 30 January 2013

References

1. Kim R, et al.: *NEJM* 2000.
2. Mekkaoui C, et al.: *ISMRM* 2011.

doi:10.1186/1532-429X-15-S1-P2

Cite this article as: Mekkaoui et al.: Myocardial infarct delineation *in vivo* using diffusion tensor MRI and the tractographic propagation angle. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):P2.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

