



Exploring the limit of metazoan thermal tolerance via comparative proteomics: thermally induced changes in protein abundance by two hydrothermal vent polychaetes

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Dilly, G. F., C. R. Young, W. S. Lane, J. Pangilinan, and P. R. Girguis. 2012. "Exploring the Limit of Metazoan Thermal Tolerance via Comparative Proteomics: Thermally Induced Changes in Protein Abundance by Two Hydrothermal Vent Polychaetes." <i>Proceedings of the Royal Society B: Biological Sciences</i> 279, no. 1741: 3347–3356.
Published Version	doi:10.1098/rspb.2012.0098
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12763599
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP

1 Exploring the limit of metazoan thermal tolerance via comparative proteomics: Thermally
2 induced expression shifts in two hydrothermal vent polychaetes

3

4 **Geoffrey F Dilly¹, C Robert Young², William S Lane³, Jasmyn Pangilinan⁴, Peter R**

5 **Girguis^{1*}**

6

7 **1 = Harvard University,**

8 **Department of Organismic and Evolutionary Biology,**

9 **16 Divinity Avenue Rm 3085,**

10 **Cambridge, MA 02138.**

11

12 **2 = CEEBE, MIT, 15 Vassar St, Cambridge, MA, 02139**

13 **3 = FAS Center for Systems Biology, Harvard University, Cambridge MA 02138**

14 **4 = Genomic Technologies, DOE JGI, 2800 Mitchell Drive Bldg, Walnut Creek, CA, 94598**

15

16 *** = send correspondence to pgirguis@oeb.harvard.edu**

17

18

18

19 **Abstract**

20 Eukaryotic thermotolerance is challenged at deep-sea hydrothermal vents, where temperatures
21 can reach 300 °C. *Paralvinella sulfincola*, an extremely thermotolerant vent polychaete, and *P.*
22 *palmiformis*, a congener with a more modest thermal tolerance, both flourish at vents along the
23 Juan de Fuca Ridge, Washington, USA. We conducted a series of shipboard, high-pressure,
24 thermotolerance experiments on both species to examine the physiological adaptations that
25 confer pronounced thermotolerance. Quantitative proteomics, a deeply sequenced EST library,
26 and glutathione (an antioxidant) assays revealed that *P. sulfincola* exhibited an upregulation in
27 the synthesis and recycling of GSH with increasing temperature, downregulated NADH and
28 succinate dehydrogenases (key enzymes in oxidative phosphorylation) with increasing
29 temperature, but maintained elevated levels of heat shock proteins (HSPs) across treatments. In
30 contrast, *P. palmiformis* exhibited more typical responses to increasing temperatures, e.g.
31 increasing HSPs at higher temperatures. These data, among the first to quantify global protein
32 and antioxidant responses to temperature in an extremely thermotolerant eukaryote, suggest that
33 *P. sulfincola*'s pronounced thermal tolerance is largely due to its capacity to mitigate oxidative
34 stress via increased synthesis of antioxidants and decreased flux through the mitochondrial
35 electron transport chain. Ultimately oxidative stress may be the key factor in limiting all
36 metazoan thermotolerance.

37 **Keywords:** Proteomics, Hydrothermal vents, Thermotolerance, Oxidative stress, *Paralvinella*

38

39 Introduction

40 Physiological adaptations to thermal stress are ubiquitous among all organisms. While
41 prokaryotes have a known upper thermal limit of at least 122 °C (1), metazoans have a much
42 lower thermal tolerance, with 45 to 47 °C as the currently accepted upper limit of homeostasis
43 (2), though unicellular eukaryotic fungi are known to grow at 60-62 °C (3). Mitochondrial
44 dysfunction (4), membrane instability, structural disintegration (2), limitations in gas transport
45 and mitochondrial dysfunction have all been implicated as possible modes of physiological
46 failure in eukaryotes (5-7).

47 There have been numerous studies to date on metazoan thermotolerance (for reviews see:
48 (8, 9)). The few that have focused on highly thermotolerant animals such as desert ants and hot
49 spring ostracods have largely examined their response to acute thermal exposure (10, 11).

50 Recently, some studies have employed proteomics to examine responses to thermal stress in
51 mesotolerant animals (6, 12); however, there remains a limited amount of biomolecular data for
52 extremely thermotolerant metazoans (13). Specifically, it remains to be determined how highly
53 thermotolerant organisms respond to chronic thermal exposure, and which physiological or
54 biochemical adaptations enable them to ameliorate physiological perturbations that arise at
55 higher temperatures.

56 Deep-sea hydrothermal vents are ideal habitats to address such questions, as these
57 environments are home to some of the most thermotolerant animals known. This includes the
58 polychaetes *Alvinella pompejana* and *Paralvinella sulfincola*. To date, numerous studies have
59 investigated the thermal tolerance of *A. pompejana*, beginning with the observation that *A.*
60 *pompejana* lives upon 81° C substratum (14). Subsequent to that, and in contrast to the in situ

Geoffrey Dilly 12/28/11 11:17 PM

Comment: Reviewer 2, Q1

Geoffrey Dilly 12/28/11 11:18 PM

Comment: Refers to Reviewer 1 Q3

61 observations, *in vitro* research on *A.pompejana* has suggested key enzymes and structural
62 components are not stable after chronic exposure to elevated temperatures (15). A recent study of
63 *A. pompejana* protein expression via 2D gel electrophoresis compared physiological responses to
64 different oxygen concentrations (13), though its response to chronic exposure remains
65 unconstrained. Notably, *A. pompejana* are not easily amenable to *in vivo* experimentation (16),
66 making it difficult to address chronic thermal tolerance in this species.

67 *Paralvinella sulfincola* is another highly thermotolerant polychaete that thrives on
68 hydrothermal sulfides in the Northwest Pacific. *P. sulfincola* are found on sulfide where
69 temperatures reach 88.5°C (17), and *in vivo* laboratory studies of *P. sulfincola* have
70 experimentally demonstrated the broadest known range of chronic thermal tolerance in
71 metazoans (5-48 °C) (18, 19) and (electronic supplementary material –ESM– Figure S1). *P.*
72 *palmiformis* - a closely related congener - is also found in these environs but exhibits markedly
73 different thermal tolerances (ESM Figure S1). Both are amenable to *in vivo* recovery and
74 laboratory experimentation, which affords the unique opportunity to elucidate the biochemical
75 responses of meso- and thermotolerant metazoans in a comparative phylogenetic context.

76 To better understand the biochemical mechanisms that underlie extreme thermal
77 tolerance, we present data from a series of *in vivo* high-pressure laboratory experiments in which
78 we examined quantitative changes in protein expression of live *P. sulfincola* and *P. palmiformis*
79 over their thermal range, including temperatures near each species' ultimate incipient lethal
80 temperature (UILT, defined here as the temperature beyond which 50% of the population cannot
81 survive indefinitely (20, 21)). These data reveal statistically significant differences in protein
82 abundance and upregulation between these two congeners, related to mitigating antioxidant
83 stress across their thermal ranges and at their respective UILTs. These data further reveal key

84 differences in antioxidant concentrations in each species. The results of this study provide the
85 first direct empirical evidence that oxidative stress may be the primary stressor at *P. sulfincola*'s
86 upper temperature limit, and illustrates the means by which *P. sulfincola* mitigates this stress.

87 **Results and Discussion**

88 The data herein comprise A) the first extensive assessment of *P. sulfincola* and *P.*
89 *palmiformis* chronic thermal tolerance; B) a thorough interrogation of their proteomes at chronic,
90 environmentally relevant temperatures using quantitative, high-throughput mass spectrometric
91 sequencing, and C) a comparison of antioxidant production between the two congeners under
92 thermal stress. A normalized expressed sequence tag (EST) library served as the database for the
93 proteomic analyses (due to the qualitative nature of these EST data, as well as the explicit focus
94 of this study on quantitative differences in expression, all data shown here are from the
95 proteomic analyses unless otherwise noted). Together these data reveal that *P. sulfincola* and *P.*
96 *palmiformis* exhibit overlap in their thermal tolerance ranges, possess markedly different
97 tolerances at their upper and lower bounds, and employ different physiological “strategies” to
98 mitigate thermal stress. Near its UILT, *P. sulfincola* maintains elevated expression of heat shock
99 proteins (HSPs) across its thermal range, rapidly resynthesizes reduced glutathione, and likely
100 decreases oxidative phosphorylation to mitigate the impact of oxygen radicals. In contrast, *P.*
101 *palmiformis* exhibited responses to chronic thermal exposure that are more similar to those
102 observed in previous studies of mesotolerant organisms, including increased representation of
103 heat shock proteins and other systems solely upon exposure to their highest chronic thermal
104 regimes.

Geoffrey Dilly 12/20/11 12:29 PM

Comment: Reviewer 1 Q1

105 While we cannot infer metabolic flux from these data (discussed below), the observed
106 systemic differences elucidate those physiological and biochemical processes most responsive to
107 thermal stress. The data suggest that the upper temperature limits of metazoan life may indeed be
108 governed by the ability of the organism to mitigate oxidative stress by managing antioxidant
109 production and vital energy yielding metabolic pathways. The sections below discuss in greater
110 detail the observed differences in protein and antioxidant expression between these two sister
111 taxa.

112 *Differences in Expression of Molecular Chaperones*

113 Molecular chaperones such as heat shock proteins (HSPs) mitigate thermal stress
114 by minimizing protein dysfunction through catalyzing nascent protein folding in the endoplasmic
115 reticulum (ER), reforming misfolded proteins, as well as other functions (22, 23). While many
116 chaperones are constitutively expressed, a large number of chaperones are up-regulated during
117 periods of cellular stress, so-called inducible forms (24). A total of 27 chaperones and co-
118 chaperones were examined in our analysis, representing members of all detected heat shock
119 proteins. Key protein families are discussed in the paragraphs below, and their representation and
120 Bayesian significance are presented in Table 1a, ESM S1 and S2). Briefly, we observed that *P.*
121 *sulfincola* exhibited elevated levels of all major chaperones, even those previously categorized as
122 inducible, over all treatments (Figure 1), while *P. palmiformis* exhibited higher chaperone
123 production primarily near the UILT.

124 Heat shock protein 70 (HSP70)

125 The 70 KDa heat shock proteins (HSP70 family) are the first characterized and best
126 understood chaperones, and are highly conserved across domains of life (23). Multiple isoforms
127 in the family are constitutive, while others are induced by heat stress (24, 25). In *P. sulfincola*,
128 GRP75 proteins, a member of the HSP70 family, exhibited the highest abundance of all
129 molecular chaperones across all *P. sulfincola* treatments. GRP75 is homologous to the human
130 HSPA9, a constitutive mitochondrial HSP (26). In contrast, GRP75 expression in *P. palmiformis*
131 was comparable across many treatments (though there was a moderate increase in expression of
132 GRP75 at 38°C relative to the cooler thermal regimes). A number of co-chaperones that interact
133 with HSP70 family were also observed in all proteomes, and though their expression varied the
134 overall trend for both species was a slight increase in the high treatments.

135 When *P. sulfincola* peptide sequences were compared against the broader NCBI non-
136 redundant protein database (ESM table S3), peptides homologous to inducible HSP70s were
137 detected, and their relative proportion to total protein remains consistent with constitutive HSP70
138 proteins. Moreover, in our *P. sulfincola* EST library, three additional HSP70 sequences with
139 human homologs were recovered, though due to their absence in our proteome data are not
140 considered in the quantitative analyses. Together these data underscore the importance of HSP70
141 proteins in thermal tolerance of both species, and the continued elevated expression in *P.*
142 *sulfincola* suggest that HSP70 proteins may be kept abundant to cope with the rapid changes in
143 temperature typically encountered by this species, which includes maintaining physiological
144 function near the organism's UILT.

145 HSP90

Geoffrey Dilly 12/20/11 12:29 PM
Comment: Next 6 lines refer to Reviewer 1 Q1

146 Although less well characterized than the HSP70 family, HSP90s are known as flexible
147 dimer ATPases that bind to a variety of cellular proteins including steroid hormone receptors,
148 transcription factors, and protein kinases (27, 28). The HSP90 protein GRP94 (a luminal protein
149 associated with the endoplasmic reticulum (29)) was detected in the *P. sulfincola* proteome,
150 exhibiting constitutive expression across all treatments (Probability of differential expression -
151 Pr(DE) 0.11). GRP94 was also observed in the *P. palmiformis* proteome, and its abundance
152 likely increased with temperature (12°C → 38°C - log 1.55, Pr(DE) 0.66). Co-chaperones such
153 as HOP, FKBP52 and others known to play a regulatory role with cytosolic HSP90s were
154 observed in both *P. sulfincola* and *palmiformis* proteomes. FKBP52 exhibited a highly
155 significant increase with temperature in both worms (Pr(DE) 1.00). HOP, which modulates
156 HSP70/90 interactions, was also upregulated with temperature in both *P. sulfincola* and *P.*
157 *palmiformis* at their highest treatments (*P.s.* 10°C → 45°C - log 0.98, Pr(DE) 0.82; *P.p.* 12°C →
158 38°C - log 0.98, Pr(DE) 0.51). The HSP90 activator AHA1 was substantially upregulated at
159 45°C in *P. sulfincola* (10°C → 45°C - log 3.58, Pr(DE) 1.00) but not in *P. palmiformis*. Notably,
160 the HSP90 inhibitor CDC37 remained constant in *P. sulfincola* and significantly decreased in *P.*
161 *palmiformis* in higher thermal regimes. The patterns observed here suggest that HSP90 is
162 constitutively expressed in *P. sulfincola*, but activity is regulated in both species through the
163 regulation of activators and inhibitors. These observations are also consistent with the
164 aforementioned hypothesis that *P. sulfincola* maintains a biochemical poise to cope with acute
165 temperature fluctuations.

166 HSP 60 and HSP27

167 HSP60 is a *mitochondrial* molecular chaperone known to confer thermal tolerance in
168 eukaryotes (30). Our analysis revealed that HSP60 was the most consistently expressed heat
169 shock protein, with high abundance across all treatments in both species. This trend was mirrored
170 in the HSP60 co-chaperone, HSP10, which assists HSP60 in protein folding during periods of
171 stress (31). These findings suggest that both species maintain pools of HSP60 and HSP10 to
172 mitigate damage to mitochondrial proteins.

173 The small 27kDa heat shock protein (sHSP), found throughout cellular compartments and
174 the cytosol, responds to both thermal and oxidative stress by binding to damaged or misfolded
175 proteins and forming reservoirs for other chaperones to correctly refold or initiate proteolytic
176 degradation (32). It is also known to upregulate key enzymes in the glutathione pathway (32, 33).
177 HSP27 was abundant across all treatments in *P. sulfincola*. However, HSP27 increased only at
178 the highest temperature in *P. palmiformis* (12°C → 38°C - log 2.09, Pr(DE) 1.00). We posit that
179 the differences observed between expression levels of HSP27 relate to oxidative stress response
180 and the glutathione pathway (discussed in detail below).

181 Foldases

182 Foldases are enzymes that catalyze rate-limiting steps in protein folding, many of which
183 play a key role in the cellular “unfolded protein response” (a stress response to an accumulation
184 of unfolded and misfolded proteins in the endoplasmic reticulum, which aims to restore normal
185 function by halting protein translation and signaling the production of molecular chaperones
186 involved in protein folding; (34)). Foldases important to the UPR were detected in both species.
187 Of note, the foldase PDIA1, a protein-thiol oxidoreductase that acts as both a chaperone and a
188 foldase (34, 35), was abundant (constitutive) across all treatment in *P. sulfincola* (Pr(DE) 0.001).

189 In *P. palmiformis*, PDIA1 abundance increased as a function of temperature (Pr(DE) 0.914),
190 reinforcing the pattern of differential response observed between these two organisms in relation
191 to thermal stress.

192 While the data on chaperones demonstrate that *P. sulfincola* maintains elevated
193 expression of chaperones across all thermal regimes, we posit that the representation and
194 abundance of chaperones does not itself explain the observed thermotolerance. Indeed, if HSP
195 abundance alone was the key factor in conferring extreme thermotolerance, then *P. palmiformis*
196 would likely have a greater thermal tolerance similar to *P. sulfincola* (with a UILT above 38°C)
197 because the representation of chaperones between these two closely related species was
198 (proportionally) equivalent at their respective highest thermal treatments. We therefore further
199 posit that elevated HSP abundances in *P. sulfincola* are more likely a reflection of its ecological
200 niche *in situ*, enabling it to survive acute, rapid shifts in temperatures caused by its proximity to
201 hot vent fluid, but does not alone explain their chronic thermal tolerance.

202 *Response to Oxidative Stress*

203 The largest shifts in protein abundance observed in both species are related to the
204 mitigation of oxidative stress. In mitochondria, the reactive oxygen species superoxide ($O_2^{\cdot-}$) is
205 generated in complexes I/III during respiration, and other ROS such as the hydroxyl radical
206 (HO^{\cdot}), and uncharged hydrogen peroxide (H_2O_2), are produced in the outer and inner membranes
207 (for review see (36, 37)). Under normative conditions, mitochondria consume more than 90% of
208 all cellular O_2 , while also producing the majority of ROS (38). However, studies have shown that
209 elevated temperatures can also increase oxidative stress in mesotolerant eukaryotes (5, 39, 40), as

210 elevated temperatures increase the metabolic demand of tissues, induce a state of functional
211 tissue hypoxia, and increase mitochondrial respiration rates (41).

212 Superoxide dismutase (SOD, EC 1.15.1.1) is a ubiquitous enzyme that is responsible for
213 catalyzing the reduction of $O_2^{\cdot-}$ to H_2O_2 . There are two forms of this metalloprotein; Cu/Zn SOD
214 (isotig03775) are primarily found in the cytosol, whereas Mn SOD (isotig06674) are located the
215 mitochondria (42). *P. sulfincola* showed no differences in the abundance of either SOD across all
216 treatments, but *P. palmiformis* exhibited significant increases in both Mn SOD (Pr(DE) 0.997)
217 and Cu/Zn SOD (Pr(DE) 0.999) (ESM tables S1, S2).

Peter Girguis 12/28/11 11:21 PM
Comment: Reviewer 1 Question 2C

218 In *P. sulfincola*, however, the production of glutathione appears to play a prominent role
219 in mitigating ROS. Glutathione (L- γ -glutamyl-L-cysteinylglycine, or GSH) is a tripeptide thiol
220 that is the primary nonprotein antioxidant in metazoans. Found in up to mM concentrations in
221 mammals, GSH mitigates oxidative stress by chemically reducing hydrogen peroxide and other
222 toxic compounds (36, 43). The enzyme glutathione peroxidase (GPx, 1.11.1.9) catalyzes this
223 reduction, yielding glutathione disulfide (GSSG). GSSG is reverted back to GSH by glutathione
224 reductase (GSR, EC 1.8.1.7). Regulation of GSH metabolism and resynthesis serves as an
225 indicator of cellular oxidative stress levels (43). As cysteine is the required peptide for *de novo*
226 GSH synthesis, and the rare amino acid selenocysteine is required for the synthesis of
227 glutathione peroxidase, increases in cysteine and in particular selenocysteine are good indicators
228 for increases in GSH cycling (Table 1b, ESM Tables S1 and S2).

229 Figure 2 depicts key steps and significant changes over temperature in the synthesis of
230 glutathione, the redox cycle of GSH and GSSG, and the catalyzing enzymes glutathione
231 peroxidase (GPx) and glutathione reductase (GSR) in *P. sulfincola* and *P. palmiformis*. Notably,

232 cystathionine beta-synthase (CBS, EC 4.2.1.22), central to both cysteine and selenocysteine
233 synthesis, exhibited the single largest fold increase with temperature of all proteins assayed in *P.*
234 *sulfincola* and nearly so for *P. palmiformis* (*P.s.* 10°C → 45°C - log 5.74, Pr(DE) 1.00; *P.p.*
235 12°C → 38°C - log 5.55, Pr(DE) 1.00). Two ATP-dependent, rate-governing steps within the
236 glutathione pathway were detected in our *P. sulfincola* and *P. palmiformis* proteomes: selenide
237 water dikinase (selD, EC 2.7.9.3), and gamma-glutamylcysteine synthetase (GCS, EC 6.3.2.2)
238 (Figure 2). SelD, essential for *de novo* synthesis of selenoproteins, increased in *P. sulfincola* at
239 both 30°C and 45°C, while no differences in abundance were observed in *P. palmiformis*. GCS,
240 the rate-limiting step in the production of GSH and subject to feedback inhibition (43), showed a
241 steady increase in abundance with temperature in *P. sulfincola*. In *P. palmiformis*, however, GCS
242 was not detected until 38°C treatment, producing a significant correlation with temperature
243 (12°C → 38°C - log 3.81). These data clearly suggest that GSH is being synthesized at higher
244 rates in response to increasing thermal stress in both species, though far more pronounced in *P.*
245 *sulfincola*.

246 Glutathione peroxidase 3 (GPx-3, cytosolic) showed significant increases in abundance at
247 both the medium and high temperature treatments in *P. sulfincola*, as well as at the highest
248 temperature treatment in *P. palmiformis*. Notably, *P. sulfincola* significantly increases its GSR
249 protein abundance while *P. palmiformis* significantly decreases it at higher temperatures. These
250 data suggest that *P. sulfincola* is continuously recycling GSH in the mitochondria. We further
251 suggest that the differences may be indicative of mitochondrial dysfunction and uncoupling in *P.*
252 *palmiformis*, possibly due to lipid peroxidation from increasing ROS activity, as has previously
253 been observed in cold-water marine mollusks exposed to heat stress and functional hypoxia (5,
254 38).

255 To further investigate the effect of thermal and oxidative stress on the pool of GSH, total
256 GSH (GSht) levels and GSH/GSSG ratios (the ratio of the reduced and oxidized forms) were
257 measured for medium and high temperature treatments in both species (ESM Figure S2). GSht
258 concentrations in *P. sulfincola* were about half those observed in *P. palmiformis*. There were no
259 measurable differences in the GSH/GSSG ratio among *P. sulfincola* worms across all thermal
260 treatments. However, in higher thermal treatments, *P. palmiformis* exhibited a 2-fold decrease in
261 the pool of GSht. Furthermore, the GSH/GSSG ratio in *P. palmiformis* exhibited more than a 3-
262 fold drop at higher thermal treatments, indicating that *P. palmiformis* were not able to effectively
263 recycle glutathione at 38°C. These trends suggest that *P. sulfincola* is well poised to sustain GSH
264 resynthesis near its UILT, allowing it to maintain functionality even under periods of high
265 oxidative stress (the limited sample size prohibited statistical analyses of these observations).

266 *Oxidative Stress and Oxidative Phosphorylation*

267 In eukaryotes, oxidative phosphorylation within the electron transport chain is
268 responsible for the majority of ATP production and ROS formation. As mentioned, research has
269 indicated that elevated temperature can lead to local tissue hypoxia (41). Here, *P. sulfincola*
270 exhibited a significant reduction in abundance of NADH dehydrogenase (10°C → 45°C - log -
271 2.01) and succinate dehydrogenase (10°C → 45°C - log -1.00), both of which are involved in the
272 mitochondrial oxidative phosphorylation (Tables S1 and S2). Indeed, a large portion of ROS is
273 generated by NADH dehydrogenase (complex I). Succinate dehydrogenase (complex II) may not
274 contribute directly to ROS formation, but it funnels electrons to complex III, which does produce
275 ROS. In *P. palmiformis*, the decrease in NADH dehydrogenase was less pronounced (12°C →
276 38°C - log -0.48), and there was an increase in abundance of succinate dehydrogenase with
277 temperature (12°C → 38°C - log 1.53). The observed patterns of NADH dehydrogenase

278 contrasts with previous studies of a heat sensitive mussel species *Mytilus galloprovincialis*, ,
279 which increased production of NADH dehydrogenase relative to its less thermotolerant congener
280 *Mytilus trossulus* (6). These data suggest that *P. sulfincola* may be actively repressing ROS
281 formation at high temperatures by lessening endogenous generation *via* the ETC, and depending
282 more heavily on anaerobic respiration at elevated temperatures.

Geoffrey Dilly 1/3/12 11:39 PM
Comment: Reviewer 2 – Question 1

283 *Global proteome responses, emerging hypotheses and future directions*

284 Quantitative mass spectrometric protein analyses reveal hundreds of differentially
285 expressed proteins per treatment, yet efforts to ally proteomic (or transcriptomic) data to
286 metabolic rate have met with limited success (44, 45). This is likely attributable to the
287 complexity of interactions among enzymes, their substrates and other factors that regulate flux
288 through a pathway. Gross changes among metabolic pathways, however, provide another –albeit
289 coarser- means of assessing organismal response to thermal stress as it reveals broad trends in
290 the abundance of proteins allied to specific systems. iPath (46) was used to map changes in
291 global protein abundance within 139 KEGG metabolic pathways, and reveal significant
292 (posterior probability <0.05), broad and complex differences in protein expression between
293 species and among treatments (ESM Figure S3a-d, S4). Protein upregulation in *P. palmiformis*
294 between the 21°and 38°C treatments is significantly higher than in other treatments. It is
295 possible that *P. palmiformis* is incapable of maintaining homeostasis at the higher temperature,
296 and is exhibiting metabolic disorder, as evident by the changes in the TCA and pentose
297 phosphate cycle, both known to respond to thermal stress (ESM Figure S5) (6, 12). At their
298 highest temperature treatments, *P. sulfincola* and *P. palmiformis* exhibited opposing patterns of
299 protein expression in the TCA cycle, with *P. sulfincola* decreasing and *P. palmiformis* increasing

Geoffrey Dilly 1/3/12 11:55 PM
Comment: I am addressing Reviewer 1 question 7 here.

300 expression of enzymes respectively. In the pentose phosphate pathway, *P. sulfincola* and *P.*
301 *palmiformis* again exhibited opposing patterns of expression, exhibiting increased and decreased
302 enzymes respectively. Depression of the TCA cycle in *P. sulfincola* may be due to thermal
303 effects on energy metabolism, or may be attributable to the oxygen concentrations in our
304 experiments, which at 130 μ M are comparable to ambient bottom water but higher than some
305 diffuse flows (discussed below). Nevertheless, these trends are consistent with a decreased
306 emphasis on aerobic respiration (TCA cycle) and the need for reducing equivalents to maintain
307 sufficient GSH for antioxidant activity (pentose can be converted into glucose 6-phosphate to
308 produce NADPH to recycle oxidized GSH (47)). Further targeted studies may better reveal
309 correlations between flux rates and protein counts, helping our understanding of the effects of
310 thermal stress on metabolic processes. In addition, future studies should also consider the impact
311 of post-translational protein modifications to physiological functions at elevated temperatures.

Peter Girguis 12/28/11 11:22 PM

Comment: Answers reviewer 2 question 2

312 **Conclusions**

313 These data lead us to conclude that *P. sulfincola* maintains a pool of both canonical
314 constitutive and inducible heat shock proteins to maintain protein function during rapid and
315 frequent exposure to high temperatures in its highly dynamic environment. Notably, enzymes
316 and pathways associated with the production of antioxidants showed the most pronounced
317 response to thermal exposure in both *P. sulfincola* and *P. palmiformis*. However, only *P.*
318 *sulfincola* increased the abundance of enzymes responsible for chemically reducing GSSG,
319 thereby maintaining its antioxidative capacity. Increasing the *de novo* synthesis of GSH from the
320 cysteine pathway (and the catalytic enzyme GPx through increases in selenocysteine) at elevated
321 temperatures further underscores this pathway's relevance in oxidative scavenging. Increased

322 production of GSR, necessary for recycling GSSG also demonstrates that *P. sulfincola* maintains
323 a sufficient pool of GSH to mitigate oxidative stress. The concurrent decreases in *P. sulfincola*
324 enzymes associated with oxidative phosphorylation may reduce the rate of oxidative radical
325 formation at high temperature.

326 In contrast, *P. palmiformis* exhibited significant increases in major molecular chaperones
327 with increasing temperature, and increases in other systems including the production of
328 antioxidants. However, *P. palmiformis* did not exhibit a comparable capacity to regenerate GSH
329 at elevated temperatures, or reduce the production of ROS from oxidative phosphorylation.
330 Indeed, *P. palmiformis* responded in a manner similar to other comparatively mesotolerant vent
331 endemics such as *Paralvinella grasslei* (16) and *Rimicaris exoculata* (48).

332 *P. sulfincola* lives on vent edifices, where they might readily encounter regions of
333 elevated temperature and varying oxygen concentration (due to radiative heating, the water
334 around vent sulfides can be warm yet exhibit a composition more similar to the ambient seawater
335 (49)). In addition, *P. sulfincola* are exposed to more sulfidic fluids as well, which might also
336 affect their thermal tolerance due to the impact of sulfide on aerobic respiration (addressing the
337 role of sulfide in thermal tolerance is beyond the scope of this study, and should be revisited in
338 future investigations). Based on the data presented here, we posit that *P. sulfincola*'s pronounced
339 thermotolerance is enabled primarily by adaptations to mitigate oxidative stress, which include
340 increasing activity of antioxidant systems and decreasing aerobic metabolism. We further
341 suggest these patterns demonstrate that managing ROS, resulting from increased mitochondrial
342 aerobic respiration at elevated temperatures, is a high priority for thermotolerant organisms.
343 Considering that all metazoans are ultimately dependent on mitochondrial aerobic respiration,
344 ROS may effectively limit them to cooler thermal regimes than thermophilic bacteria and

Geoffrey Dilly 12/20/11 12:29 PM
Comment: Next 10 lines address Reviewer 1
Question 2b

345 archaea (the most thermophilic prokaryotes are anaerobes, and exhibit a striking antioxidant
346 response when exposed to modest amounts of oxygen (50)). Although oxidative stress has been
347 implicated in previous studies on mesophilic eukaryotes (2, 5, 6, 41), this is the first study to
348 empirically derive this link between the UILT and ROS production in one of the most
349 thermotolerant metazoans on the planet, suggesting that oxidative stress -not temperature itself-
350 may limit metazoan thermal tolerance.

351 **Materials and Methods**

352 *Animal collection and experimental apparatus*

353 *Paralvinella sulfincola* and *Paralvinella palmiformis* “palm worms” were collected from
354 hydrothermal vents in the Main Endeavour field along the Juan de Fuca Ridge (47°57'N,
355 129°5'W) at a depth of 2,200m during the R/V *Atlantis* expedition 15-34 in July 2008.
356 Organisms were collected by the DSV *Alvin* on dives #4409-4423, using either a multi-chamber
357 suction sampler or an insulated sample recovery box. Upon recovery to 1 atm, worms were
358 transferred to a 4°C cold room and visually sorted based on segment number and gill
359 morphology. Aggregations of mucus and minerals were removed from the animals before
360 transfer into a flow-through high pressure aquaria system.

361 *High-pressure aquaria system*

362 *P. sulfincola* incubations were conducted in a newly designed high pressure aquaria
363 system (ESM Figure S6). *P. palmiformis* low temperature incubations (12°C) were conducted in
364 a 500 cm³ titanium flow-through system with 50 mL/min flow rate (51). Dissolved oxygen
365 concentration was measured at the inlet and outlet of each system using a polarigraphic oxygen

Geoffrey Dilly 12/20/11 12:29 PM

Comment: Reviewer 3 Q1

366 electrode (limits of detection ca. 1 μM ; YSI Inc) to verify that oxygen was always greater than
367 25 μM , which is not limiting based on known hemoglobin oxygen binding affinities of
368 alvinellids (52). Dissolved oxygen concentrations were achieved by equilibrating the seawater
369 with air, at the elevated experimental temperatures and 1 ATM. The net effect was a dissolved
370 oxygen concentration of $\sim 130 \mu\text{M}$, which is slightly higher but comparable to the ambient
371 bottom water concentration at the Juan de Fuca ridge ($\sim 120 \mu\text{M}$).

Geoffrey Dilly 12/20/11 12:29 PM

Comment: Addresses Reviewer 1 Question 2A

372 *Experimental design*

373 Though critical thermal maxima (CT_{max}) of both species and thermal preference of *P.*
374 *sulfincola* were previously examined (18, 19), we augmented these data to better establish *P.*
375 *sulfincola* and *P. palmiformis* chronic thermal tolerance (ESM Figure S1). A total of 85 *P.*
376 *sulfincola* and 108 *P. palmiformis* were utilized in this study. Chronic thermal tolerance was
377 defined as a lack of temperature-induced mortality over 12 hours of sustained exposure. On
378 occasion, $<5\%$ of individuals died during treatments, which upon further inspection we attributed
379 to recovery and handling. Based on these data, three temperatures were chosen that span the
380 chronic thermal tolerance range of each species (*P. sulfincola* = 10°C, 30°C, and 45°C; *P.*
381 *palmiformis* = 12°C, 21°C, and 38°C; ESM Figure S1). These temperatures, although not
382 identical in their ΔT , were chosen to represent the organisms' protein profiles across their
383 respective thermal tolerance ranges, including temperatures approaching their UILT. We posit
384 that the resulting data better represents protein abundance and their lower, nominal, and upper
385 temperature regimes. At each treatment, six to nine worms were maintained at constant pressure
386 and temperature for >12 hours for global protein expression analysis. To minimize the effects of
387 collection and handling, worms were first acclimated in each system at room temperature (21°C)

Peter Girguis 12/28/11 11:24 PM

Comment: Answers reviewer 1 Q4

388 for twelve hours prior to experimentation. At the conclusion of each trial, the chambers were
389 quickly depressurized, and worm health was assessed by looking for signs of embolisms, motor
390 dysfunction or other physiological damage that might have arisen from thermal exposure or other
391 experimental handling. Healthy worms were selected, and their branchiae and body tissues were
392 separated and flash frozen in liquid nitrogen for subsequent protein extraction.

393 *Transcriptome Sequencing and Analysis*

394 A *Paralvinella sulfincola* expressed sequence tag (EST) library was sequenced and built
395 by the Joint Genome Institute (Walnut Creek, CA). Briefly, mRNA was purified from total RNA
396 isolated at two different temperature conditions for two tissue types (body and gill). cDNA from
397 each was generated using an oligodT primer followed by template switching (Clontech,
398 Mountain View, CA) and subsequently normalized using the provided protocol of the Evrogen
399 Normalization kit (Evrogen, Moscow, Russia). The normalized cDNA was used to build a library
400 with the construction protocol provided in the 454 Flx Titanium Roche kit (Roche, Branford,
401 CT) and then sequenced. Four EST libraries consisting of 2,593,853 reads were filtered and
402 screened for quality and contamination to produce a filtered set of 2,382,211 reads. These reads
403 were then assembled using Newbler (v2.3-PreRelease-6/30/2009), which resulted in 80748 raw
404 contigs. Herein, contigs are single exon reads, and isotigs are contigs assembled into potential
405 gene assemblies. After a cutoff minimum length of 350 base pairs, the aforementioned sequences
406 were combined to create 24,702 sequences (24,164 isotigs and 538 contigs). The average length
407 of this library is 1,290 bp/sequence and the GC content average is 0.40 (ESM table S4). The
408 sequences were aligned using BlastX with the Swissprot database. 12,562 of the translated
409 sequences had a known BlastX match and 7,002 unique proteins were identified. Longest ORF

Geoffrey Dilly 12/20/11 12:29 PM

Comment: This refers to Rev 1 Question 13

410 translations were used as the reference library for all subsequent MS/MS oligopeptide spectra.
411 Sequences are available at the NCBI's sequence read archive (SRA;
412 <http://www.ncbi.nlm.nih.gov/>) under accession # SRA034879.

413 *Protein extraction*

414 Gill branchiae from three *P. sulfincola* and three *P. palmiformis* per treatment were
415 excised, weighed on an electronic balance (Mettler Toledo, Columbus, OH), and placed into
416 sterilized 0.5 mL glass micropestles (Wheaton, Millville, NJ) containing 24 uL of 20mM Tris pH
417 7.5 buffer and 6 uL Protease Inhibitor Cocktail (PIC) (Sigma-Aldrich, St. Louis, MO). Tissue
418 was homogenized until complete dissociation then centrifuged at 1000x g for 5 minutes. For
419 protein extraction, 0.5 mg gill branchiae were used in a modified Laemmli protein extraction
420 protocol (53). A Tris/PIC mixture at 1:1v/v and 1:20 2-mercaptoethanol/ Laemmli Buffer were
421 added, and the solution was heated at 95°C for 10 minutes. All extractions were loaded in
422 separate lanes onto 4-20% precast Precise Protein Gels (Pierce Inc) with blank lanes between
423 samples. The gels were bathed in a Tris-HEPES-SDS buffer solution and electrophoresed for 45
424 minutes at 100V. Band size and run length were assessed by including 10uL of BenchMark Pre-
425 Stained Protein Ladder 10-190 kDa (Invitrogen, Carlsbad, CA). After electrophoresis, gels were
426 rinsed and stained for three hours using the colloidal comassie blue dye Novex (Invitrogen,
427 Carlsbad, CA). Gels were visualized using a digital gel imaging system (Kodak Gel Logic 100,
428 Kodak, Rochester, NY) and sub-sectioned into six fragments according to protein size. Three
429 biological replicates from each treatment were pooled into one sample per fragment; total gel
430 surface area did not exceed 1cm². The pooled gel sub-sections were then washed with 1 mL of
431 50% acetonitrile and frozen at -20°C prior to analysis.

432 *Protein analyses by tandem mass spectrometry*

433 A total of 36 pooled samples (2 species incubated at 3 temperatures fractionated into 6
434 equal sections) were reduced, carboxyamidomethylated, and digested with trypsin. Resulting
435 peptides from each sample were analyzed over 3 technical replicates using microcapillary
436 reverse-phase HPLC directly coupled to the nano-electrospray ionization source of a
437 ThermoFisher LTQ-Orbitrap XL (replicate 1) or LTQ-Orbitrap Velos (replicates 2 and 3) hybrid
438 mass spectrometer (μ LC/MS/MS). The Orbitrap repetitively surveyed m/z range from 395-1600,
439 while data-dependent MS/MS spectra on the 20 most abundant ions in each survey scan were
440 acquired in the linear ion trap. MS/MS spectra were acquired with a relative collision energy of
441 30%, 2.5-Da isolation width, and recurring ions dynamically excluded for 60s. Preliminary
442 evaluation of peptide-spectrum matches (PSMs) was facilitated using the SEQUEST algorithm
443 with a 30 ppm mass tolerance against the *P. sulfincola* EST library and NCBI nr databases.
444 Spectral counting is a method of relative quantitation in which one compares the number of
445 MSMS spectra acquired for a particular protein across multiple LC-MS/MS datasets. Increases
446 and decreases in relative protein abundance are reflected in corresponding increases and
447 decreases in spectral counts for that protein (54, 55). PSMs were accepted with mass error <3.0
448 ppm and score thresholds to attain an estimated false discovery rate of ~1% using a reverse
449 decoy database strategy and a custom version of the Harvard Proteomics Browser Suite
450 (ThermoFisher Scientific, San Jose, CA). A total of 172,122 peptide spectra were identified with
451 an average of 14.6 amino acids/sequence, with MS/MS spectra populating 1296 referenced
452 proteins (ESM table S5).

453 *Glutathione Measurements*

454 Total GSH and GSSG levels were measured using the Glutathione Assay Kit (Cayman
455 Chem, Ann Arbor, MI) as per instructions. Spectrophotometric readings were taken kinetically
456 for 30 minutes using a Spectramax Plus³⁸⁴ (Molecular Devices, Sunnyvale, CA). Internal
457 standards were run with total GSH and GSSG experimental treatments, and standard curves were
458 built from the endpoint readings.

459 *Data Analysis and Statistics*

460 BaySeq (56) was used to determine statistically significant relative changes over
461 experimental treatments (ESM tables S1 and S2), following methods described in (57). Peptide
462 spectral counts were modeled using a negative binomial distribution to account for potential
463 overdispersion among treatment replicates. By borrowing information on replicate variance
464 among peptides over the entire dataset (ESM Figure S7), the method employed in baySeq better
465 calibrates replicate variance for individual peptides than can be achieved through standard
466 methods of modeling overdispersed count data. Using a likelihood cutoff of 0.9, Bayesian
467 analysis revealed 428 differentially expressed proteins in *Paralvinella palmiformis* and 214
468 differentially expressed proteins in *Paralvinella sulfincola*. We use the convention of a 0.9
469 likelihood cutoff throughout the analysis as in significance indicator, but it is important to note
470 that Bayesian methodology allows for the comparison of relative likelihoods that we explore
471 within the context of each protein family. Additionally, metabolic enzyme regulation was
472 examined *via* pathway analysis. The R package ShotgunFunctionalizeR (Version: 1.0-3, Date:
473 2009-10-09) was used after assigning Enzyme Commission (EC) numbers to sequences using
474 KEGG assignments and the R package BioIDMapper (Version: 2.1, Date: 2010-01-16). To
475 assess statistical support for metabolic pathway-level expression differences, ortholog data were
476 combined into KEGG pathways using methods described in (57). We assumed a binomial

477 distribution in this case, and Monte Carlo methods were used to determine the posterior
478 probability of differential expression, point estimates of pathway abundance and 95% credible
479 intervals for these estimates. Methods for iPath described in (46).

480 **Acknowledgements:** We would like to thank the captains and crew of the *R/V Atlantis* and *DSV*
481 *Alvin* for their assistance gathering samples. We would also like to thank Raymond Lee for his
482 generous donation of samples for use in this research. This material is based upon work
483 supported by the National Science Foundation under Grants# OCE 0623383 and OCE-0426109.
484 *Paralvinella sulfincola* EST sequencing was provided by JGI-DOE under CSP #796476. The
485 work conducted by the JGI-DOE is supported by the Office of Science of the U.S. Department of
486 Energy under Contract No. DE-AC02-05CH11231.

487

487 **References**

- 488 1. Takai K., Nakamura K., Toki T., Tsunogai U., Miyazaki M., Miyazaki J., Hirayama H.,
489 Nakagawa S., Nunoura T., Horikoshi K. 2008 Cell proliferation at 122 degrees C and isotopically heavy
490 CH₄ production by a hyperthermophilic methanogen under high-pressure cultivation. *Proc Natl Acad Sci*
491 *U S A* **105**(31), 10949-10954. (doi:10.1073/pnas.0712334105).
- 492 2. Pörtner H. 2001 Climate change and temperature-dependent biogeography: oxygen limitation of
493 thermal tolerance in animals. *Naturwissenschaften* **88**(4), 137-146. (doi: 10.1007/s001140100216).
- 494 3. Brock T., Libraries U.W. 1978 Thermophilic microorganisms and life at high temperatures,
495 Springer-Verlag New York.
- 496 4. O'Brien J., Dahlhoff E., Somero G. 1991 Thermal resistance of mitochondrial respiration:
497 hydrophobic interactions of membrane proteins may limit thermal resistance. *Physiological zoology*
498 **64**(6), 1509-1526.
- 499 5. Abele D., Heise K., Portner H., Puntarulo S. 2002 Temperature-dependence of mitochondrial
500 function and production of reactive oxygen species in the intertidal mud clam *Mya arenaria*. *J Exp Biol*
501 **205**(13), 1831-1841.
- 502 6. Tomanek L., Zuzow M.J. 2010 The proteomic response of the mussel congeners *Mytilus*
503 *galloprovincialis* and *M. trossulus* to acute heat stress: implications for thermal tolerance limits and
504 metabolic costs of thermal stress. *J Exp Biol* **213**(Pt 20), 3559-3574.
- 505 7. Pörtner H.O., Knust R. 2007 Climate change affects marine fishes through the oxygen limitation
506 of thermal tolerance. *Science* **315**(5808), 95-97. (doi:10.1126/science.1135471).
- 507 8. Pörtner H.O., Bennett A.F., Bozinovic F., Clarke A., Lardies M.A., Lucassen M., Pelster B.,
508 Schiemer F., Stillman J.H. 2006 Trade-offs in thermal adaptation: the need for a molecular to ecological
509 integration. *Physiol Biochem Zool* **79**(2), 295-313. (doi:10.1086/499986).
- 510 9. Somero GN. 2011 Comparative physiology: a "crystal ball" for predicting consequences of global
511 change. *Am J Physiol Regul Integr Comp Physiol*. 301(1):R1-14.
- 512 10. Gehring W., Wehner R. 1995 Heat shock protein synthesis and thermotolerance in *Cataglyphis*,
513 an ant from the Sahara desert. *Proceedings of the National Academy of Sciences of the United States of*
514 *America* **92**(7), 2994.
- 515 11. Wickstrom C., Castenholz R. 1973 Thermophilic ostracod: aquatic metazoan with the highest
516 known temperature tolerance. *Science* **181**(4104), 1063. (doi: 10.1126/science.181.4104.1063).
- 517 12. Nguyen T.T., Michaud D., Cloutier C. 2009 A proteomic analysis of the aphid *Macrosiphum*
518 *euphorbiae* under heat and radiation stress. *Insect Biochem Mol Biol* **39**(1), 20-30. (doi:
519 10.1016/j.ibmb.2008.09.014).

- 520 13. Mary J, Rogniaux H, Rees JF, Zal F. 2010 Response of *Alvinella pompejana* to variable oxygen
521 stress: a proteomic approach. *Proteomics*. **10**(12):2250-8.
- 522 14. Cary SC, Shank T, Stein J. 1998 Worms bask in extreme temperatures. *Nature* **391**(6667):545-6.
- 523 15. Chevaldonné P, Fisher C, Childress J, Desbruyères D, Jollivet D, Zal F, et al. 2000
524 Thermotolerance and the 'Pompeii worms'. *Marine Ecology Progress Series*. **208**:293-5.
- 525 16. Cottin D, Ravaux J, Leger N, Halary S, Toullec JY, Sarradin PM, et al. 2008 Thermal biology of
526 the deep-sea vent annelid *Paralvinella grasslei*: in vivo studies. *J Exp Biol* **211**(Pt 14):2196-204.
- 527 17. Sarrazin J., Levesque C., Juniper S., Tivey M. 2002 Mosaic community dynamics on Juan de
528 Fuca Ridge sulphide edifices: substratum, temperature and implications for trophic structure. *CBM-
529 Cahiers de Biologie Marine* **43**(3-4), 275-279.
- 530 18. Girguis P.R., Lee R.W. 2006 Thermal preference and tolerance of alvinellids. *Science* **312**(5771),
531 231. (doi:10.1126/science.1125286).
- 532 19. Lee R.W. 2003 Thermal tolerances of deep-sea hydrothermal vent animals from the Northeast
533 Pacific. *Biol Bull* **205**(2), 98-101.
- 534 20. Selong J.H., McMahon T.E., Zale A.V., Barrows F.T. 2001 Effect of temperature on growth and
535 survival of bull trout, with application of an improved method for determining thermal tolerance in fishes.
536 *Transactions of the American Fisheries Society* **130**(6), 1026-1037. (doi:10.1577/1548-
537 8659(2001)130<1026:EOTOGA>2.0.CO;2).
- 538 21. Kilgour D.M., McCauley R.W. 1986 Reconciling the two methods of measuring upper lethal
539 temperatures in fishes. *Environmental Biology of Fishes* **17**(4), 281-290. (doi: 10.1007/BF00001494).
- 540 22. Arndt V., Rogon C., Hohfeld J. 2007 To be, or not to be--molecular chaperones in protein
541 degradation. *Cell Mol Life Sci* **64**(19-20), 2525-2541. (doi:10.1007/s00018-007-7188-6).
- 542 23. Mayer M.P., Bukau B. 2005 Hsp70 chaperones: cellular functions and molecular mechanism.
543 *Cell Mol Life Sci* **62**(6), 670-684. (doi:10.1007/s00018-004-4464-6).
- 544 24. Feder M.E., Hofmann G.E. 1999 Heat-shock proteins, molecular chaperones, and the stress
545 response: evolutionary and ecological physiology. *Annu Rev Physiol* **61**, 243-282.
546 (doi:10.1146/annurev.physiol.61.1.243).
- 547 25. Parsell D.A., Lindquist S. 1993 The function of heat-shock proteins in stress tolerance:
548 degradation and reactivation of damaged proteins. *Annu Rev Genet* **27**, 437-496.
549 (doi:10.1146/annurev.ge.27.120193.002253).
- 550 26. Vos M.J., Hageman J., Carra S., Kampinga H.H. 2008 Structural and functional diversities
551 between members of the human HSPB, HSPH, HSPA, and DNAJ chaperone families. *Biochemistry*
552 **47**(27), 7001-7011. (doi:10.1021/bi800639z).

553 27. Terasawa K., Minami M., Minami Y. 2005 Constantly updated knowledge of Hsp90. *J Biochem*
554 **137**(4), 443-447. (doi:10.1093/jb/mvi056).

555 28. Wandinger S.K., Richter K., Buchner J. 2008 The Hsp90 chaperone machinery. *J Biol Chem*
556 **283**(27), 18473-18477. (doi:10.1074/jbc.R800007200).

557 29. Eletto D., Dersh D., Argon Y. 2010 GRP94 in ER quality control and stress responses. *Semin*
558 *Cell Dev Biol* **21**(5), 479-485. (doi: 10.1016/j.semcdb.2010.03.004).

559 30. Martin J., Horwich A.L., Hartl F.U. 1992 Prevention of protein denaturation under heat stress by
560 the chaperonin Hsp60. *Science* **258**(5084), 995-998. (doi: 10.1126/science.1359644).

561 31. Höhfeld J., Hartl F.U. 1994 Role of the chaperonin cofactor Hsp10 in protein folding and sorting
562 in yeast mitochondria. *The Journal of cell biology* **126**(2), 305. (doi: 10.1083/jcb.126.2.305).

563 32. Arrigo A.P., Virot S., Chaufour S., Firdaus W., Kretz-Remy C., Diaz-Latoud C. 2005 Hsp27
564 consolidates intracellular redox homeostasis by upholding glutathione in its reduced form and by
565 decreasing iron intracellular levels. *Antioxid Redox Signal* **7**(3-4), 414-422.

566 33. Arrigo A.P. 2001 Hsp27: novel regulator of intracellular redox state. *IUBMB Life* **52**(6), 303-307.
567 (doi:10.1080/152165401317291165).

568 34. Schroder M., Kaufman R.J. 2005 ER stress and the unfolded protein response. *Mutat Res* **569**(1-
569 2), 29-63. (doi:10.1016/j.mrfmmm.2004.06.056).

570 35. Noiva R. 1999 Protein disulfide isomerase: the multifunctional redox chaperone of the
571 endoplasmic reticulum. *Semin Cell Dev Biol* **10**(5), 481-493. (doi: 10.1006/scdb.1999.0319).

572 36. Andreyev A.Y., Kushnareva Y.E., Starkov A.A. 2005 Mitochondrial metabolism of reactive
573 oxygen species. *Biochemistry (Mosc)* **70**(2), 200-214. (doi: 10.1007/s10541-005-0102-7).

574 37. Starkov A.A. 2006 Protein-mediated energy-dissipating pathways in mitochondria. *Chem Biol*
575 *Interact* **161**(1), 57-68. (doi:10.1016/j.cbi.2006.02.009).

576 38. Abele D., Puntarulo S. 2004 Formation of reactive species and induction of antioxidant defence
577 systems in polar and temperate marine invertebrates and fish. *Comp Biochem Physiol A Mol Integr*
578 *Physiol* **138**(4), 405-415. (doi:10.1016/j.cbpb.2004.05.013).

579 39. Heise K., Puntarulo S., Pörtner H.O., Abele D. 2003 Production of reactive oxygen species by
580 isolated mitochondria of the Antarctic bivalve *Laternula elliptica* (King and Broderip) under heat stress.
581 *Comp Biochem Physiol C Toxicol Pharmacol* **134**(1), 79-90. (doi: 10.1016/S1532-0456(02)00212-0).

582 40. Keller M., Sommer A.M., Pörtner H.O., Abele D. 2004 Seasonality of energetic functioning and
583 production of reactive oxygen species by lugworm (*Arenicola marina*) mitochondria exposed to acute
584 temperature changes. *J Exp Biol* **207**(Pt 14), 2529-2538. (doi:10.1242/jeb.01050).

- 585 41. Pörtner H.O. 2002 Climate variations and the physiological basis of temperature dependent
586 biogeography: systemic to molecular hierarchy of thermal tolerance in animals. *Comp Biochem Physiol A*
587 *Mol Integr Physiol* **132**(4), 739-761. (doi: 10.1016/S1095-6433(02)00045-4).
- 588 42. Lesser M.P. 2006 Oxidative stress in marine environments: biochemistry and physiological
589 ecology. *Annu Rev Physiol* **68**, 253-278.
- 590 43. Pastore A., Federici G., Bertini E., Piemonte F. 2003 Analysis of glutathione: implication in
591 redox and detoxification. *Clin Chim Acta* **333**(1), 19-39. (doi: 10.1016/S0009-8981(03)00200-6).
- 592 44. Fraenkel D.G. 2003 The top genes: on the distance from transcript to function in yeast glycolysis.
593 *Curr Opin Microbiol* **6**(2), 198-201. (doi: 10.1016/S1369-5274(03)00023-7).
- 594 45. Fraenkel D.G. 2003 The top genes: on the distance from transcript to function in yeast glycolysis.
595 *Curr Opin Microbiol* **6**(2), 198-201. (doi: 10.1016/S1369-5274(03)00023-7).
- 596 46. Letunic I., Yamada T., Kanehisa M., Bork P. 2008 iPath: interactive exploration of biochemical
597 pathways and networks. *Trends Biochem Sci* **33**(3), 101-103. (doi:10.1016/j.tibs.2008.01.001).
- 598 47. Go Y.M., Jones D.P. 2008 Redox compartmentalization in eukaryotic cells. *Biochim Biophys*
599 *Acta* **1780**(11), 1273-1290. (doi:10.1016/j.bbagen.2008.01.011).
- 600 48. Ravaux J, Gaill F, Bris NL, Sarradin PM, Jollivet D, Shillito B. 2003 Heat-shock response and
601 temperature resistance in the deep-sea vent shrimp *Rimicaris exoculata*. *J Exp Biol* **206**(14):2345.
- 602 49. Tivey MK. 2004. Environmental conditions within active seafloor vent structures: sensitivity to
603 vent fluid composition and fluid flow. *Geophysical monograph*. **144**:137-52.
- 604 50. Kawakami R., Sakuraba H., Kamohara S., Goda S., Kawarabayasi Y., Ohshima T. 2004
605 Oxidative stress response in an anaerobic hyperthermophilic archaeon: presence of a functional
606 peroxiredoxin in *Pyrococcus horikoshii*. *Journal of biochemistry* **136**(4), 541.
- 607 51. Henry M., Childress J., Figueroa D. 2008 Metabolic rates and thermal tolerances of
608 chemoautotrophic symbioses from Lau Basin hydrothermal vents and their implications for species
609 distributions. *Deep Sea Research Part I: Oceanographic Research Papers* **55**(5), 679-695.
610 (doi: 10.1016/j.dsr.2008.02.001).
- 611 52. Hourdez S., Lallier F. 2007 Adaptations to hypoxia in hydrothermal-vent and cold-seep
612 invertebrates. *Reviews in Environmental Science and Biotechnology* **6**(1), 143-159. (doi: 10.1007/s11157-
613 006-9110-3).
- 614 53. Laemmli U. 1970 Cleavage of structural proteins during the assembly of the head of
615 bacteriophage T4. *Nature* **227**(5259), 680-685. (doi:10.1038/227680a0).
- 616 54. Liu H, Sadygov RG, Yates III JR. 2004 A model for random sampling and estimation of relative
617 protein abundance in shotgun proteomics. *Analytical Chemistry*. **76**(14):4193-201.

618 55. Zybailov B, Coleman MK, Florens L, Washburn MP. 2005 Correlation of relative abundance
619 ratios derived from peptide ion chromatograms and spectrum counting for quantitative proteomic analysis
620 using stable isotope labeling. *Analytical Chemistry*. **77**(19):6218-24.

621 56. Hardcastle T., Kelly K. 2010 baySeq: Empirical Bayesian methods for identifying differential
622 expression in sequence count data. *BMC bioinformatics* **11**(1), 422. (doi: 10.1186/1471-2105-11-422).

623 57. Ottesen EA, Marin R, 3rd, Preston CM, Young CR, Ryan JP, Scholin CA, et al. 2011
624 Metatranscriptomic analysis of autonomously collected and preserved marine bacterioplankton. *ISME J*.
625 **5**(12):1881-95.

626

627

627 **Figure Legends**

628 **Figure 1:** Molecular chaperones

629 Differences in expression between *P. sulfincola* and *P. palmiformis* in log fold-change for six
630 major molecular chaperones across their thermal range. S10 →45 = difference from *P. sulfincola*
631 maintained at 10°C to 45°C; P12 →38 = difference from *P. palmiformis* maintained at 12°C to
632 38°C. Stars (*) indicate that the log change is > 0.90 in our Bayesian analysis, indicating a
633 significant change with temperature. We assumed a binomial likelihood for the data and a Beta
634 (0.5,0.5) prior for each treatment. Monte Carlo sampling from the resulting posterior
635 distributions within each treatment was used to estimate the posterior distributions of log-fold
636 changes between treatments. We report the medians and 95% credible intervals (bars) of the
637 posterior distributions of log-fold change between treatments.

638 **Figure 2:** Representative glutathione pathway in *Paralvinella* with responses to thermal
639 exposure

640 Synthesis pathways of the antioxidant glutathione (GSH) and its catalyzing enzyme Glutathione
641 peroxidase (GPx). Ovals represent enzymes; grey rectangles indicate substrates. Grey ovals
642 represent proteins only observed in the *P. sulfincola* EST database. Color indicates significance
643 and direction of regulation. Asterisks indicate ATP-dependent enzymatic steps. Numbers in
644 diamonds correspond to protein count rows in Table 1b. Note: GPx appears twice – in synthesis
645 in the selenium pathway, and in oxidizing GSH to GSSH. DNMT is found in cysteine pathway
646 only; at present, the specific seleno-methyltransferase for Paralvinellids is unknown. Some
647 reaction cofactors omitted for simplicity. Abbreviations: AHCY, Adenosylhomocysteinase A;
648 CBS, Cystathionine β-synthase; CGL, Cystathionine γ-synthase; DNMT, DNA (cytosine-5)-

649 methyltransferase; GPx, Glutathione Peroxidase; GS, Glutathione synthetase; GSH, Glutathione;
650 GSSH, glutathione disulfide; GSR, Glutathione reductase; GSTs, Glutathione sulfur transferases;
651 MAT2, Methionine adenosyltransferase; SelD, Selenide water dikinase.

652 **Table 1a, b.** Shifts in *Paralvinella* protein abundance during thermal exposure

653 Key enzymes of *P. sulfincola* and *P. palmiformis* discussed in the text. EST refers to the (i) isotig
654 or (c) contig identifier for each enzyme. Log change refers to the shift in abundance between
655 treatments, i.e. $(\log \Delta P.p - 12 \rightarrow 21 = \text{protein log fold change between } P. palmiformis$
656 treatments 12°C and 21°C). Counts are combined between all three technical replicates and
657 normalized to treatment library sizes. Red boxes indicate a significant ($\text{Pr(DE)} > 0.9$) increase in
658 protein abundance; Blue boxes indicate a significant ($\text{Pr(DE)} > 0.9$) decrease in protein
659 abundance. **Table 1a** lists chaperones; **Table 1b** lists glutathione pathways.