



# Inhibitory single neuron control of seizures and epileptic traveling waves in humans

## Citation

Ahmed, Omar J, Mark A Kramer, Wilson Truccolo, Jason S Naftulin, Nicholas S Potter, Emad N Eskandar, Garth R Cosgrove, Andrew S Blum, Leigh R Hochberg, and Sydney S Cash. 2014. "Inhibitory single neuron control of seizures and epileptic traveling waves in humans." BMC Neuroscience 15 (Suppl 1): F3. doi:10.1186/1471-2202-15-S1-F3. <http://dx.doi.org/10.1186/1471-2202-15-S1-F3>.

## Published Version

doi:10.1186/1471-2202-15-S1-F3

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12785794>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

FEATURED TALK PRESENTATION

Open Access

# Inhibitory single neuron control of seizures and epileptic traveling waves in humans

Omar J Ahmed<sup>1\*</sup>, Mark A Kramer<sup>7</sup>, Wilson Truccolo<sup>5</sup>, Jason S Naftulin<sup>1</sup>, Nicholas S Potter<sup>4</sup>, Emad N Eskandar<sup>2</sup>, Garth R Cosgrove<sup>3</sup>, Andrew S Blum<sup>4</sup>, Leigh R Hochberg<sup>1,6</sup>, Sydney S Cash<sup>1</sup>

From The Twenty Third Annual Computational Neuroscience Meeting: CNS\*2014  
Québec City, Canada. 26-31 July 2014

Inhibitory neuronal activity is critical for the normal functioning of the brain, but is thought to go awry during neurological disorders such as epilepsy. Animal models have suggested both decreased and increased inhibition as possible initiators of epileptic activity, but it is not known if, or how, human inhibitory neurons shape seizures. Here, using large-scale recordings of neocortical single neurons in patients with secondarily generalized tonic-clonic seizures, we show that fast-spiking (FS) inhibitory activity first increases as a seizure spreads across the neocortex, impeding and altering the spatial flow of fast epileptic traveling waves. Unexpectedly, however, FS cells cease firing less than half-way through a seizure. We use biophysically-realistic computational models to show that this cessation is due to FS cells entering depolarization block as a result of extracellular potassium accumulation during the seizure and not because they are inhibited by other inhibitory subtypes. Strikingly, this absence of FS inhibitory activity is accompanied by dramatic increases in local seizure amplitude along with unobstructed traveling waves and is seen during all secondarily generalized seizures examined, independent of etiology or focus. FS cessation also leads to prominent spike-and-wave events, suggesting that FS cell dynamics control the transition between the tonic and clonic phases of these seizures. Thus, it may be possible to curtail human seizures by preventing inhibitory neurons from entering potassium-dependent depolarization block, a novel and potentially powerful therapeutic avenue in treating multiple kinds of epilepsies.

## Acknowledgements

We would like to thank the patient volunteers. This work was supported by postdoctoral fellowships from the Epilepsy Foundation (222178), MGH (2012A052031) & NINDS (F32-NS083208) awarded to OJA and by R01-NS062092 (SSC), R01-NS072023 (MAK) and R01-NS079533 (WT).

## Authors' details

<sup>1</sup>Department of Neurology, Harvard Medical School and Massachusetts General Hospital, Boston, MA 02114, USA. <sup>2</sup>Department of Neurosurgery, Harvard Medical School and Massachusetts General Hospital, Boston, MA 02114, USA. <sup>3</sup>Department of Neurosurgery, Rhode Island Hospital, Providence, RI 02903, USA. <sup>4</sup>Department of Neurology, Rhode Island Hospital, Providence, RI 02903, USA. <sup>5</sup>Department of Neuroscience, Brown University, Providence, RI 02912, USA. <sup>6</sup>School of Engineering, Brown University, Providence, RI 02912, USA. <sup>7</sup>Department of Mathematics & Statistics, Boston University, MA 02215, USA.

Published: 21 July 2014

doi:10.1186/1471-2202-15-S1-F3

Cite this article as: Ahmed et al.: Inhibitory single neuron control of seizures and epileptic traveling waves in humans. *BMC Neuroscience* 2014 15(Suppl 1):F3.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [omar.j.ahmed@gmail.com](mailto:omar.j.ahmed@gmail.com)

<sup>1</sup>Department of Neurology, Harvard Medical School and Massachusetts General Hospital, Boston, MA 02114, USA

Full list of author information is available at the end of the article