



# Body fat, energy balance and estradiol levels: a study based on hormonal profiles from complete menstrual cycles

## Citation

Ziomkiewicz, Anna, Peter T. Ellison, Susan F. Lipson, Inger Thune, and Graznya Jasienska. 2008. Body fat, energy balance, and estradiol levels: a study based on hormonal profiles from complete menstrual cycles. *Human Reproduction* 23(11): 2555-2563.

## Published Version

<http://dx.doi.org/10.1093/humrep/den213>

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:2579735>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

*Title:* Body fat accumulation, energy balance and estradiol levels: a study based on hormonal profiles from entire menstrual cycles

*Running title:* Body fat, energy balance and estradiol levels in women

*Authors:* A. Ziomkiewicz <sup>1</sup>, P. T. Ellison <sup>2</sup>, S. F. Lipson <sup>2</sup>, I. Thune <sup>3,4</sup>, G. Jasienska <sup>5</sup>

<sup>1</sup> Institute of Anthropology, Polish Academy of Science, Kuznicza 35, 50-951 Wrocław, Poland

<sup>2</sup> Department of Anthropology, Harvard University, Cambridge, USA

<sup>3</sup> Institute of Community Medicine, University of Tromsø, Norway

<sup>4</sup> The Cancer Center, Ullevål University Hospital, Oslo, Norway

<sup>5</sup> Department of Epidemiology and Population Studies, Jagiellonian University, Collegium Medicum, Krakow, Poland

*Corresponding author:* Anna Ziomkiewicz, email: [annaz@antro.pan.wroc.pl](mailto:annaz@antro.pan.wroc.pl)

*Abstract*

## BACKGROUND

It is hypothesized that female fecundity is regulated by nutritional status. Although widely cited, this hypothesis is not strongly supported by empirical data from non-obese, healthy women of reproductive age.

## METHODS

141 healthy, reproductive aged women from Southern Poland collected everyday morning saliva samples for one menstrual cycle. Levels of 17- $\beta$ -estradiol were analyzed by radioimmunoassay. Measurements of body fat and anthropometrics were taken randomly with respect to phase of the menstrual cycle. Energy balance was specified basing on changes in percentage of body fat from the beginning to the end of the observation.

## RESULTS

In women of very low to average body fat the 10% increase in fatness was associated with 5 to 7 pmol/l increase in estradiol levels. Women with high body fat had significantly lower levels of E2 comparing to women with low and average body fat.

In women with positive energy balance linear association between fatness and E2 was even stronger. Significant differences between body fat groups were also noted in estradiol profiles during menstrual cycle. In contrast, no such relationship was found in women with negative energy balance.

## CONCLUSIONS

In healthy women body fat was positively associated with estradiol levels although

this effect was restricted only to women with positive energy balance. These results point to the importance of negative energy balance as a factor affecting the relationship between fatness and estradiol levels.

*Key words:* 17- $\beta$ -estradiol /body fat percentage/energy balance/menstrual cycle

### *Introduction*

Even a relatively small decline in reproductive ovarian hormone levels during menstrual cycle has a marked effect on woman's fecundity (Ellison, 1991, Lipson & Ellison 1996). Estradiol, as the major and most active ovarian form of estrogen, is involved in ovarian follicle development, quality of the ovum and ovulation process. A small reduction in the levels of estradiol during follicular phase, without any significant change in the length of menstrual cycle, may lead to disturbed development of ovarian follicle, resulting in decreased probability of conception (Yoshimura and Wallach, 1987). Higher levels of follicular estradiol were observed in cycles resulting in conception comparing to cycles without conceptions in healthy subjects (Baird et al., 1997, 1999, Lipson and Ellison, 1996; Li et al., 2001, Lu et al., 1999; Venners et al 2006). In women undergoing in vitro fertilization higher levels of estradiol after the time of embryo transfer (Chen et al., 2003) or hCG administration (Blazar et al., 2004) were also strongly positively related to pregnancy success after the in vitro procedure.

Inter-individual variation in levels of steroid ovarian hormones can be attributed

to several factors, including those related to energy availability and metabolism. While the influence of physical activity on ovarian hormone levels is relatively well recognized (Burrows and Bird, 2000; Chen and Brzyski, 1999; De Souza and Williams, 2004; Elias and Wilson, 1993; Jasienska, 2003; Jasienska and Ellison, 1998, 2004; Jasienska et al., 2006; Warren and Perlroth, 2001), the role of nutritional status is still controversial. In particular there is no agreement if the nutritional status can affect reproductive hormone levels independently from the influence of physical activity and energy balance, which were hypothesized to have predominant effect on reproductive physiology (Ellison, 2001, 2003, Jasienska, 2001, Jasienska and Ellison, 2004, Schneider, 2004, Wade and Jones, 2004, Wade et al., 1996).

The regulatory role of nutritional status on reproductive ability was first hypothesized by Frisch (Frisch, 1984) and the evidence on the relationship between nutritional status and ovarian hormone levels comes from studies on women with anorexia nervosa (Kirchengast and Huber, 2004; Miller et al., 2004, van Binsbergen et al., 1990, Warren, 1983). In these women, low body mass or low body mass index (BMI) is related to low levels of estradiol and inhibition of menstrual cycles. However, results of the studies on healthy, relatively well-nourished women are contradictory. Studies report no relationship (Dorgan et al., 1995; Ukkola et al., 2001, Ivandi et al., 1998; Panter-Brick et al., 1993), positive relationship (Barnett et al., 2001, 2002; Brunning et al., 1992; Furberg et al., 2005; Zanker and Swaine, 1998), or negative relationship (Howard et al., 1987, Potischman et al., 1996, Thomas et al., 1997a, b; Westhoff et al., 1996) between nutritional status and ovarian reproductive hormone levels, therefore it is not clear if

biological variation in the reproductive hormone levels can be attributed to the influence of nutritional status.

However, several authors suggested that nutritional status cannot be considered as the factor regulating reproductive functioning independently from the effect of physical activity and energy balance (Ellison, 2001, 2003, Jasienska, 2001, Jasienska and Ellison, 2004, Wade and Jones, 2004). It is hypothesized that energy balance and physical activity have predominant role in reproductive function regulation. As energy balance affects nutritional status a relationship between nutritional status and estradiol can be in fact a by-product of the association between energy balance and reproductive hormone levels. In none of the cited studies the effect of physical activity and/or energy balance was accounted for thus it is impossible to determine if the observed changes in the levels of reproductive hormones are independently modified by the nutritional status.

Although recently several molecular models of the association between nutritional status and fecundity were proposed (Budak et al., 2006, Caprio et al., 2001, Rexford and Flier, 2000, Smith et al., 2002, Tatarani, 1997), conclusive results confirming this association are still lacking. Here, we report results of a study investigating an association between ovarian estradiol levels measured during entire menstrual cycle and nutritional status indicated by body fat percentage in 141 healthy, well-nourished, but not obese women from Southern Poland. To account for the confounding effect of energy balance separate analysis were conducted in sub-sample of 131 women for whom energy balance was estimated on the basis of changes in the body fat concentration. Our results suggest that in women with positive energy balance

nutritional status has independent, regulatory effect on the levels of estradiol during menstrual cycles whereas in women with negative energy balance no such an effect can be observed.

## *Methods*

### Study Participants

The participants of the study were 141 women from urban and rural areas of Southern Poland. Women were recruited for the study by local media advertisement in the urban area, and through their parish in the rural area between June 2001 and June 2003. Selection of women for the study was based on the following criteria: age between 24 and 37 years, self-assessed regular menstrual cycles not shorter than 25 and not longer than 35 days, no fertility problems, no gynecological and endocrinological disorders, not taking hormonal oral contraceptives or other hormonal medications for the period of six months prior to the recruitment, and not being pregnant or lactating during the six months prior to the recruitment. The research protocol was approved by the Bioethical Committee of Jagiellonian University.

### General questionnaire and physical activity assessment

General questionnaire collecting information about the place of birth, age, birth weight and birth length, age of menarche, education, marital status, reproductive history, use of hormonal medication, and tobacco consumption was distributed to the study participants.

Physical activity was assessed on the base of a pre-set daily log completed by

women every day during their menstrual cycle. It collected data about hours of sleep, waking-up time, and time spent during the day on physical activities in five categories. Detailed description of the methods was published elsewhere (Jasienska et al 2006).

#### Anthropometric measurements and energy balance adjustment

Anthropometric measurements were taken from each woman twice, randomly with respects to the beginning and to the end of menstrual cycle. The maximal time elapsing from the first measurement to the beginning of the menstrual cycle and from the end of menstrual cycle to the second measurement was less then 65 days.

Measurements of height, weight, body fat percentage, breasts, underbreast, waist and hips circumferences were taken by the trained anthropologist. Body fat percentage was measured by bioimpedance using TANITA scale; model TBF 551 with measurement accuracy of 0.1%. Body mass was measured using the same Tanita scale, with the measurement accuracy of 0.1 kg. Body mass index (BMI) was calculated as ratio of height [m] to body mass [kg] squared. Waist and hips circumferences measurements were used to calculate waist-to-hip ratio (WHR). Breast and underbreast circumference measurements were used to calculate breast-to-underbreast ratio (BUR).

Energy balance was determined based on changes in the percentage of body fat between the first and the second measurement. A woman was classified as having positive energy balance when the difference in body fat percentage between the first and second measurement was equal or greater then -1%. When this difference was smaller then -1% a woman was classified as having negative energy balance

#### Salivary estradiol assay procedure and estradiol indices



Packages containing plastic vials and laboratory-tested chewing gum were distributed to women before the beginning of their menstrual cycle. During one entire menstrual cycle, every day in the morning after waking-up, women collected saliva samples to plastic tubes pretreated with sodium azide following published protocols (Lipson and Ellison, 1989). Saliva samples from 20 days (-5 to -24 reversed cycle days) were analyzed for E2 concentrations using RIA I-125-based kit (#39100, Diagnostic Systems Laboratories, Webster, Texas, USA) with published modifications to the manufacturer's protocol (Jasienska et al 2004). Average intra-assay variability was 9% and inter-assay variability varied from 23% for lower (15 pmol/l) to 13% for higher (50 pmol/l) values. The sensitivity of estradiol assay was 4 pmol/l. Before statistical analysis, days of cycles were aligned on the basis of midcycle drop day (day 0) identification, which provides a reasonable estimate of the day of ovulation (Lipson and Ellison, 1996). Values of E2 concentration from 18 consecutive days were used in the analysis, and the following estradiol indices were calculated: "mean E2" (mean of days from -9 to 8), "mid-cycle E2" (mean of days from -2 to 2), "day -1 E2" (day -1 E2 value, day before the midcycle drop day), "day 0 E2" (day 0 E2 value, midcycle drop day), "mean follicular E2" (mean of days from -9 to -1) and "mean luteal E2" (mean of days from 0 to 8).

#### Statistical analysis

Women were divided into four groups based on the quartiles of the distribution of body fat percentage. The established designation of fatness quartiles was maintained in all analyses to allow comparisons between women groups (all women, positive

energy balance women, negative energy balance women) and thus produced differences in sample sizes. Differences among very low, low, average and high body fat groups in basic characteristics i.e. age, size at birth, age at menarche, menstrual cycle length, body anthropometrics, body composition, physical activity and number of cigarettes smoked per day were tested by one-way factorial analysis of variance (ANOVA) followed by Duncan tests. Separate comparisons in basic characteristic were also made in body fat quartiles of women with negative and positive energy balance. The same procedure was applied to test the significance of the differences in E2 indices between body fat quartiles.

Additionally, simple regression models tested the effect of body fat on E2 levels, with each E2 index as the dependent variable and body fat as the independent predictor. Separate analyses were conducted for women with positive and negative energy balance.

Repeated measure analysis of variance was used to test the differences in E2 profiles among the body fat groups in positive energy balance women. Separate models for values of E2 levels during -9 to 8 days (whole cycle), -9 to -1 days (follicular phase) and 0 to 8 (luteal phase) as dependent variables and body fat group as the independent were tested. The contrast analyzes indicated statistical significance of differences among four body fat groups.

## *Results*

### BODY FAT AND ESTRADIOL

#### General characteristics

General descriptive statistics of all women categorized with respect to their fatness are presented in Table I. Women characterized by very low, low, average and high body fat did not differ significantly with respect to age, birth weight and birth height, age of first menstruation, number of cigarettes smoked per day and physical activity. They differed significantly with respect to usual length of menstrual cycle ( $F_{3,134}=2.73$ ,  $p<0.05$ ) and anthropometric traits. Women with average body fat had significantly shorter usual, self-reported length of menstrual cycles as compared to women with very low body fat (28.1 days for average body fat group vs 30.2 days for very low body fat group). As expected, women characterized by higher body fat content had significantly higher body mass, BMI, and higher waist-to-hip ratio (all within groups comparisons were statistically significant at  $p<0.05$ ), but they did not differ significantly with respect to their breast-to-underbreast ratio.

#### Estradiol levels in body fat groups

Mean levels of 17- $\beta$ -estradiol in subsequent body fat groups are presented in Table II. Comparing to women characterized by average body fat women with very low and high body fat percentage had significantly lower levels of estradiol during follicular phase (21.1 vs 17.0 and 15.9 pmol/l,  $F_{3,124}=3.22$ ,  $p=0.025$ ), midcycle (25.4 pmol/l vs 19.7 and 18.2 pmol/l,  $F_{3,126}=4.03$ ,  $p=0.009$ ), and on day -1 (38.8 pmol/l vs 29.1 and 26.6

pmol/l,  $F_{3,123}=4.60$ ,  $p=0.004$ ), day 0 (20.4 pmol/l vs 15.2 and 13.4 pmol/l,  $F_{3,122}= 3.35$ ,  $p=0.021$ ). Similar differences were also observed in comparisons between women with low body fat and women with very low and high body fat.

Additional evidence for the relationship between nutritional status and estradiol levels came from simple regression analysis restricted to a group of non-overweight women (body fat <31%). In these women, we observed linear, positive association between the amount of accumulated body fat and mean E2 ( $R^2=0.085$ ,  $p=0.003$ ), follicular E2 ( $R^2=0.075$ ,  $p=0.007$ ), mid-cycle E2 ( $R^2=0.094$ ,  $p=0.002$ ) (Figure 1) and luteal E2 ( $R^2=0.076$ ,  $p=0.006$ ). Statistically significant relationship was also observed for E2 on day -1 ( $R^2=0.097$ ,  $p=0.002$ ) and on day 0 ( $R^2=0.085$ ,  $p=0.004$ ). The 10% increase in percentage of body fat in the range of 9.1 to 30.8% was associated with 7 pmol/l increase in mid-cycle E2 and 5 pmol/l increase in follicular phase E2, luteal phase E2 and mean E2 during the menstrual cycles.

## BODY FAT ENERGY BALANCE AND ESTRADIOL LEVELS

### General characteristics

To test the effect of energy balance on the relationship between body fat and estradiol separate analyzes were conducted in 131 women characterized by positive or negative energy balance. General characteristics of women from those groups are presented in Table III. Women with negative energy balance as compared to those with positive energy balance had significantly longer usual cycle length (30.0 days vs 28.8 days respectively) and lower body weight at birth (3.38 kg vs 3.07 kg). However, they

did not differ significantly in any of the anthropometric or life style parameters.

In contrast significant differences in anthropometrics were observed within both negative and positive energy balance group between women characterized by very low, low, average and high body fat content. As expected, higher weight, body fat, body mass index and waist-to-hip ratio were observed in women with higher body fat content comparing to women with lower body fat content (all between group comparisons significant at  $p < 0.05$ ) in positive energy balance group. In negative energy balance the same trend was observed for weight, body mass index and body fat. Additionally, in positive energy balance group women with very low and high body fat content had significantly higher birth weight comparing to women with low and average body fat content (3.58 kg for very low body fat content group and 3.57 kg for high body fat content group vs 3.18 kg for low body fat content group and 3.21 kg for average body fat content group).

Estradiol levels in body fat and energy balance groups

Average values of consecutive E2 indices in body fat groups of women with positive and negative energy balance group are presented in Table IV. Anova analyzes of estradiol levels in these groups revealed significant differences in all estradiol indices between women differing with respect to body fat content in positive but not in negative energy balance group. Significantly lower levels of mean E2, mid-cycle E2, day -1 E2 and follicular phase E2 were observed in women with very low and high body fat content comparing to women with low and average body fat content. Additionally, levels of day 0 E2 and luteal phase E2 were significantly lower in women with very low

and high body fat group in comparison with women with average body fat group.

Repeated measure analysis of variance revealed significant differences in estradiol profiles during menstrual cycle between body fat groups ( $F_{3,66}=3.81$ ,  $p<0.05$ ) in women with positive energy balance. Women with very low and high body fat had significantly lower levels of E2 during whole menstrual cycle comparing to women with average and high fat (Figure 2). Similar differences between body fat groups was also observed with respect to levels of E2 during follicular phase ( $F_{3,66}=3.69$ ,  $p<0.05$ ) and levels of E2 during luteal phase ( $F_{3,66}=3.17$ ,  $p<0.05$ ).

The relationship between body fat and E2 levels was further confirmed by regression analyzes in non-overweight women with positive but not in those with negative energy balance (Table V). In women with positive energy balance we observed linear association between amount of accumulated body fat and mean E2 ( $R^2=0.13$ ,  $p<0.01$ ), mid-cycle E2 ( $R^2=0.15$ ,  $p<0.01$ ), follicular E2 ( $R^2=0.11$ ,  $p<0.05$ ) and luteal E2 ( $R^2=0.14$ ,  $p<0.01$ ) (Figure 3). The same trend was also observed for estradiol on day -1 ( $R^2=0.14$ ,  $p<0.01$ ) and on day 0 ( $R^2=0.14$ ,  $p<0.01$ ). In the range of 9.1 to 30.8% of body fat, the 10% increase in body fat content was associated with about then 7 pmol/l increase in mean E2, luteal E2 and follicular E2. Also 10% increase in body fat content was associated with about 9 pmol/l increase in E2 on day 0, 10 pmol/l increase in midcycle E2 and almost 15 pmol/l increase in E2 on day -1.

### *Discussion*

To our knowledge this is the first study which clearly demonstrates the

relationship between the amount of fat accumulated in woman's body and the entire cycle estradiol profile in a relatively large sample of healthy, well-nourished but not obese women. This is also the first study, which demonstrates how this relationship can be modified by energy balance.

In healthy non-obese women, we found non-linear association between body fat percentage and levels of estradiol during menstrual cycle. Women with very low (below 22%) and high body fat (above 31%) had 25 to 35% lower levels of follicular and mid-cycle estradiol. Lower levels were also observed for days 0 and -1. Furthermore, when the analysis was restricted to women characterized by body fat percentage lower than 31%, we found a positive linear relationship between fat accumulation and estradiol levels. The 10% increase in accumulated body fat was associated with 5 to 7 pmol/l increase in estradiol levels.

Interesting differences in the association between body fat and estradiol levels were demonstrated between groups of women with positive and negative energy balance. Whereas in women with negative energy balance no relationship between body fat and estradiol levels was found, in women with positive energy balance the association was even stronger than in case of the whole sample of women. Significant differences between body fat groups were observed in all indices of estradiol. Women with very low and high body fat had 30 to 45% lower levels of E2 than women with low and average body fat. Additionally in non-overweight, positive energy balance women 10% increase in body fat was associated with 7 to even 15 pmol/l increase in estradiol levels.

Levels of estradiol during menstrual cycle and especially during its follicular phase are related to follicular diameter, oocyte quality and endometrium morphology and thickness (Cahill et al., 2000, Ohno and Fujimoto, 1998). Lower levels of E2 during the menstrual cycles and during ovulation co-occur very frequently with lower pregnancy rate both in healthy, naturally conceiving women (Li et al., 2001, Lipson and Ellison, 1996, Lu et al., 1999; Venners et al 2006) and women undergoing in vitro fertilization procedure (Blazar et al., 2004, Chen et al., 2003). It can be estimated that 25 to 35% difference in estradiol levels in very low and high body fat group observed in our study can be translated into 2 to 3 times lower probability of conception (Lipson and Ellison, 1996).

Results of our study clarify and extend the accumulated evidence on the association between nutritional status and levels of reproductive steroids. Several studies conducted on different groups of premenopausal women (diabetic, obese, dieting, very lean and normal weight) demonstrated contradictory results (Barnett et al., 2001, 2002; Brunning et al., 1992; Furberg et al., 2005; Zanker and Swaine, 1998, Howard et al., 1987, Potischman et al., 1996, Thomas et al., 1997a, b; Westhoff et al., 1996). This inconsistency partly results from methodological limitations, especially calculations of mean E2 levels based on no more than a few samples from each menstrual cycle. Due to substantial intra-cycle variation in E2 levels, such sampling is vastly insufficient and can lead to errors in estimating mean E2 levels for individual women (Jasienska & Jasienski in press). In most of the studies estradiol levels were analyzed from a single blood sample, whereas Williams et al. (2002), relying on repeated measurements of



reproductive steroids in 34 women, showed that at least eight samples taken from a single subject are necessary to detect about 80% of biological variation in estradiol levels during a particular menstrual cycle. In our study, the number of samples taken from a single subject considerably exceeded this requirement.

Another limitation of other studies is using BMI as the indicator of nutritional status. BMI does not represent sufficient information about nutritional status and accumulated body fat (Kyle et al., 2003, Piers et al., 2000). Frequently, individuals classified as overweight on the basis on the BMI criteria are of normal adiposity, especially when characterized by high muscle mass (Hortobagyi et al., 1994, Nevill et al., 2005, Wit and Bush, 2005). Conversely, women classified as normal using BMI criteria frequently have increased adiposity (Frankenfield et al. 2001). In our study, this limitation was omitted by direct measurement of body fat percentage. Lack of relationship between estradiol and BMI noted in our study and positive relationship between E2 and percentage of body fat observed at the same time, provides further evidence that BMI may be a poor indicator of nutritional status.

Additionally to the careful measurements of estradiol levels and nutritional status assessment we were able to estimate the energy balance of women based on the changes in body fat during the observational period. This estimation allowed us to investigate the interactions between nutritional status, energy balance and the entire cycle estradiol levels, which to our knowledge were never reported before. Energy balance was shown to influence ovarian steroids profiles in several studies (Ellison et al., 1989, Lager and Ellison, 1990, Panter-Brick, 1993). Our results indicate that negative

energy balance caused by increased physical activity and/or inadequate calories intake has confounding effect on the association between nutritional status and reproductive hormone levels. This fact can explain lack of the relationship between nutritional status and levels of reproductive hormones demonstrated in populations or groups of women characterized by high physical activity (Jasienska, 1998, Jasienska and Ellison, 2004, Lager and Ellison, 1990), women losing weight due to voluntarily caloric restriction (Lager and Ellison, 1990) and women from hunter gatherer and horticulturalist groups in Africa and Nepal (Ellison et al, 1989, Panter-Brick, 1993) who experienced periods of restricted caloric intake and high physical activity. In contrast, association between body fat and reproductive hormones frequently was demonstrated in women from western population who generally have higher energy intake and lower physical activity, none of these studies however controlled energy balance (Barnett et al., 2001, 2002; Brunning et al., 1992; Furberg et al., 2005; Zanker and Swaine, 1998).

In our study, no linear association between E2 and body fat was found in a group of overweight women, but the levels of E2 in these women were significantly lower than in normal adiposity women. This finding corresponds with results of other studies concerned with levels of E2 in overweight and obese women. Drop in E2 surge was noted in studies by Grenman et al. (1986), Kopelman et al. (1980), Leenen et al. (1994) and recently by Tworoger et al. (2006). Several authors demonstrated that in women increased adiposity and obesity are related to high androgenic activity (Evans et al., 1983, Hauner et al., 1988, Norman and Clark 1998, Wabitsch et al., 1995). This may explain the inversion in the pattern of the association between body fat percentage and

estradiol levels observed in our study. In accordance with our results, Wang et al. (2000) showed that in underweight (BMI <20) and overweight (BMI>25) women, the probability of pregnancy during assisted reproduction treatment was about 20% lower comparing to women of normal range of BMI. Our high body fat group corresponded closely to overweight category in study by Wang et al. (see body fat % to BMI recalculation in Gallagher et al, 2000). Additionally, recently Gesink Law et al. (2006) demonstrated significantly reduced fecundity in overweight and obese women of reproductive age. Since the estradiol levels are highly related both to endometrial thickness and ovum quality, our results suggest that lower fecundity, conception and pregnancy rates in underweight, overweight and obese women can be mediated by unfavorable estradiol environment.

Decreased levels of estradiol in overweight and obese women found in our study have significant clinical implications for breast cancer research. Recently, Baer et al. (2006) in the large cohort of premenopausal women demonstrated the significant relationship between the energy status during childhood and adolescence and the risk of breast cancer. In women from the most overweight group during the childhood and the adolescence the risk of premenopausal breast cancer was almost 50% lower than in women from the leanest group. The same pattern was also shown in study by Michels et al. (2006), in which BMI at the age of 18 was the strongest predictor of premenopausal breast cancer risk. Our results point to the possible hormonal mechanism of the observed relationship. Since the high estradiol level is regarded the major mediator of breast cancer and the body mass index highly correlates with the amount of

accumulated body fat it can be hypothesized that the reduction in the premenopausal breast cancer risk is mediated through the reduction of the estradiol levels in overweight and obese women.

Our results confirm the hypothesis about the regulatory role of nutritional status on potential fertility in reproductive age women, but only in those with positive energy balance. In women of low and normal adiposity, we observed a positive association between the amount of accumulated body fat and levels of estradiol during menstrual cycle. Similar effect was demonstrated in several different studies (Barnett et al., 2001, 2002; Brunning et al., 1992; Zanker and Swaine, 1998), but in these studies nutrition status was assessed by BMI and they did not control for the effect of energy balance

In women of reproductive age, energetic resources are partitioned in order to maintain normal both physiological processes and reproduction. Increased energy requirements of reproductive processes can be partially supported by energetic reserves stored in women's bodies in the form of fat. This fat depot is formulated in favorable environmental conditions when energy intake is high, energy expenditure of physical activity is low and total energy balance is positive. Consequences of inadequate nutritional status of women during pregnancy are highly adverse both to the child and to the mother. Low body mass of a woman prior to and during pregnancy is associated with high risk of preterm labor, intrauterine growth retardation, low birth weight of an infant and maternal depletion syndrome (Ehrenberg et al., 2003; Jelliffe and Maddocks, 1964; Kramer, 2003; Winkvist et al., 1992). Short-term reproductive suppression in women with low energy reserves may function as an evolutionary adaptation protecting

against these risks and improving chances of successful reproduction in the future (Ellison 1990, Ellison, 2001, Jasienska 2003).

### *Funding*

Norwegian Cancer Society (TP 49 257 project no. 05087 to I.T.); Foundation for the Norwegian Health and Rehabilitation Organisations (59010-2000, 2001, 2002 to I.T.)

### *Acknowledgements*

This project was supported by a grant nr 2 P05D 017 28 from the Polish Committee for Scientific Research to Grazyna Jasienska and from the Norwegian Cancer Society. We are grateful to women who participated in the study, and to students of the Faculty of Public Health, Jagiellonian University who worked as research assistants, Kolastyna SA and Yves Rocher Poland.

*References*

Baird DD, Wilcox AJ, Weinberg CR, Kamel F, McConnaughey DR, Musey PI, Collins DC (1997) Preimplantation hormonal differences between the conception and non-conception menstrual cycles of 32 normal women. *Hum Reprod* 12, 2607-2613.

Baird DD, Weinberg CR, Zhou H, Kamel F, McConnaughey DR, Kesner JS, Wilcox AJ (1999) Preimplantation urinary hormone profiles and the probability of conception in healthy women. *Fertil Steril* 71, 40-49

Barnett JB, Woods MN, Rosner B, McCormack C, Longcope C, Houser RF, Gorbach SL (2001) Sex Hormone Levels in Premenopausal African-American Women With Upper and Lower Body Fat Phenotypes. *Nutr Cancer* 41, 47-56.

Barnett JB, Woods MN, Rosner B, McCormack C, Floyd L, Longcope C, Gorbach SL (2002) Waist-to-Hip Ratio, Body Mass Index and Sex Hormone Levels Associated with Breast Cancer Risk in Premenopausal Caucasian Women. *J Med Sci* 2, 170-176.

Bear HJ, Colditz GA, Rosner B (2005) Body fatness during childhood and adolescence and incidence of breast cancer in premenopausal women: a prospective cohort study. *Breast Cancer Res* 7, R314-R325.

Blazar AS, Hogan JW, Frankfurter D, Hackett R, Keefe DL (2004) Serum estradiol positively predicts outcomes in patients undergoing in vitro fertilization. *Fertil Steril* 81, 1707-1709.

Brunning PF, Bonfrer JMG, Hart AAM, van Noord PAH, van der Hoeven H, Collette HJA, Battermann JJ, de Jong-Bakker M, Nooijen WJ, de Waard F (1992) Body Measurements, Estrogen Availability and the Risk of Human Breast Cancer: a Case-Control Study. *Int J Cancer* 51, 14-19.

- Budak E, Fernandez Sanchez M, Bellver J, Cervero A, Simon C, Pellicer A (2006) Interactions of the hormones leptin, ghrelin, adiponectin, resistin, and PYY3-36 with the reproductive system. *Fertil Steril* 85, 1563-1581.
- Burrows M, Bird S (2000) The Physiology of the Highly Trained Female Endurance Runner. *Sports Med* 30, 281-300.
- Cahill DJ, Wardle PG, Harlow CR, Hunt LP, Hull MG (2000) Expected contribution to serum oestradiol from individual ovarian follicles in unstimulated cycles. *Hum Reprod* 15, 1909-12.
- Caprio M, Fabbrini E, Isidori AM, Aversa A, Fabbri A (2001) Leptin in reproduction. *Trends in Endocrin Met* 12, 65-72.
- Chen EC, Brzyski RG (1999) Exercise and reproductive dysfunction. *Fertil Steril* 71, 1-6.
- Chen CH, Zhang X, Barnes R, Confino E, Milad M, Puscheck E, Kazer RR (2003) Relationship between peak serum estradiol levels and treatment outcome in vitro fertilization cycles after embryo transfer on day 3 or day 5. *Fertil Steril* 80, 75-79.
- De Souza MJ, Williams NI (2004) Physiological aspects and clinical sequel of energy deficiency and hypoestrogenism in exercising women. *Hum Reprod Update* 10, 433-448.
- Dorgan JF, Reichman ME, Judd JT, Brown C, Longcope C, Schatzkin A, Albanes D, Campbell WS, Franz C, Kahle L, Taylor PR (1995) The relation of body size to plasma levels of estrogens and androgens in premenopausal women (Maryland, United States). *Cancer Cause Control* 6, 3-8.
- Ehrenberg HM, Dierker L, Milluzzi C, Mercer BM (2003) Low maternal weight, failure to thrive in pregnancy, and adverse pregnancy outcomes. *Am J Obstet and Gynecol* 189,

1726-1730.

Elias AN, Wilson AF (1993) Exercise and gonadal function. *Hum Reprod* 8, 1747-1761.

Ellison PT (1988) Human Salivary Steroids: Methodological Considerations and Applications in Physical Anthropology. *Yearb Phys Anthropol* 31, 115-142.

Ellison PT (1990) Human Ovarian Function and Reproductive Ecology: New Hypotheses. *Am Anthropol* 92, 933-952.

Ellison PT (1991) Reproductive ecology and human fertility. In Lasker GW, Mascie-Taylor CGN (eds) *Applications of biological anthropology to human affairs*. New York: Cambridge University Press, p 14-54.

Ellison PT (1999) Reproductive ecology and reproductive cancers. In Panter-Brick C, C.M. Worthman CM (eds) *Hormones, Health, and Behavior. A Socio-ecological and Lifespan Perspective*. New York: Cambridge University Press, pp. 184-209.

Ellison, P.T.: *On Fertile Ground*. Cambridge: Harvard University Press, 2001.

Ellison PT, Peacock NR, Lager C (1989) Ecology and ovarian function among Lese women of the Ituri Forest, Zaire. *Am J Phys Anthropol* 78, 519-526.

Evans DJ, Hoffmann RG, Kalkhoff RK, Kissebah AH (1983) Relationship of androgenic activity to body fat topography, fat cell morphology, and metabolic aberrations in premenopausal women. *J Clin Endocr Metab* 57, 304-310.

Frankenfield DC, Rowe WA, Cooney RN, Smith JS, Becker D (2001) Limits of body mass index to detect obesity and predict body composition.

*Nutrition* 17, 26-30.

Frish RE (1984) Body Fat, Puberty and Fertility. *Biol Rev* 59, 161-188.



Furberg A, Jasienska G, Bjurstam N, Torjesen PA, Emaus A, Lipson SF, Ellison PT, Thune I (2005) Metabolic and hormonal profiles: HDL cholesterol as a plausible biomarker of breast cancer risk. The Norwegian EBBA Study. *Cancer Epidem Biomar* 14, 33-40.

Gallagher D, Heymsfield SB, Heo M, Jebb SA, Murgatroyd PR, Sakamoto Y (2000) Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *Am J Clin Nutr* 72, 694-701.

Gesink Law DC, Maclehose RF, Longnecker MP (2006) Obesity and time to pregnancy. *Hum Reprod* 22, 414-420.

Grenman S, Rönnemaa T, Irjala K, Kaihola HL, Grönroos M (1986) Sex Steroid, Gonadotropin, Cortisol, and Prolactin Levels in Healthy Massively Obese Women: Correlation with Abdominal Fat Cell Size and Effect of Weight Reduction. *J Clin Endocr Metab* 63, 1257-1261.

Hauer H, Ditschuneit HH, Pal SB, Moncayo R, Pfeiffer EF (1988) Fat distribution, endocrine and metabolic profile in obese women with and without hirsutism. *Metabolism* 37, 281-286.

Hortobagyi T, Israel RG, O'Brien KF (1994) Sensitivity and specificity of the Quetelet index to assess obesity in men and women. *Eur J Clin Nutr* 48, 369-375.

Howard BV, Pan XR, Harper I, Kuusi T, Taskinen MR (1987) Lack of sex differences in high density lipoproteins in Pima Indians. Studies of obesity, lipase activities, and steroid hormones. *Arteriosclerosis* 7, 292-300.

Ivandi A, Prpi -Kri evac, Su i M, Juri M (1998) Hyperinsulinemia and Sex

Hormones in Healthy Premenopausal Women: Relative Contribution of Obesity, Obesity Type, and Duration of Obesity. *Metabolism* 47, 13-19.

Jasienska G, Ellison PT (1998) Physical work causes suppression of ovarian function in women *P Roy Soc Lond B Bio* 265, 1847-1851

Jasienska G (2001) Why Energy Expenditure Causes Reproductive Suppression in Women. An Evolutionary and Bioenergetic Perspective. In Ellison P.T. (ed) *Reproductive Ecology and Human Evolution*. Aldine de Gruyter, New York, USA, pp. 59-84.

Jasienska G (2003) Energy metabolism and the evolution of reproductive suppression in the human female. *Acta Biotheor* 51, 1-18.

Jasienska G, Ellison PT (1998) Physical work causes suppression of ovarian function in women. *P R Soc B* 265 1847-1851.

Jasienska G, Ellison PT (2004) Energetic factors and seasonal changes in ovarian function in women from rural Poland. *Am J Hum Biol* 16, 563-580.

Jasienska G, Jasienski M (2007) Inter-population, inter-individual, inter-cycle, and intra-cycle natural variation in progesterone levels: a quantitative assessment and implications for population studies. *Am J Hum Biol* (in press).

Jasienska G, Ziolkiewicz A, Thune I, Lipson SF, Ellison P T (2006) Habitual physical activity and estradiol levels in women of reproductive age. *Eur J Cancer Prev* 15, 439-445.

Jelliffe DB, Maddocks I (1964) Notes on Ecologic Malnutrition In the New Guinea Highlands. *Clin Pediatr* 35, 432-438.

Kirchengast S, Huber J (2004) Body composition characteristics and fat distribution patterns in young infertile women *Fertil Steril* 81, 539-544.

Kopelman PG, Pilkington TR, White N, Jeffcoate SL (1980) Abnormal sex steroid secretion and binding in massively obese women *Clin Endocrinol* 12, 363-369.

Kramer MS (2003) The Epidemiology of Adverse Pregnancy Outcomes: An Overview *J Nutr* 133, 1952S-1956S.

Kyle U, Schutz Y, Dupertuis YM, Pichard C (2003) Body Composition Interpretation: Contributions of the Fat-Free Mass Index and the Body Fat Mass Index *Nutrition* 19, 597-604.

Lager C, Ellison PT (1990) Effect of moderate weight loss on ovarian function assessed by salivary progesterone measurements *Am J Hum Biol* 2, 303-312.

Leenen R, van der Kooy K, Seidell JC, Deurenberg P, Koppeschaar HPF (1994) Visceral Fat Accumulation in Relation to Sex Hormones in Obese Men and Women Undergoing Weight Loss Therapy *J Clin Endocr Metab* 78, 1515-1520.

Li H, Nakajima ST, Chen J, Todd HE, Overstreet JW, Lasley BL (2001) Differences in hormonal characteristics of conceptive versus nonconceptive menstrual cycles *Fertil Steril* 75, 549-553.

Lipson SF, Ellison PT (1996) Comparison of salivary steroid profiles in naturally occurring conception and non-conception cycles. *Hum Reprod* 11, 2090-2096.

Lu Y, Bentley GR, Gann PH, Hodges KR, Chatterton RT (1999) Salivary estradiol and progesterone levels in conception and nonconception cycles in women: evaluation of a new assay for salivary estradiol *Fertil Steril* 71, 863-868.

Michels KB, Terry KL, Willet WC (2006) Longitudinal study on the role of body size in premenopausal breast cancer *Archiv Intern Med* 166, 2395-2402.

Miller KK, Grinspoon S, Gleysteen S, Grieco KA, Ciampa J, Breu J, Herzog DB, Klibanski A (2004) Preservation of Neuroendocrine Control of Reproductive Function Despite Severe Undernutrition. *J Clin Endocr Metab* 89, 4434-4438.

Nevill AM, Stewart AD, Olds T, Holder R (2005) Relationship between adiposity and body size reveals limitations of BMI *Am J Phys Anthropol* 129, 151-156.

Norman RJ, Clark AM (1998) Obesity and reproductive disorders: a review *Reprod Fert Develop* 10, 55-63.

Ohno Y, Fujimoto Y (1998) Endometrial oestrogen and progesterone receptors and their relationship to sonographic appearance of the endometrium. *Hum Reprod* 4, 560-564.

Panter-Brick C, Lotstein DS, Ellison PT (1993) Seasonality of reproductive function and weight loss in rural Nepali women *Hum Reprod* 8, 684-690.

Piers LS, Soares MJ, Frandsen SL, O'Dea K (2000) Indirect estimates of body composition are useful for groups but unreliable in individuals. *Int J Obes Relat Metab Disord* 24, 1145-1152.

Potischman N, Swanson CA, Siiteri P, Hoover RN (1996) Reversal of relation between body mass and endogenous estrogen concentrations with menopausal status. *J Natl Cancer I* 88, 756-758.

Rexford SA, Flier JS (2000) Leptin. *Annu Rev Physiol* 62, 413-437.

Schneider JE (2004) Energy balance and reproduction. *Physiol Behav* 81, 289-317.

Smith GD, Jackson LM, Foster DL (2002) Leptin regulation of reproductive function and

fertility. *Theriogenology* 57, 73-86.

Tataranni PA, Monroe MB, Dueck CA, Traub SA, Nicolson M, Manore MM, Matt KS, Ravussin E (1997) Adiposity, plasma leptin concentration and reproductive function in active and sedentary females. *Int J Obes Relat Metab Disord* 21, 818-821.

Thomas HV, Key TJ, Allen DS, Moore JW, Dowsett M, Fentiman IS, Wang DY (1997) Reversal of Relation Between Body Mass and Endogenous Estrogen Concentrations With Menopausal Status. *J Natl Cancer I* 89, 396-397.

Thomas HV, Key TJ, Allen DS, Moore JW, Dowsett M, Fentiman IS, Wang DY (1997) A prospective study of endogenous serum hormone concentrations and breast cancer risk in premenopausal women on island of Guernsey *Brit J Cancer* 76, 401-405.

Twooroger SS, Eliassenn AH, Missmer SA, Baer H, Rich-Edwards J, Michels KB, Barbieri RL, Dowsett M, Hankinson SE (2006) Birth weight and body size through life in relation to sex hormones and prolactin concentrations in premenopausal women *Cancer Epidem Biomar* 15, 2494-2501.

Ukkola O, Gagnon J, Rankinen T, Thompson PA, Hong Y, Leon AS, Rao DC, Skinner JS, Wilmore JH, Bouchard C (2001) Age, body mass index, race and other determinants of steroid hormone variability: the HERITAGE Family Study *Eur J Endocrinol* 145, 1-9.

van Binsbergen CJM, Coelingh Bennink HJT, Odink J, Haspels AA, Koppeschaar HPF (1990) A Comparative and Longitudinal Study on Endocrine Changes Related to Ovarian Function in Patients with Anorexia Nervosa. *J Clin Endocr Metab* 71: 705-711.

Venners SA, Liu X, Perry MJ, Korrick SA, Li ZP, Yang F, Yang JH, Lasley BL, Xu XP, Wang XB (2006) Urinary estrogen and progesterone metabolite concentrations in

menstrual cycles of fertile women with non-conception, early pregnancy loss or clinical pregnancy. *Hum Reprod* 21, 2272-2280.

Wabitsch M, Hauner H, Heinze E, Böckmann Benz R, Mayer H, Teller W (1995) Body Fat Distribution and Steroid Hormone Concentrations in Obese Adolescent Girls before and after Weight Reduction. *J Clin Endocr Metab* 80, 3469-3475.

Wade GN, Jones JE (2004) Neuroendocrinology of nutritional infertility. *Am J Physiol* 287, R1277-R1296.

Wade GN, Schneider JE, Li H-Y (1996) Invited review: Control of fertility by metabolic cues. *Am J Physiol* 270, E1-E19.

Wang JX, Davies M, Norman RJ (2000) Body mass and probability of pregnancy during assisted reproduction treatment: retrospective study. *BMJ* 321, 1320-1321.

Warren MP (1983) Effects of undernutrition on reproductive function in the human. *Endocr Rev* 4, 363-377.

Warren MP, Perlroth NE (2001) The effects of intense exercise on the female reproductive system. *J Endocrinol* 170, 3-11.

Westhoff C, Gentile G, Lee J, Zacur H, Helbig D (1996) Predictors of Ovarian Steroid Secretion in Reproductive-Age Women. *Am J Epidemiol* 144, 381-388.

Williams AE, Maskarinec G, Franke AA, Stanczyk FZ (2002) The temporal reliability of serum estrogens, progesterone, gonadotropins, SHBG and urinary estrogen and progesterone metabolites in premenopausal women *BMC Women's Health* 2, 13

Winkvist A, Rasmussen KM, Habicht J-P (1992) A new definition of the maternal depletion syndrome. *Am J Pub Health* 82, 691-694.

Wit K, Bush EA (2005) College Athletes with an Elevated Body Mass Index Often Have a Higher Upper Arm Muscle Area, but Not Elevated Triceps and Subscapular Skinfolde. J Am Diet Assoc 105, 599-602.

Yoshimura Y, Wallach EE (1987) Studies of the mechanism(s) of mammalian ovulation. Fertil Steril 47, 22-34.

Zanker CL, Swaine IL (1998) The Relationship Between Serum Oestradiol Concentration and Energy Balance in Young Women Distance Runners. Int J Sports Med 19, 104-108.

Table I

General characteristics of study participants. Mean and standard deviation (in parentheses) for four consecutive body fat groups compared by one-way ANOVA tests.

	All women N=141	Very low body fat ≤22% N=36	Low body fat ≤26.5% N= 35	Average body fat ≤30.8% N= 35	High body fat >30.8 N=35
Age [years]	29.8 (3.34)	29.6 (3.07)	29.7 (3.75)	29.4 (3.25)	30.6 (3.25)
Age of menarche [years]	13.4 (1.31)	13.6 (1.37)	13.2 (1.26)	13.3 (1.35)	13.6 (1.28)
Usual cycle length [days]*	29.2 (3.13)	30.2 (3.82)	29.6 (3.52)	28.1 (2.16)	29.1 (2.46)
Birth weight [kg]*	3.27 (0.611)	3.34 (0.626)	3.22 (0.502)	3.24 (0.500)	3.28 (0.819)
Birth length [cm]	53.2 (4.38)	53.1 (3.74)	53.7 (2.90)	52.6 (4.37)	53.4 (6.37)
Height [cm]	163.1 (6.51)	161.7 (6.12)	162.1 (6.30)	164.2 (7.11)	164.3 (6.26)
Weight [kg]*	60.1 (8.37)	51.4 (3.67)	56.3 (3.10)	62.5 (4.40)	70.3 (6.15)
BMI [kg/m <sup>2</sup> ]*	22.6 (2.84)	19.7 (1.25)	21.4 (1.47)	23.2 (1.65)	26.0 (1.92)
Body fat% *	26.4 (6.52)	17.9 (3.19)	24.4 (1.19)	28.8 (1.22)	34.6 (2.31)
BUR	1.16 (0.037)	1.15 (0.037)	1.15 (0.036)	1.16 (0.036)	1.17 (0.037)
WHR *	0.72 (0.043)	0.70 (0.029)	0.71 (0.037)	0.72 (0.041)	0.75 (0.051)
Physical activity [MET*hour/day]	5.8 (2.95)	4.7 (1.47)	6.1 (3.09)	5.9 (2.95)	6.4 (3.68)
Number of cigarettes	1.8 (4.44)	1.1 (3.65)	1.9 (4.30)	2.1 (5.20)	2.0 (4.59)

\* differences between body fat groups at p<0.05



Table II

Estradiol indices in consecutive body fat groups. Mean and standard deviation (in parentheses) for four consecutive body fat groups compared by one-way ANOVA tests.

	All	Very low	Low	Average	High
E2 [pmol/l]	women	body fat	body fat	body fat	body fat
	N=130	N=34	N=34	N=31	N=31
mean E2	18.5 (8.99)	17.0 (8.45)	20.3 (9.70)	20.7 (9.32)	16.1 (7.87)
mid-cycle E2*	22.1 (11.43)	19.7 (10.54)	25.4 (11.77)	25.2 (12.51)	18.2 (9.20)
day -1 E2 *	33.3 (18.27)	29.1 (15.38)	39.2 (17.75)	38.8 (22.26)	26.6 (14.22)
day 0 E2 *	17.1 (11.36)	15.2 (10.33)	19.4 (11.74)	20.4 (13.16)	13.4 (8.70)
mean follicular E2 *	18.7 (8.97)	17.0 (8.08)	20.9 (10.27)	21.1 (9.71)	15.9 (6.56)
mean luteal E2	18.3 (9.84)	17.1 (9.60)	19.6 (10.01)	20.2 (9.39)	16.3 (10.26)

\*differences between body fat groups significant at  $p < 0.05$

Table III

General characteristics in consecutive body fat quartiles of women with positive and negative energy balance. Mean and standard deviation (in parentheses) for four consecutive body fat groups compared by one-way ANOVA tests.

	All women N=131	Positive Energy Balance N=75	Negative Energy Balance N=56
<b>Age [years]</b>	29.9 (3.36)	30.1 (3.49)	29.6 (3.17)
Very low fat	29.4 (3.01)	29.9 (3.04)	28.9 (2.91)
Low fat	29.9 (3.77)	30.7 (3.83)	28.9 (3.58)
Average fat	29.1 (3.34)	28.7 (3.35)	29.8 (3.33)
High fat	30.9 (3.13)	31.3 (3.42)	30.5 (2.83)
<b>Age of menarche [years]</b>	13.4 (1.34)	13.3 (1.22)	13.5 (1.48)
Very low fat	13.7 (1.42)	13.5 (1.64)	13.9 (0.88)
Low fat	13.1 (1.29)	13.1 (1.20)	13.1 (1.46)
Average fat	13.2 (1.31)	13.3 (1.10)	13.0 (1.60)
High fat	13.7 (1.27)	13.5 (0.80)	13.9 (1.63)
<b>Usual cycle length [days]**</b>	29.3 (3.25)	28.8 (3.02)**	30.0 (3.45)**
Very low fat	30.3 (4.04)	29.3 (3.18)	32.3 (4.90)
Low fat	29.6 (3.64)	29.3 (3.97)	29.9 (3.23)
Average fat	28.3 (2.23)	27.8 (2.04)	29.0 (2.35)
High fat	29.0 (2.56)	28.6 (2.37)	29.4 (2.77)
<b>Birth weight [kg]**</b>	3.26 (0.627)	3.38 (0.500)**	3.07 (0.748)**
Very low fat	3.38 (0.641)	3.58 (0.547)	3.01 (0.702)
Low fat	3.25 (0.478)	3.18 (0.477)	3.34 (0.484)

Average fat	3.12 (0.519)	3.21 (0.493)	3.01 (0.554)
High fat	3.29 (0.847)	3.57 (0.349)	2.86 (1.177)
<b>Birth length [cm]</b>	53.3 (4.59)	53.8 (4.08)	52.5 (5.36)
Very low fat	53.3 (3.99)	53.7 (4.67)	52.5 (1.76)
Low fat	53.8 (2.66)	53.9 (2.94)	53.6 (2.32)
Average fat	52.9 (4.73)	52.9 (5.49)	53.0 (3.71)
High fat	53.0 (6.76)	54.7 (2.22)	50.5 (10.20)
<b>Height [cm]</b>	163.1 (6.53)	163.3 (6.71)	162.7 (6.33)
Very low fat	161.9 (6.11)	162.4 (6.62)	160.5 (5.04)
Low fat	162.6 (6.18)	162.3 (6.16)	162.9 (6.42)
Average fat	163.7 (7.29)	163.6 (7.07)	163.9 (7.84)
High fat	164.2 (6.53)	165.1 (7.21)	163.2 (5.80)
<b>Weight [kg]*</b>	60.0 (8.40)	60.0 (8.51)	60.1 (8.39)
Very low fat	51.5 (3.64)	51.4 (3.72)	51.1 (3.45)
Low fat	55.9 (3.03)	56.1 (3.31)	55.6 (2.70)
Average fat	62.3 (4.61)	62.6 (3.81)	61.9 (5.64)
High fat	70.1 (6.39)	71.1 (6.44)	69.2 (6.38)
<b>BMI [kg/m<sup>2</sup>]*</b>	22.4 (3.43)	22.5 (2.90)	22.3 (4.07)
Very low fat	19.7 (1.30)	19.5 (1.33)	19.9 (1.34)
Low fat	21.2 (1.34)	21.3 (1.35)	21.0 (1.36)
Average fat	22.6 (4.20)	23.5 (1.57)	21.5 (6.13)
High fat	26.0 (2.00)	26.1 (2.10)	26.0 (1.95)
<b>Body fat% *</b>	26.3 (6.49)	25.9 (6.57)	26.9 (6.29)
Very low fat	17.6 (3.04)	17.2 (2.84)	18.4 (3.40)

Low fat	24.1 (1.23)	24.2 (1.19)	24.0 (1.32)
Average fat	28.6 (1.31)	28.7 (1.19)	28.4 (1.48)
High fat	34.4 (2.30)	34.4 (2.38)	34.4 (2.30)
<b>BUR</b>	1.16 (0.037)	1.16 (0.034)	1.16 (0.039)
Very low fat	1.15 (0.039)	1.14 (0.027)	1.17 (0.052)
Low fat	1.16 (0.037)	1.15 (0.032)	1.16 (0.042)
Average fat	1.16 (0.035)	1.16 (0.038)	1.15 (0.029)
High fat	1.17 (0.036)	1.17 (0.038)	1.17 (0.035)
<b>WHR *</b>	0.72 (0.044)	0.72 (0.046)	0.71 (0.042)
Very low fat	0.70 (0.029)	0.70 (0.027)	0.71 (0.034)
Low fat	0.70 (0.035)	0.71 (0.028)	0.70 (0.044)
Average fat	0.72 (0.043)	0.73 (0.044)	0.71 (0.039)
High fat	0.75 (0.053)	0.76 (0.055)	0.73 (0.046)
<b>Physical activity [MET*hour/day]*</b>	6.0 (3.00)	6.4 (3.22)	5.4 (2.59)
Very low fat*	4.7 (1.47)	5.1 (1.37)	3.9 (1.28)
Low fat	5.9 (2.90)	6.6 (3.15)	5.0 (2.33)
Average fat	6.6 (3.10)	6.7 (3.25)	6.5 (2.97)
High fat	6.6 (3.71)	7.2 (4.42)	6.0 (2.76)
<b>Number of cigarettes</b>	1.3 (4.01)	1.2 (3.46)	1.6 (4.79)
Very low fat	0.7 (2.46)	1.2 (3.10)	0.0 (0.00)
Low fat	1.4 (3.98)	1.3 (2.68)	1.6 (5.43)
Average fat	1.3 (4.23)	0.6 (2.42)	2.2 (5.94)
High fat	2.0 (5.15)	1.9 (5.39)	2.2 (5.07)

---

\* differences between body fat quartiles significant at  $p < 0.001$

\*\* differences between positive and negative energy balance significant at  $p < 0.05$

Table IV

Estradiol indices in consecutive body fat quartiles of women with positive and negative energy balance. Mean and standard deviation (in parentheses) for four consecutive body fat groups compared by one-way ANOVA tests.

	All Women N=120	Positive Energy Balance N=70	Negative Energy Balance N=50
<b>mean E2</b>	18.3 (8.92)	17.3 (9.16)	19.6 (10.86)
Very low fat	16.8 (8.58)	14.0 (7.52)*	20.8 (9.06)
Low fat	20.7 (9.61)	20.2 (10.86)	21.3 (7.81)
Average fat	19.7 (8.96)	20.7 (9.80)	18.0 (7.52)
High fat	15.7 (7.86)	13.4 (4.43)*	18.2 (9.96)
<b>mid-cycle E2</b>	21.9 (11.27)	20.2 (11.86)	24.4 (10.12)
Very low fat	19.5 (10.83)*	16.0 (10.32)*	24.3 (10.34)
Low fat	26.1 (11.45)	24.4 (12.48)	28.7 (9.64)
Average fat	24.2 (12.12)	24.7 (13.3)	23.4 (10.45)
High fat	17.7 (8.74)*	14.4 (6.37)*	21.2 (9.76)
<b>day -1 E2</b>	33.1 (17.85)	30.9 (19.1)	35.7 (15.71)
Very low fat	29.4 (15.78)*	24.9 (16.00)*	34.4 (13.49)
Low fat	39.8 (17.67)	39.1 (19.02)	40.77 (16.21)
Average fat	37.1 (21.41)	36.0 (21.87)	38.8 (21.65)
High fat	25.8 (12.93)*	22.1 (14.59)*	29.7 (9.93)
<b>day 0 E2</b>	16.8 (11.21)	14.7 (11.10)	19.9 (10.86)
Very low fat	14.6 (10.24)	11.4(8.10)*	18.7 (12.03)
Low fat	20.0 (11.52)	15.8 (11.25)	26.5 (8.89)
Average fat	19.6 (12.90)	20.8 (14.72)	17.7 (9.81)

High fat	13.1 (8.61)*	9.9 (4.48)*	16.7 (10.80)
<b>mean follicular E2</b>	18.6 (8.94)	17.7 (9.46)	19.7 (8.24)
Very low fat	17.2 (8.33)	14.6 (7.47)*	21.1 (8.68)
Low fat	21.1 (10.30)	21.1 (11.63)	21.1 (8.44)
Average fat	20.3 (9.33)	20.9 (9.93)	19.4 (8.78)
High fat	15.6 (6.53)	13.8 (4.96)*	17.4 (7.62)
<b>mean luteal E2</b>	17.9 (9.67)	16.7 (9.54)	19.5 (9.79)
Very low fat	16.4 (9.16)	13.4 (7.91)*	20.6 (9.78)
Low fat	20.2 (9.79)	19.2 (11.13)	21.6 (7.59)
Average fat	19.0 (9.17)	20.4 (10.36)	16.7 (6.64)
High fat	15.8 (10.34)	13.0 (5.32)*	18.9 (13.44)

---

\* differences between body fat quartiles significant at  $p < 0.05$

Table V

The association between estradiol indices and body fat percentage in women with positive and negative energy balance and body fat below 31%. Results of simple regression analysis.

	Positive Energy Balance			Negative Energy Balance		
	N	R <sup>2</sup>	p	N	R <sup>2</sup>	p
Mean E2	55	0.13	0.006	36	0.001	NS
Follicular E2	53	0.11	0.017	36	0.003	NS
Mid-cycle E2	55	0.15	0.003	36	0.010	NS
Day -1 E2	53	0.14	0.006	36	0.039	NS
Day 0 E2	53	0.14	0.005	35	0.010	NS
Luteal E2	55	0.14	0.005	36	0.002	NS

Figure 1




Figure 1. The association between body fat percentage and E2 levels in women with the body fat percentage below 31%.



Figure 2




Figure 2. E2 profiles in very low, low, average and high body fat groups of women with positive energy balance.

Figure 3




Figure 3. The association between body fat percentage and E2 levels in women with positive energy balance and body fat percentage below 31%.