



Does Sleep Play a Role in Memory Consolidation? A Comparative Test

Citation

Capellini, Isabella, Patrick McNamara, Brian T. Preston, Charles L. Nunn, and Robert A. Barton. 2009. Does sleep play a role in memory consolidation? A comparative test. PLoS One 4(2): e4609. doi:10.1371/journal.pone.0004609.

Published Version

<http://dx.doi.org/10.1371/journal.pone.0004609>

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:2703476>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Does Sleep Play a Role in Memory Consolidation? A Comparative Test

Isabella Capellini^{1*}, Patrick McNamara², Brian T. Preston³, Charles L. Nunn^{3,4,5}, Robert A. Barton¹

1 Department of Anthropology, Durham University, Durham, United Kingdom, **2** Department of Neurology, Boston University School of Medicine and Boston VA Healthcare System, Boston, Massachusetts, United States of America, **3** Department of Anthropology, Harvard University, Cambridge, Massachusetts, United States of America, **4** Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany, **5** Department of Integrative Biology, University of California, Berkeley, California, United States of America

Abstract

Sleep is a pervasive characteristic of mammalian species, yet its purpose remains obscure. It is often proposed that 'sleep is for the brain', a view that is supported by experimental studies showing that sleep improves cognitive processes such as memory consolidation. Some comparative studies have also reported that mammalian sleep durations are higher among more encephalized species. However, no study has assessed the relationship between sleep and the brain structures that are implicated in specific cognitive processes across species. The hippocampus, neocortex and amygdala are important for memory consolidation and learning and are also in a highly activated state during sleep. We therefore investigated the evolutionary relationship between mammalian sleep and the size of these brain structures using phylogenetic comparative methods. We found that evolutionary increases in the size of the amygdala are associated with corresponding increases in NREM sleep durations. These results are consistent with the hypothesis that NREM sleep is functionally linked with specializations of the amygdala, including perhaps memory processing.

Citation: Capellini I, McNamara P, Preston BT, Nunn CL, Barton RA (2009) Does Sleep Play a Role in Memory Consolidation? A Comparative Test. PLoS ONE 4(2): e4609. doi:10.1371/journal.pone.0004609

Editor: Olaf Sporns, Indiana University, United States of America

Received: September 19, 2008; **Accepted:** January 15, 2009; **Published:** February 25, 2009

Copyright: © 2009 Capellini et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by NIMH grant number 1R01MH070415-01A1 to PM, RB and CN and the Max Planck Society (CN, BP). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Isabella.Capellini@durham.ac.uk

Introduction

It has been suggested that sleep is of particular importance to brain processes such as memory consolidation and learning [1–4]. Experimental studies have supported this 'memory consolidation' hypothesis of sleep function by showing that sleep-deprived human and animal subjects perform poorly in learning tasks when compared to individuals that are well rested [2,5]. However, the approach and conclusions of these studies are often criticized, due to the stress associated with sleep deprivation experiments [6,7] and because memory consolidation can also occur in the absence of sleep.

The comparative study of sleep variation offers a complementary approach to investigating potential adaptive functions of sleep [8,9] and most comparative research has focused on sleep durations. The importance of sleep times is reflected by the observation that when sleep deprived, experimental human and animal subjects exhibit a 'sleep rebound' proportional to the amount of sleep lost [10], indicating that the amount of sleep, or of some specific component of sleep, is physiologically relevant. Previous comparative studies have suggested that the great interspecific variation in sleep durations observed in mammals may reflect either functional benefits or ecological constraints, or both [e.g. 9,11,12–15]. Recent analyses on mammalian sleep durations have reported a positive relationship between rapid-eye-movement (REM) sleep and mammalian whole brain volume, which has been taken as support for a cognitive function of sleep [8,12 but see 11]. While these reports are consistent with a memory related function of sleep, the brain is a complex organ, and comparative evidence suggests that functionally specific

regions have changed in size independently of whole brain size [16]. Measures of total brain size or of encephalization are therefore too coarse to substantiate the idea of a functional association between sleep and specific cognitive processes. A potentially more targeted approach is to examine sleep parameters in relation to specific brain regions that play a role in memory consolidation.

Ecologically-imposed needs for increased memory capacity should be reflected by an increase in the size of brain structures that are responsible for memory processing and consolidation [17]. For example, spatial memory is important in animals that hoard food because it improves their ability to retrieve stored food at a later time, which in turn enhances fitness. The hippocampus is one of the most important brain structures involved in spatial memory processing and memory retention, and studies in birds have shown that hippocampal volumes and hippocampal neuron numbers are higher in species and populations that cache food relative to those that do not exhibit such behaviour [17–19]. Similarly, if sleep serves a specific function with regard to memory consolidation and learning, we expect that greater memory-related demands result in a greater need for sleep. Those brain structures that are devoted to memory processing and learning therefore should be positively associated with sleep durations.

Many brain regions are involved in the diverse aspects of memory formation, but those hypothesised to have prominent roles in forming adaptively relevant associations in mammals include the hippocampus, amygdala and neocortex [17,20–24]. During sleep a variety of brain structures are in a highly activated state, including those specifically linked to memory consolidation and learning.

Interactions of the amygdala and hippocampus with one another and the neocortex during NREM sleep are well documented [25], while the outflow from the hippocampus to the neocortex is inhibited during REM but not NREM sleep [22,25]. Recent studies converge on the conclusion that procedural and emotional forms of memory benefit from both REM and NREM sleep, while episodic memory benefits only from NREM sleep [4,26–28].

Here we test the hypothesis that sleep is involved in memory consolidation and learning processes, predicting that evolutionary increases in the relative size of mammalian neocortical, hippocampal, and amygdalar regions will be associated with increased durations of REM and NREM sleep. We test these predictions by conducting phylogenetically controlled analyses of mammalian sleep durations and brain structures.

Materials and Methods

We constructed a dataset of mammalian sleep durations (REM and NREM sleep times in hours/day) from an exhaustive search of the published literature [29; data available at <http://www.bu.edu/phylogeny/index.html>]. In previous analyses of this dataset we found that when sleep was recorded for less than 12 hours, sleep times were significantly underestimated, and that EEG studies tended to have lower estimates of sleep durations relative to non-EEG behavioural studies [11]. We thus restricted our analyses to studies that recorded sleep durations with EEG equipment for at least 12 hours. We excluded monotremes and aquatic mammals because their peculiar sleep architecture may not be comparable to that of terrestrial mammals [9,30].

We extracted data on overall brain volume and the total volumes of individual brain components from a paper that employed uniform measurement procedures across species [31]. Data on neocortical and hippocampal volumes were available for 14 species in our sleep dataset, and data on amygdalar volumes for 13 species. Our final dataset comprised eight primates, one tree shrew, three ‘insectivores’, and two rodents (see Appendix S1). All variables were log-transformed to achieve normality.

Because closely related species tend to exhibit similar sleep durations [11], we implemented statistical methods that explicitly incorporate phylogeny to account for the lack of statistical independence in the data due to common ancestry [32–35]. Specifically we used the program BayesTraits [36,37] to perform a multiple regression analysis with the method of phylogenetic generalized least squares (PGLS). PGLS converts the phylogeny into a variance-covariance matrix of species relationships, which is then used to weight the parameters of regression analysis estimated with maximum likelihood [ML, 36,37]. We based our tests on the mammalian phylogenetic tree by Bininda-Emonds et al. [38] with updated branch lengths [39].

For each brain structure volume we calculated the volume of the remaining brain as a log-transformed difference from total brain volume, and used these volumes to account for the scaling of brain components with total brain volume [40]. We used ‘the rest of the brain’ for each individual structure instead of total brain volume when controlling for scaling effects because total brain volume includes the volume of the structure of interest. Our procedure thus ensured that the brain structure of interest was not represented on the X and Y axes simultaneously. REM and NREM sleep times were tested against each structure volume and corresponding volume of the remaining brain with multiple regression in PGLS. This allowed us to control for both phylogenetic relatedness of species and for scaling effects. We controlled for multiple testing using the false discovery rate test [FDR, 41,42]. All tests were two-tailed with $\alpha = 0.05$.

Results

After controlling for scaling effects, NREM sleep increased with amygdala volume (amygdala: $t_{10} = 4.60$, $p = 0.001$, rest of the brain: $t_{10} = -4.74$, $p < 0.001$, model- $R^2 = 0.70$; Figure 1) while the correlations with neocortex and hippocampus volumes were not significant (neocortex: $t_{11} = -0.77$, $p = 0.458$, rest of the brain: $t_{11} = 0.65$, $p = 0.527$, model- $R^2 = 0.06$; hippocampus: $t_{11} = 1.89$, $p = 0.086$, rest of the brain: $t_{11} = -1.93$, $p = 0.080$, model- $R^2 = 0.26$). We found no significant association between REM sleep durations and any of the brain structures we used (amygdala: $t_{10} = 1.27$, $p = 0.232$, rest of the brain: $t_{10} = -1.39$, $p = 0.195$, model- $R^2 = 0.16$; neocortex: $t_{11} = -1.71$, $p = 0.115$, rest of the brain: $t_{11} = 1.53$, $p = 0.153$, model- $R^2 = 0.22$; hippocampus: $t_{11} = 0.11$, $p = 0.918$, rest of the brain: $t_{11} = -0.26$, $p = 0.793$, model- $R^2 = 0.02$). After controlling for multiple testing, NREM sleep remained significantly correlated with amygdalar volume (FDR estimated threshold of significance: $\alpha = 0.008$).

Discussion

We found that evolutionary increases in NREM sleep durations were correlated with evolutionary increases in the size of the amygdala, and this effect was independent of both scaling effects and phylogeny. We found no evidence of positive relationships between REM sleep and brain structures implicated in memory consolidation and learning. The hippocampus showed a tendency to increase with NREM sleep, although this relationship was not statistically significant ($p < 0.09$). Sample sizes are however




Figure 1. NREM sleep time and amygdala. NREM sleep durations increase with relative amygdalar volumes after accounting for scaling effects [(NREM sleep time) = $1.50 + 0.66 * (\text{amygdala volumes}) - 0.45 * (\text{rest of the brain})$; see text]. The plot shows relative amygdalar volumes, which were calculated with a phylogenetically corrected regression of amygdalar volumes on the rest of the brain, using ML in PGLS (see methods). Species number: (1) *Microcebus murinus*, (2) *Rattus norvegicus*, (3) *Nannospalax ehrenbergi*, (4) *Tupaia glis*, (5) *Callithrix jacchus*, (6) *Pan troglodytes*, (7) *Saimiri sciureus*, (8) *Papio hamadryas*, (9) *Erythrocebus patas*, (10) *Macaca mulatta*, (11) *Tenrec ecaudatus*, (12) *Erinaceus europaeus*, (13) *Aotus trivirgatus*. doi:10.1371/journal.pone.0004609.g001

relatively limited in our analyses, and it would be worthwhile to reassess this relationship once more data on sleep and brain regions have accumulated.

Our results are broadly consistent with electrophysiological, computational and neuroimaging studies that functionally link NREM slow wave sleep with information flow from amygdalar-hippocampal structures to neocortical sites during sleep [3,22,25,26,28,43], but suggest that the amygdala is the key locus of anatomical change in sleep-regulated memory enhancement. These studies converge on a two-step model of memory consolidation that suggests an initial information transfer from the hippocampal-amygdalar complexes to neocortical sites during NREM slow wave sleep, and then a later integration of this information into existing semantic memory networks in the neocortex during REM sleep. Thus both the amygdala and the hippocampus are active during NREM sleep and only the amygdala is active during both sleep states. This may explain why we could detect a stronger link between the amygdala and NREM sleep than between the hippocampus and NREM sleep with currently available data. Insufficient comparative data exist to determine which part of the amygdala correlates most strongly with NREM sleep. The amygdala is a heterogeneous structure comprising nuclei with divergent projection systems [44]. Despite such differences, however, these nuclei have strong reciprocal connections and evolved together in a closely coordinated fashion [45].

We found no association between neocortical volumes and sleep durations. Although this result is surprising, the neocortex includes different subdivisions that undertake different functions [17] and the lack of association with sleep in our results might reflect this composite nature of the neocortex as well as small sample sizes. At present, however, sample sizes on volumes of neocortical subdivisions are not sufficient to investigate this possibility. An alternative hypothesis is that the neocortex might respond more strongly to sleep intensity than to sleep durations (see below).

Our results conflict with interpretations from some previous comparative studies of brain size and sleep durations. For example, Lesku et al. [12] found that mammals with relatively larger brains engage in relatively more REM sleep, and they concluded that this association indicates that REM sleep is important for memory consolidation, learning or other cognitive functions. In contrast, we found that NREM sleep durations, but not REM sleep durations, were linked with brain regions involved in memory processing. Although it is possible that Lesku et al. [12]

identified a more generalized relationship between REM sleep and overall brain function, a subsequent study was unable to confirm this pattern when using more comparable data and different methods [11]. Thus, while our analyses provide some evidence for a memory-related function of sleep in relation to specific brain structures, there is conflicting evidence regarding the involvement of sleep in whole brain function.

Other aspects of sleep architecture are likely to be important in sleep function and evolution, particularly sleep intensity. Slow-wave activity during NREM sleep is considered to be a measure of sleep intensity and is implicated in the homeostatic regulation of sleep; sleep deprived subjects experience increases in slow-wave activity as well as the duration of subsequent sleep periods [46,47]. Thus, memory related demands of sleep might also be met by an increase in sleep intensity. Unfortunately, there are insufficient comparative data currently available to investigate how sleep intensity may interact with sleep durations and brain structure in generating the benefits of sleep.

Our analyses were limited by the availability of data on both brain structures and sleep durations and require confirmation when larger sample sizes become available. Our study nonetheless reveals the first evidence of correlated evolution between sleep and specific brain structures, and provides support for the idea that memory consolidation may be among the ultimate functions of sleep.

Supporting Information

Appendix S1 Sleep & Brain data

Found at: doi:10.1371/journal.pone.0004609.s001 (0.01 MB XLS)

Acknowledgments

We wish to thank Erica Harris and Nikita Patel for their help with the database of sleep traits and logistic support, Chris Venditti and Chris Orban for their advice with the analysis in BayesTraits.

Author Contributions

Conceived and designed the experiments: PM CLN RB. Analyzed the data: IC CLN. Wrote the paper: IC PM BTP CLN RB. Helped collect the data: IC PM BTP CLN RB. Analyzed the data: BTP CLN RB.

References

- Wagner U, Gais S, Born J (2001) Emotional memory formation is enhanced across sleep intervals with high amounts of rapid eye movement sleep. *Learn Memory* 8: 112–119.
- Smith C (1995) Sleep states and memory processes. *Behav Brain Res* 69: 137–145.
- Stickgold R, Walker MP (2007) Sleep-dependent memory consolidation and reconsolidation. *Sleep Med* 8: 331–343.
- Walker MP, Stickgold R (2006) Sleep, memory and plasticity. *Ann Rev Psychol* 57: 139–166.
- Bonnet M (2005) Sleep fragmentation. In: Kushida C, ed. *Sleep deprivation: basic science, physiology and behavior*. New York: Marcel Dekker. pp 103–120.
- Siegel JM (2001) The REM sleep-memory consolidation hypothesis. *Science* 294: 1058–1063.
- Vertes RP (2004) Memory consolidation in sleep: dream or reality. *Neuron* 44: 135–148.
- Zepelin H (1989) Mammalian sleep. In: Kryger MH, Roth T, Dement WC, eds. *Principles and practices of sleep medicine*. Philadelphia: Saunders. pp 30–49.
- Zepelin H, Siegel JM, Tobler I (2005) Mammalian sleep. In: Kryger MH, Roth T, Dement WC, eds. *Principles and practices of sleep medicine*. 4th ed. New York: Saunders. pp 91–100.
- Tobler I (2000) Phylogeny and sleep regulation. In: Kryger MH, Roth T, Dement WC, eds. *Principles and practices of sleep medicine*. Philadelphia: Saunders. pp 72–81.
- Capellini I, Barton RA, McNamara P, Preston BT, Nunn CL (2008) Phylogenetic analysis of the ecology and evolution of mammalian sleep. *Evolution* 62: 1764–1776.
- Lesku JA, Roth TC, Amlaner CJ, Lima SL (2006) A phylogenetic analysis of sleep architecture in mammals: the integration of anatomy, physiology and ecology. *Am Nat* 168.
- Allison T, Cicchetti DV (1976) Sleep in mammals: ecological and constitutional correlates. *Science* 194: 732–734.
- Elgar MA, Pagel MD, Harvey PH (1988) Sleep in mammals. *Anim Behav* 36: 1407–1419.
- Preston BT, Capellini I, McNamara P, Barton RA, Nunn CL (2009) Parasite resistance and the adaptive significance of sleep. *BMC Evolutionary Biology* 9: 7.
- Barton RA, Harvey PA (2000) Mosaic evolution of brain structure in mammals. *Nature* 405: 1055–1057.
- Striedter GF (2005) *Principles of brain evolution*. Sunderland, Massachusetts (USA): Sinauer Associates, Inc. Publishers.
- Krebs JR (1990) Food-storing birds: adaptive specialization in brain and behaviour? *Phil Trans R Soc Lond B* 329: 153–160.
- Roth TCI, Pravosudov VV (2008) Hippocampal volumes and neuron numbers increase along a gradient of environmental harshness: a large-scale comparison. *Proc R Soc, Lond B* 276: 401–405.
- Squire LR, Zola SM (1996) Structure and function of declarative and nondeclarative memory systems. *PNAS* 93: 13515–13522.

21. Paré D, Collins DR, Pelletier JG (2002) Amygdala oscillations and the consolidation of emotional memories. *Trends Cogn Sci* 6: 306–314.
22. Paré D, Royer S, Smith Y, Lang EJ (2003) Contextual inhibitory gating of impulse traffic in the intra-amygdaloid network. *Ann NY Acad Sci* 985: 78–91.
23. Isaacson RL (2008) The hippocampus. In: Squire LR, ed. *Encyclopedia of Neuroscience*. London: Elsevier Science.
24. Squire LR, Schacter DL, eds. *Neuropsychology of memory*. 3rd Edition ed. New York: The Guilford Press.
25. Buzsáki G (1998) Memory consolidation during sleep: a neurophysiological perspective. *J Sleep Res Suppl* 7: 17–23.
26. Sterpenich V, Albouy G, Boly M, Vandewalle G, Darsaud A, et al. (2007) Sleep-related hippocampo-cortical interplay during emotional memory recollection. *PLoS Biology* 5: 1–14.
27. Rauchs G, Desgranges B, Foret J, Eustache F (2005) The relationship between memory systems and sleep stages. *J Sleep Res* 14: 123–140.
28. Yoo SS, Gujar N, Hu P, Jolesz FA, Walker MP (2007) The human emotional brain without sleep - a prefrontal amygdala disconnect. *Current Biology* 17: R877–878.
29. McNamara P, Capellini I, Harris E, Nunn CL, Barton RA, et al. (2008) The phylogeny of sleep database: a new resource for sleep scientists. *The Open Sleep Journal* 1: 11–14.
30. Mukhametov LM (1984) Sleep in marine mammals. *Exp Brain Res Suppl* 8: 227–238.
31. Stephan H, Frahm H, Baron G (1981) New and revised data on volumes of brain structures in insectivores and primates. *Folia Primatol* 35: 1–29.
32. Harvey PA, Pagel M (1991) *The comparative method in evolutionary biology*. Oxford: Oxford University Press.
33. Nunn CL, Barton RA (2001) Comparative methods for studying primate adaptation and allometry. *Evol Anthropol* 10: 81–98.
34. Garland T, Bennett AF, Rezende EL (2005) Phylogenetic approaches in comparative physiology. *J Exp Biol* 208: 3015–3035.
35. Freckleton RP, Harvey PH, Pagel M (2002) Phylogenetic analysis and comparative data: a test and review of evidence. *Am Nat* 160: 712–726.
36. Pagel M (1997) Inferring evolutionary processes from phylogenies. *Zoologica Scripta* 26: 331–348.
37. Pagel M (1999) Inferring the historical patterns of biological evolution. *Nature* 401: 877–884.
38. Bininda-Emonds ORP, Cardillo M, Jones KE, MacPhee RDE, Beck RMD, et al. (2007) The delayed rise of present-day mammals. *Nature* 446: 507–512.
39. Bininda-Emonds ORP, Cardillo M, Jones KE, MacPhee RDE, Beck RMD, et al. (2008) Corrigendum. The delayed rise of present-day mammals. *Nature* 456: 274.
40. Barton RA (1998) Visual specialization and brain evolution in primates. *Proc R Soc, Lond B* 265: 1933–1937.
41. Benjamini Y, Hochberg Y (1995) Controlling for the false discovery rate: a practical and powerful approach to multiple testing. *J Roy Stat Soc B* 57: 289–300.
42. Verhoeven KJF, Simonsen KL, McIntyre LM (2005) Implementing the false discovery rate control: increasing your power. *Oikos* 108: 643–647.
43. Sejnokski TJ, Destexhe A (2000) Why do we sleep? *Brain Res Suppl* 886: 208–223.
44. Swanson LW, Petrovich GD (1998) What is the amygdala? *Trends Neurosci* 21: 323–331.
45. Barton RA, Aggleton JP, Grenyer R (2003) Evolutionary coherence of the mammalian amygdala. *Proc R Soc, Lond B* 270: 539–543.
46. Achermann P, Borbély AA (2003) Mathematical models of sleep regulation. *Frontiers in Bioscience* 8: s683–693.
47. Borbély AA, Achermann P (2005) Sleep homeostasis and models of sleep regulation. In: Kryger MH, Roth T, Dement WC, eds. *Principles and practice of sleep medicine*. Philadelphia: Saunders. pp 405–417.