



# The Association between Hypertension and Depression and Anxiety Disorders: Results from a Nationally-Representative Sample of South African Adults

## Citation

Grimsrud, Anna, Dan J. Stein, Soraya Seedat, David Williams, Landon Myer, and J. Jaime Miranda. 2009. The association between hypertension and depression and anxiety disorders: Results from a nationally-representative sample of South African adults. PLoS ONE 4(5): e5552.

## Published Version

doi:10.1371/journal.pone.0005552

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4453999>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

# The Association between Hypertension and Depression and Anxiety Disorders: Results from a Nationally-Representative Sample of South African Adults

Anna Grimsrud<sup>1</sup>, Dan J. Stein<sup>2</sup>, Soraya Seedat<sup>3</sup>, David Williams<sup>4</sup>, Landon Myer<sup>1\*</sup>

**1** School of Public Health & Family Medicine, University of Cape Town, Cape Town, South Africa, **2** Department of Psychiatry & Mental Health, University of Cape Town, Cape Town, South Africa, **3** Department of Psychiatry, University of Stellenbosch, Stellenbosch, South Africa, **4** Department of Society, Human Development and Health, Harvard School of Public Health, Boston, Massachusetts, United States of America

## Abstract

**Objective:** Growing evidence suggests high levels of comorbidity between hypertension and mental illness but there are few data from low- and middle-income countries. We examined the association between hypertension and depression and anxiety in South Africa.

**Methods:** Data come from a nationally-representative survey of adults (n = 4351). The Composite International Diagnostic Interview was used to measure DSM-IV mental disorders during the previous 12-months. The relationships between self-reported hypertension and anxiety disorders, depressive disorders and comorbid anxiety-depression were assessed after adjustment for participant characteristics including experience of trauma and other chronic physical conditions.

**Results:** Overall 16.7% reported a previous medical diagnosis of hypertension, and 8.1% and 4.9% were found to have a 12-month anxiety or depressive disorder, respectively. In adjusted analyses, hypertension diagnosis was associated with 12-month anxiety disorders [Odds ratio (OR) = 1.55, 95% Confidence interval (CI) = 1.10–2.18] but not 12-month depressive disorders or 12-month comorbid anxiety-depression. Hypertension in the absence of other chronic physical conditions was not associated with any of the 12-month mental health outcomes (p-values all < 0.05), while being diagnosed with both hypertension and another chronic physical condition were associated with 12-month anxiety disorders (OR = 2.25, 95% CI = 1.46–3.45), but not 12-month depressive disorders or comorbid anxiety-depression.

**Conclusions:** These are the first population-based estimates to demonstrate an association between hypertension and mental disorders in sub-Saharan Africa. Further investigation is needed into role of traumatic life events in the aetiology of hypertension as well as the temporality of the association between hypertension and mental disorders.

**Citation:** Grimsrud A, Stein DJ, Seedat S, Williams D, Myer L (2009) The Association between Hypertension and Depression and Anxiety Disorders: Results from a Nationally-Representative Sample of South African Adults. PLoS ONE 4(5): e5552. doi:10.1371/journal.pone.0005552

**Editor:** J. Jaime Miranda, London School of Hygiene and Tropical Medicine, Peru

**Received:** January 14, 2009; **Accepted:** April 20, 2009; **Published:** May 14, 2009

**Copyright:** © 2009 Grimsrud et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The South African Stress and Health study was carried out in conjunction with the World Health Organization World Mental Health (WMH) Survey Initiative. We thank the WMH staff for assistance with instrumentation, fieldwork, and data analysis. These activities were supported by the United States National Institute of Mental Health (R01MH070884), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (FIRCA R01-TW006481), the Pan American Health Organization, Eli Lilly and Company, Ortho-McNeil Pharmaceutical, Inc., GlaxoSmithKline, and Bristol-Myers Squibb. The South Africa Stress and Health study was funded by grant R01-MH059575 from the National Institute of Mental Health and the National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. Dan Stein and Soraya Seedat are also supported by the Medical Research Council (MRC) of South Africa. A complete list of WMH publications can be found at <http://www.hcp.med.harvard.edu/wmh/>. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: [landon.myer@uct.ac.za](mailto:landon.myer@uct.ac.za)

## Introduction

Chronic forms of morbidity, including mental disorders and hypertension, play a central role in shaping the burden of disease in the developing world. In South Africa there is a high prevalence of mental disorders, with an estimated 16% of adults living with an anxiety disorder and 10% with major depression [1]. Hypertension is a leading risk factor for mortality and morbidity worldwide, accounting for approximately 6% of global deaths [2–4]. The prevalence of hypertension in South Africa is over 20% [5].

A number of studies from Europe and North America have described an increased prevalence of chronic physical conditions among those with mental disorders [6,7]. There have been mixed findings for an association between hypertension and anxiety disorders in developed countries, with conflicting results from studies using the same design, and using the same measurements. Some studies have shown a positive association between hypertension and anxiety in both crude and multivariate analyses [8–12]. Conversely, there are studies that show no crude or adjusted association between hypertension and anxiety [13–15]. Several studies have observed a positive crude association between

hypertension and anxiety disorders that does not persist after adjustment [16–18]. Evidence is inconclusive and does not always include adjustment for relevant confounding variables, particularly traumatic life experiences.

Less evidence is available on the relationship between hypertension and depressive disorders. A handful of studies have suggested that depression may be more common among individuals with hypertension [9,19–20] while a much larger body of evidence shows no association between hypertension and depression [12–15,21–22]. Additional research found no crude or adjusted association between hypertension and depression [12–15,18,22–24].

Most of the literature investigates the relationship between hypertension and either depression or anxiety alone and few studies have examined comorbid depression and anxiety [12]. The relationship between comorbid anxiety-depression and chronic physical conditions was examined in the data from 17 countries that completed World Mental Health surveys [25]. Those with non-comorbid depressive disorder, non-comorbid anxiety disorder, and comorbid depression-anxiety were all more likely to have hypertension compared to persons with neither a depressive nor an anxiety disorder.

Limited evidence is available from developing countries. Furthermore, analyses need to consider important confounders such as trauma and other chronic conditions. Trauma has not previously been adjusted for, despite its potential to be associated with both hypertension and mental health disorders. Hypertension is often a risk factor for other chronic conditions, and therefore an observed association between hypertension and a mental health disorder may not persist after adjustment for other chronic conditions as the true association may be between more severe chronic conditions and the mental health disorder.

Despite the prevalence of mental disorders and chronic diseases such as hypertension in South Africa and other developing countries, few studies examine the relationship between these conditions in this setting. This paper investigates the associations between self-reported hypertension diagnosis and Diagnostic and Statistical Manual of Mental Disorders 4<sup>th</sup> edition (*DSM-IV*) defined 12-month a) anxiety disorders b) depressive disorders and c) comorbid anxiety-depression. We hypothesized that hypertension and anxiety and depressive disorders will be associated even after adjusting for potential confounding variables including demographic and socioeconomic risk factors, substance use disorders, traumatic life events and other chronic physical conditions.

## Methods

This study was conducted according to the principles expressed in the Declaration of Helsinki. The study was approved by the Institutional Review Board of the University of Stellenbosch (Project Number “N06/07/134”). All patients provided written informed consent for the collection of samples and subsequent analysis.

The South African Stress and Health (SASH) study is a cross-sectional survey of mental health in the South African adult population. The study is part of the World Health Organization World Mental Health Survey Initiative [26–28]. Its rationale and design have been described in detail previously [26,29].

SASH sampling and data collection were conducted between 2002 and 2004. The survey population included all resident South Africans 18 years and older who lived in households and hostels during the field period of the study. A three-stage probability sample design included a primary stage of stratification by

enumerator area (EA, a unit of census administration), a second stage of sampling dwelling units within each EA and a final stage random sub-selection of a single adult respondent from each of the selected households.

The EA units provided the primary stage sampling units for the SASH sampling. The approximately 85,000 EAs were then assigned to one of 53 strata. These strata were defined based on the province, urban/rural status and the majority population group (African, Coloured, White or Indian). A random sample of 5 dwelling units was selected from each EA to be included in the study population and was subsequently contacted for interviews. Within each selected dwelling unit, a single adult respondent was selected to participate. Fieldworkers made up to three attempts to contact the selected participants. The survey had a high initial response rate of 87%.

Interviews were conducted in the homes of the selected participants by trained field interviewers. The interviewers were trained in psychiatric interviewing and conducted a structured diagnostic interview which was translated into seven of the national languages.

## Measures

**Hypertension and other chronic conditions.** Hypertension was assessed as self-reported lifetime diagnosis in the section on *Chronic conditions* in the questionnaire. Participants responded to the following question: “Did a doctor or other health professional ever tell you that you had any of the following illnesses: high blood pressure”. In addition to this binary measure, a four-level categorical variable combining hypertension and other chronic physical conditions was created. This variable was divided into none (no chronic physical condition), hypertension only, hypertension plus another chronic physical condition, and other chronic physical condition only. Chronic physical conditions other than hypertension included the following: arthritis or rheumatism, chronic back or neck problems, any other chronic pain, a stroke, a heart attack, heart disease, asthma, any other chronic lung disease like COPD (chronic obstructive pulmonary disease) or emphysema, diabetes or high blood sugar, an ulcer in your stomach or intestine, thyroid disease, a neurological problem, like multiple sclerosis, Parkinson’s, or seizures, epilepsy or seizures, and cancer.

**Mental Disorders.** The World Mental Health pencil and paper (PAPI) version of the WHO Composite International Diagnostic Interview Version 3.0 (CIDI-3.0) was used to assess mental disorders according to the definitions and criteria of the *DSM-IV* [8]. Lifetime and 12-month *DSM-IV* diagnosis of the following anxiety disorders were included in this analysis: (i) panic disorder without agoraphobia, (ii) generalized anxiety disorder (GAD), (iii) social phobia, (iv) agoraphobia without panic disorder, and (v) posttraumatic stress disorder (PTSD). Lifetime and 12-month *DSM-IV* diagnosis of major and minor depression was also assessed. Substance use disorders were treated as a potential confounder. The variable for substance use disorders included lifetime *DSM-IV* diagnosis of alcohol abuse (with and without depression), alcohol dependence with abuse, drug abuse (with and without depression) and drug dependence with abuse. Earlier versions of the CIDI have been found to have good inter-rater and test-retest reliability [30] and in other WMH methodological studies there has been strong validity of CIDI diagnoses when compared to diagnoses based on blinded clinical reappraisal interviews [31].

**Additional Variables.** Socioeconomic status (SES) was measured using estimated household income, years of participant education, and participant employment. In addition, an asset index based on 17 items of individual and household

wealth was used to capture variation of socioeconomic position in the developing-country setting [32]. These measures of asset ownership were used to construct an aggregate asset score, which was categorized into categories for 0–5, 6–12 and 13–17 assets.

In addition to reporting measures of SES (household income, education, employment status and asset index) individually, we standardized the raw income, education and asset index values, and summed them to create an aggregate measure of SES. This standardized measure was divided into quartiles for analysis corresponding to low, low-moderate, moderate-high and high SES [33].

Alcohol use was described based on the years when the participants drank the most and analysed a categorical variable of more than once a week, once a week to once a month, and less than once a month. Lifetime smokers were defined as those who had smoked more than 100 cigarettes in their lifetime, and current smokers were those who self-reported currently smoking. The measurement of traumatic life events was based on the Life Event schedule of the World Mental Health Survey and included experiences of crime, war, natural disasters, accidents and illness.

**Analysis.** Data were analysed using Stata Version 10.0 (Stata Corporation, College Station, USA). All analyses reported accounted for the complex survey design based on person-level weights that incorporated sample selection, non-response and post-stratification factors. All statistical tests were two-sided at  $\alpha = 0.05$ .

Descriptive statistics were generated with continuous variables reporting means and categorical data reporting proportions. The prevalence and proportions of the demographic and socioeconomic variables, as well as other variables (alcohol use, smoking, chronic physical conditions and traumatic life events), were examined among the total sample, and stratified by hypertension, anxiety disorders (lifetime and 12-month), depressive disorders (lifetime and 12-month), and comorbid anxiety-depression (lifetime and 12-month). The odds ratios (OR) and 95% confidence intervals (CI) of anxiety disorders, depressive disorders, and comorbid anxiety-depression by demographics, socioeconomic and other variables were generated with those with no mental disorder (no anxiety or depression) as the reference group.

The prevalence of hypertension and other chronic physical conditions in those with mental disorders (anxiety, depression and comorbid anxiety-depression), or the physical-mental comorbidity prevalences, were calculated for both lifetime and 12-month mental disorders. For each crude association, Pearson's chi-squared test used to compare proportions was calculated with the reference group being those without anxiety or depression. In addition, age adjusted odds ratios, with the reference group being those without anxiety or depression, with 95% confidence intervals and Pearson's chi-squared test for comparison of proportions were calculated for the physical-mental comorbidities.

We developed a series of multiple logistic regression models to assess the independent association between, 1) hypertension and anxiety disorders, 2) hypertension and depressive disorders, and 3) hypertension and comorbid anxiety-depression. Models for both hypertension as a binary variable (yes/no) and categorical variable including other chronic physical conditions (no chronic physical condition, hypertension only, hypertension and another chronic physical condition, and another chronic physical condition) were built. The models included adjustment for demographics, an aggregate measure of SES, lifetime smoking and alcohol use, substance use disorders and traumatic life events.

## Results

The demographic characteristics and socioeconomic characteristics of the study sample are presented in Table 1. The mean age was 37 years and half of the sample was married (50%). 76% of the sample was Black, 10% Coloured, 10% White and 3% Indian/Asian. Less than a third of the sample was employed (31%) and 74% had completed at least some high school. Of the total sample, 8% had a 12-month anxiety disorder, 5% a 12-month depressive disorder and 1% had 12-month comorbid anxiety-depression (Table 2).

Overall, 17% of participants reported being told that they had hypertension. More than forty percent of the sample (43%) had at least one chronic physical condition, and 79% of those with hypertension had at least one chronic condition (Table 1). Strong crude associations were found between hypertension diagnosis and age, sex, marriage, education, employment status, smoking (current and lifetime), chronic physical conditions, and traumatic life events ( $p$ -values all  $< 0.001$ ). Women were 2.6 times more likely to report a lifetime diagnosis of hypertension compared to men ( $p < 0.001$ ). People with another chronic condition were 6.7 times more likely to report a lifetime diagnosis of hypertension compared to those without a hypertension diagnosis ( $p < 0.001$ ). Among measures of socioeconomic status, those with a hypertension diagnosis were less educated and more likely to be unemployed. The aggregate measure of SES was strongly associated with hypertension; those with a hypertension diagnosis had a lower SES than those without a hypertension diagnosis (Table 1).

Table 2 presents some of the demographic and other characteristics of those with and without 12-month anxiety and depressive disorders. Females were more than twice as likely to have 12-month anxiety or depressive disorders (OR = 2.2, 95% CI = 1.6–2.9; OR = 2.3, 95% CI = 1.6–3.2), and more than seven times more likely to have comorbid anxiety-depression compared to men (OR = 7.4, 95% CI = 3.4–16.0). Those with mental disorders also reported an increased prevalence of chronic conditions other than hypertension with significantly more reporting being diagnosed as hypertensive. Those with a 12-month anxiety disorder were twice as likely to report a lifetime diagnosis of hypertension and twice as likely to report another chronic condition (OR = 2.0, 95% CI = 1.4–2.8; OR = 2.2, 95% CI = 1.7–2.8). Traumatic life events were also highly associated with all 12-month mental disorders. Those with a 12-month depressive disorder were 2.9 times more likely to report 1 or 2 traumatic life events, 4.2 times more likely to report 3 or 4 traumatic life events and 5.8 times more likely to report 5 or more traumatic life events compared to those without a 12-month depressive disorder (Table 2).

There was variability in the association between a recent diagnosis of hypertension and specific mental disorders during the previous 12 months (Table 3). Hypertension was more commonly reported in participants with agoraphobia, generalized anxiety disorder and panic disorder, but was not increased in individuals with PTSD or social phobia. Major depression was associated with an increased prevalence of hypertension. Overall, those with a 12-month mental disorder were 1.68 times more likely to report being diagnosed as hypertensive compared to those without a 12-month mental disorder.

Table 4 presents the association between hypertension and mental health disorders in models that do and do not adjust for other chronic conditions as well as adjusting for a range of other variables. Without adjusting for other chronic conditions, self-reported hypertension is associated with 12-month anxiety disorders (OR = 1.55) but not with 12-month depressive disorders ( $p$ -value) or 12-month comorbid anxiety-depression ( $p$ -value)

**Table 1.** Description of sample, stratified by hypertension diagnosis.

	Total	Hypertension diagnosis	No hypertension diagnosis	p-value
	%	%	%	
N (%)	4351	767 (16.7)	3584 (83.3)	
<b>Demographic Characteristics</b>				
Mean Age	37.0	50.3	34.3	-
Age categories: 18–29	39.1	9.0	45.1	<0.001
30–39	22.1	13.5	23.8	
40–49	18.1	24.2	16.9	
50+	20.7	53.3	14.2	
Sex: Male	46.3	28.0	50.0	<0.001
Female	53.7	72.0	50.0	
Race: Black	76.2	76.1	76.3	0.271
Coloured	10.4	11.7	10.2	
White	10.0	10.2	9.9	
Indian/Asian	3.4	2.0	3.6	
Currently married	50.1	54.0	47.1	<0.001
Location: Rural	38.4	41.6	37.7	0.074
Urban	61.6	58.5	62.3	
<b>Socioeconomic status</b>				
Education: None	6.8	12.7	5.6	<0.001
Grade 1–7	19.1	33.9	16.2	
Grade 8–11	35.4	32.9	35.9	
Completed high school	23.5	11.4	25.9	
Post-high school education	15.3	9.2	16.5	
Employed	31.0	26.0	32.0	0.020
Household income (mean)	59403.9			
R0	13.7	13.3	13.7	0.244
R1–5000	29.5	32.9	28.9	
R5001–25000	15.4	16.7	15.1	
R25000–100000	19.6	17.4	20.1	
R100001+	21.8	19.7	22.3	
Assets owned by household: 0–5	39.3	38.3	39.5	
6–12	37.4	41.0	36.7	
13–17	23.3	20.7	23.8	0.177
SES Quartile: 0	22.5	36.0	19.8	<0.001
1	24.5	22.1	24.9	
2	25.0	20.8	25.9	
3	28.0	21.0	29.4	
<b>Other</b>				
Alcohol: Lifetime – None	77.8	79.1	77.5	0.157
Rare	5.8	7.0	5.5	
Moderate	7.5	5.7	7.9	
Heavy	8.9	8.2	9.1	
Alcohol: Current – None	65.7	71.8	64.5	0.053
Rare	10.2	8.7	10.5	
Moderate	16.8	13.5	17.4	
Heavy	7.3	6.0	7.6	
Smoking				
Lifetime smoker	30.0	26.1	30.8	0.033
Current smoker	23.8	15.9	25.4	<0.001
Chronic physical conditions	42.6	78.5	35.4	<0.001

**Table 1.** Cont.

	Total	Hypertension diagnosis	No hypertension diagnosis	p-value
	%	%	%	
Vascular	11.2	40.2	5.4	<0.001
Other	39.4	69.9	33.3	<0.001
Traumatic life events				
None	26.6	17.3	28.5	<0.001
1–2	49.9	52.5	49.4	
3–4	10.8	11.9	10.6	
5 or more	12.7	18.3	11.5	

doi:10.1371/journal.pone.0005552.t001

(Table 4). When other chronic conditions are included in the final model, there is no association between hypertension (in the absence of another chronic condition) are mental health disorders. However, hypertension and other chronic condition are associated with 12-month anxiety disorders (OR = 2.25). Furthermore, other chronic conditions are associated with 12-month anxiety (OR = 1.74), depressive (OR = 1.56), and comorbid anxiety-depressive disorder (OR = 2.35) (Table 4).

Table 5 presents the multivariate models examining the association between 12-month mental health disorders and

participant demographic, socioeconomic and health characteristics. After adjustment, females were 3 times more likely to have a 12-month anxiety disorder or depressive disorder, and more than 12 times more likely to have 12-month comorbid anxiety-depressive disorder compared to men. Furthermore, independent of hypertension status, those with 5 or more traumatic life events were more likely to have a 12-month anxiety disorder, depressive disorder, and co-morbid anxiety-depressive disorder compared to those without a traumatic life event (p-values all <0.05).

**Table 2.** Description of sample by 12-month *DSM-IV* Anxiety & Major Depression.

	No depression or anxiety	Anxiety	OR (95% CI)	Major Depression	OR (95% CI)	Comorbid anxiety-depression	OR (95% CI)
	%	%		%		%	
Prevalence	87.8	8.1		4.9		1.4	
Mean Age	36.8	38.4		37.6		39.5	
Age categories:18–29	39.8	33.3	1.0	32.2	1.0	22.9	1.0
30–39	21.9	23.2	1.3 (1.0–1.7)	24.4	1.4 (0.9–2.1)	30.8	2.4 (1.0–6.0)
40–49	17.6	21.2	1.4 (1.0–2.1)	23.3	1.6 (1.1–2.5)	20.9	2.1 (1.0–4.2)
50+	20.7	22.3	1.3 (0.8–2.0)	20.2	1.2 (0.8–1.8)	25.4	2.1(1.0–4.5)
Sex: Male	48.4	30.1	1.0	29.2	1.0	11.3	1.0
Female	51.7	69.9	2.2 (1.6–2.9)	70.8	2.3 (1.6–3.2)	88.7	7.4 (3.4–16.0)
Race: Black	76.2	79.9	1.0	71.8	1.0	79.8	1.0
Coloured	10.1	11.5	1.1 (0.7–1.7)	13.4	1.4 (0.9–2.1)	9.7	0.9 (0.3–2.4)
White	10.4	6.5	0.6 (0.2–1.6)	7.6	0.8 (0.3–2.1)	7.7	0.7 (0.1–5.9)
Indian/Asian	3.2	2.1	0.6 (0.3–1.3)	7.2	2.4 (1.1–5.0)	2.8	0.8 (0.2–3.1)
Currently married	50.1	48.6	0.9 (0.7–1.2)	53.1	1.1 (0.9–1.5)	50.9	1.0 (0.6–1.7)
Location: Urban	61.2	66.1	1.2 (0.9–1.6)	65.2	1.2 (0.9–1.7)	76.2	2.0 (1.1–2.6)
Hypertension diagnosis	15.7	26.9	2.0 (1.4–2.8)	21.2	1.4 (1.0–2.0)	31.6	2.4 (1.2–5.1)
Lifetime smoker	29.0	34.6	1.3 (1.0–1.6)	38.9	1.6 (1.1–2.2)	27.0	0.9 (0.5–1.7)
Current smoker	23.2	24.6	1.1 (0.9–1.4)	31.1	1.5 (1.0–2.2)	19.0	0.8 (0.4–1.6)
Other chronic physical condition	40.3	59.8	2.2 (1.7–2.8)	59.8	2.2 (1.6–3.1)	67.8	3.1 (1.6–6.2)
Vascular	10.2	20.0	2.2 (1.5–3.3)	16.3	1.7 (1.1–2.6)	21.8	2.5 (1.0–5.8)
Other	37.1	57.8	2.3 (1.8–3.0)	56.2	2.2 (1.5–3.1)	64.6	3.1 (1.6–6.1)
Traumatic life events: None	28.3	16.8	1.0	10.0	1.0	10.2	1.0
1–2	50.2	45.0	1.5 (1.0–2.2)	51.5	2.9 (1.7–4.9)	40.6	2.3 (0.9–5.6)
3–4	10.4	13.0	2.1 (1.3–3.5)	15.7	4.2 (2.4–7.6)	19.9	5.3 (1.9–14.7)
5 or more	11.1	25.2	3.8 (2.6–5.7)	22.8	5.8 (3.6–9.4)	29.4	7.4 (2.7–20.3)

doi:10.1371/journal.pone.0005552.t002

**Table 3.** Prevalence and relative odds of 12-month mental disorders by past hypertension diagnosis.

	Total % (n)	Hypertension diagnosis %	No hypertension diagnosis	p-value	OR (95% CI)
Any anxiety disorder	8.23 (375)	26.92	15.8	<0.001	1.97 (1.41–2.75)
Agoraphobia	5.13 (232)	22.7	16.4	0.042	1.50 (1.01–2.22)
Generalized anxiety disorder	1.91 (90)	41.0	16.2	<0.001	3.59 (2.13–6.05)
Panic disorder	0.79 (37)	39.0	16.5	0.002	3.23 (1.51–6.93)
PTSD	0.58 (27)	25.6	16.6	0.232	1.72 (0.69–4.31)
Social phobia	1.88 (87)	19.4	16.6	0.475	1.21 (0.71–2.05)
Any depressive disorder	5.4 (249)	21.2	16.4	0.060	1.37 (0.99–1.90)
Major depressive disorder	4.8 (223)	23.0	16.4	0.020	1.52 (1.07–2.17)
Minor depressive disorder	0.6 (26)	6.7	16.8	0.076	0.36 (0.11–1.18)
Comorbid anxiety-depression	1.4 (67)	31.6	16.5	0.019	2.34 (1.14–4.83)
Any anxiety or depressive disorder	12.3 (557)	23.9	15.8	<0.001	1.68 (1.31–2.16)

doi:10.1371/journal.pone.0005552.t003

## Discussion

This study suggests that self-reported diagnosis of hypertension is more common among South Africans with 12-month anxiety disorders, depressive disorders and comorbid anxiety-depression compared to those without a mental disorder. However, hypertension diagnosis without another chronic physical condition (hypertension only) was not associated with any of the mental health outcomes (anxiety, depression, and comorbid anxiety-depression) in multivariate models. Hypertension diagnosis and another chronic condition were consistently associated with anxiety disorders, depressive disorders and comorbid anxiety-depression after adjustment for various confounding variables. Therefore, adjusting for other chronic conditions explains the observed crude associations between mental health disorders and hypertension.

This analysis makes significant advances on previous studies of this association. This is the first population-based study from sub-Saharan Africa to document an association between anxiety and depressive disorders and hypertension. Our analysis included adjustment for traumatic life events which has previously never been done; given the strong associations between life events and

both hypertension and common mental disorders, accounting for this covariation is an important consideration. In addition, few other studies of the association between hypertension and mental disorders have included other chronic conditions as a potential confounder.

Our evidence for an association between hypertension and anxiety disorders after adjusting for demographics, SES, lifetime smoking and alcohol behaviour, substance use disorders and traumatic life events is similar to the majority of studies in the literature [8,12]. The observed odds ratio of 1.6 may be smaller than those found in previous studies as we adjusted for a larger number of potential cofounders. Our principle confounding factor appeared to be traumatic life events, a variable that has previously never been considered in this association. The findings add to the previous evidence of an association between hypertension and anxiety disorders as well as providing evidence that the association exists in a developing country context.

Major depressive disorder demonstrated a positive crude association with hypertension diagnosis that did not persist in multivariate analysis when adjustment was made for demographics and traumatic life events. These results are similar to the majority of studies in the literature which suggests no association, or a crude

**Table 4.** Multivariate models examining the association between 12-month mental disorders and hypertension\*.

Model	Hypertension variable(s)	12-month anxiety disorders	12-month depressive disorders	12-month comorbid anxiety-depression
		OR (95% CI)	OR (95% CI)	OR (95% CI)
Model A – Hypertension as a binary variable	Hypertension diagnosis	1.55 (1.10–2.18)	1.01 (0.68–1.50)	1.38 (0.60–3.18)
Model B – Hypertension as a categorical variable including other chronic physical conditions	Hypertension diagnosis only	1.58 (0.74–3.38)	0.81 (0.35–1.87)	2.74 (0.81–9.22)
	Hypertension and another chronic physical condition	2.25 (1.46–3.45)	1.44 (0.91–2.29)	2.14 (0.82–5.63)
	Another chronic physical condition	1.74 (1.28–2.37)	1.56 (1.07–2.26)	2.35 (1.10–5.01)

\*Models A and B are both adjusted for demographic variables (age, sex, race marriage, location), SES, lifetime smoking and alcohol use, substance use disorders and traumatic life events.

doi:10.1371/journal.pone.0005552.t004

**Table 5.** Summary of multivariate models examining the association between hypertension and 12-month mental health outcomes\*.

	Anxiety		Depression		Comorbid anxiety-depression	
	OR	95% CI	OR	95% CI	OR	95% CI
Hypertension variable						
None	1.0		1.0		1.0	
Hypertension diagnosis only	1.58	0.74–3.38	0.81	0.35–1.87	2.74	0.81–9.22
Hypertension diagnosis and chronic	2.25	1.46–3.45	1.44	0.91–2.29	2.14	0.82–5.63
Chronic only	1.74	1.28–2.37	1.56	1.07–2.26	2.35	1.10–5.01
Age: 18–29						
30–39	1.09	0.80–1.47	1.17	0.74–1.85	1.95	0.82–4.62
40–49	1.03	0.68–1.56	1.27	0.78–2.07	1.30	0.68–2.47
50+	0.94	0.56–1.56	0.97	0.62–1.52	1.76	0.83–3.70
Sex: Male						
Female	3.24	2.25–4.67	3.47	2.33–5.15	12.50	4.59–34.04
Race: Black						
Coloured	0.74	0.44–1.24	1.00	0.65–1.54	0.49	0.18–1.32
White	0.43	0.17–1.09	0.57	0.27–1.24	0.41	0.06–2.68
Indian/Asian	0.48	0.21–1.07	1.96	0.95–4.05	0.52	0.13–2.15
Currently married	0.87	0.66–1.13	0.98	0.72–1.33	0.86	0.51–1.44
Location: Rural						
Urban	1.36	1.00–1.85	1.09	0.75–1.58	2.08	1.18–3.65
SES Quartile – 0						
1	0.95	0.64–1.43	1.19	0.77–1.81	1.35	0.71–2.59
2	0.85	0.56–1.28	0.91	0.58–1.41	1.24	0.54–2.85
3	1.01	0.63–1.63	1.04	0.61–1.78	1.28	0.43–3.78
Alcohol: Lifetime – None						
Rare	1.29	0.65–2.54	1.96	1.00–3.84	2.75	0.53–14.40
Moderate	1.25	0.67–2.32	1.91	1.11–3.29	2.91	0.89–9.55
Heavy	1.47	0.81–2.67	1.14	0.61–2.11	3.23	1.28–8.17
Lifetime smoker	1.57	1.20–2.06	2.03	1.34–3.08	1.28	0.46–3.52
Substance use disorder	2.32	1.32–4.09	1.44	0.83–2.52	1.73	0.67–4.47
Traumatic life events – None						
1–2	1.15	0.77–1.70	2.44	1.44–4.14	1.52	0.61–3.75
3–4	1.44	0.88–2.37	3.36	1.91–5.91	3.18	1.16–8.76
5 or more	2.62	1.70–4.03	4.27	2.49–7.32	4.49	1.59–12.65

\*All models are adjusted for demographic variables (age, sex, race marriage, location), SES, lifetime smoking and alcohol use, substance use disorders and traumatic life events.

doi:10.1371/journal.pone.0005552.t005

association that does not persist in multivariate analysis between hypertension and depressive disorders.

In our study, the association with hypertension and another chronic physical condition was attenuated after adjustment for demographics and traumatic life events. However, failure to show an adjusted association between hypertension and comorbid anxiety-depression may be the result of a lack of statistical power as only 67 people had comorbid anxiety-depression.

The study sample is nationally representative which gives it strong generalisability to the country and meaningful insight into the burden of hypertension and mental health in the country. Furthermore, with the exception of one study from Zimbabwe [35], this is the first large scale study of the association between hypertension and mental health in an African and developing country context.

There are several notable limitations to these results. The cross-sectional design limits the ability to ascertain temporality between hypertension diagnosis and mental health outcomes. The use of self-reported lifetime diagnosis of hypertension as the hypertension measure is a limitation of the study. Self-reporting may introduce measurement bias, where those who have a mental health disorder are more likely to recall a diagnosis of hypertension thus inflating the true association between hypertension and mental health disorders. Furthermore, self-reported hypertension is generally underreported [36]. Previous research suggests that measured blood pressure is modestly associated with self-reported hypertension status. However, reassurance is obtained from previous data which indicate that self-reported hypertension is moderately associated with objective data on hypertensive status [37–38].



The prevalence of self-reported hypertension from this data was 17% which is less than the 21% reported by the 1998 South African Demographic and Health Survey which measured hypertension at the time of administering the questionnaire [4].

Mental health was assessed using the CIDI 3.0, an instrument designed in the context of developing countries [39]. A significant body of literature is available on the cross-cultural challenges in measuring mental health. The literature highlights two primary issues; one on the linguistic issues regarding comprehension and understanding of mental health constructs and one on the socio-cultural nature of mental health and how the symptoms may manifest differently in different societies [40–41]. As an example, it has been recognized that depression for black Africans may present with different linguistic phrases and symptoms may both occur in different places and in different ways to those of Western culture [40]. South Africa is a heterogeneous society, with significant diversity in cultures. The CIDI 3.0 has not been validated for South Africa, with research on the concordance of the CIDI compared to standardized clinical interviews for *DSM-IV* diagnosis (SCID) done only in France, Italy, Spain and the United States. The results show the agreement to be good, with CIDI 12-month prevalences generally conservative relative to SCID [31]. There is potential for this measurement to introduce bias to the results, both through random and systematic misclassification. Random misclassification may have happened if the misclassification

of mental disorders occurred to the same degree among those with and without hypertension diagnosis.

A strength of the dataset is that information is available on a number of potential confounders that have not been accounted for in previous studies of this association. We also had information on a large number of chronic physical conditions, both conditions in the past 12-months and lifetime diagnoses from health care professionals. Adjustment for other chronic physical conditions proved highly important, demonstrating that hypertension on its own has no association with anxiety, depression or comorbid anxiety-depression in crude or adjusted analyses. We were able to adjust for demographics and socioeconomics, as well as lifestyle risk factors (smoking and alcohol use) and substance use disorders.

There is a high prevalence of hypertension and mental health disorders in South Africa. The comorbidity observed in developing countries between chronic physical conditions and mental disorders is also present in the South African context. Multivariate analysis adjusting for other chronic physical conditions and traumatic life events show a smaller than previously presented association between hypertension and mental health disorders.

### Author Contributions

Conceived and designed the experiments: DJS DRW. Performed the experiments: DJS SS DRW. Analyzed the data: ATG LM. Wrote the paper: ATG LM.

### References

- Stein DJ, Seedat S, Herman A, Moomal H, Heeringa SG, et al. (2008) Lifetime prevalence of psychiatric disorders in South Africa. *The British Journal of Psychiatry* 192: 112–117.
- Murray CJL, Lopez AD (1997) Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *The Lancet* 349: 1498–504.
- Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJL (1997) Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *The Lancet* 367: 1747–57.
- Department of Health South Africa (1998) South Africa Demographic and Health Survey - 1998. South Africa.
- Steyn KA, Gaziano TABE, Bradshaw DC, Laubscher RD, Fourie JA, et al. (2001) Hypertension in South African adults: results from the Demographic and Health Survey, 1998. *Journal of Hypertension* 19: 1717–25.
- Jones DR, Macias C, Barreira PJ, Fisher WH, Hargreaves WA, et al. (2004) Prevalence, Severity, and Co-occurrence of Chronic Physical Health Problems of Persons With Serious Mental Illness. *Psychiatric Services* 55: 1250–7.
- Evans DL, Charney DS, Lewis L, Golden RN, Gorman JM, et al. (2005) Mood Disorders in the Medically Ill: Scientific Review and Recommendations. *Biological Psychiatry* 58: 175–89.
- Harter MC, Conway KP, Merikangas KR (2003) Associations between anxiety disorders and physical illness. *European Archives of Psychiatry and Clinical Neuroscience* 253: 313–20.
- Coelho R, Hughes AM, Fernandes da Fonseca A, Bond MR (1989) Essential hypertension: The relationship of psychological factors to the severity of hypertension. *Journal of Psychosomatic Research* 33: 187–96.
- Tie-Min WEI, Lexin W (2006) Anxiety symptoms in patients with hypertension: a community-based study. *International Journal of Psychiatry in Medicine* 36: 315.
- Dorn T, Joris Yzermans C, Guigt H, van der Zee J (2007) Disaster-related stress as a prospective risk factor for hypertension in parents of adolescent fire victims. *American Journal of Epidemiology* 165: 410–7.
- Paterniti SMD, Alperovitch AMD, Ducimetiere PP, Dealberto M-JMDP, Lepine J-PMD, et al. (1999) Anxiety But Not Depression Is Associated With Elevated Blood Pressure in a Community Group of French Elderly. *Psychosomatic Medicine* 61: 77–83.
- Yan LL, Liu K, Matthews KA, Daviglus ML, Ferguson TF, Kiefe CI (2003) Psychosocial Factors and Risk of Hypertension: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. *JAMA* 290: 2138–48.
- Shim EH, Poston WSC, Kimball KT, St. Jeor ST, Foreyt JP (2001) Blood pressure and symptoms of depression and anxiety: a prospective study. *American Journal of Hypertension* 14: 660–4.
- Jones-Webb R, Jacobs DR, Flack JM, Liu K (1996) Relationships between Depressive Symptoms, Anxiety, Alcohol Consumption, and Blood Pressure: Results from the CARDIA Study. *Alcoholism: Clinical and Experimental Research* 20: 420–7.
- Kadam UT, Jordan K, Croft PR (2005) Clinical comorbidity was specific to disease pathology, psychological distress, and somatic symptom amplification. *Journal of Clinical Epidemiology* 58: 909–17.
- Markovitz JH, Matthews KA, Kannel WB, Cobb JL, D'Agostino RB (1993) Psychological predictors of hypertension in the Framingham Study. Is there tension in hypertension? *JAMA* 270: 2439–43.
- Scott K, Oakley Browne M, McGee M, Wells J, et al. (2006) Mental-physical comorbidity in Te Rau Hinengaro: the New Zealand Mental Health Survey. *Australian and New Zealand Journal of Psychiatry* 40: 7.
- Davidson K, Jonas BS, Dixon KE, Markovitz JH (2000) Do Depression Symptoms Predict Early Hypertension Incidence in Young Adults in the CARDIA Study? *Arch Intern Med* 160: 1495–500.
- Jonas BS, Franks P, Ingram DD (1997) Are symptoms of anxiety and depression risk factors for hypertension? Longitudinal evidence from the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Archives of Family Medicine* 6: 43–9.
- Levenstein S, Smith MW, Kaplan GA (2001) Psychosocial Predictors of Hypertension in Men and Women. *Arch Intern Med* 161: 1341–6.
- Wiehe M, Fuchs S, Moreira L, Moraes R, Pereira G, et al. (2006) Absence of association between depression and hypertension: results of a prospectively designed population-based study. *Journal of Human Hypertension* 20: 434–9.
- Scherrer JFMA, Xian HP, Buchholz KKP, Eisen SAMDM, Lyons MJP, et al. (2003) A Twin Study of Depression Symptoms, Hypertension, and Heart Disease in Middle-Aged Men. *Psychosomatic Medicine* 65: 548–57.
- Lowe B, Grafe K, Ufer C, Kroenke K, Grunig E, et al. (2004) Anxiety and Depression in Patients With Pulmonary Hypertension. *Psychological Medicine* 34: 831–6.
- Scott KM, Bruffaerts R, Tsang A, Ormel J, Alonso J, et al. (2007) Depression-anxiety relationships with chronic physical conditions: Results from the World Mental Health surveys. *Journal of Affective Disorders*, in press.
- Consortium TWWMHS (2004) Prevalence, Severity, and Unmet Need for Treatment of Mental Disorders in the World Health Organization World Mental Health Surveys. *JAMA* 291: 2581–90.
- Ustun TB (1999) The global burden of mental disorders. *American Journal of Public Health* 89: 1315–8.
- Medicine HSo (2007) The World Mental Health Survey Initiative. vol. 2005.
- Williams DR, Herman A, Kessler RD, Sonnega J, Seedat S, et al. (2004) The South Africa Stress and Health Study: Rationale and Design. *Metabolic Brain Disease* 19: 135–47.
- Andrew G, Peters L (1998) The psychometric properties of the Composite International Diagnostic Interview. *Social Psychiatry and Psychiatric Epidemiology* 33: 8.
- Haroo J, Arbabzadeh-Bouchez S, Brugha T, de Girolamo G, Guyer M, et al. (2006) Concordance of the Composite International Diagnostic Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health Surveys. *International Journal of Methods in Psychiatric Research* 13.

32. Myer L, Ehrlich RI, Susser ES (2004) Social Epidemiology in South Africa. *Epidemiology Reviews* 26: 112–23.
33. Myer L, Stein DJ, Grimsrud A, Seedat S, Williams DR (2008) Social determinants of psychological distress in a nationally-representative sample of South African adults. *Social Science & Medicine* 66: 1828–40.
34. Williams D, Herman A, Stein D, Heeringa S, Jackson P, et al. (2007) Twelve-month mental disorders in South Africa: prevalence, service use and demographic correlates in the population-based South African Stress and Health Survey. *Psychological Medicine* 33: 211–20.
35. Somova LI, Connolly C, Diara K (1995) Psychosocial predictors of hypertension in Black and White Africans. *Journal of Hypertension* 13: 193–9.
36. Goldman N, Lin IF, Weinstein M, Lin Y-H (2003) Evaluating the quality of self-reports of hypertension and diabetes. *Journal of Clinical Epidemiology* 56: 148–54.
37. Alonso A, Beunza JJ, Delgado-Rodriguez M, Martinez-Gonzalez MA (2005) Validation of self reported diagnosis of hypertension in a cohort of university graduates in Spain. *BMC Public Health* 5.
38. Tormo MJ, Navarro C, Chirlaque MD, Barber X, Spain tEGo (2000) Validation of self diagnosis of high blood pressure in a sample of the Spanish EPIC cohort: overall agreement and predictive values. *Journal of Epidemiology and Community Health* 54: 221–6.
39. Wittchen HU (1994) Reliability and validity studies of the WHO-composite international diagnostic interview (CIDI): a critical review. *J Psychiatr Res* 28: 57–84.
40. Smit J, van den Berg C, Bekker L-G, Seedat S, Stein D (2006) Translation and cross-cultural adaptation of a mental health battery in an African setting. *Afr Health Sci* 6(4): 215–222.
41. Ellis C (2003) Cross-cultural aspects of depression in general practice. *South African Medical Journal* 93: 342–45.