Shenandoah Cloud and Photochemistry Experiment (SCAPE): Overview

Citation

Keene, William C., Daniel J. Jacob, Robert W. Talbot, and J. William Munger. 1995. "Shenandoah Cloud and Photochemistry Experiment (SCAPE): Overview." *Journal of Geophysical Research* 100 (D5): 9313. doi:10.1029/95jd00848.

Published Version

doi:10.1029/95JD00848

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:14121767>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Shenandoah Cloud and Photochemistry Experiment (SCAPE): Overview

William C. Keene,¹ Daniel J. Jacob,² Robert W. Talbot,³ and J. William Munger²

Introduction

The Shenandoah Cloud and Photochemistry Experiment (SCAPE) investigated the chemistry of the atmosphere at a high-elevation forested site in central Virginia with particular focus on the factors controlling concentrations of ozone, carbonyl compounds, and carboxylic acids. Carbonyl compounds and carboxylic acids are ubiquitous chemical constituents of the global troposphere, but many aspects of their biogeochemical cycling are poorly understood. Carbonyls are generated from the oxidation of hydrocarbons; formaldehyde (HCHO) produced via this pathway is an important precursor for OH. Thus the carbonyls are intermediates in and tracers of tropospheric photochemistry [e.g., *Carlier et al.*, 1988]. Carboxylic acids are major natural sources of atmospheric acidity thereby partially controlling pH-dependent chemical transformations in clouds [e.g., *Keene and Gallo-way*, 1988]. The aqueous-phase oxidation of HCHO in clouds has been hypothesized as a significant source of formic acid (HCOOH) and the aqueous-phase oxidation of HCOO⁻ a potentially important in-cloud sink for OH [*Chameides and Davis*, 1983; *Jacob*, 1986]. Under many conditions, clouds should be a net sink for tropospheric HCOOH [*Jacob and Wofsy*, 1988]. Specific research objectives were (1) to assess the relative importance of NO_x and hydrocarbons in limiting regional ozone production, (2) to investigate sources and sinks for carbonyl compounds and carboxylic acids, and (3) to examine the droplet-size dependence of solute concentrations and the thermodynamics of phase partitioning for carbonyl compounds and carboxylic acids in clouds.

Experimental Design

SCAPE was a collaborative effort involving investigators from several institutions. Measurement and modeling activities are summarized in Table 1; readers are referred to the accompanying papers for additional details.

The field experiment was conducted between September 1 and October 1, 1990. Most of the chemical species and physical parameters were measured from a 15-m tower on a mountain ridge at Pinnacles (1037-m elevation) in the north

central section of the Shenandoah National Park, Virginia (38°37'N, 78°21'W). The top of the tower was 3 m above the surrounding forest canopy. O₃, CO, NO, NO_y, and peroxides were continuously measured through a 10-m sampling mast on a portable laboratory van in a clearing approximately 20 m from the base of the tower.

When clouds enveloped the site, coarse (>18-μm diameter) and fine (19- to 5.5-μm diameter) droplets and liquid-water content were sequentially sampled. Cloud-water pH and aqueous-phase and vapor-phase concentrations of carbonyls and carboxylic acids are measured in parallel over 1.0- to 1.5-hour intervals. Volatile carbonyls and carboxylic acids were also sampled over diel cycles during noncloud periods.

Overview of Results

On the basis of measured O₃, CO, NO, NO_y, H₂O₂, and HCHO, *Jacob et al.* [this issue] found evidence for a seasonal transition from NO_x-limited to hydrocarbon-limited O₃ production during September. They argue that such a seasonal transition in photochemical regime should be a general phenomenon over the eastern United States.

Munger et al. [this issue] report that the oxidation of isoprene was a major source for ambient HCHO and CHOCHO. They observed maximum concentrations (HCHO > 2 parts per billion by volume; CHOCHO > 100 parts per trillion by volume (pptv) during stagnation episodes in association with warm temperatures and elevated O₃ and H₂O₂. Concentrations of CH₃C(O)CHO were generally below the detection limit of 50 pptv.

Talbot et al. [this issue] infer that direct emissions from vegetation and soils were the dominant sources of carboxylic acids; model calculations and correlation analyses using various tracer compounds suggest that the decomposition of hydrocarbons and combustion of biomass or fossil fuels were relatively less important.

Both *Munger et al.* [this issue] and *Keene et al.* [this issue] found that aqueous-phase concentrations of carbonyls and carboxylic acids in clouds did not differ significantly between fine- and coarse-droplet-size fractions. Aqueous HCHO was supersaturated relative to the gas phase during a daytime cloud event; *Munger et al.* [this issue] hypothesize the hydrolysis of HOCH₂OOH as a possible explanation.

The phase partitioning for CH₃COOH measured by *Keene et al.* [this issue] in clouds was within the uncertainty of thermodynamic data. However, temporal variability in gaseous HCOOH was damped relative to equilibria with the aqueous phase suggesting that liquid water on the forest canopy may have buffered HCOOH in near-surface clouds. No evidence was found for significant HCOOH production in clouds; cloud-water deposition to the forest canopy appeared to be an important sink for HCOOH.

¹Department of Environmental Sciences, University of Virginia, Charlottesville.

²Division of Applied Sciences, Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts.

³Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham.

Table 1. Sampling, Measurement, and Modeling Components of SCAPE

Species	Media	Institution	Investigator
Carbonyl compounds ^a	vapor, ^b cloud water, ^c precipitation ^d	Harvard University	J.W. Munger, B.C. Daube, D.J. Jacob
Carboxyl acids ^e	vapor ^b	University of New Hampshire	B.W. Mosher, R.W. Talbot
Carboxyl acids, major inorganic ions ^f	cloud water, ^c precipitation ^d	University of Virginia	W.C. Keene, J.R. Maben
O ₃ , CO, NO, NO _y , H ₂ O ₂ , organic peroxides, UV radiation ^{g,h}	vapor	University of Rhode Island	B.G. Heikes, M. Lee
Hydrocarbons ⁱ	vapor	University of California, Santa Barbara	D. Pierotti
Liquid-water content ^j	cloud	University of Virginia	W.C. Keene, J.R. Maben
Temperature, pressure, RH, wind speed and direction	---	University of Virginia	P. Thompson, J. Sigmon
Meteorology, trajectory analyses	---	NOAA ARL, University of Virginia	R.S. Artz, J.L. Moody
Photochemical modeling	---	Harvard University	D.J. Jacob, L.W. Horowitz

^aFormaldehyde (HCHO), glyoxal (CHOCHO), and methylglyoxal (CH₃C(O)CHO).

^bSampled using mist chambers; an inertial separator developed by B. C. Daube was positioned at the inlet to remove cloud droplets and coarse aerosol (> 1-μm diameter) from the sample air.

^cSampled using size-segregating collector developed by B. C. Daube and operated by W. C. Keene and J. R. Maben.

^dManually sampled by W. C. Keene and J. R. Maben during precipitation events; sampling intervals varied as a function of sample volume.

^eFormic (HCOOH), acetic (CH₃COOH), and pyruvic (CH₃(CO)COOH) acids.

^fHCOOH_t (HCOOH_{aq} + HCOO⁻), CH₃COOH_t (CH₃COOH_{aq} + CH₃COO⁻), H⁺, SO₄²⁻, NO₃⁻, Cl⁻, NH₄⁺, Ca²⁺, Mg²⁺, Na⁺, and K⁺.

^gMeasured continuously.

^hInstrumentation to measure O₃, CO, NO, and NO_y was provided by R. R. Dickerson, B. G. Doddridge, and K. Civerolo at the University of Maryland.

ⁱGrab samples were collected between September 4 and 14, 1990.

^jMeasured using two independent techniques.

Acknowledgments. Detailed acknowledgments appear in each of the SCAPE papers, but we thought it appropriate to briefly mention here the major contributors without whose support this effort would not have been possible. Financial support for the program was provided by the National Science Foundation, the NOAA Air Resources Laboratory, the Merck Foundation, and the Electric Power Research Institute. The Mountain Cloud Chemistry Project allowed us to use their field facilities and the Shenandoah National Park provided on-site housing and laboratory space. Several individuals not explicitly mentioned in this introduction contributed substantively to the effort and are so acknowledged in the accompanying papers. We thank Anne Thompson who served as editor and capably guided the SCAPE manuscripts through peer review. We also acknowledge the editorial assistance provided by Paul Ginsburg. In closing, we wish to thank all the participants for contributing to an intellectually stimulating and thoroughly enjoyable investigation.

References

- Carlier, P., H. Hannachi, and G. Mouvier, The chemistry of carbonyl compounds in the atmosphere—A review, *Atmos. Environ.*, **20**, 2079–2099, 1988.
- Chameides, W. L., and D. D. Davis, Aqueous phase source of formic acid in clouds, *Nature*, **304**, 427–429, 1983.
- Jacob, D. J., Chemistry of OH in remote clouds and its role in the production of formic acid and peroxymonosulfate, *J. Geophys. Res.*, **91**, 9807–9826, 1986.
- Jacob, D. J., and S. C. Wofsy, Photochemical production of carboxylic acids in a remote continental atmosphere, in *Acid Deposition at High Elevation Sites*, edited by M. H. Unsworth and D. Fowler, Kluwer Academic, Norwell, Mass., 1988.
- Jacob, D. J., L. W. Horowitz, B. G. Heikes, R. R. Dickerson, R. S. Artz, and W. C. Keene, Seasonal transition from NO_x- to hydrocarbon-limited ozone production over the eastern United States in September, *J. Geophys. Res.*, this issue.
- Keene, W. C., and J. N. Galloway, The biogeochemical cycling of formic and acetic acids through the troposphere: An overview of current understanding, *Tellus*, **40B**, 322–334, 1988.
- Keene, W. C., B. W. Mosher, D. J. Jacob, J. W. Munger, R. W. Talbot, R. S. Artz, J. R. Maben, B. R. Daube, and J. N. Galloway, Carboxylic acids in clouds at a high-elevation forested site in central Virginia, USA, *J. Geophys. Res.*, this issue.
- Munger, J. W., D. J. Jacob, B. C. Daube, L. W. Horowitz, W. C. Keene, and B. G. Heikes, Formaldehyde, glyoxal, and methylglyoxal in air and cloud water at a rural mountain site in central Virginia, *J. Geophys. Res.*, this issue.
- Talbot, R. W., B. W. Mosher, B. G. Heikes, D. J. Jacob, J. M. Munger, B. C. Daube, W. C. Keene, J. M. Maben, and R. S. Artz, Carboxylic acids in the rural continental atmosphere of the eastern United States, *J. Geophys. Res.*, this issue.
- D. J. Jacob and J. W. Munger, Division of Applied Sciences, Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA 02138. emails: djj or jwm@io.harvard.edu
- W. C. Keene, Department of Environmental Sciences, University of Virginia, Clark Hall, Charlottesville, VA 22903. email: wck@virginia.edu
- R. W. Talbot, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, NH 03824. email: rwt@christa.unh.edu

(Received March 10, 1995; revised March 10, 1995; accepted March 10, 1995.)