Hybrid zones and the speciation continuum in *Heliconius* butterflies

Citation

Mallet, James, and Kanchon K. Dasmahapatra. 2012. "Hybrid Zones and the Speciation Continuum in *Heliconius* Butterflies ." *Mol Ecol* 21 (23) (November 21): 5643–5645. doi:10.1111/mec.12058.

Published Version

doi:10.1111/mec.12058

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:25288054>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Hybrid zones and the speciation continuum
in *Heliconius* butterflies

James Mallet^{1,2}

and

Kanchon K. Dasmahapatra¹

1. Galton Laboratory, Department of Genetics, Evolution and
Environment, University College London,
Gower St., LONDON WC1E 6BT, UK

and

2. Department of Organismic and Evolutionary Biology,
16 Divinity Avenue, CAMBRIDGE, MA 02138, USA

Corresponding author: James Mallet (j.mallet@ucl.ac.uk)
tel: +44 - (0)20 7679 7412

2012. Molecular Ecology 21 (in press).

**Commentary on: Arias CF, Rosales C, Salazar C et al. (2012) Sharp
genetic discontinuity across a unimodal *Heliconius* hybrid zone.
Molecular Ecology, 21 (in press).**

25 Tropical butterflies in the genus *Heliconius* have long been models in
26 the study of the stages of speciation. *Heliconius* are unpalatable to
27 predators, and many species are notable for multiple geographic
28 populations with striking warning colour pattern differences associated
29 with Müllerian mimicry. There is a speciation continuum evident in
30 *Heliconius* hybrid zones, across which mimicry patterns are often
31 different, but where hybrids are common and little else differs, through
32 to 'bimodal' hybrid zones with strongly marked molecular differences
33 with few hybrids, through to 'good' sympatric species. Now Arias et al.
34 (2012) have found an intermediate case in Colombian *Heliconius cydno*
35 showing evidence for assortative mating and molecular differences, but
36 where hybrids are abundant.

38 Hybrid zones in neotropical *Heliconius* butterflies have long fascinated
39 evolutionary biologists, and informed about stages of speciation. Henry
40 Walter Bates was the first to study *Heliconius* on a scientific basis, and he
41 was also the first to explain mimicry among unrelated species via natural
42 selection. Together with Fritz Müller, Bates was able to explain the
43 selective advantage of this mimicry between edible and unpalatable
44 species (Batesian mimicry), and among unpalatable species (Müllerian
45 mimicry): birds, having learnt to avoid the warning colours of one inedible
46 species, thereafter shun other species with convergent or 'mimetic' colour
47 patterns (Bates 1862; Müller 1879).

48
49 It is less well known that Bates, who was particularly interested in the
50 origin of species after reading Darwin's works, was also the first to

document a hybrid zone and assess its implications for understanding the continuum of speciation (Bates 1863). He carried out this work on a hybrid zone between colour pattern races of *Heliconius melpomene* in the Brazilian Amazon. Recent work with heliconiines has shown how mimicry and speciation are closely intertwined across the adaptive radiation in this group (McMillan *et al.* 1997; Linares 1997; Jiggins *et al.* 2001; Arias *et al.* 2008). Now a new paper closes the gap between species and geographic races still further (Arias *et al.* 2012).

Speciation appears to be more of a problem for humans to understand than it is for the organisms that do it. Learned books tell us that speciation was a conundrum not solved by Darwin, and one that is not clearly understood even today. Nonetheless, many intermediate stages of ongoing speciation are evident both below the species level (ecotypes, ecological races, and host races) and above the species level (involving hybridization and gene flow between species, and hybrid speciation). Speciation clearly exists as a continuum across the species boundary, and assessing these intermediate stages is key to understanding the process.

Studies involving the genus *Heliconius* have been major recent contributors to our knowledge of the speciation continuum. An early study used hybrid zone dynamics to measure the strong natural selection on mimicry, and showed how this selection could stabilize narrow zones of hybridization between Peruvian geographic races of *H. erato* differing in mimetic colour pattern. Similar results are also evident from the coincident hybrid zone between races of its co-mimic *H. melpomene* (Mallet *et al.* 1990). Recent studies have shown that virtually no genomic differences exist across these Peruvian hybrid zones, except at fixed differences in a handful of genomic regions controlling the colour pattern and mimicry shift (Countermand *et al.* 2010; Nadeau *et al.* 2012), suggesting that in many cases mimetic shifts may be a first stage of divergence towards speciation. However, mating of the geographic races within each species is random in the centre of each hybrid zone, so the contribution of the evolution of geographic races like those in Peru to speciation was still unclear. *Heliconius cydno* is a species with widespread mimicry polymorphisms, such as in *H. cydno weymeri* studied here in the southern Cauca Valley (Fig. 1). In an Ecuadorean population of this species, colour pattern contributes to assortative mating among mimetic morphs in one direction of cross (Chamberlain *et al.* 2009).

90 How such racial divergence might contribute to speciation was further
91 studied along a wet-dry ecotone in Ecuador. A narrow hybrid zone
92 between *H. (erato) himera* and *H. erato cyrbia*, which were previously
93 regarded as conspecific, was rather different in character than the Peruvian
94 hybrid zones in *H. erato* and *H. melpomene*. In particular, there were strong
95 differences in mtDNA and nuclear markers across the hybrid zone. Within
96 the hybrid zone multilocus genotypes, including colour pattern loci,
97 formed a bimodal distribution. Phenotypic hybrids between the taxa,
98 although present, formed only about 10% of the population in the centre of
99 the hybrid zone. This was largely due to assortative mating and ecological
100 differences between the taxa, because no hybrid inviability or sterility was
101 found (Jiggins *et al.* 1997; McMillan, Jiggins, and Mallet 1997). Clearly,
102 considerable progress towards speciation has been achieved because the
103 bimodality of the genotypic distribution shows that two clusters of
104 genotypes coexist without collapsing, albeit in a narrow zone of contact.
105 As a result, *H. himera* and *H. erato* were thereafter classified as separate
106 species.

107
108 In Colombia there is a somewhat comparable hybrid zone along a wet-dry
109 ecotone between another two taxa also formerly considered geographic

110 races of *H. erato*: *H. (e.) chestertonii* and *H. e. venus*. Although considerable
111 hybridization is reported in the hybrid zone, hybrids again remain rare
112 (25%) compared to "pure" colour pattern phenotypes, and both mtDNA
113 and nuclear markers also differ between the taxa (Arias *et al.* 2008).
114
115 It is extremely difficult for foreigners to obtain permits to collect wild
116 species for DNA-based work in Colombia, and there have been, at least
117 until recently, public order difficulties across much of that country.
118 Mauricio Linares, the senior Colombian author of the current paper, has
119 endured a number of life-threatening situations to obtain valuable data on
120 Colombian *Heliconius* hybrid zones. For a long time his research team has
121 been studying a major hybrid zone between two mimetic races, *H. cydno*
122 *cydnides* and *H. c. weymeri* in the Cauca Valley, sandwiched between the
123 Central and Western Cordilleras of the Andes (Fig. 1). The current paper,
124 from an all-Latin American group, shows some extremely interesting
125 patterns from this hybrid zone. The two races show asymmetrical
126 assortative courtship, and there is also clear mtDNA as well as some
127 evidence for nuclear genetic differences across the zone. Yet hybrids in the
128 centre of the zone appear abundant, giving no evidence for phenotypic
129 bimodality: hence the authors' designation of the hybrid zone as

"unimodal" (Arias *et al.* 2012). Along the continuum of speciation, this *Heliconius cydno* hybrid zone therefore sits between bimodal hybrid zones like those of *H. himera* and *H. erato*, and classical clinal hybrid zones like those between Peruvian races of *H. erato* or *H. melpomene* (Table 1).

The authors also find that the mtDNA haplotypes of the southern race (*H. c. weymeri*) are closely related to those of the Pacific coastal race *H. cydno zelinde*, whereas the northern *H. c. cydnides* groups with races further east, from the Magdalena Valley. According to the authors, this argues for the current hybrid zone forming by secondary contact as a result of the spread of *H. c. weymeri* accompanied by colour pattern divergence across passes in the Western Andes into the southern Cauca Valley. Here, it met *H. c. cydnides* which itself spread from the east. As with most phylogeographic hypotheses, this is hard to prove, especially as it has been shown that some colour patterns in *Heliconius erato* seem to have spread behind moving contact zones while in parapatric contact with another race (Blum 2002), so potentially explaining the existence of multiple, disjunct but otherwise very similar mimicry races up and down the Andes (Hines *et al.* 2011). Nonetheless, the existence of strong mtDNA and some nuclear divergence across this *Heliconius cydno* contact zone argues for greater population

restructuring than just a moving colour pattern cline within a continuous geographic range; the authors rightly argue that this is good evidence that the hybrid zone was achieved via secondary contact.

What is not yet quite clear is whether speciation can take place during complete overlap in sympatry. The existence of regular albeit low-level hybridisation among many sympatric *Heliconius* species (Mallet *et al.* 2007), and the high degree of sympatry among sister species (Rosser *et al.* 2012) argues that sympatric speciation may be common (Neil Rosser *et al.* in prep.). Formation of the hybrid species *Heliconius heurippa* and *H. elevatus* were almost certainly also explained by sympatric transfer of colour pattern genes between species (Mavárez *et al.* 2006; Heliconius Genome Consortium 2012). On the other hand, the existence of a continuum of speciation in parapatric hybrid zones like those studied by Arias *et al.* (2012) (Table 1) suggests that at least some of the speciation is completed in parapatric contact or perhaps, sometimes even in complete geographic isolation.

Funding is getting more difficult for evolutionary studies in "developed" countries. Meanwhile, strong economic growth in emerging industrial

170 powers like Colombia is contributing to greatly improved strength of their
171 own science base. Further work on problems like these will be done
172 increasingly by indigenous teams like this Colombian research group, with
173 their ready access to high biodiversity in the complex topography of the
174 Andes and adjacent tropical rainforests. This work shows great potential to
175 clarify remaining questions about speciation in *Heliconius* soon.

FIGURE LEGEND:

Figure 1. Mimetic races of *Heliconius cydno* and its Müllerian comimics in the Cauca Valley of Colombia. Left side, top to bottom: *Heliconius cydno cydnides*, *H. cydno weymeri* form 'weymeri,' *H. cydno weymeri* form 'gustavi.' Right side, comimics: *H. eleuchia eleusinus*, *Elzunia humboldt*, *H. (erato) chestertonii*. Photo by Carlos Arias.

Table 1. The speciation continuum in *Heliconius* butterflies: examples of hybrid zones among species and geographic races.

Taxon 1	Taxon 2	Locality	Unimodal /Bimodal /Sympatry	mtDNA divergence	Nuclear divergence	Taxonomic status	References
<i>H. erato favorinus</i>	<i>H. erato emma</i>	NE Peru	Unimodal	No	Colour pattern loci only	Geographic races	Mallet <i>et al.</i> 1990; Countermand <i>et al.</i> 2010
<i>H. melpomene amaryllis</i>	<i>H. melpomene aglaope</i>	NE Peru	Unimodal	No	Colour pattern loci only	Geographic races	Mallet <i>et al.</i> 1990; Nadeau <i>et al.</i> 2012
<i>H. cydno cydnides</i>	<i>H. cydno weymeri</i>	Cauca, Colombia	Unimodal	Yes	Some divergence	Geographic races	Arias <i>et al.</i> 2012; this paper
<i>H. erato cyrbia</i>	<i>H. (erato) himera</i>	S Ecuador	Bimodal, 10% hybrids	Yes	Strong	Species	Jiggins <i>et al.</i> 1997
<i>H. erato venus</i>	<i>H. (erato) chestertonii</i>	W Colombia	Bimodal, 25% hybrids	Yes	Yes	Species	Arias <i>et al.</i> 2008

Reference List

- Arias CF, Muñoz AG, Jiggins CD *et al.* (2008) A hybrid zone provides evidence for incipient ecological speciation in *Heliconius* butterflies. *Molecular Ecology*, **17**, 4699-4712.
- Arias CF, Rosales C, Salazar C *et al.* (2012) Sharp genetic discontinuity across a unimodal *Heliconius* hybrid zone. *Molecular Ecology*, **21**, xxx-xxx.
- Bates HW (1862) Contributions to an insect fauna of the Amazon valley. Lepidoptera: Heliconidae. *Transactions of the Linnean Society of London*, **23**, 495-566.
- Bates HW (1863) *A Naturalist on the River Amazons*. John Murray, London.
- Blum MJ (2002) Rapid movement of a *Heliconius* hybrid zone: evidence for phase III of Wright's shifting balance theory? *Evolution*, **56**, 1992-1998.
- Chamberlain NL, Hill RI, Kapan DD, Kronforst MR (2009) Polymorphic butterfly reveals the missing link in ecological speciation. *Science*, **326**, 847-850.
- Heliconius Genome Consortium (2012) Butterfly genome reveals promiscuous exchange of mimicry adaptations among species. *Nature (London)*, **487**, 94-98.
- Hines HM, Counterman BA, Papa R *et al.* (2011) Contrasted modes of evolution in the same genome: wing patterning genes redefine the mimetic history of *Heliconius* butterflies. *Proceedings of the National Academy of Sciences, USA*, **108**, 19666-19671.
- Jiggins CD, McMillan WO, King P, Mallet J (1997) The maintenance of species differences across a *Heliconius* hybrid zone. *Heredity*, **79**, 495-505.
- Jiggins CD, Naisbit RE, Coe RL, Mallet J (2001) Reproductive isolation caused by colour pattern mimicry. *Nature (London)*, **411**, 302-305.
- Linares M (1997) The ghost of mimicry past: laboratory reconstitution of an extinct butterfly 'race'. *Heredity*, **78**, 628-635.

Mallet J, Barton N, Lamas G *et al.* (1990) Estimates of selection and gene flow from measures of cline width and linkage disequilibrium in *Heliconius* hybrid zones. *Genetics*, **124**, 921-936.

Mallet J, Beltrán M, Neukirchen W, Linares M (2007) Natural hybridization in heliconiine butterflies: the species boundary as a continuum. *BMC Evolutionary Biology*, **7**, 28.

Mavárez J, Salazar C, Bermingham E *et al.* (2006) Speciation by hybridization in *Heliconius* butterflies. *Nature (London)*, **441**, 868-871.

McMillan WO, Jiggins CD, Mallet J (1997) What initiates speciation in passion-vine butterflies? *Proceedings of the National Academy of Sciences, USA*, **94**, 8628-8633.

Müller F (1879) *Ituna* and *Thyridia*; a remarkable case of mimicry in butterflies. *Proceedings of the Entomological Society of London*, **1879**, xx-xxix.

Rosser N, Phillimore AB, Huertas B, Willmott KR, Mallet J (2012) Testing historical explanations for gradients in species richness in heliconiine butterflies of tropical America. *Biological Journal of the Linnean Society*, **105**, 479-497.