



A Meningococcal Outer Membrane Vesicle Vaccine Incorporating Genetically Attenuated Endotoxin Dissociates Inflammation from Immunogenicity

Citation

Dowling, D. J., H. Sanders, W. K. Cheng, S. Joshi, S. Brightman, I. Bergelson, C. Pietrasanta, et al. 2016. "A Meningococcal Outer Membrane Vesicle Vaccine Incorporating Genetically Attenuated Endotoxin Dissociates Inflammation from Immunogenicity." *Frontiers in Immunology* 7 (1): 562. doi:10.3389/fimmu.2016.00562. <http://dx.doi.org/10.3389/fimmu.2016.00562>.

Published Version

doi:10.3389/fimmu.2016.00562

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:29739048>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



A Meningococcal Outer Membrane Vesicle Vaccine Incorporating Genetically Attenuated Endotoxin Dissociates Inflammation from Immunogenicity

David J. Dowling^{1,2}, Holly Sanders³, Wing Ki Cheng^{1,2,4}, Sweta Joshi^{1,4}, Spencer Brightman^{1,4}, Ilana Bergelson¹, Carlo Pietrasanta^{1,2,4,5}, Simon D. van Haren^{1,2,4}, Sandra van Amsterdam³, Jeffrey Fernandez⁶, Germie P. J. M. van den Dobbelsteen³ and Ofer Levy^{1,2,4*}

OPEN ACCESS

Edited by:

Joseph M. Bliss,
Women & Infants Hospital
of Rhode Island, USA

Reviewed by:

Katie Louise Flanagan,
Monash University, Australia
Juliana Cassataro,
National Scientific and Technical
Research Council, Argentina
Patricia Talamás-Rohana,
CINVESTAV, Mexico

*Correspondence:

Ofer Levy
ofer.levy@childrens.harvard.edu

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Immunology

Received: 10 October 2016

Accepted: 22 November 2016

Published: 08 December 2016

Citation:

Dowling DJ, Sanders H, Cheng WK, Joshi S, Brightman S, Bergelson I, Pietrasanta C, van Haren SD, van Amsterdam S, Fernandez J, van den Dobbelsteen GPJM and Levy O (2016) A Meningococcal Outer Membrane Vesicle Vaccine Incorporating Genetically Attenuated Endotoxin Dissociates Inflammation from Immunogenicity. *Front. Immunol.* 7:562. doi: 10.3389/fimmu.2016.00562

¹Department of Medicine, Division of Infectious Diseases, Boston Children's Hospital, Boston, MA, USA, ²Harvard Medical School, Boston, MA, USA, ³Janssen Vaccines and Prevention B.V., Leiden, Netherlands, ⁴Precision Vaccine Program, Division of Infectious Diseases, Boston Children's Hospital, Boston, MA, USA, ⁵Neonatal Intensive Care Unit, Department of Clinical Sciences and Community Health, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, University of Milan, Milan, Italy, ⁶Janssen Research and Development, LLC, Spring House, PA, USA

Background: Group B *Neisseria meningitidis*, an endotoxin-producing Gram-negative bacterium, causes the highest incidence of group B meningococcus (MenB) disease in the first year of life. The Bexsero vaccine is indicated in Europe from 8 weeks of age. Endotoxin components of outer membrane vesicles (OMVs) or soluble lipopolysaccharide (LPS) represent a potential source of inflammation and residual reactogenicity. The purpose of this study was to compare novel candidate MenB vaccine formulations with licensed vaccines, including Bexsero, using age-specific human *in vitro* culture systems.

Methods: OMVs from wild type- and inactivated *lpxL1* gene mutant-*N. meningitidis* strains were characterized in human neonatal and adult *in vitro* whole blood assays and dendritic cell (DC) arrays. OMVs were benchmarked against licensed vaccines, including Bexsero and whole cell pertussis formulations, with respect to Th-polarizing cytokine and prostaglandin E2 production, as well as cell surface activation markers (HLA-DR, CD86, and CCR7). OMV immunogenicity was assessed in mice.

Results: Δ *lpxLI* native OMVs (nOMVs) demonstrated significantly less cytokine induction in human blood and DCs than Bexsero and most of the other pediatric vaccines (e.g., PedvaxHib, EasyFive, and bacillus Calmette–Guérin) tested. Despite a much lower inflammatory profile *in vitro* than Bexsero, Δ *lpxLI* nOMVs still had moderate DC maturing ability and induced robust anti-*N. meningitidis* antibody responses after murine immunization.

Conclusion: A meningococcal vaccine comprised of attenuated LPS-based OMVs with a limited inflammatory profile *in vitro* induces robust antigen-specific immunogenicity *in vivo*.

Keywords: group B meningococci, outer membrane vesicles, vaccine, newborn, dendritic cells

INTRODUCTION

Human newborns and infants suffer a high frequency of infection compared to older children and adults (1), in part due to distinct early life immunity with impaired host defense (2). Early life immunization is desirable, but vaccine-induced responses of newborns and young infants demonstrate slow initiation, low immunogenicity, and reduced persistence of functional antibodies and cell-mediated responses (2). Although the majority of global immunization schedules are focused on the pediatric age group, development of early life vaccines has been hampered by this distinct immunity and an *ad hoc* approach to developing vaccines, relying on adult-derived results that may inaccurately predict infant responses (3).

Neisseria meningitidis (meningococcus) is a Gram-negative, endotoxin-producing organism that is a normal commensal of the human nasopharynx and is an important cause of invasive bacterial infection in children worldwide (4). Recent immunization programs with capsular-polysaccharide vaccines have dramatically reduced the incidence of serogroup C and A meningococcal disease in North America, Europe, and Africa (5). However, meningitis and septicemia caused by serogroup B meningococci remain a major health concern in young children, as similar capsule-based vaccines cannot be developed against this serogroup (6). Outer membrane vesicle (OMV)-based vaccines have historically been used, with some success, to control outbreaks of disease caused by serogroup B meningococci. OMVs are produced by the blebbing of membranes of clinical-derived live Gram-negative bacteria during *in vitro* growth and are useful vaccine components, as immuno-stimulatory membrane components [lipids, proteins, lipopolysaccharide (LPS), etc.] from meningococci are represented (7). As OMV yield from culture alone is too low for vaccine production, detergent extraction is used to force vesiculation and increase yield and has the added advantage of reducing LPS content to prevent overt reactogenicity. Novartis' 4-component aluminum hydroxide (Alum)-adjuvanted group B meningococcus (MenB) vaccine (4CMenB, Trade name: Bexsero) has been licensed by both the European Medicines Agency and the U.S. Food and Drug Agency and comprised OMVs and three recombinant immunogenic *N. meningitidis* proteins identified by reverse vaccinology (8). Another meningococcal serogroup B vaccine, Trumenba, contains two recombinant proteins with no OMV component, is approved in the U.S. for use in individuals 10 through 25 years of age, but lacks the potentially broader antigen repertoire inherent to OMVs.

The currently licensed meningococcal vaccines have the potential to reduce mortality and morbidity associated with MenB infections, but do have some limitations. Although detergent extraction of OMVs removes the majority of the LPS, the remaining endotoxin, particularly the soluble LPS, may still result in residual reactogenicity, necessitating the use of the Alum to ameliorate excess toxicity (9). Even so, immunization with the OMV-containing Bexsero may correlate with rates of reactogenicity (e.g., fever $\geq 38^{\circ}\text{C}$, tenderness at injection site) as common as 10% in infants less than 1 years of age (10), and severe reactogenicity is reported in some cases (11). Bexsero

may also enhance reactogenicity when given together with other vaccines (12), prompting some primary care physicians to prescribe anti-inflammatory agents (13) to prevent potential reactions. Furthermore, as conventional assays for preclinical and release testing of vaccines, such as the rabbit pyrogenicity test and *Limulus* amoebocyte lysate (LAL) assay, were developed for vaccines containing no or negligible amounts of endotoxin, predicting the reactogenicity of LPS-containing OMV vaccines is difficult and requires more sophisticated models (14).

To overcome the potential reactogenicity of wild-type (WT) OMV-based vaccines while maintaining the inherent adjuvant activity of vesicles, a vaccine has previously been developed using a *N. meningitidis* strain with genetically attenuated endotoxin (15). This enabled the use of detergent-free manufacturing processes to produce native OMVs (nOMVs) (16). To assess the potential of such OMV vaccines, we reasoned that it would be important to better understand their interactions with human leukocytes, including dendritic cells (DCs). DCs are professional antigen-presenting cells (APCs) that play a vital role in shaping adaptive immunity. DC maturation begins when endogenous or exogenous danger molecules are recognized by pattern recognition receptors [e.g., Toll-like receptors (TLRs)] triggering upregulation of costimulatory molecules and production of immune polarizing cytokines (17). Of note, in addition to interactions via TLR4, *N. meningitidis* LPS also directly interacts with DCs through the C-type lectin receptor DC-SIGN and modulates their function (18).

In this study, we benchmarked novel candidate MenB vaccine formulations against licensed vaccines, including Bexsero, using physiologically relevant human neonatal and adult whole blood (WBA) and monocyte-derived dendritic cell (MoDC) *in vitro* culture systems, both of which employ age-specific autologous plasma, a rich source of age-specific immunomodulatory molecules (2, 19). We assessed vaccine-induced production of Th-polarizing cytokines, upregulation of surface activation markers (20), and production of potential vaccine reactogenicity biomarkers such as IL-1 β and prostaglandin E2 (PGE₂) (3, 21, 22). We found that a meningococcal vaccine comprised of attenuated LPS-based OMVs with a relatively low inflammatory profile toward human leukocytes *in vitro* induced robust antigen-specific immunogenicity *in vivo*. Our observations, may inform translational development of MenB vaccines in high-risk populations, such as newborns and infants.

MATERIALS AND METHODS

Ethics Statement

Peripheral blood samples were collected after written informed consent from healthy adult volunteers with approval from the Ethics Committee of Boston Children's Hospital, Boston, MA, USA (X07-05-0223). Non-identifiable cord blood samples were collected immediately after elective cesarean section delivery (epidural anesthesia) of the placenta. Births to HIV-positive or febrile mothers were excluded. Human experimentation guidelines of the U.S. Department of Health and Human Services, the Brigham and Women's Hospital, Beth Israel Medical Center, and Boston Children's Hospital were observed, with approval

from the Ethics Committee of The Brigham and Women's Hospital, Boston, MA, USA (protocol number 2000-P-000117) and Beth Israel Deaconess Medical Center Boston, MA, USA (2011P-000118). Mice were obtained from Envigo (Indianapolis, IN, USA), and studies were approved by the Janssen Research and Development, LLC Institutional Animal Care and Use Committee.

Human Blood Processing and *In Vitro* Stimulation

Human blood was anti-coagulated with 20 U/ml pyrogen-free sodium heparin (American Pharmaceutical Partners, Inc., Schaumburg, IL, USA). All blood products were kept at room temperature and processed within 4 h from collection. Whole blood (WB) was mixed 1:1 with pre-warmed (37°C) RPMI 1640 medium (Invitrogen, Carlsbad, CA, USA) and 180 μ l plated in a 96-well U-bottom plate (Becton Dickinson, Franklin Lakes, NJ, USA) containing 20 μ l of 10 \times freshly prepared treatment, as described previously (23).

Human MoDCs

Blood was layered onto Ficoll-Hypaque gradient (GE Healthcare, Waukesha, WI, USA) to collect cord blood mononuclear cells (CBMCs) or peripheral blood mononuclear cells (PBMCs). Monocytes were isolated by positive CD14 selection with magnetic microbeads (Miltenyi Biotec, Auburn, CA, USA). Preparations were routinely >95% pure as assessed by flow cytometry (23–26). Monocytes were cultured in tissue culture dishes at 0.4×10^6 cells/ml in RPMI media containing fresh 10% autologous platelet-poor plasma, supplemented with recombinant human (rh) IL-4 (50 ng/ml) and rh granulocyte-macrophage colony-stimulating factor (rhGM-CSF) (100 ng/ml) (R&D Systems, Minneapolis, MN, USA) with one replenishment of fresh media/cytokines at day 3. After 5–6 days, immature MoDCs were handled, as described previously (23, 25).

Vaccines, TLR Agonists, and Multi-Analyte Assays

Licensed vaccines were commercially obtained. Ultrapure LPS (*Salmonella minnesota* R595, List Biological Laboratories, Campbell, CA, USA) was used at 100 ng/ml. Supernatants were assayed by ELISA for TNF (BD Biosciences, San Jose, CA, USA), IL-1 β (eBiosciences, San Diego, CA, USA) and by competitive monoclonal enzyme immunoassay (EIA) for PGE₂ (Cayman Chemical, Ann Arbor, MI, USA). Additionally, assay supernatants were analyzed by magnetic bead multiplex cytokine/chemokine assay (Millipore, Billerica, MA, USA) and analyzed on the Luminex[®] 100/200™ System employing xPOTENT[®] software (Luminex, Austin, TX, USA) and Millipore Milliplex Analyst (version 3.5.5.0). PGE₂ concentrations were determined using the analysis tool at www.myassays.com.

Flow Cytometry

Monocyte-derived dendritic cells were treated with blocking agent for 10 min (Sigma-Aldrich), transferred to staining buffer

[1 \times PBS, 0.5% (v/v) human serum albumin] and stained for 30 min at 4°C in the dark (1×10^5 cells/staining) with fluorophore-labeled antibodies [CD80/Fluorescein isothiocyanate/Clone L307.4, CD86/phycoerythrin/Clone 2331, CCR7/V450/Clone 150503, and HLA-DR/PerCP-Cy5.5/Clone L243 (BD Biosciences)]. Cells were then centrifuged ($500 \times g$, 10 min), washed, fixed [1% (v/v) paraformaldehyde], and acquired using a LSRII flow cytometer employing FACSDiva software (BD Biosciences). Data were analyzed using the FlowJo (Tree Star, Inc., Ashland, OR, USA). Typically, 10,000 events per sample were acquired.

Production of OMVs and Characterization

Outer membrane vesicles were provided by Janssen Vaccines and Prevention B.V. Briefly, OMVs were produced from *N. meningitidis* H44/76 (B:P1.7,16;F3-3) WT or derivatives. nOMVs were produced from a H44/76 Δ R Δ L (15, 16, 27), resulting in a pentaacylated meningococcal LPS from the *lpxL1* mutant *N. meningitidis* strain, lacking the secondary C12:0 acyl chain at the 2'-position of the lipid A. Detergent-extracted OMVs were produced from the WT strain or the Δ R Δ L mutant (28). Adsorption to aluminum hydroxide (Sigma-Aldrich) was performed, as described elsewhere (29). Before use, OMVs were diluted in 10 mM Tris pH 7.4/3% (w/v) sucrose to 50 μ g total protein/ml. OMV size was determined by dynamic light scattering (DLS) using the Zetasizer nanoseries (Malvern Nano-ZS, 1 1/4 532 nm, Westborough, MA, USA).

In Vivo Immunogenicity

Adult Balb/c mice (6–8 weeks, female, 10/group) were immunized with two doses of OMVs (2.5 μ g total protein/dose), two doses of Bexsero [at 2.5 μ g total protein/dose (equivalent to 1/10th human dose)], or buffer control, subcutaneously in the scruff of the neck. Doses were administered 4 weeks apart, with terminal bleed taken 2 weeks after the second dose. Serum bactericidal assays (SBAs) were performed, as described previously (30), against the WT H44/76 strain with sera from individual mice using baby rabbit sera (Cedarlane, Burlington, VT, USA) as an exogenous complement source. The H44/76 strain was chosen as our OMV formulations were produced from a *N. meningitidis* H44/76 strain and since H44/76 is used as an immunogenicity indicator strain for Bexsero approval. Reported titers are the reciprocal of the serum dilution giving 50% killing. If 50% killing was not achieved the sample was assigned a titer of 4 (half of the lower limit of detection).

Statistical Analyses and Graphics

Statistical significance and graphs were generated using Prism v. 5.0b (GraphPad Software, La Jolla, CA, USA) and Microsoft Excel (Microsoft Corporation, Redmond, WA, USA). For data analyzed by normalization to control values, column statistics were conducted using the two-tailed Wilcoxon signed-rank test or unpaired Mann–Whitney test as appropriate. Results were considered significant at $p < 0.05$ and indicated as follows: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, # $p < 0.05$, ## $p < 0.01$, ### $p < 0.001$, + $p < 0.05$, ++ $p < 0.01$, and +++ $p < 0.001$.

RESULTS

OMVs Incorporating Genetically Attenuated Endotoxin Have Reduced Cytokine Induction Potential in Human Newborn and Adult Blood


Wild-type OMVs were produced from *N. meningitidis* H44/76 (B:P1.7,16;F3-3), while nOMVs (Figure 1A), incorporating a genetically attenuated version of endotoxin, were produced from a H44/76 $\Delta R\Delta L$ mutant strain (16, 27). All OMVs were characterized by DLS for particle size (Figure 1B; Figure S1 in Supplementary Material), with all formulations falling within a range of 80–120 nm in diameter. Next, we tested the ability of (1) the commercially available Bexsero vaccine, (2) detergent-extracted WT OMV (WT dOMV) (i.e., a Bexsero-like OMV formulation), (3) $\Delta R\Delta L$ mutant nOMV strain ($\Delta lpxLI$ nOMV) (28), and (4) a $\Delta lpxLI$ dOMV formulation (Figure 2A), to induce titration-dependent cytokine production in human neonatal and adult blood (Figures 2A,B). Non-detergent treated WT OMVs containing WT LPS were not tested as these are known

to be highly cytotoxic. As indicated by measurement of TNF and IL-1 β , both Bexsero and the WT dOMV robustly activated neonatal cord and adult peripheral blood in a titration-dependent manner in all concentrations tested, where levels of these two mediators were significantly increased over baseline ($p < 0.001$). The mutant OMVs also significantly induced production of TNF and IL-1 β , but only at the highest concentration in newborn blood ($p < 0.05$) (Figures 2A,B; Figure S2 in Supplementary Material). When compared to Bexsero at the highest equivalent volume-to-volume (v/v) dilution tested (1:10), $\Delta lpxLI$ nOMV demonstrated reduced TNF production in neonatal ($p < 0.001$) and adult ($p < 0.01$) blood. $\Delta lpxLI$ nOMV were also less effective in inducing IL-1 β production in newborn ($p < 0.05$) and WBA ($p < 0.01$). Similar trends were observed when comparing the $\Delta lpxLI$ nOMV to WT dOMV.

We next broadened our characterization of the ability of WT dOMV, $\Delta lpxLI$ dOMV, and $\Delta lpxLI$ nOMV (at v/v 1:1000–1:10) to induce concentration-dependent cytokine production from newborn and WBA using multiplexing assays (Figures 2C,D; Figure S3 in Supplementary Material). When compared head-to-head, the $\Delta lpxLI$ dOMV and $\Delta lpxLI$ nOMV induced markedly lower cytokine production profiles as compared to the WT dOMV, especially at the lowest concentration tested. The addition of Alum, the most commonly used adjuvant worldwide, and a component of the Bexsero vaccine to the OMV formulations altered the OMV (both wild type and mutant) induced innate cytokine production profiles (Figures S3 and S4 in Supplementary Material). Most notably, Alum reduced the ability of the WT dOMV to induce IFN γ and the chemokine CXCL10 (interferon inducible 10) (Figures S5A,B in Supplementary Material).

$\Delta lpxLI$ nOMV Mature Human Dendritic Cells without Bexsero-Associated Inflammatory Profile

To further characterize the innate immune effects of OMV formulations, 96-well human MoDC-based arrays were generated after culturing CD14 selected monocytes with IL-4 and GM-CSF in the presence of autologous plasma, a rich source of age-specific soluble immunomodulatory factors (19), as described previously (23, 25). In response to Bexsero, a strong upregulation of CD80, HLA-DR (MHCII), and CCR7 and a modest increase in CD86 were observed in both newborn and adult DCs (Figures 3A,B). Both the $\Delta lpxLI$ nOMV and WT dOMV (with or without Alum) produced slightly reduced DC maturation profiles as compared to the Bexsero. Interestingly, the Alum-adjuvanted $\Delta lpxLI$ nOMV formulation induced a similar upregulation profile to Bexsero in adult but not newborn MoDCs. Titration-dependent induced MoDC cytokine induction (Figure 3C) supported the flow cytometry results and mostly mirrored those observed in WB. When compared head-to-head, the mutant OMV formulations induced lower cytokine production profiles as compared to the WT formulations, especially at the lowest concentration tested. Interestingly, as seen in the WBA, the addition of Alum selectively enhanced WT dOMV-induced IL-1 β production from human DCs (Figure 3D). Moreover, the addition of Alum to the $\Delta lpxLI$






FIGURE 2 | OMVs containing genetically attenuated endotoxin demonstrate relatively low cytokine induction in human newborn and adult blood. Human neonatal and adult blood cultured *in vitro* for 6 h with buffer control (RPMI) or with increasing concentrations of wild type (WT) and mutant OMV formulations (1:1000–10 v/v). Supernatants were collected for ELISA (A,B) and multiplex assay [(C) newborn, (D) adult]. Results represent means ± SEM of N = 7. For analyses at individual treatments (e.g., control RPMI vs. Bexero 1:10), unpaired Mann–Whitney test was applied at each concentration, and statistical significances are denoted as follows: *p < 0.05, **p < 0.01, and ***p < 0.001. For comparison of ΔlpxLI nOMV to Bexsero, *p < 0.05, **p < 0.01, and ***p < 0.001. For comparison of ΔlpxLI nOMV to WT dOMV, *p < 0.05, **p < 0.01, and ***p < 0.001.

nOMV formulation only induced a slight increase in IL-1β from newborn (Figure 3D; Figure S6A in Supplementary Material) and adult DCs (Figures 3E,F; Figure S6B in Supplementary Material) at the highest concentration tested (1:10 v/v).

ΔlpxLI nOMV Are Less Inflammatory toward Human Leukocytes than the Majority of Pediatric Vaccines

To gain further insight into the inflammatory potential of the ΔlpxLI nOMV formulation, we next benchmarked it against a number of conventionally licensed pediatric vaccines (Table S1 in Supplementary Material) in both WB and MoDC assays. We have previously demonstrated that several of these licensed vaccines, such as the EasyFive (DTwP-HepB-Hib), induce distinct reactivity biomarker profiles *in vitro* (31, 32). When tested at equivalent v/v treatment concentrations, ΔlpxLI nOMV conversely induced a lower cytokine response for most

innate cytokines (Figures 4A,B) and chemokines (Figure S7 in Supplementary Material) tested, grouping closer to pediatric vaccines such as PCV13 and HBV, than the more inflammatory (Alum + TLR agonist)-containing pediatric vaccines (i.e., PedvaxHIB and EasyFive).

Next, immature newborn (Figure 5A) and adult DCs (Figure 5B) were assessed for vaccine-induced production of PGE₂, a molecule whose *in vitro* production has been correlated with reactogenicity *in vivo* (31–33). Of note, newborn DCs demonstrated significantly reduced ΔlpxLI nOMV-mediated PGE₂ responses as compared to both Bexsero (p < 0.01) and WT dOMV (with Alum) (p < 0.05) (Figure 5A). A similar pattern was observed for adult DCs, but with significance only observed between Bexsero and ΔlpxLI nOMV (p < 0.01) (Figure 5B).

Newborn (Figure 5C) and adult DCs (Figure 5D) responses to ΔlpxLI nOMV were also benchmarked against conventional licensed pediatric vaccines (at 1:10 v/v, Tables S1 and S2 and Figure S8 in Supplementary Material) in the MoDC array. Here,




FIGURE 3 | $\Delta lpxLI$ nOMV mature human dendritic cells without the high *in vitro* inflammatory profile induced by Bexsero. (A) Human newborn and adult MoDCs were cultured in 10% autologous plasma (v/v) and stimulated for 24 h with OMV vaccines (each at 1:10 v/v). Surface expression of costimulatory molecules and HLA was determined by flow cytometry. **(B)** Average MFI of newborn and adult MoDCs. **(C,D)** MoDCs cultured 24 h with increasing concentrations of OMV vaccines (each at 1:1000, 1:100, and 1:10 v/v). **(E,F)** Multiplex cytokine production from treated DCs. Mean \pm SEM, $n = 5-6$. For analyses at individual treatments (e.g., control RPMI vs. Bexsero), unpaired Mann-Whitney test was applied at each concentration, and statistical significances are denoted as follows: * $p < 0.05$, ** $p < 0.01$, and not significant (NS).

we focused on the correlation of both PGE₂ and IL-1 β production, as co-production of both in human monocytic assays may predict rabbit pyrogenicity (i.e., fever) *in vivo* (33). Overall, a common trend was observed. The bacillus Calmette–Guérin (BCG) vaccine, Bexsero and WT dOMV consistently induced the highest production of both PGE₂ and IL-1 β . $\Delta lpxLI$ nOMV conversely induced a lower MoDC PGE₂/IL-1 β profile than WT

OMV-based vaccines, suggesting similarity to low reactivity pediatric vaccines such as PCV13 and HBV (Figures 5C,D).

$\Delta lpxLI$ nOMV Dissociates Inflammation from Immunogenicity

Having characterized the relatively low reactivity potential of $\Delta lpxLI$ nOMV toward human leukocytes *in vitro*, we next




FIGURE 4 | $\Delta lpxLI$ nOMV are less inflammatory *in vitro* than licensed combination vaccines. Human (A) neonatal and (B) adult blood cultured *in vitro* for 6 h with buffer control (RPMI) or with increasing concentrations of WT and $\Delta lpxLI$ OMV formulations (each at 1:1000, 1:100, and 1:10 v/v) as well as the vaccines PedvaxHib, EasyFive, BCG, Cervarix, PCV13, and HBV (1:1000–10 v/v). Results for TNF, IL-1 β , IL-6, IL-10, IL-12p40, and IL-12p70 are shown and represent means \pm SEM of $N = 7$.

assessed the ability of $\Delta lpxLI$ nOMV to induce immunogenicity in mice *in vivo*. Four cohorts of 10 mice each (Figure 6) were immunized subcutaneously with Tris/Sucrose (buffer control), $\Delta lpxLI$ nOMV or WT dOMV (2.5 μ g total protein/dose, dOMVs formulated with Alum), or a 1/10th dose of Bexsero (equivalent to 2.5 μ g dOMV). Two doses were administered 4 weeks apart, with blood collected 2 weeks after the second dose to obtain serum for SBA. The buffer control alone failed to induce meningococcal bactericidal antibody responses. In marked contrast, Bexsero ($p < 0.05$), WT dOMV ($p < 0.01$), and $\Delta lpxLI$ nOMV ($p < 0.01$) all induced robust and significant antibody responses as compared to baseline (Figure 6). Remarkably, even though $\Delta lpxLI$ nOMVs demonstrated reduced inflammation potential toward human leukocytes *in vitro*, there was no significant difference in immunogenicity as compared to either Bexsero or WT dOMV.

DISCUSSION

The majority of global immunization schedules are pediatric, with particular focus on newborns and young infants, yet most vaccine discovery programs do not rationally design vaccine formulations for use in humans in early life. Indeed, even when employed, *in vitro* modeling of human responses often occurs in late stage

vaccine development. Such current pre-clinical approaches may contribute to vaccine formulations inducing sub-optimal responses in the very young (3). To more completely evaluate an OMV-based *N. meningitidis* vaccine, we took a rational vaccine design approach, in which we combined a relevant *in vivo* model and age-specific human *in vitro* culture systems that together may better model OMV-induced innate immunomodulatory capacity and immunogenicity potential. Overall, when compared to the WT detergent-extracted OMV (WT dOMV), mutant detergent-extracted OMV ($\Delta lpxLI$ dOMV), and multiple licensed vaccines with respect to innate signaling toward human newborn and adult leukocytes *in vitro*, a prototype $\Delta lpxLI$ nOMV vaccine generated a pattern of response suggestive of low reactogenicity potential while still demonstrating robust immunogenicity in mice *in vivo*. Demonstration of non-inferior *in vivo* immunogenicity of the $\Delta lpxLI$ nOMV vaccine, as compared to Bexsero, is noteworthy, especially as the *in vitro* DC maturation profiles are divergent. Such a result may indicate that the inclusion of native endotoxin molecules into traditionally designed OMV-based vaccines may not be as essential for immunogenicity as often assumed.

A key concern regarding modern vaccine development is reactogenicity (33), the propensity of a formulation to cause acute inflammatory events either locally, such as erythema or tenderness




FIGURE 5 | $\Delta lpxLI$ nOMV-induced lower MoDC-PGE₂ and -IL-1 β than Alum-adjuvanted WT OMV and the licensed Bexsero vaccine. Human newborn and adult MoDCs were cultured in 10% autologous plasma (v/v) and stimulated for 24 h with increasing concentrations of OMV vaccines (each at 1:1000, 1:100, and 1:10 v/v). (A) Newborn and (C) adult PGE₂ production, as measured by ELISA. Potential reactogenicity biomarker correlation of PGE₂ with IL-1 β for newborn (B) and adult (D) MoDCs after treatment with wild WT and mutant OMV formulations (at 1:10 v/v) as well as the vaccines PedvaxHib, EasyFive, BCG, Cervarix, PCV13, and HBV (at 1:10 v/v). Mean \pm SEM, $n = 5-6$. PGE₂ levels are depicted with untreated basal levels subtracted. For analyses at individual treatments (e.g., Bexsero 1:10 vs. $\Delta lpxLI$ nOMV 1:10), Wilcoxon test was applied at each concentration and statistical significances are denoted as follows: * $p < 0.05$ and ** $p < 0.01$.

at the injection site, or systemically, such as fever. In this context, *in vitro* assays that provide potential reactogenicity biomarker data may be highly advantageous in de-risking formulation selection for *in vivo* use (21). By employing an *in vitro* human DC array, we benchmarked the immunomodulatory abilities of an $\Delta lpxLI$ nOMV-based vaccine against several licensed pediatric vaccines. DCs are logical targets for such *in vitro* studies as they efficiently process antigens for induction of immunity against pathogens and their components (34). The few published studies that assessed the activating effects of OMV on human leukocytes *in vitro* (35–40) evaluated fixed concentrations with limited cytokine measurements. One study evaluated concentration-dependent OMV responses in human WB and PBMC assays, solely from adult donors (39). Accordingly, our study is the first to comprehensively investigate (a) the ontogeny of OMV-induced immune responses and (b) the responses of human DCs to OMVs as benchmarked against licensed vaccines. $\Delta lpxLI$ nOMV induced cytokine responses significantly lower than those induced by either Bexsero or licensed vaccines (i.e., PedvaxHIB and EasyFive). As demonstrated by the stronger induction of costimulatory molecule expression by the $\Delta lpxLI$ nOMV with Alum in adult but not newborn MoDCs, the dramatically enhanced human DC IL-1 β production in response to Alum-adjuvanted WT dOMV, as well as our prior studies demonstrating distinct ontological effects of




FIGURE 6 | $\Delta lpxLI$ nOMV induce robust immunogenicity *in vivo*. Mouse functional antibodies levels were determined by serum bactericidal assays (SBAs) after subcutaneous immunization with $\Delta lpxLI$ nOMV or WT dOMV (2.5 μ g total protein/dose) formulations, as well as the vaccines Bexsero (1/10th human dose) or buffer control. Two doses were administered 4 weeks apart, with terminal bleed taken 2 weeks after the second dose. Mutant OMV demonstrated enhanced immunogenicity over buffer control treatment, and equivalent induction of functional antibodies to the WT formulations, $n = 10$. Horizontal line indicates geometric mean. Wilcoxon test was applied between buffer and each OMV formulation, or between OMV formulations as indicated. Statistical significance is denoted as follows: * $p < 0.05$, ** $p < 0.01$, and not significant (NS).

Alum on *in vitro* DC IL-1 β production (24), it is reasonable to suggest that our age-specific *in vitro* systems may be useful tools with respect to characterizing candidate vaccine formulations with or without Alum.

Our study has multiple strengths including (a) robust age-specific human *in vitro* modeling, (b) extensive benchmarking to multiple licensed vaccines, and (c) confirmation of immunogenicity *in vivo*. That said, our study also has some limitations including (a) the unavoidable use of different species for *in vitro* (human) and *in vivo* (mouse) studies, (b) that *in vitro* systems, no matter how carefully developed, are imperfect models for responses *in vivo*, (c) the lack of full concordance in the antigens profile found in Bexsero- which contains added recombinant protein- and our $\Delta lpxLI$ nOMV formulations that do not, and (d) the lack of non-detergent treated- WT strain-derived OMVs that contain WT LPS. While such a formulation was initially not evaluated as part of our current study due to the known cytotoxic effects in humans, based on our results, such WT OMVs may provide a benchmark in future studies employing human *in vitro* systems. Of note, our experiences to date with *in vitro* benchmarking suggest that our *in vitro* studies provide insights relevant *in vivo* (2). Additionally, studies evaluating OMV vaccine-based reactogenicity *in vivo* using similar strategies to ours in mice (41) and humans (42) support these conclusions.

Several aspects of our results will prompt future investigations. For example, nucleotide-binding oligomerization domain (NOD)-like receptor-mediated IL-1 β production is greater in human newborn cord than adult peripheral blood (43, 44) and slowly decreases to adult levels over the first years of life (45). A similar phenomenon for basal PGE₂ levels in WB has been

observed (32). Therefore, further study of the IL-1 β /PGE₂ axis as a predictive marker of vaccine reactogenicity may need to take ontogeny into account. Recent use of proteomics (secretomics) has profiled hundreds of proteins released by human monocytes when stimulated *in vitro* with Alum or TLR agonists, several of which have been validated upon study of licensed adjuvanted vaccines, providing opportunities for further refinement of future adjuvanticity and reactogenicity biomarkers (31). Finally, the evaluation of different pentaacyl lipid A mutants (46) and comparison of their modified agonist properties using age-specific human WB and DC assays should also be considered.

The persistently high global burden of meningococcal disease in the very young infants and adolescents provides a compelling rationale for developing additional safe and effective early life vaccines (47). Overall, two key aspects of our study deserve particular emphasis: (a) human *in vitro* systems model age-specific biomarker responses that may correspond to reactogenicity and immunogenicity as benchmarked to licensed vaccines and (b) these *in vitro* systems may enable systematic comparison of formulations with and without candidate adjuvants early in the design process to inform translational development. Overall, if further validated, such an approach of *in vitro* assay system-informed age-specific vaccine development may open new paths to more precise vaccine development for distinct vulnerable populations.

AUTHOR CONTRIBUTIONS

DD, GD, and OL designed the study. DD, WC, SJ, SB, IB, CP, and SH conducted the *in vitro* experiments. HS, SA, and JF produced OMV vaccines and conducted the *in vivo* experiments. DD wrote the manuscript. OL provided overall mentorship and assisted in writing the manuscript. HS, WC, SJ, SB, IB, CP, SH, SA, JF, and GD contributed to helpful discussions and the careful approval of the final manuscript. All the authors have given final approval for the version submitted for publication.

REFERENCES

- Prabhudas M, Adkins B, Gans H, King C, Levy O, Ramilo O, et al. Challenges in infant immunity: implications for responses to infection and vaccines. *Nat Immunol* (2011) 12:189–94. doi:10.1038/ni0311-189
- Dowling DJ, Levy O. Ontogeny of early life immunity. *Trends Immunol* (2014) 35:299–310. doi:10.1016/j.it.2014.04.007
- Dowling DJ, Levy O. Pediatric vaccine adjuvants: components of the modern vaccinologist's toolbox. *Pediatr Infect Dis J* (2015) 34:1395–8. doi:10.1097/INF.0000000000000893
- Plotkin SA, Orenstein WA, Offit PA. *Vaccines*. Sixth ed. Philadelphia, PA: Saunders (2012).
- Ladhani SN, Ramsay M, Borrow R, Riordan A, Watson JM, Pollard AJ. Enter B and W: two new meningococcal vaccine programmes launched. *Arch Dis Child* (2016) 101:91–5. doi:10.1136/archdischild-2015-308928
- Andrews SM, Pollard AJ. A vaccine against serogroup B *Neisseria meningitidis*: dealing with uncertainty. *Lancet Infect Dis* (2014) 14:426–34. doi:10.1016/S1473-3099(13)70341-4
- Sanders H, Kaaik P, Van Den Dobbelen GP. Preclinical evaluation of MenB vaccines: prerequisites for clinical development. *Expert Rev Vaccines* (2013) 12:31–42. doi:10.1586/erv.12.137

ACKNOWLEDGMENTS

The authors thank the members of the OL's Laboratory for assistance with phlebotomy as well as helpful discussions. The authors thank their donors and acknowledge the assistance of the Labor and Delivery staff at both The Brigham and Women's Hospital, Boston and Beth Israel Deaconess Medical Center, Boston. The authors are grateful for the mentorship and support of Drs. Michael Wessels and Gary R. Fleisher.

FUNDING

This study was supported by Janssen Vaccines and Prevention B.V. (formerly Crucell Holland B.V.). OL's laboratory is supported by U.S. National Institutes of Health (NIH) grants 1R01AI100135-01, and 3R01AI067353-05S1, the National Institutes of Allergy and Infectious Diseases (NIAID), NIH, Department of Health and Human Services, NIH UO1 award Molecular Mechanisms of Combination Adjuvants (1U01AI124284-01), Adjuvant Discovery Program Contract No. HHSN272201400052C as well as Global Health (OPPGH5284) and Grand Challenges Explorations (OPP1035192) awards from the Bill & Melinda Gates Foundation. CP was supported by the scholarship “J. Miglierina,” Fondazione Comunitaria del Varesotto, Varese, Italy. OL's Laboratory has received sponsored research support from VentiRx Pharmaceuticals, 3M Drug Delivery Systems, MedImmune, Crucell (Johnson & Johnson), Shire, and an internal Boston Children's Hospital award to the Precision Vaccines Program.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://journal.frontiersin.org/article/10.3389/fimmu.2016.00562/full#supplementary-material>.

- Serruto D, Bottomley MJ, Ram S, Giuliani MM, Rappuoli R. The new multicomponent vaccine against meningococcal serogroup B, 4CMenB: immunological, functional and structural characterization of the antigens. *Vaccine* (2012) 30(Suppl 2):B87–97. doi:10.1016/j.vaccine.2012.01.033
- Fredriksen JH, Rosenqvist E, Wedege E, Bryn K, Bjune G, Froholm LO, et al. Production, characterization and control of MenB-vaccine “Folkehelsa”: an outer membrane vesicle vaccine against group B meningococcal disease. *NIPH Ann* (1991) 14:67–79; discussion 79–80.
- CHMP. *Annex I – Summary of Product Characteristics*. European Medicines Agency European Public Assessment Report (EPAR). Bexsero meningococcal group-B vaccine, EMEA/H/C/002333, London (2015).
- Tenenbaum T, Niessen J, Schroten H. Severe upper extremity dysfunction after 4CMenB vaccination in a young infant. *Pediatr Infect Dis J* (2016) 35:94–6. doi:10.1097/INF.0000000000000917
- Vesikari T, Esposito S, Prymula R, Ypma E, Kohl I, Toneatto D, et al. Immunogenicity and safety of an investigational multicomponent, recombinant, meningococcal serogroup B vaccine (4CMenB) administered concomitantly with routine infant and child vaccinations: results of two randomised trials. *Lancet* (2013) 381:825–35. doi:10.1016/S0140-6736(12)61961-8
- Prymula R, Esposito S, Zuccotti GV, Xie F, Toneatto D, Kohl I, et al. A phase 2 randomized controlled trial of a multicomponent meningococcal serogroup

- B vaccine (I). *Hum Vaccin Immunother* (2014) 10:1993–2004. doi:10.4161/hv.28666
14. Vipond C, Findlay L, Feavers I, Care R. Limitations of the rabbit pyrogen test for assessing meningococcal OMV based vaccines. *ALTEX* (2016) 33:47–53. doi:10.14573/altex.1509291
 15. Van Der Ley P, Steeghs L, Hamstra HJ, Ten Hove J, Zomer B, Van Alphen L. Modification of lipid A biosynthesis in *Neisseria meningitidis* lpxL mutants: influence on lipopolysaccharide structure, toxicity, and adjuvant activity. *Infect Immun* (2001) 69:5981–90. doi:10.1128/IAI.69.10.5981-5990.2001
 16. Van De Waterbeemd B, Streefland M, Van Der Ley P, Zomer B, Van Dijken H, Martens D, et al. Improved OMV vaccine against *Neisseria meningitidis* using genetically engineered strains and a detergent-free purification process. *Vaccine* (2010) 28:4810–6. doi:10.1016/j.vaccine.2010.04.082
 17. Dowling D, Hamilton CM, O'Neill SM. A comparative analysis of cytokine responses, cell surface marker expression and MAPKs in DCs matured with LPS compared with a panel of TLR ligands. *Cytokine* (2008) 41:254–62. doi:10.1016/j.cyto.2007.11.020
 18. Steeghs L, Van Vliet SJ, Uronen-Hansson H, Van Mourik A, Engering A, Sanchez-Hernandez M, et al. *Neisseria meningitidis* expressing lgtB lipopolysaccharide targets DC-SIGN and modulates dendritic cell function. *Cell Microbiol* (2006) 8:316–25. doi:10.1111/j.1462-5822.2005.00623.x
 19. Pettengill MA, Van Haren SD, Levy O. Soluble mediators regulating immunity in early life. *Front Immunol* (2014) 5:457. doi:10.3389/fimmu.2014.00457
 20. Levitz SM, Golenbock DT. Beyond empiricism: informing vaccine development through innate immunity research. *Cell* (2012) 148:1284–92. doi:10.1016/j.cell.2012.02.012
 21. Mastelic B, Garcon N, Del Giudice G, Golding H, Gruber M, Neels P, et al. Predictive markers of safety and immunogenicity of adjuvanted vaccines. *Biologicals* (2013) 41:458–68. doi:10.1016/j.biologicals.2013.08.006
 22. Lewis DJ, Lythgoe MP. Application of “systems vaccinology” to evaluate inflammation and reactogenicity of adjuvanted preventative vaccines. *J Immunol Res* (2015) 2015:909406. doi:10.1155/2015/909406
 23. Dowling DJ, Tan Z, Prokopowicz ZM, Palmer CD, Matthews MA, Dietsch GN, et al. The ultra-potent and selective TLR8 agonist VTX-294 activates human newborn and adult leukocytes. *PLoS One* (2013) 8:e58164. doi:10.1371/journal.pone.0058164
 24. Philbin VJ, Dowling DJ, Gallington LC, Cortes G, Tan Z, Suter EE, et al. Imidazoquinoline toll-like receptor 8 agonists activate human newborn monocytes and dendritic cells through adenosine-refractory and caspase-1-dependent pathways. *J Allergy Clin Immunol* (2012) 130(195–204):e199. doi:10.1016/j.jaci.2012.02.042
 25. Palmer CD, Ninkovic J, Prokopowicz ZM, Mancuso CJ, Marin A, Andrianov AK, et al. The effect of stable macromolecular complexes of ionic polyphosphazene on HIV Gag antigen and on activation of human dendritic cells and presentation to T-cells. *Biomaterials* (2014) 35:8876–86. doi:10.1016/j.biomaterials.2014.06.043
 26. Ganapathi L, Van Haren S, Dowling DJ, Bergelson I, Shukla NM, Malladi SS, et al. The imidazoquinoline toll-like receptor-7/8 agonist hybrid-2 potently induces cytokine production by human newborn and adult leukocytes. *PLoS One* (2015) 10:e0134640. doi:10.1371/journal.pone.0134640
 27. Van De Waterbeemd B, Zomer G, Kaaijk P, Ruiterkamp N, Wijffels RH, Van Den Dobbelen GP, et al. Improved production process for native outer membrane vesicle vaccine against *Neisseria meningitidis*. *PLoS One* (2013) 8:e65157. doi:10.1371/journal.pone.0065157
 28. Sanders H. *Investigation of the Potential of PorA and FetA as Meningococcal Vaccine Components*. Oxford: University of Oxford (2012).
 29. Norheim G, Aase A, Caugant DA, Hoiby EA, Fritszone E, Tangen T, et al. Development and characterisation of outer membrane vesicle vaccines against serogroup A *Neisseria meningitidis*. *Vaccine* (2005) 23:3762–74. doi:10.1016/j.vaccine.2005.02.021
 30. Maslanka SE, Gheesling LL, Libutti DE, Donaldson KB, Harakeh HS, Dykes JK, et al. Standardization and a multilaboratory comparison of *Neisseria meningitidis* serogroup A and C serum bactericidal assays. The Multilaboratory Study Group. *Clin Diagn Lab Immunol* (1997) 4:156–67.
 31. Oh DY, Dowling DJ, Ahmed S, Choi H, Brightman S, Bergelson I, et al. Adjuvant-induced human monocyte secretome profiles reveal adjuvant- and age-specific protein signatures. *Mol Cell Proteomics* (2016) 15(6):1877–94. doi:10.1074/mcp.M115.055541
 32. Van Haren SD, Ganapathi L, Bergelson I, Dowling DJ, Banks M, Samuels RC, et al. In vitro cytokine induction by TLR-activating vaccine adjuvants in human blood varies by age and adjuvant. *Cytokine* (2016) 83:99–109. doi:10.1016/j.cyto.2016.04.001
 33. Zaitseva M, Romantseva T, Blinova K, Beren J, Sirota L, Drane D, et al. Use of human MonoMac6 cells for development of in vitro assay predictive of adjuvant safety in vivo. *Vaccine* (2012) 30:4859–65. doi:10.1016/j.vaccine.2012.05.002
 34. Bachmann MF, Jennings GT. Vaccine delivery: a matter of size, geometry, kinetics and molecular patterns. *Nat Rev Immunol* (2010) 10:787–96. doi:10.1038/nri2868
 35. Koeberling O, Seubert A, Granoff DM. Bactericidal antibody responses elicited by a meningococcal outer membrane vesicle vaccine with overexpressed factor H-binding protein and genetically attenuated endotoxin. *J Infect Dis* (2008) 198:262–70. doi:10.1086/589308
 36. Steeghs L, Keestra AM, Van Mourik A, Uronen-Hansson H, Van Der Ley P, Callard R, et al. Differential activation of human and mouse toll-like receptor 4 by the adjuvant candidate LpxL1 of *Neisseria meningitidis*. *Infect Immun* (2008) 76:3801–7. doi:10.1128/IAI.00005-08
 37. Koeberling O, Giuntini S, Seubert A, Granoff DM. Meningococcal outer membrane vesicle vaccines derived from mutant strains engineered to express factor H binding proteins from antigenic variant groups 1 and 2. *Clin Vaccine Immunol* (2009) 16:156–62. doi:10.1128/CVI.00403-08
 38. Franssen F, Hamstra HJ, Boog CJ, Van Putten JP, Van Den Dobbelen GP, Van Der Ley P. The structure of *Neisseria meningitidis* lipid A determines outcome in experimental meningococcal disease. *Infect Immun* (2010) 78:3177–86. doi:10.1128/IAI.01311-09
 39. Stoddard MB, Pinto V, Keiser PB, Zollinger W. Evaluation of a whole-blood cytokine release assay for use in measuring endotoxin activity of group B *Neisseria meningitidis* vaccines made from lipid A acylation mutants. *Clin Vaccine Immunol* (2010) 17:98–107. doi:10.1128/CVI.00342-09
 40. Koeberling O, Seubert A, Santos G, Colaprico A, Uguzzoli M, Donnelly J, et al. Immunogenicity of a meningococcal native outer membrane vesicle vaccine with attenuated endotoxin and over-expressed factor H binding protein in infant rhesus monkeys. *Vaccine* (2011) 29:4728–34. doi:10.1016/j.vaccine.2011.04.095
 41. Daniels-Treffandier H, De Nie K, Marsay L, Dold C, Sadarangani M, Reyes-Sandoval A, et al. Impact of reducing complement inhibitor binding on the immunogenicity of native *Neisseria meningitidis* outer membrane vesicles. *PLoS One* (2016) 11:e0148840. doi:10.1371/journal.pone.0148840
 42. Keiser PB, Biggs-Cicatelli S, Moran EE, Schmiel DH, Pinto VB, Burden RE, et al. A phase 1 study of a meningococcal native outer membrane vesicle vaccine made from a group B strain with deleted lpxL1 and synX, over-expressed factor H binding protein, two PorAs and stabilized OpcA expression. *Vaccine* (2011) 29:1413–20. doi:10.1016/j.vaccine.2010.12.039
 43. Kollmann TR, Crabtree J, Rein-Weston A, Blimkie D, Thommai F, Wang XY, et al. Neonatal innate TLR-mediated responses are distinct from those of adults. *J Immunol* (2009) 183:7150–60. doi:10.4049/jimmunol.0901481
 44. Reikie BA, Adams RC, Ruck CE, Ho K, Leligowicz A, Pillay S, et al. Ontogeny of toll-like receptor mediated cytokine responses of South African infants throughout the first year of life. *PLoS One* (2012) 7:e44763. doi:10.1371/journal.pone.0044763
 45. Burl S, Townend J, Njie-Jobe J, Cox M, Adetifa UJ, Touray E, et al. Age-dependent maturation of toll-like receptor-mediated cytokine responses in Gambian infants. *PLoS One* (2011) 6:e18185. doi:10.1371/journal.pone.0018185
 46. Pupo E, Hamstra HJ, Meiring H, Van Der Ley P. Lipopolysaccharide engineering in *Neisseria meningitidis*: structural analysis of different pentaacyl lipid A mutants and comparison of their modified agonist properties. *J Biol Chem* (2014) 289:8668–80. doi:10.1074/jbc.M114.554345
 47. Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE, et al. Global, regional, and national causes of child mortality: an updated systematic analysis for

2010 with time trends since 2000. *Lancet* (2012) 379:2151–61. doi:10.1016/S0140-6736(12)60560-1

Conflict of Interest Statement: HS, SA, JF, and GD are employees of Janssen Vaccines and Prevention B.V., part of the Janssen pharmaceutical companies of Johnson & Johnson. This study was funded by Janssen Vaccines and Prevention B.V. (formerly Crucell B.V.). All the other authors report no potential conflicts.

Copyright © 2016 Dowling, Sanders, Cheng, Joshi, Brightman, Bergelson, Pietrasanta, van Haren, van Amsterdam, Fernandez, van den Dobbsteijn and Levy. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.