Atom—Atom Collision Processes in Astrophysics: Theoretical Studies

Citation

Dalgarno, A. 1967. "Atom—Atom Collision Processes in Astrophysics: Theoretical Studies." *Reviews of Modern Physics* 39 (4) (October 1): 850–857. doi:10.1103/revmodphys.39.850.

Published Version

doi:doi.org/10.1103/RevModPhys.39.850

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:29914192>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Atom-Atom Collision Processes in Astrophysics: Theoretical Studies*

A. DALGARNO†

School of Physics and Applied Mathematics, The Queen's University, Belfast, Northern Ireland

Theoretical studies of atom-atom collision processes which may be relevant to the interpretation of astrophysical phenomena are reviewed. Work on excitation, ionization, and electron capture in atomic hydrogen by fast protons and He^+ ions is summarized. Processes involving slow collisions that are listed include radiative charge transfer and association, associative detachment, Penning ionization, spin-change, collision-induced deactivation, mutual neutralization, and rotational excitation.

Atom-atom collision processes which are relevant to astrophysical problems appear to occur either at thermal energies or at very high energies. Low-energy collisions require special considerations, but, with the exception of rearrangement processes, high-energy collisions can be described by the Born approximation. It follows that proton-impact cross sections for excitation and ionization of the target species are comparable to, but larger than, those for electron impact for equal *velocities* of the proton and the electron. Excitation and ionization by heavy particles other than bare nuclei is complicated by the possibility of excitation and ionization of the incident heavy particle; and collisions in which the incident particle and the target particle both undergo transitions may be more probable at high energies than collisions in which only one of the colliding pair undergoes a transition. However, high-energy neutral particles are rarely of interest in astrophysics since they are readily ionized, and the reverse process of charge transfer or electron capture is improbable. Although the high-energy behavior of the cross section for charge transfer

is still obscure (see Nikolaev 1965 and Bransden 1966 for reviews), it is clear that the cross section ultimately decreases rapidly. The expected rapid decrease may not emerge from laboratory investigations since capture of inner shell electrons is important up to very high energies (Mapleton 1966).

Except for bare nuclei, heavy particles can induce transitions involving a change of spin multiplicity. In such transitions the core of the heavy particle behaves as a spectator in the high-velocity limit, and the active electron may be regarded as a free electron. Explicit calculations for the reactions

have been carried out by Bates and Crothers (1967).

Comprehensive accounts of atom-atom collisions are

† Present address: Smithsonian Astrophysical Observatory and Harvard College Observatory, Cambridge, Mass.

* Paper prepared for International Astronomical Union Colloquium on Atomic Collision Processes, Boulder, Colorado, July 1966.

given in *Atomic and Molecular Processes* edited by Bates (1962), and a collection of more recent papers and reviews comprises *Atomic Collision Processes* edited by McDowell (1964).

HIGH-ENERGY COLLISIONS

Because of the occurrence of cosmic rays, high-energy collisions of protons and of neutral hydrogen atoms with hydrogen and possibly helium assume a special importance, and we present a brief discussion of the available data.

Comprehensive data are available only within the Born approximation. Calculations on a few processes have been performed using more refined approximations which are useful in assessing the accuracy and the range of validity of the Born approximation results (Bates 1959, 1961, Mittleman 1961, Bates and Williams 1964, Coleman and McDowell 1965, Lovell and McElroy 1965, Ingber 1965, Fulton and Mittleman 1965, Cheshire 1965, Coleman and McDowell 1966, Wilets and Gallaher 1966, Callaway and Dugan 1966, Basu, Sil, and Bhattacharya 1966, McCarroll and Salin 1966).

EXCITATION AND IONIZATION OF H BY H^+

Bates and Griffing (1953) (see also Bates 1958) have presented the results of Born approximation calculations of cross sections $Q(n_i l_i; n_f l_f)$ for the excitation processes

and the ionization process

(See also Peach 1965.) For the ionization process, they present the velocity distributions of the ejected electron. The ionization cross section agrees closely with the measurements of Gilbody and Ireland (1964) above 40 keV, but lie above the measurements of Fite, Stebbings, Hummer, and Brackman (1960) below 40 keV. The cross section for excitation into the $2p$ state is about twice that measured by Stebbings, Young, Oxley, and Ehrhardt (1965) between 5 and 30 keV but falls below it for energies below 3 keV.

At high energies (E measured in keV),

$$Q(1s; 2s) \sim (11.1/E)[1 - (7.8/E)]$$

and

$$Q(1s; 2p) \sim (128/E)[\log E - 1.185 + (4.1/E)].$$

Cross sections are expressed in units of πa_0^2 . Thus excitation of p states dominates in the asymptotic region. Cross sections for other values of n_i , l_i , n_f , and l_f may be derived from electron-impact cross sections (Bates and Griffing 1953, Carew and Milford 1963). May (1965a) has obtained a simple formula for excitation into all the substates corresponding to a particular value of n when n is large. He shows that

$$Q(1s; \sum_l nl) = [I(E)/n^3](1 + O(n^{-2})), \quad (3)$$

where $I(E)$ is given in Table I. Cross sections for excitation into all the discrete levels

$$\sum_{nl} Q(1s; nl)$$

have been calculated by Butler and Parcell (1965), and their results are reproduced in Table II.

Cross sections for transitions in which the target atom is excited have been computed by Carew and Milford (1963), who give results for the target atom in states with principal quantum numbers $n_i = 2, 3, 4, 5$, and 10 for transitions such that $n_f - n_i = 1$ or 2. With decreasing threshold energy, the cross-section maximum increases to lower energy and, as n_i increases, proton impact becomes more efficient than electron impact in causing transitions in the thermal energy range.

EXCITATION AND IONIZATION OF H BY H

Bates and Griffing (1954, 1955) (see also Bates 1958) have presented the results of Born approximation calculations of cross sections $Q(1s - n_1 l_1; 1s - n_2 l_2)$ for the excitation processes

and for ionization processes

TABLE I. Values of the function $I(E)$ appearing in Eq. (3).

Impact energy E (keV)	$I(E)$
1.6	0.58
3.1	2.5
6.3	6.1
12.5	8.9
25	8.5
50	6.5
100	4.4
200	2.8

TABLE II. Cross sections Q in units of a_0^2 for $H^+ + H(1s) \rightarrow H^+ + H(nl)$.

Impact energy (keV)	$\sum_l \{Q(1s, 2l) + Q(1s, 3l)\}$	$\sum_n Q(1s; nl)$
6.24	2.26	2.53
7.71	2.49	2.83
9.76	2.67	3.04
12.74	2.74	3.12
17.35	2.75	3.15
25.0	2.58	2.95
39.0	2.24	2.57
69.4	1.72	1.95
156.1	1.08	1.23
277.6	0.74	0.83
624.5	0.41	0.44

At high impact energies, E (keV),

$$Q(1s - 2s; 1s - \sum) \sim 4.3/E \quad (7)$$

$$Q(1s - 2p; 1s - \sum) \sim 21/E, \quad (8)$$

where $\sum$ is the sum of all possible final states, including the continuum. For ionization at high energies,

$$Q(1s - C; 1s - \sum) \sim 128/E.$$

Collisions in which the incident atom remains in the $1s$ state contribute only about 13% of the total.

May (1965b) has derived further asymptotic formulas for

$$Q(1s - \sum_l nl; 1s - 1s) \quad \text{and} \quad Q(1s - \sum_l nl; 1s - \sum),$$

showing that for large n they decrease as n^{-3} .

Bates and Griffing (1955) have obtained the velocity distributions of the electrons ejected in ionizing collisions of stationary hydrogen atoms by hydrogen-atom impact. To compute the velocity distributions of electrons ejected from the moving projectile atom, it is necessary to calculate also the angular distributions of the ejected electrons, and this calculation has been carried out by Dalgarno and Griffing (1958).

Collisions in which the target atom is excited have been investigated by Bouthiette, Healey, and Milford (1964) and collisions in which both the projectile and the target atom are excited by Pomilla and Milford (1966).

The charge transfer process

has been investigated by Mapleton (1965).

CHARGE TRANSFER OF H^+ IN H

Cross sections for electron capture by protons from atomic hydrogen

have been computed using the Brinkman-Kramer form of the Born approximation for final states up to $4f$ by Bates and Dalgarno (1953), and their work has been

extended to higher states of excitation and to capture from excited atoms by Butler and Johnston (1964), by May (1964), by Hiskes (1965), and by May and Lodge (1965). The accuracy of the absolute cross sections is open to question, but the relative values for capture into different excited states appear to be satisfactory for impact energies above 40 keV (cf. Hiskes 1965). For large values of n , the cross sections decrease as n^{-3} .

Comparisons of different first-order approximations (cf. Bates 1962) have been made by Bates and Dalgarno (1952), Jackson and Schiff (1953), Bassel and Gerjuoy (1960), McCarroll (1961), and Mapleton (1962).

The cross section for capture into the $2p$ state is about twice that measured by Stebbings *et al.* (1965) between 5 and 30 keV, but falls below it for energies below 3 keV.

STOPPING POWER OF H^+ IN H

The efficiency with which a beam of protons is slowed down in a gas of ground-state hydrogen atoms has been examined in detail by Dalgarno and Griffing (1955) using the Born approximation to the cross sections for the various processes. The mean energy expended in producing an ion pair in hydrogen gas has also been calculated as a function of impact energy (Dalgarno and Griffing 1958). The neutralization of the ion beam appears to extend the range of impact energies over which the mean energy per ion pair is essentially constant at a value of about 32 eV.

STOPPING POWER OF H^+ IN AN H^+ , e PLASMA

The energy loss of fast protons in an ionized plasma has been calculated by Hayakawa and Kitao (1956) and by Butler and Buckingham (1962). The presence of free electrons in a gas increases markedly the rate of energy loss.

EXCITATION AND IONIZATION OF He BY H^+ AND H

Theoretical calculations of the excitation and ionization of He are much less comprehensive than for H. The following processes have been studied:

(Moiseiwitsch and Stewart 1954, Bell 1961, Bell and Skinner 1962);

(Bell 1961, Bell and Skinner 1962);

(Mapleton 1958, Peach 1965);

(Dalgarno and McDowell 1955, Mapleton 1958);

(Adler and Moiseiwitsch 1957);

(Bates and Williams 1957).

The predicted ionization cross sections agree well with the measurements of Hooper, Harmer, Martin, and McDaniel (1962) at impact energies above 400 keV but lie above the experimental cross sections (Federenko, Afrosimov, Il'in, and Solov'ev 1960, Hooper *et al.* 1962) at lower energies.

CHARGE TRANSFER OF H^+ IN He

Mapleton (1961, 1963) has used the Born approximation to calculate cross section for electron capture by protons from helium,

for various excited states of the end products. The total charge-transfer cross section is in good agreement with the measured cross section (Stier and Barnett 1956, Barnett and Reynolds 1958) above 40 keV, and the cross section for capture into the $3s$ state of atomic hydrogen is in good agreement with that measured by Hughes, Dawson, Doughty, Kay, and Stiger (1966) above 100 keV, but lies above it at lower energies. Hughes *et al.* also obtained evidence suggesting that the cross sections for capture into the n th state of hydrogen decreases as n^{-3} for impact energies above about 20 keV. The theoretical cross sections for capture into the $2s$ and $2p$ states of hydrogen lie well above those measured by Pretzer, Van Zyl, and Geballe (1964), by Colli, Cristofori, Frigerio, and Sona (1962) and by Jaacks, Van Zyl, and Geballe (1965) at energies below 20 keV.

The electron capture process for the case when the hydrogen atom and helium atom are produced in their ground states has been studied using higher-order approximations by Bransden and Cheshire (1963), by Green, Stanley, and Chang (1965), and by Bransden and Sin Fai Laim (1966).

LOW-ENERGY COLLISIONS

Excitation by proton impact

The cross section for direct excitation by electron impact is broadly similar to that by proton impact if the electron and proton have the same velocity. The electron impact excitation cross section rises rapidly from threshold to a maximum, which occurs at an impact velocity corresponding to about twice the threshold energy, and then decreases asymptotically as $E^{-1} \ln E$

for optically allowed transitions, or as E^{-1} for optically forbidden transitions. Thus electron impact excitation is usually more efficient than proton impact excitation except when the threshold energy is only a small fraction of the thermal energy of the plasma.

An example of an excitation process for which proton impact is more efficient than electron impact is the $2s-2p$ transition in atomic hydrogen

The threshold energies are, respectively, 0.0354 and 0.327 cm⁻¹. The reaction is important in reducing the intensity of the two-quantum emission from H(2s).

The cross section has been calculated by Seaton (1955) using a partial wave analysis in which he ensured that no partial wave violated the conservation laws. The derived rate coefficients at 10 000° and at 20 000°K are reproduced in Table III. They are an order of magnitude larger than those for electrons. The charge transfer process

is negligible compared to the direct reaction (Boyd and Dalgarno 1958).

Collisions which lead merely to a redistribution of angular momentum

will usually proceed very rapidly.

Similar reactions which may be of astrophysical importance are the proton impact excitation of molecular rotations

The particular case

has been investigated by Thaddeus and Clauser (1966) in connection with the cosmic microwave radiation at $\lambda = 2.63$ mm (see also Field and Hitchcock 1966). Their calculations appear to be based upon an approximation analogous to the Bethe form of the Born approximation (cf. Seaton 1962). There occurs a strong long-range interaction between the proton and the permanent dipole moment of CN, and the derived cross section for 1-eV protons is about 10^{-12} cm². Thaddeus and Clauser suggest that the process may be important in HII regions.

TABLE III. Rate coefficient k for $\text{H}^+ + \text{H}(2s) \rightarrow \text{H}^+ + \text{H}(2p_i)$.

$T^\circ\text{K}$		10 000	20 000
k (cm ³ sec ⁻¹)	$j=\frac{1}{2}$	2.5×10^{-4}	2.1×10^{-4}
	$j=\frac{3}{2}$	2.2×10^{-4}	2.2×10^{-4}

TABLE IV. Diffusion coefficients D^+ of H^+ in H.

$T^\circ\text{K}$	1000	2000	5000	10 000	20 000	50 000
$D^+_{\text{n(H)}} \times 10^{17}$	1.7	2.6	4.6	7.1	11.1	20.0

Because of the strong coupling between different rotational levels during the collision, the cross sections for exciting higher rotational levels will be of comparable magnitude.

CHARGE TRANSFER

The symmetrical resonance charge-transfer process

does not involve an electronic transition, and its cross section can be predicted reliably. At 1 eV the cross section is about 4×10^{-15} cm² decreasing slowly with increasing energy (Dalgarno and Yadav 1953, Ferguson 1961, Peek 1966).

Resonance charge transfer significantly modifies the diffusion of H^+ in H. Values of the ion diffusion coefficient have been calculated by Dalgarno (1961a), and they are reproduced in Table IV.

Asymmetric charge transfer processes

involve an electronic transition, and they will, in most cases, proceed very slowly at thermal energies. Exceptions occur, and one such is

which is accidentally resonant. Chamberlain (1956) has suggested that the reaction may affect the ratio of the 0 to 0⁺ densities in the Cassiopeia radio source.

There is no satisfactory theory of asymmetric charge transfer reactions, but an approximate analysis by Rapp (1963) of the measurements at high energies by Fite, Smith, and Stebbings (1962) shows that it behaves like a symmetric charge transfer reaction down to very low energies. The predicted cross section at 1 eV is 2×10^{-15} cm², a value consistent with the cross section of 8×10^{-15} cm² at 1000°K derived by Hanson, Patterson, and Degaonkar (1963) from upper atmosphere data.

RADIATIVE CHARGE TRANSFER

Radiative charge-transfer processes in which a photon is emitted may proceed more rapidly than ordinary charge transfer at thermal energies. Calculations have been carried out for two simple cases using a theory in which the probability of a transition of the quasi-molecule formed by the colliding pair is integrated along the classical path (Bates 1951). The reactions which have been investigated are

(Arthurs and Hyslop 1957) and

(Allison and Dalgarno 1965). The first reaction has a rate coefficient of $1.5 \times 10^{-13} \text{ sec}^{-1}$ at $20\,000^\circ\text{K}$ and the second a rate coefficient of $4 \times 10^{-15} \text{ cm}^3 \text{ sec}^{-1}$.

RADIATIVE ASSOCIATION

Radiative association

provides a mechanism for the formation of molecules in gases at low densities. It may be described by a theory analogous to that appropriate to radiative charge transfer. Bates (1951) has computed the rate coefficient for

obtaining a value of $5 \times 10^{-18} \text{ cm}^3 \text{ sec}^{-1}$ at 1000°K increasing to $4 \times 10^{-16} \text{ cm}^3 \text{ sec}^{-1}$ at $20\,000^\circ\text{K}$. Bates also tabulates the rate coefficient for the free-free transition

For the reactions

Bates (1951) has given tentative estimates of $2 \times 10^{-18} \text{ cm}^3 \text{ sec}^{-1}$, but he notes that the rate coefficient of the former may be $6 \times 10^{-18} \text{ cm}^3 \text{ sec}^{-1}$ if the carbon atoms are all in the 3P_0 state, and that the rate coefficient of the latter may vanish if the carbon atoms are all in the $^2P_{1/2}$ state.

The rate of formation of HeH^+ has not been calculated, though the molecule undoubtedly exists (Michels 1966, Harris 1966).

The free-bound absorption

has been investigated by Erkovich (1960), Soshnikov (1964), and Solomon (1964) for the transition ($1s\sigma 2p\sigma^3\Sigma_u^+ - 1s\sigma 2s\sigma^3\Sigma_g^+$). The process does not seem to be a significant source of absorption.

The radiative association of two hydrogen atoms

is a very slow process since the stabilizing transition is highly forbidden. Malville (1964) has given an estimate of $4 \times 10^{-27} \text{ cm}^3 \text{ sec}^{-1}$ at 100°K for its rate coefficient.

COLLISION-INDUCED RADIATIVE DEACTIVATION

Allison and Dalgarno (1963) have calculated the rate coefficient for

Because of the large polarizability of $\text{He}(2^1S)$, the reaction proceeds quite rapidly with a rate coefficient of $10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ at $20\,000^\circ\text{K}$.

A similar study of the reaction

has been completed by Allison, Browne, and Dalgarno (1966), who obtain a rate coefficient of $3 \times 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$ at $20\,000^\circ\text{K}$.

ASSOCIATIVE DETACHMENT

Associate detachment

is an efficient mechanism for the destruction of negative ions. It may be regarded as proceeding through the formation of a quasi-molecule XY^- which can undergo autodetachment. Since the mean time for autodetachment is usually much less than the mean time the nuclei remain in the autodetaching region, the probability of autodetachment is high. This simple picture led Dalgarno (1961a) to advocate a rate coefficient of up to $10^{-9} \text{ cm}^3 \text{ sec}^{-1}$ for

Developments in the theory of resonating states should yield a more precise estimate (see in particular Bardsley, Herzenberg, and Mandl 1966).

The process is also of interest in that it leads to a production of hydrogen molecules (McDowell 1961).

MUTUAL NEUTRALIZATION

Mutual neutralization

usually proceeds rapidly at thermal energies because of the long range attraction in the initial channel and because intersecting potential energy surfaces are available connecting the initial and final channels. (Bates and Massey 1954, Bates and Boyd 1956). The case

has been studied in detail by Bates and Lewis (1955), who have shown that the rate coefficient may be as large as $10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ and varies approximately as $T^{-1/2}$.

PENNING IONIZATION

Thermal ionization

can also occur efficiently in many instances because of the existence of intersecting potential energy surfaces (Bates and Massey 1954). The colliding systems may become bound forming XY^+ , a process known as associative ionization. The processes are analogous to associative detachment in that a radiationless transition (autoionization) occurs.

Quantal studies have been made by Katsuura (1965), Smirnov and Firsov (1965), Mori (1966), Sheldon (1966), and Bates, Bell, and Kingston (1967). Bates *et al.* have emphasized that the long-range interactions can lead to large rate coefficients. For

Bates *et al.* estimate a rate coefficient at temperature T of $2 \times 10^{-8} T^{-1/6} \text{ cm}^3 \text{ sec}^{-1}$ and for

a rate coefficient of $7 \times 10^{-10} T^{1/6} \text{ cm}^3 \text{ sec}^{-1}$. They note that Penning ionization processes, which can take place only through exchange and which involve the transfer of an electron between compact orbitals around unlike centers, such as

will be less efficient. Rate coefficients for the ionization of alkali atoms by metastable inert gas atoms have been computed by Sheldon (1966) and for the ionization of Li and Cs by Smirnov and Firsov (1965). The values are of the order of $10^{-9} - 10^{-10} \text{ cm}^3 \text{ sec}^{-1}$.

EXCITATION BY NEUTRAL HYDROGEN ATOM IMPACT

The excitation of rotational levels of molecular hydrogen by the impact of hydrogen atoms is an important cooling mechanism in interstellar space. Cross sections for

have been calculated by Takayanagi and Nishimura (1960) using an approximate version of the distorted wave method. The distorted wave method has been applied without further approximation by Dalgarno, Henry, and Roberts (1966) with results in close agreement with the earlier calculations. Their cooling rates are reproduced in Table V.

Recently Allison and Dalgarno (1967) have solved the set of coupled equations that describes the collision, obtaining cross sections which are somewhat lower at high temperatures than those derived from the distorted-wave approximation. The corresponding rate coefficient is included in Table V. However, the major uncertainty in the predicted cooling rates lies in the interaction potential between H and H_2 .

Takayanagi and Nishimura (1960) and Dalgarno *et al.* (1966) have also tabulated excitation rates for other rotational transitions in H_2 and D_2 .

The excitation of H_2 by H_2

is less important. The most accurate cross sections are those computed by Allison and Dalgarno (1967).

Takayanagi and Nishimura (1960) have roughly estimated the rate coefficient for excitation of the 0-1

TABLE V. Rate coefficient $\text{H}-\text{H}_2(j-j')$ ($\text{cm}^3 \text{ sec}^{-1}$).

$T^\circ\text{K}$	0-2		1-3 (a)	2-4 (a)
	(a)	(b)		
20	3.4(-23)	4.5(-23)		
30	2.8(-19)	4.0(-19)	7.21(-25)	
40	2.8(-17)	3.5(-17)	1.3(-21)	
50	4.4(-16)	5.1(-16)	1.3(-19)	4.3(-23)
60	2.9(-15)	3.1(-15)	2.7(-18)	3.1(-21)
80	3.2(-14)	3.1(-14)	1.3(-16)	6.8(-19)
100	1.4(-13)	1.3(-13)	1.5(-15)	1.9(-17)
200	3.8(-12)	3.1(-12)	2.6(-13)	2.1(-14)
300	1.4(-11)	1.1(-11)	1.9(-12)	2.9(-13)
500	5.3(-11)	3.7(-11)	1.3(-11)	3.4(-12)
1000	2.2(-10)		7.8(-11)	3.4(-11)
2000	7.1(-10)		3.1(-10)	1.7(-10)
3000	1.4(-9)		6.4(-10)	3.7(-10)
4000	2.2(-9)		1.0(-9)	6.2(-10)
5000	3.1(-9)		1.5(-9)	9.2(-10)

^a From Dalgarno, Henry, and Roberts (1966).

^b From Allison and Dalgarno (1967).

rotational transition in CH, CN, and OH to be $3.5 \times 10^{-10} \text{ cm}^3 \text{ sec}^{-1}$, corresponding to a cross section of about 10^{-15} cm^2 .

SPIN-CHANGE PROCESSES

Purcell and Field (1956) drew attention to the role of the spin-change process

in controlling the spin population of atomic hydrogen and gave a rough estimate of its cross section. The process has been further studied by Dalgarno (1961b) and by Dalgarno and Henry (1964) using a theory which assumes the collision to be elastic and which ignores the effect of molecular rotation during the collision. The derived cross sections, averaged over a Maxwellian velocity distribution, oscillate about a value of $8 \times 10^{-15} \text{ cm}^2$ for $T < 100^\circ\text{K}$ and decrease slowly at higher temperatures.

A similar process

was advanced by Burgess, Field, and Michie (1960) as a possible cooling mechanism in interstellar space, and Dalgarno and Rudge (1964) proposed the additional mechanisms

giving a rough estimate of the cross section. Smith (1966b) has computed the corresponding cooling rates, and he concludes that the O-H reaction is more efficient than the C-H and Si-H reactions (despite the long-range attraction), and that below 300°K the O-H reaction is a more efficient cooling mechanism than excitation of the rotational levels of H_2 by hydrogen atom impact.

The theoretical description used by Smith is an extension of that presented by Dalgarno and Rudge (1965) for collision-induced changes in the hyperfine structure of alkali metal vapors, and it may not be adequate for collision-induced changes in fine structure (cf. Nikitin 1965, Callaway and Bauer 1965).

QUENCHING COLLISIONS

The quenching of radiation through the destruction process

has been studied theoretically by Nikitin and Bykhovskii (1964), but the lack of adequate wave functions for the quasi-molecule XNa has prevented quantitative predictions. For inert gas atoms, the measured cross sections are about 10^{-16} cm². Larger cross sections occur when X is a diatomic molecule (cf. Starr and Shaw 1966).

CHEMICAL REACTIONS

Herzberg (1955) has listed a number of exothermic reactions which produce molecular hydrogen:

and Stecher and Williams (1967) have listed other exothermic reactions which produce CH, OH, NH, CN, CO, and N₂. It is conventional practice to adopt the Arrhenius form for the rate coefficients

$$K = A \exp(-K/kT)$$

and to assume that the activation energy E is given by

$$A = 0.055D,$$

where D is the dissociation energy of the initial molecule (Polanyi 1962). A review of some of the reactions involving atomic hydrogen has been presented by Kaufman (1964). Recent work on OH radical reactions is described by Skinner and Ringrose (1965) (see also Carroll and Salpeter 1966).

TRANSPORT PROPERTIES

The thermal conductivity and the viscosity of atomic hydrogen have been calculated to high accuracy by Dalgarno and Smith (1962) at high temperatures and by Buckingham, Fox, and Gal (1965) at low temperatures. The diffusion coefficient of protons in atomic hydrogen gas is given in Table IV. Formulas for the transport coefficients appropriate to a fully ionized plasma have been derived by Spitzer (1956) (see Devoto 1966).

ACKNOWLEDGMENT

The preparation of this review was partly supported by the GCA Corporation, Bedford, Massachusetts.

REFERENCES

- Adler, J., and Moiseiwitsch, B. L., 1957, *Proc. Phys. Soc. (London)* **A70**, 117.
 Allison, D. C., and Dalgarno, A., 1963, *Proc. Phys. Soc. (London)* **81**, 23.
 Allison, D. C., and Dalgarno, A., 1965, *Proc. Phys. Soc. (London)* **85**, 845.
 Allison, A. C., and Dalgarno, A., 1967, *Proc. Phys. Soc. (London)* **90**, 609.
 Allison, D. C., Browne, J. C., and Dalgarno, A., 1966, *Proc. Phys. Soc. (London)* **89**, 41.
 Arthurs, A. M., and Hyslop, J., 1957, *Proc. Phys. Soc. (London)* **A70**, 849.
 Bardsley, J. N., Herzenberg, A., and Mandl, F., 1966, *Proc. Phys. Soc. (London)* **89**, 305 and 321.
 Barnett, C. F., and Reynolds, H. K., 1958, *Phys. Rev.* **109**, 355.
 Bassel, R. H., and Gerjuoy, E., 1960, *Phys. Rev.* **117**, 749.
 Basu, D., Sil, N. C., and Bhattacharya, D. M., 1966, *Phys. Rev.* **144**, 56.
 Bates, D. R., 1951, *Monthly Notices Roy. Astron. Soc.* **111**, 303.
 Bates, D. R., 1958, *Proc. Roy. Soc. (London)* **A245**, 299.
 Bates, D. R., 1959, *Proc. Roy. Soc. (London)* **A73**, 227.
 Bates, D. R., 1961, *Proc. Phys. Soc. (London)* **A77**, 59.
 Bates, D. R., 1962, *Atomic and Molecular Processes* (Academic Press Inc., New York).
 Bates, D. R., Bell, K., and Kingston, A. E., 1967, *Proc. Phys. Soc. (London)* **91**, 288.
 Bates, D. R., and Boyd, T. J. M., 1956, *Proc. Phys. Soc. (London)* **A69**, 910.
 Bates, D. R., and Crothers, D. S. F., 1967, *Proc. Phys. Soc. (London)* **90**, 73.
 Bates, D. R., and Dalgarno, A., 1952, *Proc. Phys. Soc. (London)* **A65**, 919.
 Bates, D. R., and Dalgarno, 1953, *Proc. Phys. Soc. (London)* **A66**, 972.
 Bates, D. R., and Griffing, G. W., 1953, *Proc. Phys. Soc. (London)* **A66**, 961.
 Bates, D. R., and Griffing, G. W., 1954, *Proc. Phys. Soc. (London)* **A67**, 663.
 Bates, D. R., and Griffing, G. M., 1955, *Proc. Phys. Soc. (London)* **A68**, 90.
 Bates, D. R., and Lewis, J. T., 1955, *Proc. Phys. Soc. (London)* **A68**, 173.
 Bates, D. R., and Massey, H. S. W., 1954, *Phil. Mag.* **45**, 111.
 Bates, D. R., and Williams, A., 1957, *Proc. Phys. Soc. (London)* **A70**, 117.
 Bates, D. R., and Williams, D. A., 1964, *Proc. Phys. Soc. (London)* **83**, 425.
 Bell, R. J., 1961, *Proc. Phys. Soc. (London)* **78**, 903.
 Bell, R. J., and Skinner, B. G., 1962, *Proc. Phys. Soc. (London)* **80**, 404.
 Bouthilliet, D. B., Healey, J. A., and Milford, S. N., 1964, *Atomic Collision Processes* (North-Holland Publ. Co., Amsterdam), p. 1081.
 Boyd, T. J. M., and Dalgarno, A., 1958, *Proc. Phys. Soc. (London)* **72**, 694.
 Bransden, B. H., 1966, *Advan. Atomic Mol. Phys.* **1**, 85.
 Bransden, B. H., and Cheshire, I., 1963, *Proc. Phys. Soc. (London)* **81**, 820.
 Bransden, B. H., and Sin Fai Lam, L. T., 1966, *Proc. Phys. Soc. (London)* **87**, 653.
 Buckingham, R. A., Fox, J. W., and Gal, E., 1965, *Proc. Roy. Soc. (London)* **A284**, 237.
 Burgess, A., Field, G. B., and Michie, R. W., 1960, *Astrophys. J.* **131**, 529.
 Butler, S. T., and Buckingham, M. J., 1962, *Phys. Rev.* **126**, 1.
 Butler, S. T., and Johnston, I. D. S., 1964, *Nucl. Fusion* **4**, 196.
 Callaway, J., and Bauer, E., 1965, *Phys. Rev.* **140**, A1072.
 Callaway, J., and Dugan, A. F., 1966, *Phys. Letters* **21**, 295.
 Carew, J., and Milford, S. N., 1963, *Astrophys. J.* **138**, 772.
 Carroll, T. O., and Salpeter, E. E., 1966, *Astrophys. J.* **143**, 609.
 Chamberlain, J. W., 1956, *Astrophys. J.* **124**, 390.

- Cheshire, I., 1965, *Phys. Rev.* **138**, A992.
- Coleman, J. P., and McDowell, M. R. C., 1965, *Proc. Phys. Soc. (London)* **85**, 1097.
- Coleman, J. P., and McDowell, M. R. C., 1966, *Proc. Phys. Soc. (London)* **87**, 879.
- Colli, L., Cristofori, F., Frigerio, G. E., and Sona, P. G., 1962, *Phys. Letters* **3**, 1962.
- Dalgarno, A., 1961a, *Ann. Geophys.* **17**, 16.
- Dalgarno, A., 1961b, *Proc. Roy. Soc. (London)* **A262**, 132.
- Dalgarno, A., and Griffing, G. W., 1955, *Proc. Roy. Soc. (London)* **A232**, 423.
- Dalgarno, A., and Griffing, G. W., 1958, *Proc. Roy. Soc. (London)* **A248**, 415.
- Dalgarno, A., and Henry, R. J. W., 1964, *Proc. Phys. Soc. (London)* **83**, 157.
- Dalgarno, A., Henry, R. J. W., and Roberts, C. S., 1966, *Proc. Phys. Soc. (London)* **88**, 611.
- Dalgarno, A., and McDowell, M. R. C., 1955, *The Airglow and the Aurorae* (Pergamon Press, London).
- Dalgarno, A., and Rudge, M. J., 1964, *Astrophys. J.* **140**, 800.
- Dalgarno, A., and Rudge, M. J., 1965, *Proc. Roy. Soc. (London)* **A286**, 519.
- Dalgarno, A., and Smith, F. J., 1962, *Proc. Roy. Soc. (London)* **A267**, 417.
- Dalgarno, A., and Yadav, H. N., 1953, *Proc. Phys. Soc. (London)* **A66**, 173.
- Devoto, R. S., 1966, *Phys. Fluids* **9**, 1230.
- Erkovich, C., 1960, *Opt. Spectry.* **8**, 162.
- Federenko, N. V., Afrosimov, V. V., Il'in, R. N., and Solov'ev, E. S., 1960, *Proceedings of the IVth International Conference on Ionization Phenomena in Gases* (North-Holland Publ. Co., Amsterdam), p. 47.
- Ferguson, A. F., 1961, *Proc. Roy. Soc. (London)* **A246**, 540.
- Field, G. B., and Hitchcock, J. L., 1966, *Phys. Rev. Letters* **16**, 817.
- Fite, W. L., Smith, A. C. H., and Stebbings, R. F., 1962, *Proc. Roy. Soc. (London)* **A268**, 527.
- Fite, W. L., Stebbings, R. F., Hummer, D. G., and Brackman, R. T., 1960, *Phys. Rev.* **119**, 663.
- Fulton, M. J., and Mittleman, M. H., 1965, *Ann. Phys. (N. Y.)* **33**, 65.
- Gilbody, H. B., and Ireland, J. V., 1964, *Proc. Roy. Soc. (London)* **A277**, 137.
- Green, T. A., Stanley, H. E., and Chang, Y. C., 1965, *Helv. Phys. Acta* **38**, 109.
- Hanson, W. B., Patterson, T. L. N., and Degaonkar, S. J., 1963, *J. Geophys. Res.* **68**, 6213.
- Harris, F. E., 1966, *J. Chem. Phys.* **44**, 3636.
- Hayakawa, S., and Kitao, K., 1956, *Progr. Theoret. Phys. (Kyoto)* **16**, 139.
- Herzberg, G., 1955, *Mem. Soc. Roy. Sci. Liège* (4) **15**, 291.
- Hiskes, J. R., 1965, *Phys. Rev.* **137**, A361.
- Hooper, J. W., Harmer, D. S., Martin, D. W., and McDaniel, E. W., 1962, *Phys. Rev.* **125**, 2000.
- Hughes, R. H., Dawson, H. R., Doughty, B. M., Kay, D. B., and Stiger, C. A., 1966, *Phys. Rev.* **146**, 53.
- Ingber, L., 1965, *Phys. Rev.* **139**, A35.
- Jackson, J. D., and Schiff, H. I., 1953, *Phys. Rev.* **89**, 359.
- Jaacks, D., Van Zyl, B., and Geballe, R., 1965, *Phys. Rev.* **137**, A340.
- Katsuura, K., 1965, *J. Chem. Phys.* **42**, 3771.
- Kaufman, F., 1964, *Ann. Geophys.* **20**, 106.
- Lovell, S. E., and McElroy, M. B., 1965, *Proc. Roy. Soc. (London)* **A283**, 100.
- McCarroll, R. W., 1961, *Proc. Roy. Soc. (London)* **A246**, 547.
- McCarroll, R. W., and Salin, A., 1966, *Ann. Phys. (Paris)* **1**, 283.
- McDowell, M. R. C., 1961, *Observatory* **81**, 240.
- McDowell, M. R. C., 1964, *Atomic Collision Processes* (North-Holland Publ. Co., Amsterdam).
- Malville, J. M., 1964, *Astrophys. J.* **139**, 198.
- Mapleton, R. A., 1958, *Phys. Rev.* **109**, 1166.
- Mapleton, R. A., 1961, *Phys. Rev.* **122**, 528.
- Mapleton, R. A., 1962, *Phys. Rev.* **126**, 1477.
- Mapleton, R. A., 1963, *Phys. Rev.* **130**, 1839.
- Mapleton, R. A., 1965, *Proc. Phys. Soc. (London)* **85**, 841.
- Mapleton, R. A., 1966, *Phys. Rev.* **145**, 25.
- May, R. M., 1964, *Nucl. Fusion* **4**, 207.
- May, R. M., 1965a, *Phys. Letters* **14**, 98.
- May, R. M., 1965b, *Phys. Letters* **14**, 198.
- May, R. M., and Lodge, J. G., 1965, *Phys. Rev.* **137**, A699.
- Michels, H. H., 1966, *J. Chem. Phys.* **44**, 3834.
- Mittleman, M. H., 1961, *Phys. Rev.* **122**, 499.
- Moiseiwitsch, B. L., and Stewart, A. L., 1954, *Proc. Phys. Soc. (London)* **A67**, 1069.
- Mori, M., 1966, *J. Phys. Soc. Japan* **21**, 979.
- Nikitin, E. E., 1965, *J. Chem. Phys.* **43**, 744; and *Opt. Spectry.* **19**, 91.
- Nikitin, E. E., and Bykhovskii, V. K., 1964, *Opt. Spectry.* **17**, 444.
- Nikolaev, V. S., 1965, *Soviet Phys.—Usp.* **8**, 269.
- Parcell, L., and Butler, S. T., 1965, *Phys. Letters* **14**, 110.
- Peach, G., 1965, *Proc. Phys. Soc. (London)* **85**, 709.
- Peek, J. M., 1966, *Phys. Rev.* **143**, 33.
- Polanyi, J. C., 1962, *Atomic and Molecular Processes* (Academic Press Inc., New York), p. 807.
- Pomilla, F. R., and Milford, S. N., 1966, *Astrophys. J.* **144**, 1174.
- Pretzer, D., Van Zyl, B., and Geballe, R., 1964, *Atomic Collision Processes* (Academic Press Inc., New York), p. 618.
- Purcell, E. M., and Field, G. B., 1956, *Astrophys. J.* **124**, 542.
- Rapp, D., 1963, *J. Geophys. Res.* **68**, 1773.
- Seaton, M. J., 1955, *Proc. Phys. Soc. (London)* **A68**, 457.
- Seaton, M. J., 1962, *Atomic and Molecular Processes* (Academic Press Inc., New York), p. 374.
- Sheldon, J. W., 1966, *J. Appl. Phys.* **37**, 2928.
- Skinner, G. B., and Ringrose, G. H., 1965, *J. Chem. Phys.* **42**, 2190.
- Smirnov, B. M., and Firsov, O. B., 1965, *JETP Letters* **2**, 297.
- Smith, F. J., 1963, *Planet. Space Sci.* **11**, 126.
- Smith, F. J., 1966a, *Planet. Space Sci.* **14**, 929.
- Smith, F. J., 1966b, *Planet. Space Sci.* **14**, 937.
- Solomon, P. M., 1964, *Astrophys. J.* **139**, 999.
- Soshnikov, V. N., 1964, *Opt. Spectry.* **17**, 186.
- Spitzer, L., 1956, *Physics of Fully Ionized Gases* (Interscience Publishers, Inc., New York).
- Starr, W. L., and Shaw, T. M., 1966, *J. Chem. Phys.* **44**, 4181.
- Stebbing, R. F., Young, R. A., Oxley, C. L., and Ehrhardt, H., 1965, *Phys. Rev.* **138**, A1312.
- Stecher, T. P., and Williams, D. A., 1967, *Astrophys. J.* **146**, 88.
- Stier, P. M., and Barnett, C. F., 1956, *Phys. Rev.* **103**, 896.
- Takayanagi, K., and Nishimura, S., 1960, *Publ. Astron. Soc. Japan* **12**, 77.
- Thaddeus, P., and Clauser, J. F., 1966, *Phys. Rev. Letters* **16**, 819.
- Wilets, L., and Gallaher, D. F., 1966, *Phys. Rev.* **147**, 13.