C2absorption-line diagnostics of diffuse interstellar clouds

Citation

Cecchi-Pestellini, Cesare, and Alexander Dalgarno. 2002. "C2absorption-Line Diagnostics of Diffuse Interstellar Clouds." *Monthly Notices of the Royal Astronomical Society* 331 (3) (April): L31–L34. doi:10.1046/j.1365-8711.2002.05418.x.

Published Version

doi:10.1046/j.1365-8711.2002.05418.x

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:30403734>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

C₂ absorption-line diagnostics of diffuse interstellar clouds

Cesare Cecchi-Pestellini^{1,2★} and Alexander Dalgarno¹

¹*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA*

²*Dipartimento di Astronomia e Scienza dello Spazio, Università di Firenze, Largo E. Fermi 2, 50125 Firenze, Italy*

Accepted 2002 February 11. Received 2002 February 11; in original form 2002 January 14

ABSTRACT

The excitation of the diatomic carbon molecule in diffuse interstellar clouds is discussed for a cloud with large density fluctuations of small linear filling factor along the line of sight. Possible implications for the chemistry of the diffuse cloud in front of Cygnus OB2 No. 12 are discussed.

Key words: molecular processes – ISM: clouds – ISM: individual: Cygnus OB2 No. 12 – ISM: molecules.

1 INTRODUCTION

C₂ radicals play an important role in the study of the structure, properties and evolution of diffuse interstellar clouds. The homonuclear molecule C₂ was originally thought to be an effective optical probe of kinetic temperatures in interstellar clouds (Souza & Lutz 1977). However, radiative pumping effects, similar to those affecting rotational populations in H₂, have been shown to be competitive with the collisional processes (van Dishoeck & Black 1982), reducing the value of C₂ as a direct probe of kinetic temperature but providing a new source of information on the radiation field and molecular processes that determine rotational excitation.

Several absorption lines of C₂ arising from different rotational levels belonging to the ground electronic and vibrational state X¹Σ_g⁺, *v* = 0, mainly in the Phillips and Mulliken systems, have been observed in interstellar clouds. Interstellar C₂ was discovered towards ζ Oph by Chaffee & Lutz (1978) who detected one line of the 2–0 band of the Phillips system at 10 148 Å. Snow (1978) reported a detection of a line of the 0–0 Mulliken system at 2313 Å in a *Copernicus* ultraviolet spectrum of the same object. The 0–0 band has been studied in detail by Lambert, Sheffer & Federman (1995), using a high-resolution spectrum of ζ Oph from the *Hubble Space Telescope*. They also detected the weaker F¹Π_u–X¹Σ_g⁺ 0–0 band at 1342 Å. The 3–0 Phillips band at 7720 Å, weaker by a factor of 2 than the 2–0 band, was discovered by van Dishoeck & Black (1986).

The C₂ molecule rotational ladder, in sharp contrast to those of heteronuclear molecules like CN or CO, has a significant population of high-*J* levels. The high-*J* level populations are extremely sensitive to the ambient gas density, and if accurate column densities for levels *J* > 4 are available, the gas density can be inferred. In the uniform-density model of van Dishoeck & Black (1982), the distribution of the rotational levels is determined by the ratio $n_c\sigma_0/I$, where n_c is the density of the collision partners, σ_0 is

the cross-section for *J* = 2 → 0 quenching, and *I* is the incident interstellar radiation field.

The observed rotational level distribution does not, however, exclude the presence of high-density gas as in the model of the cloud in front of Cyg OB2 No. 12 (Cecchi-Pestellini & Dalgarno 2000). At the temperatures characteristic of interstellar clouds, an increase in density reduces significantly the populations of high rotational levels as the collisions drive them towards thermal equilibrium. Then if most of the column density of C₂ in the low-lying *J* levels along the line of sight resides in the diffuse gas, the high-density component may be present but undetectable. The C₂ rotational distribution arising in a mixed-phase medium inferred by absorption measurements may mimic the response from a diffuse homogeneous gas at some *average* density and kinetic temperature.

In this Letter, we investigate the extent of the C₂ rotational ladder perturbation arising from the presence of high-density regions randomly mixed with a more tenuous background medium. In Section 2 we briefly survey the molecular processes that govern the excitation of C₂, while in Section 3 we present models of the equilibrium of C₂ in diffuse interstellar clouds with embedded dense clumps. The results of the modelling procedure are applied to the line of sight towards Cyg OB2 No. 12 (Section 4), where Gredel, Black & Yan (2001) observed a large number of C₂ rotational lines. Recently, along this apparently diffuse line of sight, Geballe et al. (1999) detected the molecular ion H₃⁺ with an abundance comparable to that found in dense clouds (Geballe & Oka 1996). This detection has raised questions about the processes responsible for the production of such large amounts of H₃⁺ in the diffuse gas, and about what physical conditions prevail in the medium (Geballe et al. 1999; Cecchi-Pestellini & Dalgarno 2000; Gredel et al. 2001).

2 THE C₂ MOLECULE

In interstellar clouds, C₂ molecules exist mainly in the X¹Σ_g⁺ state. The production of vibrationally and rotationally excited C₂ is

★E-mail: cecchi-pestellini@cfa.harvard.edu

initiated by absorption of diffuse starlight in the allowed electronic transitions to the singlet $A^1\Pi_u$ (Phillips system), $D^1\Sigma_u^+$ (Milliken system) and $F^1\Pi_u$ states. These absorptions are followed by fluorescence to the electronic ground state. Collisions with the ambient gas species will contribute to establish a steady-state distribution of level populations. During the fluorescence cascade any change Δv in the vibrational quantum number is possible so that the molecule is left in an excited vibrational level in the $X^1\Sigma_g^+$ state. Because of the homonuclear nature of the molecule, no electric dipole transitions among vibrational or rotational levels are allowed within an electronic state. The main de-excitation processes are singlet–triplet (intercombination) transitions to levels of the $a^3\Pi$ state, followed by transitions back to the $X^1\Sigma_g^+$ state. These intercombination transitions cool down the rotational and vibrational Boltzmann temperatures of the C_2 radicals. This process dominates direct quadrupole transitions when $v \neq 0$ (Le Bourlot & Roueff 1986).

We solve the balance equations by means of the cascade efficiency formalism (Black & Dalgarno 1976) in the form given by Le Bourlot, Roueff & Viala (1987). The final system for the fractional population of the first 10 rotational states ($J_{\max} = 18$) of the ground vibrational state, f_J , reads

$$f_J \left(\sum_{J' \neq J} C_{J,J'} n_{H_2} + A_{J,J-2} + \sum_{k \in A,D} \zeta_{J,k} \right) = \sum_{J' \neq J} f_{J'} C_{J',J} n_{H_2} + f_{J+2} A_{J+2,J} + \sum_{J' \neq J} f_{J'} \sum_{k \in A,D} \zeta_{J',k} D_{k,J}, \quad (1)$$

where $C_{J,J'}$ are the (de)excitation rates for H_2 collisions, $A_{J,J-2}$ the (quadrupole) rotational transition probabilities (van Dishoeck & Black 1982), and $\zeta_{J,k}$ the total radiative pumping out of the state J to the vibro-rotational levels of the excited electronic states $A^1\Pi_u$ and $D^1\Sigma_u^+$, A and D respectively,

$$\zeta_{J,k} = \frac{\pi e^2}{mc} f_{J,k} I(\nu_{k,J}) \mathcal{X}. \quad (2)$$

In this equation $I(\nu_{k,J})$ is the interstellar radiation field at the transition frequency, $\mathcal{X}$ is the radiation field scaling factor, and $f_{J,k}$

is the oscillator strength

$$f_{J,k} = 1.499 \frac{2k+1}{2J+1} \left(\frac{\nu_{k,J}}{cm^{-1}} \right)^2 A_{k,J}, \quad (3)$$

where $A_{k,J}$ is the radiative transition probability. $D_{k,J}$ is the probability that a cascade from $A^1\Pi_u$ or $D^1\Sigma_u^+$ leads to $X^1\Sigma_g^+(0,J)$ via all possible paths. Its definition is given by Le Bourlot et al. (1987). The fractional level populations are subject to the normalization condition $\sum_i f_i = 1$.

References to the molecular constants are given by van Dishoeck & Black (1982), Le Bourlot et al. (1987) and Rousselot et al. (2000).

3 EQUILIBRIUM OF C_2

As discussed by van Dishoeck & Black (1982), the observed C_2 rotational population distribution provides information on the temperature and density along the line of sight. The gas kinetic temperature, T_k , is usually determined from the rotational excitation temperature, T_{ex} , of the lowest rotational levels, because the population density in these levels is not significantly affected by radiative decays. Rotational diagrams are constructed with the level populations normalized to the population of the level $J = 2$, forcing agreement for its column density, N_2 (cf. van Dishoeck & Black 1982).

If dense gas is present, the measured column density of C_2 in a rotational level J is the superposition of the contributions from both the diffuse and dense phases, $(N_J)_{\text{obs}} = N_{C_2} f_J + N_{C_2}^* f_J^* = N_J (1 + \eta f_J^* / f_J)$, where the superscript * indicates quantities in local thermal equilibrium (LTE) and $\eta = N_{C_2}^* / N_{C_2}$ is the ratio between the total C_2 column densities of the dense and diffuse phases. The relative populations as ratios of column densities are then given by

$$\left(\frac{N_J}{N_2} \right)_{\text{obs}} = \left(\frac{N_J}{N_2} \right) \frac{1 + \eta f_J^* / f_J}{1 + \eta f_2^* / f_2}. \quad (4)$$

When $\eta = 0$ we recover the ratio for homogeneous diffuse clouds.

We have computed the relative populations of the C_2 rotational ladder (equation 4) as functions of the excitation energy $\Delta E(J)$ of level J for several assumptions about the gas physical state. In Fig. 1 we show the results for two kinetic temperatures, $T_k = 20$ and

Figure 1. Relative rotational populations as functions of excitation energy $\Delta E(J)$. The distributions are shown for gas densities of 100, 500 and 1000 cm^{-3} . Solid lines: $\eta = 0$; dotted lines: $\eta = 0.1$; dashed lines: $\eta = 0.5$. (a) $T_k = 20 \text{ K}$; (b) $T_k = 45 \text{ K}$.

Figure 2. Relative rotational populations as functions of excitation energy $\Delta E(J)$ at $T_k = 50$ K. Dotted line: $n_H = 150 \text{ cm}^{-3}$; solid line: $n_H = 200 \text{ cm}^{-3}$; dashed line: $n_H = 400 \text{ cm}^{-3}$; dot-dashed line: $n_H = 600 \text{ cm}^{-3}$. Points represent mixed-phase models – triangles: $n_H = 90 \text{ cm}^{-3}$, $\eta = 0.1$; diamonds: $n_H = 150 \text{ cm}^{-3}$, $\eta = 0.1$; squares: $n_H = 320 \text{ cm}^{-3}$, $\eta = 0.15$; circles: $n_H = 260 \text{ cm}^{-3}$, $\eta = 0.3$; asterisks: $n_H = 400 \text{ cm}^{-3}$, $\eta = 0.5$; pluses: $n_H = 560 \text{ cm}^{-3}$, $\eta = 0.1$.

45 K, and three different gas densities, $n_H = 100, 500$ and 1000 cm^{-3} . The assumed interstellar radiation field is given by van Dishoeck & Black (1982). We set $\mathcal{X} = 1$. The fractions of gas in LTE that we consider are $\eta = 0, 0.1$ and 0.5 . As expected, the changes introduced in the rotational distribution by the presence of packets of dense gas along the line of sight increase with decreasing density of the diffuse component. At low densities, $n_H \leq 100 \text{ cm}^{-3}$, and temperatures, $T_k \leq 20$ K, intermediate rotational levels ($4 \leq J \leq 8$) are considerably underpopulated with respect to LTE. Thus the excitation temperatures reflect the presence of very small fractions of dense gas, as low as a few per cent along the line of sight. For larger densities of the dominant diffuse component, C_2 rotational levels are insensitive to density fluctuations of relatively small linear filling factor.

The average excitation temperatures derived from homogeneous models can be obtained from mixed-phase descriptions with suitable kinetic temperatures and densities of the two phases. The rotational diagrams corresponding to constant low-density models are well reproduced by two-phase models with a range of linear filling factors for the dense component and densities of the diffuse background (Fig. 2). Departures from the relative population ratios derived from the standard homogeneous model are very small and well inside the characteristic observational uncertainties. In all models the fraction of dense gas exceeds $\eta = 0.1$. Thus C_2 absorption-line diagnostics appear unable to detect the presence of small-scale density fluctuations, unless the gas along the line of sight has a low density $n_H \leq 100 \text{ cm}^{-3}$ and a low temperature $T_k \leq 20$ K.

4 THE CASE OF CYGNUS OB2 NO. 12

The recent discovery of H_3^+ towards the heavily reddened star Cygnus OB2 No. 12 (Geballe et al. 1999) has renewed interest in the diffuse cloud in front of the star. Geballe et al. (1999) constructed a model of the line of sight towards Cyg OB2 No. 12 with a hydrogen density of 10 cm^{-3} that, as they noted, led to an excessively long path length and did not explain the observations of

Figure 3. C_2 excitation diagram of Cyg OB2 No. 12. The three solid lines drawn in the diagram represent model population distributions at $T_k = 35$ K and $n_H = 400, 600$ and 800 cm^{-3} . Points correspond to individual line detections in the (2–0) band (triangles: R branch; diamonds: Q branch; squares: P branch). Dotted line: $n_H = 540 \text{ cm}^{-3}$, $\eta = 0.1$; dashed line: $n_H = 440 \text{ cm}^{-3}$, $\eta = 0.2$; dot-dashed line: $n_H = 300 \text{ cm}^{-3}$, $\eta = 0.4$; long-dashed line: $n_H = 540 \text{ cm}^{-3}$, $\eta = 0.1$, $T_k = 15$ K in the dense phase.

$\text{CO}(2-1)$ emission (Geballe et al. 1999) and C_2 absorption (Gredel & Münch 1994). The low-density model of Geballe et al. (1999) is consistent with the *ISO* observations that have shown the presence of the $3.4\text{-}\mu\text{m}$ hydrocarbon feature, and the absence of the 3.0- and $4.27\text{-}\mu\text{m}$ features of H_2O and CO_2 ices, respectively (Whittet et al. 1997). The model was questioned by Cecchi-Pestellini & Dalgarno (2000) because of the assumption that the fractional abundance of H_2 was 0.25, and they presented a model of the material in front of the star Cyg OB2 No. 12 in which dense cold cores are proposed to exist embedded in diffuse clumps of gas. The model is consistent with *ISO* data and the absence of ice features. It predicts the presence of HCO^+ in a detectable amount. The recent detection of $\text{HCO}^+(1-0)$ emission along this line of sight by Scappini et al. (2000) provides support for this scenario. Yet another model was put forward by Gredel et al. (2001) in which the molecular abundances are a response to an X-ray ionization rate $\zeta \approx 10^{-15} \text{ s}^{-1}$.

To constrain the physical conditions of density and temperature of the cloud molecular material in front of Cyg OB2 No. 12, Gredel et al. (2001) performed a detailed analysis of C_2 interstellar absorption lines up to $J = 18$ in the (2,0) and (3,0) bands of the Phillips system. The large number of rotational lines constrains the gas kinetic temperature and the density to $T_k = 35$ K and $n_H = 600 \pm 100 \text{ cm}^{-3}$. These conditions do not exclude the clumpy model of Cecchi-Pestellini & Dalgarno (2000). In Fig. 3 we show some theoretical population distributions computed for a temperature of $T_k = 35$ K using both homogeneous and mixed-phase models. Only the (2–0) band rotational transitions are shown. We did not perform any averages for individual rotational levels. The spread in the populations arising from different branches may be taken as a rough indication of the measurement errors. Assuming the distribution profile derived from a homogeneous cloud at $n_H = 600 \text{ cm}^{-3}$ as the best fit to the observations (Gredel et al. 2001), we find that the mixed-phase model with $n_H = 540 \text{ cm}^{-3}$ and $\eta = 0.1$ is in excellent agreement with the data. More *extreme* models with $\eta = 0.2$ and 0.4 fit the observations moderately well. Since temperature variations are

likely to follow density fluctuations, we have also computed the level distribution profile assuming a lower kinetic temperature, $T_k = 15$ K, in the dense phase. We find no significant differences.

These results imply that the observed C_2 rotational distribution is not inconsistent with the presence of high-density gas of linear filling factor $\eta \leq 0.1$ along the line of sight. The above upper limit is derived assuming a similar C_2 fractional abundance in both diffuse and dense phases, and it could be larger because, according to Gredel & Münch (1994), chemical models predict a sudden decrease in C_2 abundance at densities above $10^{3.5} \text{ cm}^{-3}$. Since the total hydrogen column density along the line of sight is $N_H = 2 \times 10^{22} \text{ cm}^{-2}$ (Geballe et al. 1999), the fraction of thermal gas yields $N_H^* \leq 2 \times 10^{21} \text{ cm}^{-2}$. Assuming in the high-density regime an almost total conversion of C to CO, and adopting a carbon cosmic abundance of 1×10^{-4} (Sofia, Fitzpatrick & Meyer 1998), we obtain an upper limit for the CO column density produced in the dense phase of $N_{CO} = 2 \times 10^{17} \text{ cm}^{-2}$, consistent with the observed value of $3 \times 10^{16} \text{ cm}^{-2}$ (Geballe et al. 1999). Indeed, observations of the CO (2–1) emission by Scappini et al. (2000) imply a gas density $n_H \geq 10^4 \text{ cm}^{-3}$.

5 CONCLUSION

We have shown that C_2 absorption observations are not sensitive to the presence of dense gas of small linear filling factor along the line of sight. This result has implications for our understanding of the elusive chemistry of diffuse interstellar clouds. Millimetre-wave absorption spectra of molecules like CO, CS, SO, CN, HCN, HNC, H_2S , C_2H and H_2CO (Lucas & Liszt 1999) suggest large concentrations of these species in environments that are poorly shielded from the ambient ultraviolet radiation. In conventional chemical evolution models the physical characteristics of interstellar diffuse clouds are assumed to vary slowly on subparsec scales. While the clumpy structure of dense molecular clouds is widely accepted, the diffuse interstellar medium is usually considered a more homogeneous environment. The incorporation of density fluctuations, length-scale and filling factor in theoretical models could be crucial for the interpretation of diffuse cloud chemistry.

The chemistry of the cloud in front of Cyg OB2 No. 12 may be explained by a model in which dense cores are embedded in diffuse clumps of gas (Cecchi-Pestellini & Dalgarno 2000). We have

shown that C_2 absorption measurements are consistent with this scenario. Since the H_3^+ molecular ion has now been definitely detected in at least four diffuse lines of sight (Geballe 2000), the chemistry of the gas is due not to the peculiarities of a particular region, but to a general feature of the diffuse interstellar medium.

Excitation analyses of other molecules in the framework of the mixed-phase model would be instructive.

ACKNOWLEDGMENTS

This research has been supported by the Division of Astronomy of the US National Science Foundation. CC-P thanks the Italian MURST.

REFERENCES

- Black J. H., Dalgarno A., 1976, *ApJ*, 203, 132
- Cecchi-Pestellini C., Dalgarno A., 2000, *MNRAS*, 317, L6
- Chaffee F. H., Lutz B. L., 1978, *ApJ*, 221, L91
- Geballe T. R., 2000, *Phil. Trans. R. Soc. A*, 358, 2503
- Geballe T. R., Oka T., 1996, *Nat*, 384, 334
- Geballe T. R., McCall B. J., Hinkle K. H., Oka T., 1999, *ApJ*, 510, 251
- Gredel R., Münch G., 1994, *A&A*, 285, 640
- Gredel R., Black J. H., Yan M., 2001, *A&A*, 375, 553
- Lambert D. L., Sheffer Y., Federman S. R., 1995, *ApJ*, 438, 740
- Le Boulart J., Roueff E., 1986, *J. Mol. Spectrosc.*, 120, 157
- Le Boulart J., Roueff E., Viala Y., 1987, *A&A*, 188, 137
- Lucas R., Liszt H. S., 1999, in Ossenkopf V., Stutzki J., Winnewisser G., eds, *The Physics and the Chemistry of the Interstellar Medium*. GCA-Verlag, Herdecke, p. 112
- Rousselot P., Hill S. M., Burger M. H., Brain D. A., Laffont C., Moreels G., 2000, *Icarus*, 146, 263
- Scappini F., Cecchi-Pestellini C., Codella C., Dalgarno A., 2000, *MNRAS*, 317, L6
- Snow T. P., 1978, *ApJ*, 220, L93
- Sofia U. J., Fitzpatrick E., Meyer D. M., 1998, *ApJ*, 499, 951
- Souza S. P., Lutz B. L., 1977, *ApJ*, 216, L49
- van Dishoeck E. F., Black J. H., 1982, *ApJ*, 258, 533
- van Dishoeck E. F., Black J. H., 1986, *ApJ*, 307, 332
- Whittet D. C. B. et al., 1997, *ApJ*, 490, 729

This paper has been typeset from a $\text{\LaTeX}$ file prepared by the author.