PBDEs (polybrominated diphenyl ethers) pose a risk to captive giant pandas

Citation

Chen, Yi-ping, Ying-juan Zheng, Qiang Liu, Aaron M. Ellison, Yan Zhao, and Qing-yi Ma. 2017. "PBDEs (polybrominated Diphenyl Ethers) Pose a Risk to Captive Giant Pandas." *Environmental Pollution* 226 (July): 174–181. doi:10.1016/j.envpol.2017.04.023.

Published Version

10.1016/j.envpol.2017.04.023

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:32686919>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

PBDEs (polybrominated diphenyl ethers) pose a risk to captive giant pandas

Yi-ping Chen^{a, b*}, Ying-juan Zheng^a, Qiang Liu^a, Aaron M. Ellison^c, Yan Zhao^a, Qing-yi Ma^d

^a State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, CAS, Xi'an, 710075, China

^b College of Life Science, Northwest Normal University, Lanzhou 730070, China

^c Harvard University, Harvard Forest, Petersham, Massachusetts, 01368, USA

^d Shaanxi Wild Animal Research Center, Zhouzhi, Xi'an, 710402, China

*Corresponding author: Dr YP Chen, Institute of Earth Environment, Chinese Academy of Science, No. 97,

Yan-Xiang Road, Xi'an, 710061, China; Tel: +86-29-62336269; Fax: +86-29 -62336269; E-mail:

chenyp@ieecas.cn

Abstract

The Qinling subspecies of giant panda (*Ailuropoda melanoleuca qinlingensis*) is highly endangered; fewer than 350 individuals still inhabit the Qinling Mountains. Previous research revealed that captive pandas were exposed to bromine, so we hypothesized that captive pandas also were exposed to, and affected by, polybrominated diphenyl ethers (PBDEs). To test this hypothesis, we sampled blood and feces of captive and wild pandas, their drinking water, food (bamboo leaves) from the Shaanxi Wild Animal Research Center (SWARC) and the Foping National Nature Reserve (FNNR), and supplemental feedstuff fed to captive pandas at SWARC. We found 13 congeners of PBDEs in fecal samples, of which BDE47, BDE66, BDE71, BDE99, and BDE154 predominated; total PBDE concentration in feces of captive pandas was 255%

higher than in wild pandas. We found nine PBDEs congeners in blood samples: BDE153 and BDE183 predominated, and concentrations of PBDEs in blood from captive pandas also were significantly higher than in wild pandas. The primary source of PBDEs appears to be the bamboo fed to the pandas: total concentration of PBDEs were 5473 and 4835 $\text{pg}\cdot\text{g}^{-1}$ in the bamboo *Fargesia qinlingensis*; 2192 and 1414 $\text{pg}\cdot\text{g}^{-1}$ in the bamboo *Bashannia fargesii*; 0.066, 0.038 $\text{pg}\cdot\text{mL}^{-1}$ in drinking water; and 28.8 $\text{pg}\cdot\text{g}^{-1}$ in supplemental feedstuff for captive and wild pandas, respectively. BDE99 and BDE47 could threaten the health of captive pandas, whereas other PBDE congeners may pose additional health risks to the captive pandas. In the short term, this risk may be ameliorated by strict control of food quality. In the long term, however, reducing air, water and soil contamination to improve environmental quality will best reduce these risks.

Keywords: PBDEs; Captive Panda; Feces and Blood; Food; Health risk

Capsule: Captive pandas were exposed to higher concentrations of toxic PBDEs than wild pandas. PBDEs are most prevalent in the bamboo fed to the pandas, highlighting the need for quality control on the food supply of captive pandas.

Introduction

The giant panda (*Ailuropoda melanoleuca*) is one of the rarest animals in the world. Approximately 1800 individuals remain in anthropogenically fragmented habitats (SFA, 2015), of which < 350 individuals are of the Qinling subspecies (*A. melanoleuca qinlingensis*) living in the

Qinling Mountains of China (SFA, 2015). In the last several decades, two strategies have been used to protect this species. One strategy is *ex-situ* breeding in, for example, the Beijing Zoo, the Wolong Breeding Center, and the Shaanxi Wild Animal Research Center (SWARC). The other strategy is the establishment of natural conservation zones to preserve panda habitat. In the last several decades, 67 conservation zones, with a total area > 43,600 km², have been established (SFA, 2015).

It is generally assumed that captive breeding centers can effectively protect giant pandas from the adverse impacts of human activities. However, canine distemper virus has killed at least four pandas at SWARC (Mara, 2015) suggesting that new measures are needed to protect captive individuals of this iconic endangered species. Environmental pollution further stresses captive animals. For example, we have shown that captive pandas are exposed to heavy metals including cadmium, zinc, chromium, arsenic and lead (Chen et al., 2016). We also found that chlorine and bromine were 690% and 330% higher in feces of captive pandas than in those of wild pandas (Chen and Ma, 2017), and we therefore hypothesized that captive pandas may be exposed to and affected by polybrominated diphenyl ethers (PBDEs).

PBDEs are brominated flame retardants that are used in electronic equipment, textiles, cabinets for television and computers, and in many plastic products (WHO, 1994; Darnerud et al., 2001; Kim et al., 2012). PBDEs are lipophilic, are released slowly into the environment, can bioaccumulate in tissues of humans and other mammals, and are toxic to them (Hooper and McDonald, 2000; De Wit, 2002; Hu et al., 2008). Exposure of laboratory animals to high concentrations of PBDEs can suppress production of antibodies and proliferation of lymphocytes

(Darnerud and Thuvander, 1998), decrease thymic weights (Fowles et al., 1994), cause immunomodulatory turbulence, and lead to hormonal deficits (Eriksson et al., 2001; Branchi et al., 2003). Modulating effects of PBDE exposure on endocrine systems of wild animal also have been documented (Legler and Brouwer, 2003; Darnerud, 2003). Recent research showed that giant pandas were exposed to PCDDs, PCDFs, PCBs, and heavy metals from the bamboo they eat (*Fargesia qinlingensis* and *Bashania fargesii*) in both captive breeding centers and *in situ* in conservation areas (Chen et al., 2016). However, there has been no research on exposure of captive or wild pandas to PBDEs, or their concentrations in panda feces and blood.

The objective of this study was to (1) test whether captive or wild pandas are exposed to PBDEs. (2) document and compare the concentrations of PBDEs in wild and captive pandas; and (3) identify possible sources of PBDEs contamination. Feces, drinking water, and food (bamboo) were collected from SWARC and the Foping National Nature Reserve (FNNR), and blood samples and supplemental feedstuff were sampled at SWARC (Fig.1).

Figure 1. Sample Collection Sites. The Shaanxi Wild Animal Research Center (SWARC) is located at $34^{\circ} 04' N$, $108^{\circ} 19' E$ in Zhouzhi County, Shaanxi province. The Foping National Nature Reserve (green shaded area) is located in the area bounded by $33^{\circ} 33' - 33^{\circ} 44' N$, $107^{\circ} 40' - 107^{\circ} 55' E$ within the Qinling Mountains (blue shaded area).

79 Materials and methods

80 Sample collection

81 Giant pandas are protected in China, capturing them is illegal, and so samples from wild
 82 animals must be collected non-invasively. Fecal samples (droppings: Fig. 1) were collected from
 83 16 different locations within FNNR. Sampling locations were spaced 10-km apart and every four
 84 independent samples were pooled into a mixed sample. Droppings of 16 captive pandas were

collected from SWARC, which was established in 1987 to conserve the Qinling panda. Each set of 16 fecal samples were pooled for analysis into four samples each consisting of four independent samples.

Fresh leaves of living plants (500 g) of the two bamboo species (*Fargesia qinlingensis*, *Bashania fargesii*) that are the primary food of the panda were collected in proximity to where the droppings were collected at FNNR and around SWARC. Water samples (500 mL) were collected into Pyrex borosilicate amber glass bottles from streams at FNNR near where we collected the droppings, and from the SWARC water supply. At both FNNR and SWARC, 12 samples of each bamboo species and of freshwater were collected. They subsequently were pooled to produce four mixed samples each consisting of three independent samples from each site. In addition, four samples of mixed feedstuff, provided as a nutrient supplement for captive pandas, also were collected from SWARC.

Finally, blood samples were obtained from three similarly-aged pandas rescued from the Qinling Mountains and three captive pandas bred at SWARC. These blood samples were residuals from regular, routine physical examinations of the individual pandas. Prior to examination, the pandas were anesthetized with 25% ketamine (dosage: 8 mg·kg⁻¹). After collection, the blood was placed in EDTA tubes and frozen at -80 °C for analysis of PBDEs.

Sample preparation and extraction

PBDE congeners were analyzed using US Environmental Protection Agency (EPA) method 1614 with minor modifications (Li et al., 2008). Bamboo, feces, and feedstuff samples

were freeze dried and then homogenized by passing them through a stainless steel sieve (0.5-mm mesh). Each 3-g homogenized sample was spiked with a ^{13}C -labeled surrogate standard (EPA methods 1613B and 1668A) and extracted using accelerated solvent extraction (ASE) for 24 h with dichloromethane (150 mL) and hexane (150 mL) at 55 °C. After ASE, acidic silica (15 g, 30% w/w) was added to the sample to remove lipids. Then, 5 g of anhydrous sodium sulfate was added to the extract. The extract sample was rotary evaporated to 2 ml and then passed through a multi-layered silica-gel column that had been pre-cleaned by hexane (100 mL). After the sample was loaded, the PBDE congeners were eluted with 70 ml hexane followed by 70 mL dichloromethane. The eluant was then concentrated to 2 ml on the rotary evaporator. Its volume was further reduced with a gentle nitrogen flow and the solvent was changed to 20 μL nonane in a minivial.

PBDEs in water samples were extracted using US EPA method 1614. Prior to extraction, 1 L of the liquid samples were filtered using 0.45- μm microporous membranes to remove the particle phase and then spiked with a ^{13}C -labeled BDE-LCS standard. Organic halogen pollutants were adsorbed by siphon, 50 mL of acetone was used to flush the pillars and a 100-mL mixture of acetone and water was collected. Resins were extracted using ASE for 24 h with dichloromethane (300 mL) and hexane (300 mL) at 55 °C. After ASE, a 15-mL acetone-cleaned soxhlet extractor and zeolite was used to obtain a 430-mL solution, which was transferred to a 1000-mL separatory funnel, washed with 100 mL of ultrapure water for 3 times, and extracted with 30-mL n-hexane for 3 times to obtain a clear and transparent organic phase solution. This organic phase solution was evaporated to 2–3 mL and purified on the silica gel column. After

adding 50 ml n-hexane and evaporated to 2–3 mL, we added 15 mL isooctane, evaporated to 2–3 mL and used 6-mL isooctane to clean. The solution was purged by nitrogen stream and diluted to 1 mL in a brown bottle at –20 °C for further analysis.

Instrumental analysis

BDEs 17, 28, 47, 66, 71, 85, 99, 100, 138, 153, 154, 183, and 190 were analyzed using gas chromatography (Agilent 6890, USA) coupled with a high-resolution mass spectrometer (HRMS). The HRMS (Waters Micromass, Manchester, UK) operated in selected ion monitoring (SIM) mode with resolution >10,000. The m/z ratio of all the PBDE congeners was 79 and 81 except for BDE47 (m/z 325 and 327), BDE99 (m/z 404 and 406) and BDE183 (m/z 562 and 564). Exactly 1 µL of the sample was injected with a CTC PAL autosampler in splitless mode into an HB-5 (30 × 250-µm i.d. × 0.1-µm film thickness) capillary column for separation. The flow rate of the carrier gas was 1.2 mL/min and the carrier gas was Helium. The program was as follows: the injector was temperature programmed to ramp from 60 °C to 320 °C at 150 °C/min. The oven started at 80 °C held for 1 min, increased to 200 °C at 10 °C/min, held at 200 °C for 1 min, increased to 300 °C at 20 °C/min, and then held at 300 °C for 5 min. The temperature of the ion source was 150 °C.

Quality assurance and quality control

All solvents were pesticide-residue grade and were purchased from Fisher (Hampton, NH, USA). Silica gel was obtained from Merck (silica gel 60, Darmstadt, Germany). ¹³C labeled

surrogate and labeled injection standards were purchased from Wellington Laboratories (Guelph, Canada).

All analytical procedures were checked by the strict quality assurance and control measures to avoid sample- and cross-contamination. Reference material and 3 blank control samples (ultrapure water) were analyzed using the same methods as described above. Triplicate samples were analyzed to determine repeatability and reproducibility. To monitor analytic losses, all samples were spiked with internal standards of ^{13}C -labeled BDE47, 99, and 153. The mean recoveries of ^{13}C -labeled surrogate PBDE congeners 47, 99 and 153 were in the range of $54.2 \pm 12.1\%$, $66.0 \pm 10.1\%$, $102.2 \pm 20.1\%$, respectively, which were well in the limits according to US EPA Method 1614; all the content of PBDEs in the control (blank) samples were below the limit of detection (LOD), which for BDEs 17, 28, 47, 66, 71, 85, 99, 100, 138, 153, 154, 183, and 190 are 5, 4, 3, 15, 6, 7, 7, 7, 4, 1, 12, and 4 $\text{pg}\cdot\text{g}^{-1}$, respectively). If concentrations of actual samples fell below LOD, 1/2 of the LOD values were used in subsequent statistical analysis.

Data analysis

Correlation analysis (CA) and principal components analysis (PCA) were used to analyze the association between 13 PBDE congeners in different samples. Paired samples were analyzed using *t*-tests. All statistical analyses were done using the IBM statistical package SPSS 20.0 (IBM Corp., USA).

Evaluation methods

The giant panda's health risk evaluation was calculated using following equations detailed in the Exposure Factors Handbook (US EPA 1997). Assuming that giant pandas only feed on bamboo leaves, average daily dose (ADD) was calculated as:

$$ADD = \frac{C \times IR_s \times EF \times ED}{BW \times AT}$$

where C is concentration of PBDEs (mg/kg), IR_s is the ingestion rate of bamboo, EF is the exposure frequency (350 day/year), ED is the exposure duration (10.36 years), BW is the average body weight (105 kg, from mid-range of 80-130 kg [Zhang and Wei 2006]), and AT is the averaging time (3781.4 days).

Noncancer toxic risk was determined by the model hypothesis of HQ (Hazard Quotient):

$$HQ = \frac{ADD}{RfD_o}$$

where RfD_o is the reference dose of PBDEs (US EPA, 1997). Risk increases with HQ (Hang et al. 2009). If $HQ \leq 1$, risk exposure is relatively safe. If $1 < HQ \leq 10$, considerable threat is suggested, Finally, if $HQ \geq 10$, high chronic risk is suggested.

Results

Concentrations of PBDEs

Total PBDEs concentrations were consistently and significantly greater in captive pandas and their food supply than in wild pandas and their food and water supply (Fig. 2). In the fecal samples, $\Sigma 13$ PBDE of captive pandas was 2.55 times greater than in wild pandas (Fig. 2A). $\Sigma 13$ PBDE of *Fargesia qinlingensis* was 1.13 times higher, and of *Bashania fargesii* 1.55 times

189 higher, in leaves eaten by captive pandas (Figs. 2B, 2C). Water samples had low concentrations
190 of PBDEs (Fig. 2D).

191 Thirteen congeners of PBDEs were found in fecal samples; BDE47, BDE66, BDE71,
192 BDE99 and BDE154 predominated in captive pandas (Fig. 3A). Of the dozen congeners found in
193 the two bamboo species eaten by captive pandas, BDE47 and BDE99 predominated in *Fargesia*
194 *qinlingensis* and *Bashania fargesii*, respectively (Figs. 3B, 3C). Although captive pandas were
195 exposed to somewhat higher concentrations of PBDEs in their water supply (Fig. 2D), the
196 concentrations of each congener were quite low and none predominated (Fig. 3D). Ten PBDE
197 congeners were found in the supplemental feedstuff provided for captive pandas, with BDE28
198 and BDE183 predominating (Fig. 3E). Finally, nine PBDE congeners were found in the blood
199 samples collected from SWARC. BDE153 and BDE183 were the predominant congeners in
200 captive panda blood samples, and occurred in significantly higher concentrations than in blood
201 sampled from wild pandas (Fig. 3F).

Fig. 2. Total concentrations of PBDEs ($\Sigma 13\text{PBDEs}$) in (A) fecal samples; (B) leaves of *Fargesia qinlingensis*; (C) leaves of *Bashania fargesii*; (D) drinking water; (E) supplemental feedstuff; and (F) blood sample of wild (gray bars) and captive (cross-hatched bars) giant pandas. The concentrations of PBDEs in A, B, C and E are based on dry weight. In (F), the wild pandas were three 17-year old individuals rescued from Qingling and the captive pandas were 8–9-years old. Bars (means ± 1 SE of the mean from $n = 4$ independent replicates comprising three or four pooled samples). Different letters indicate significant differences between the wild and captive pandas identified using Tukey HSD test (all $P < 0.01$). $\text{pg} \cdot \text{g}^{-1} \text{lw}^{-1}$ = nanograms per gram lipid weights.

Fig. 3. Concentrations of individual PBDE congeners in (A) feces; (B) leaves of *Fargesia qinlingensis*; leaves of *Bashania fargesii*; (D) drinking water; (E) supplemental feedstuff; and (F) blood from wild (cross-hatched bars) and captive (black bars) giant pandas. Numbers 1 – 13 on the x-axis denote different congeners, respectively: BDE17; BDE28; BDE47; BDE66; BDE71; BDE85; BDE99; BDE100; BDE138; BDE153; BDE154; BDE183; BDE190. Different letters indicate significant differences between captive and wild pandas ($P < 0.05$).

Statistical results

The highest positive correlations in $\Sigma 13\text{PBDEs}$ were detected for feces and blood vs. bamboos ($r > 0.88$), and blood vs. feedstuff ($r = 0.88$). Concentration of PBDEs in water samples were not significantly correlated with any of the other samples (Table 1).

Three principal axes accounted for most of the variance in the groupings of PBDE congeners (Fig. 4). In leaves of *Fargesia qinlingensis*, the first three principal axes accounted for 87.8% of the variance (Fig. 4A), whereas they accounted for 62.5% for *Bashania fargesii* (Fig. 4B), and 77.4% in feedstuff (Fig. 4C). Only one grouping was found for PBDEs in drinking water (19.8% of the variance; Fig. 4D). The PCA identified clusters of the congeners BDE47, BDE66, BDE71, BDE99, and BDE154 in leaves of *Bashania fargesii* that matched the predominant congeners found in fecal samples (compare Fig. 4B with Fig. 3A).

Table 1 Spearman correlation matrix for PBDEs measured in captive samples

	<i>Fargesia qinlingensis</i>	<i>Bashania fargesii</i>	Blood	Water	Feces
<i>Bashania fargesii</i>	0.89**				
Blood	0.92**	0.92**			
Water	-0.11	-0.10	0.15		
Feces	0.90**	0.88**	0.93**	-0.09	
Feedstuff	0.73**	0.75**	0.88**	0.309	0.79**

**, $P < 0.01$ level (2-tailed test).

Fig. 4. Principal component triplot (principal axes 1–3) for 13 PBDE congeners in (A) leaves of *Fargesia qinglingensis*; (B) leaves of *Bashania fargesii*; (C) supplemental feedstuff; and (D) drinking water of samples taken at SWARC.

Health risk assessment

The ordering of hazard quotients (HQ) of the top five PBDE congeners in captive pandas was BDE99 (1.60) > BDE47 (1.10) > BDE100 (1.07) > BDE153 (0.97) > BDE154 (0.55), and was BDE47 (1.20) > BDE100 (0.99) > BDE154 (0.75) > BDE153 (0.57) > BDE99 (0.55) in wild pandas. with HQ values >1, BDE99 and BDE47 appear to pose a significant health risk to captive giant pandas, whereas BDE47 is congener putting wild pandas at risk.

Discussion

The first objective of this study was to test this hypothesis that giant pandas were exposed to PBDEs. Our previous research had found that Br in fecal samples of captive pandas were 3.3 times higher than in wild pandas (Chen et al., 2017), and data reported here supported the resulting hypothesis that captive pandas were exposed to PBDEs (Figs. 2A, 3A). In captive pandas, BDE47, BDE66, BDE71, BDE99, BDE154 were the major PBDE congeners in fecal samples, which matched the profile of PBDEs in leaves of the bamboo *Bashania fargesii* (Fig. 4B). This result suggests that the bamboo that is the primary food supply for pandas is the main source of PBDE exposure. Plants can accumulate high concentrations of toxicants in tissues that subsequently can accumulate and in the animals (Burreau et al. 2006; Voorspoels et al. 2007; McKinney et al. 2011; Krieger et al. 2016). Once in their body, PBDEs can lead to deficiencies in neural responses, thyroid hormone disorders, and carcinogenicity (McDonald 2002; Staskal et al. 2005; Lee et al. 2014).

Many studies have shown that PBDEs bioaccumulate in aquatic (Law et al., 2003) and terrestrial (Huwe et al., 2002; Jaspers et al., 2005; Pirard et al., 2007; Voorspels et al., 2006, 2007; Da Chen et al., 2012, 2013; Crosse et al., 2012; Andersen et al., 2015) species. However, we are aware of few studies that sampled PBDEs in fecal samples (Zheng et al., 2015), which are an appropriately non-invasive method to sample PBDEs in a rare species like the giant panda.

The second objective of our research was to compare exposure of wild and captive pandas to PBDEs. Not surprisingly, given the anthropogenic origin of PBDEs, the $\Sigma 13$ PBDE was significantly higher in samples taken from captive pandas. Plants often reflect the content of

organic pollutants in the environment (Collins and Finnegan, 2010) because the soil-air-plant pathway (pollutants volatilized from soils into the atmosphere are deposited onto plants) describes uptake of organic pollutants from contaminated soils (e.g., Paterson et al., 1991; Trapp and Matthies., 1997; Harrad et al., 2006; Collins and Finnegan., 2010; Ding et al., 2014). We found that the $\Sigma 13$ PBDE of in *Fargesia qinlingensis* and *Bashania fargesii* growing around the captive breeding center was significantly higher than that growing in the nature reserve (Fig. 2B and 2C). The panda captive breeding center at SWARC is located close to the large city of Xi'an and also to several potential sources of PBDE contamination: waste incinerators, electronic-waste processing facilities, and industrial discharges (He et al., 2014; Kosior et al., 2015; Wang et al., 2015).

The final objective of our research was to identify possible sources of exposure for PBDE pollution by pandas. This study is the first to measure PBDE concentrations in bamboo, so we can compare our results only with those from unrelated plant species. The $\Sigma 13$ PBDEs of *Fargesia qinlingensis* and *Bashania fargesii* growing at FNNR and SWARC in our study were higher than that of the moss *Pleurozium schreberi* in uncontaminated (755.6 pg/g dry mass) or urban (3062.9 pg/g dry mass) areas. Pandas eat little besides bamboo. They can consume on average 30 kg/day (Tuanmu et al., 2013) and bamboo accounts for > 99% of their diet (Hu, 1991, 2000), yet only 25% of the nutrients in bamboo can be assimilated (Zhou et al., 2008). This diet, together with the similarities in congener profiles of panda fecal samples (Fig. 3A) and the clustering in the PCA, (Fig. 4B) suggest that bamboo is the primary source of PDBE exposure for pandas.

The very low concentrations of PBDEs in water (Figs. 2D, 3D, 4D) suggest that it is an unlikely source of PBDEs for pandas. Likewise, the feedstuff appears an unlikely route of exposure. In captivity, pandas are fed a steamed bread supplement (“feedstuff”) that includes additional ingredients, including milk powder, apple, carrot, steamed bran, rice flour, maize flour, bean flour, fishmeal, bone meal, and mineral additives that provide supplemental nutrients essential for successful breeding programs (Chen and Ma, 2016). The $\Sigma 13$ PBDE in feedstuff (28.8 pg·g⁻¹; Fig. 2E) exceeded PBDEs concentrations in some farmland grains (13.7 pg·g⁻¹: Luo et al., 2009) but not others (30–440 pg·g⁻¹: Zheng et al., 2015). This might be because the feedstuff used in our research was purchased from local markets instead of having been made on site from locally-grown ingredients.

To our knowledge, this is the first investigation of exposure of pandas to PBDEs, and one of only a very few studies of PBDE exposure and bioaccumulation in a terrestrial species (Hoshi et al., 1998; Christensen et al., 2005). There are few comparables, but the *HQ* model for exposure risk suggests that BDE99, BDE47, and BDE153, each of which has an *HQ* >1, could threaten the health of captive giant pandas.

Pandas in captive breeding centers generally are thought to be better protected from human activities than are wild pandas in nature conservation zones. However, our previous research has shown that captive pandas are exposed to a variety of environmental pollutants (Chen et al. 2016). The data presented here provide further evidence that the habitat and captive breeding centers of giant pandas are polluted by PBDEs, and that exposure to PBDEs is significantly greater in captive conditions.

PBDEs in pandas most likely come from the bamboo they eat. Every giant panda consumes 30 kg of shoots and leaves of bamboo, on average, every day (Tuanmu et al. 2013). Therefore, even relatively low concentration of PBDEs in bamboo can still lead to a high dietary exposure that threatens the health of the giant pandas. For mammals, PBDEs can be transferred to nursing offspring via mother's milk (Travis and Hattermer-Frey, 1991; Darnerud, 2003; Beineke et al., 2005; Beineke et al., 2007). PBDEs also have immunotoxicity and can be immunosuppressants (Arkoosh et al., 2010; Frouin et al., 2010; Lv et al., 2015) that make pandas more vulnerable to bacterial and viral infections.

Conclusions and Recommendations

Our data suggest that pandas are exposed to high levels of PDBEs in captive breeding centers, and may represent a significant health risk for pandas in captivity. We recommend that managers of these centers and captive breeding programs, including the Chinese State Forestry Administration (SFA), seek strategies to minimize PDBE exposure by pandas lest decades of successful *ex situ* conservation efforts become compromised by the increasing pollution associated with Chinese economic development. A short-term solution to addressing this issue is to reduce the supply of contaminated bamboo and to grow uncontaminated bamboo strictly for captive pandas. In the long term, however, sustaining a successful captive breeding program for pandas will require reduction of air, water, and soil pollution that will lead to improvements in the environmental quality of the giant panda's natural habitat.

Acknowledgements

This research was supported by a project from SKLLQQG (State Key Laboratory of Loess and Quaternary Geology) and IEECAS (Institute of Earth Environment, Chinese Academy of Sciences (ZZBS1303)). AME's participation in this project was supported by the Chinese Academy of Sciences (CAS) Presidential International Fellowship Initiative for Visiting Scientists, Grant no. 2016VB074.

References

- Arkoosh, M.R., Boylen, D., Dietrich, J., Anulacion, B.F., Ylitalo, G., Bravo, C.F., Johnson, L.L., Loge, F.J., Collier, T.K., 2010. Disease susceptibility of salmon exposed to polybrominated diphenyl ethers (PBDEs). *Aquat. Toxicol.* 98, 51–59.
- Andersen, M.S., Fuglei, E., König, M., Lipasti, I., Pedersen, Å., Polder, A., Yoccoz, N.G., Routti, H., Levels and temporal trends of persistent organic pollutants (POPs) in arctic foxes (*Vulpes lagopus*) from Svalbard in relation to dietary habits and food availability. *Sci. Total Environ.* 472, 112–122.
- Beineke, A., Siebert, U., McLachlan, M., Bruhn, R., Thron, K., Failing, K., Müller, G., Baumgärtner, W., 2005. Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbor porpoises (*Phocoena phocoena*). *Environ. Sci. Technol.* 39, 3933–3938.
- Beineke, A., Siebert, U., Stott, J., Müller, G., Baumgärtner, W., 2007. Phenotypical characterization of changes in thymus and spleen associated with lymphoid depletion in

347 free-ranging harbor porpoises (*Phocoena phocoena*). Vet. Immunol. Immunop. 117, 254–
 348 265.

349 Branchi, I., Capone, F., Alleva, E., Costa, L.G., 2003. Polybrominated diphenyl ethers
 350 neurobehavioral effects following developmental exposure. Neurotoxicol. 24, 449–462.

351 Burreau, S., Zebuhr, Y., Broman, D., Ishaq, R., 2006. Biomagnification of PBDEs and PCBs in
 352 food webs from the Baltic Sea and the northern Atlantic Ocean. Sci. Total Environ. 366.
 353 659–672.

354 Chen, D., Letcher, R.J., Martin, P., 2012. Flame retardants in eggs of American kestrels and
 355 European starlings from southern Lake Ontario region (North America). J. Environ.
 356 Monit. 14. 2870–2876.

357 Chen, D., Martin, P., Burgess, N.M., Champoux, L., Elliott, J.E., Forsyth, D.J., Idrissi, A., and
 358 Letcher, R.J., 2013. European starlings (*Sturnus vulgaris*) suggest that landfills are an
 359 important source of bioaccumulative flame retardants to Canadian terrestrial ecosystems.
 360 Environ. Sci. Technol. 47. 12238–12247.

361 Chen, Y.P., Maltby, L., Liu, Q., Song, Y., Zheng, Y.J., Ellison, A.M., Ma, Q.Y., Wu, X.M., 2016.
 362 Captive pandas are at risk from toxic chemicals. Front. Ecol. Environ. 14. 363-367.

363 Chen, Y.P., Ma, Q.Y., 2017. Supplement nutrition has potential risk for captive pandas. J. Earth
 364 Environ. (in press).

365 Christensen, J.R., MacDuffee, M., Macdonald, R.W., Whitticar, M., Ross, P. 2005. Persistent
 366 organic pollutants in British Columbia grizzly bears: consequence of divergent diets.
 367 Environ. Sci. Technol. 39, 6952–6960.

368 Collins, C.D., Finnegan, E., 2010. Modeling the plant uptake of organic chemicals, including the
369 soil–air–plant pathway. *Environ. Sci. Technol.* 44, 998–1003.

370 Crosse, J.D., Shore, R.F., Wadsworth, R.A., Jones, K.C., Pereira, G., 2012. Long-term trends in
371 pbdes in sparrowhawk (*Accipiter nisus*) eggs indicate sustained contamination of UK
372 terrestrial ecosystems. *Environ. Sci. Technol.* 46, 13504–13511.

373 Darnerud, P.O., 2003. Toxic effects of brominated flame retardants in man and in wildlife.
374 *Environ. Int.* 29, 841–853.

375 Darnerud, P.O., Thuvander, A., 1998. Studies on immunological effects of polybrominated
376 diphenyl ethers (PBDE) and polychlorinated biphenyl (PCB) exposure in rat and mice.
377 *Organohalogen Compd.* 35, 415–418.

378 Darnerud, P.O., Eriksen, G.S., Johannesson, T., Larsen, P.B., Viluksela, M., 2001.
379 Polybrominated diphenyl ethers: occurrence, dietary exposure, and toxicology. *Environ.*
380 *Health Persp.* 109, 49–68.

381 De Wit, C.A., 2002. An overview of brominated flame retardants in the environment.
382 *Chemosphere* 46, 583–624.

383 Ding, C., Chang, W.J., Zeng, H., Ni, H.G., 2014. Field and modeling study of PBDEs uptake by
384 three tree species. *Sci. Total Environ.* 472, 923–92.

385 Eriksson, P., Jakobsson, E., Fredriksson, A., 2001. Articles brominated flame retardants: a novel
386 class of developmental neurotoxicants in our environment? *Environ. Health Persp.* 109,
387 903–908.

388 Fowles, J.R., Fairbrother, A., Baechersteppan, L., Kerkvliet, N.I., 1994. Immunological and
 389 endocrine effects of the flame-retardant pentabromodiphenyl ether de-71 in c57blr6j mice.
 390 Toxicology 86, 49–61.

391 Frouin, H., Lebeuf, M., Hammill, M., Masson, S., Fournier, M., 2010. Effects of individual
 392 polybrominated diphenyl ethers (PBDE) congeners on harbour seal immune cells in vitro.
 393 Mar. Pollut. Bull. 60, 291–298.

394 Hang, X.S., Wang, H.Y., Zhou, J.M., Ma, C.L., Du, C.W., Chen, X.Q., 2009. Risk assessment of
 395 potentially toxic element pollution in soils and rice (*Oryza sativa*) in a typical area of the
 396 Yangtze River Delta. Environ. Pollut. 157, 2542–2549.

397 Harrad, S., Ren, J., Hazrati, S., Robson, M., 2006. Chiral signatures of PCB# s 95 and 149 in
 398 indoor air, grass, duplicate diets and human faeces. Chemosphere 63, 1368–1376.

399 He, W., Qin, N., He, Q.S., Kong, X.Z., Liu, W.X., Wang, Q.M., Yang, C., Jiang, Y.J., Yang, B.,
 400 Bai, Z.L., Wu, W.J., Xu, F.L. 2014. Atmospheric PBDEs at rural and urban sites in
 401 central China from 2010 to 2013: residual levels, potential sources and human exposure.
 402 Environ. Pollut. 192, 232–243.

403 Hooper, K., McDonald, T.A., 2000. The PBDEs: An emerging environmental challenge and
 404 another reason for breast-milk monitoring programs. Environ. Health Persp. 108, 387.

405 Hoshi, H., Minamoto, N., Iwata, H., Shiraki, K., Tatsukawa, R., Tanabe, S., Fujita, S., Hirai, K.,
 406 Kinjo, T. 1998. Organochlorine pesticides and polychlorinated biphenyl congeners in
 407 wild terrestrial mammals and birds from Chubu Region, Japan: Interspecies Comparison
 408 of the Residue Levels and Compositions. Chemosphere 36, 211–3221.

409 Hu, G.C., Luo, X.J., Dai, J.Y., Zhang, X.L., Wu, H., Zhang, C.L., Guo, W., Xu, M.Q., Mai, B.X.,
 410 Weit, F.W. 2008. Brominated flame retardants, polychlorinated biphenyls, and
 411 organochlorine pesticides in captive giant panda (*Ailuropoda melanoleuca*) and red panda
 412 (*Ailurus fulgens*) from China. Environ. Sci. Technol. 42: 4704–4709.

413 Hu, J.C., 1991. Habitating environment of giant pandas and food base. A special topic of
 414 zoology. Beijing University Press, Beijing, in Chinese.

415 Hu, J.C., 2000. Review on the classification and population ecology of the giant panda.
 416 Zool. Res. 21, 28–34.

417 Huwe, J.K., Lorentzsen, M., Thuresson, K., Bergman, A., 2002. Analysis of mono- to
 418 deca-brominated diphenyl ethers in chickens at the parts per billion level. Chemosphere
 419 46, 635–640.

420 Jaspers, V., Covaci, A., Maervoet, J., Dauwe, T., Voorspoels, S., Schepens, P., Eens, M., 2005.
 421 Brominated flame retardants and organochlorine pollutants in eggs of little owls (*Athene*
 422 *noctua*) from Belgium. Environ. Pollut. 136, 81–88.

423 Kim, T.H., Bang, D.Y., Lim, H.J., Won, A.J., Ahn, M.Y., Patra, N., Chung, K.K., Kwack, S.J.,
 424 Park, K.L., Han, S.Y., Choi, W.S., Han, J.Y., Lee, B.M., Oh, J.E., Yoon, J.H., Lee, J.,
 425 Kim, H.S. 2012. Comparisons of polybrominated diphenyl ethers levels in paired South
 426 Korean cord blood, maternal blood, and breast milk samples. Chemosphere 87, 97–104.

427 Kosior, G., Klánová, J., Vanková, L., Kukucka, P., Chropenová, M., Brudzinska-Kosior, A.,
 428 Samecka-Cymerman, A., Kolon, K., Kempers, A.J., 2015. *Pleurozium schreberi* as an

ecological indicator of polybrominated diphenyl ethers (PBDEs) in a heavily industrialized urban area. *Ecol. Indic.* 48, 492–497.

Krieger, L.K., Szeitz, A., Bandiera, S.M. 2016. Evaluation of hepatic biotransformation of polybrominated diphenyl ethers in the polar bear (*Ursus maritimus*). *Chemosphere* 146, 555–564.

Lee, H.J., An, S., Kim, G.B. 2014. Background level and composition of polybrominated diphenyl ethers (PBDEs) in creek and subtidal sediments in a rural area of Korea. *Sci Total Environ* 470–47, 1479–1484

Legler, J., Brouwer, A., 2003. Are brominated flame retardants endocrine disruptors? *Environ. Int.* 29, 879–885.

Li, Y.M., Jiang, G.B., Wang, Y.W., Wang, P., Zhang, Q.H., 2008. Concentrations, profiles and gas-particle partitioning of PCDD/Fs, PCBs and PBDEs in the ambient air of an E-waste dismantling area, Southeast China. *Chin. Sci. Bull.* 53, 521–528.

Law, R.J., Alaee, M., Allchin, C.R., Boon, J.P., Lebeuf, M., Lepom, P., Stern, G.A., 2003. Levels and trends of polybrominated diphenylethers and other brominated flame retardants in wildlife. *Environ. Int.* 29, 757–770.

Luo, X.J., Liu, J., Luo, Y., Zhang, X.L., Wu, J.P., Lin, Z., Chen, S.J., Mai, B.X., Yang, Z.Y. 2009. Polybrominated diphenyl ethers (PBDEs) in free-range domestic fowl from an e-waste recycling site in South China: levels, profile and human dietary exposure. *Environ. Int.* 35, 253–258.

449 Lv, Q.Y., Wan, B., Guo, L.H., Zhao, L., Yang, Y., 2015. In vitro immune toxicity of
 450 polybrominated diphenyl ethers on murine peritoneal macrophages: apoptosis and
 451 immune cell dysfunction. *Chemosphere* 120, 621–630.

452 Mara, H., 2015. Captive pandas succumb to killer virus. *Science* 347, 700–701.

453 McDonald, T.A. 2002. A perspective on the potential health risks of PBDEs. *Chemosphere* 46,
 454 745–755.

455 McKinney, M.A., Dietz, R., Sonne, C., Guise, S.D., Skirnisson, K., Karlsson, K., Steingrimsen,
 456 E., Letcher, R.J. 2011. Comparative hepatic microsomal biotransformation of selected
 457 PBDEs, including decabromodiphenyl ether, and decabromodiphenyl ethane flame
 458 retardants in Arctic marine-feeding mammals. *Environ. Toxicol. Chem.* 30, 1506–1514.

459 Paterson, S., Mackay, D., Bacci, E., Calamari, D., 1991. Correlation of the equilibrium and
 460 kinetics of leaf–air exchange of hydrophobic organic chemicals. *Environ. Sci. Technol.*
 461 25, 866–871.

462 Pirard, C., DePauw, E., 2007. Absorption, disposition and excretion of poly-brominated
 463 diphenylethers (PBDEs) in chicken. *Chemosphere* 66, 320–325.

464 SFA (State Forestry Administration)., 2015. The 4th national survey report on giant panda in
 465 China. Science Press, Beijing (in Chinese).

466 Staskal, D.F., Diliberto, J.J., DeVito, M.J., Birnbaum, L.S. 2005. Toxicokinetics of BDE 47 in
 467 female mice: effect of dose, route of exposure, and time. *Toxicol. Sci.* 83, 215–223.

468 Trapp, S., Matthies, M., 1997. Modeling volatilization of PCDD/F from soil and uptake into
 469 vegetation. *Environ. Sci. Technol.* 31, 71–74.

470 Travis, C.C., Hattermer-Frey, H.A., 1991. Human exposure to dioxin. *Sci. Total Environ.* 104,
 471 97–127.

472 Tuanmu, M.N., Vina, A., Winkler, J.A., Li, Y., Xu, W.H., Ouyang, Z.Y., Liu, J.G., 2013.
 473 Climate-change impacts on understorey bamboo species and giant pandas in China's
 474 Qinling Mountains. *Nat. Clim. Change* 3, 249–253.

475 US EPA. 1997. Exposure factors handbook; EPA/600/P-95/002Fa, b, c; Environmental
 476 Protection Agency. Office of Research and Development, Washington, DC.

477 Voorspoels, S., Covaci, A., Jaspers, V.L.B., Neels, H., Schepens, P. 2007. Biomagnification of
 478 PBDEs in three small terrestrial food chains. *Environ. Sci. Technol.* 41, 411–416.

479 Wang, J.X., Liu, L.L., Wang, J.F., Pan, B.S., Fu, X.F., Zhang, G., Zhang, L., Lin, K.F., 2015.
 480 Distribution of metals and brominated flame retardants (BFRs) in sediments, soils and
 481 plants from an informal e-waste dismantling site, South China. *Environ. Sci. Pollut. Res.*
 482 22, 1020–1033.

483 WHO, World Health Organization., 1994. Brominated Diphenyl Ethers. IPCS, Environmental
 484 Health Criteria. Geneva, PP.162.

485 Zheng, X.B., Luo, X.J., Zheng, J., Zeng, Y.H., Mai, B.X., 2015. Contaminant sources,
 486 gastrointestinal absorption, and tissue distribution of organohalogenated pollutants in
 487 chicken from an e-waste site. *Sci. Total Environ.* 505, 1003–1010.

488 Zhou, S.Q., Huang, J.Y., Liu, B., Zhang, Y.H., Tan, Y.C., Zhou, X.P., Huang, Y., Li, D.S.,
 489 Zhang, G.Q., Wei, R.P., Tang, C.X., Wa, P.Y., Zhang, H.M., 2008. A preliminary study

490 on the food utilization ratio of wildness training panda. Sichuan Forest Exploration

491 Design. 30, 17.