



# Arginine dependence of tumor cells: targeting a chink in cancer's armor

## Citation

Patil, M D, J Bhaumik, S Babykutty, U C Banerjee, and D Fukumura. 2016. "Arginine Dependence of Tumor Cells: Targeting a Chink in Cancer's Armor." *Oncogene* 35 (38) (April 25): 4957–4972. doi:10.1038/onc.2016.37.

## Published Version

doi:10.1038/onc.2016.37

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:37133883>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

1           **Arginine Dependence of Tumor Cells: Targeting a Chink in**  
2   **Cancer's Armor**

3  
4  
5           **Mahesh D. Patil,<sup>1</sup> Jayeeta Bhaumik,<sup>1</sup> Suboj Babykuty,<sup>2</sup>**  
6   **Uttam Chand Banerjee<sup>1</sup> and Dai Fukumura<sup>2\*</sup>**

7  
8  
9  
10          <sup>1</sup>*Department of Pharmaceutical Technology (Biotechnology), National Institute of*  
11               *Pharmaceutical Education and Research, Sector-67, S.A.S. Nagar – 160062*  
12   *Punjab, India*

13  
14          <sup>2</sup>*Edwin L. Steele Laboratory, Department of Radiation Oncology, Massachusetts General*  
15               *Hospital and Harvard Medical School, 100 Blossom Street, Boston, MA 02114, USA*

16  
17   \*Corresponding author's e-mail:

18   [dai@steele.mgh.harvard.edu](mailto:dai@steele.mgh.harvard.edu)

19

1 **Abstract**

2 Arginine, one among the twenty most common natural amino acids, plays a pivotal  
3 role in cellular physiology as it is being involved in numerous cellular metabolic and  
4 signaling pathways. Dependence on arginine is diverse for both tumor and normal cells. Due  
5 to decreased expression of argininosuccinate synthetase (ASS) and/or ornithine  
6 transcarbamoylase (OTC), several types of tumor are auxotrophic for arginine. Deprivation of  
7 arginine exploits a significant vulnerability of these tumor cells and leads to their rapid  
8 demise. Hence, enzyme-mediated arginine depletion is a potential strategy for the selective  
9 destruction of tumor cells. Arginase, arginine deiminase (ADI) and arginine decarboxylase  
10 (ADC) are potential enzymes that may be used for arginine deprivation therapy. These  
11 arginine catabolizing enzymes not only reduce tumor growth but also make them susceptible  
12 to concomitantly administered anti-cancer therapeutics. Most of these enzymes are currently  
13 under clinical investigations and if successful will potentially be advanced as anti-cancer  
14 modalities.

15

16 **Keywords: cancer, arginine deprivation, arginase, arginine deiminase, arginine**  
17 **decarboxylase**

18

## 1           **Introduction**

2           Amino acids play a major role in regulating important cellular events in both normal  
3           and malignant cells. Besides their role in the synthesis of hormones and peptides, amino acids  
4           also function as cell signaling molecules, playing a modulatory role in gene expression.<sup>1</sup>  
5           Amino acids regulate RNA synthesis by diverse mechanisms ranging from regulating  
6           transcription factors assembly,<sup>2</sup> to total mRNA turnover.<sup>3,4</sup> Amino acids are major  
7           determinants of a normal cellular physiology, therefore potential signaling pathways such as  
8           amino acid response (AAR) pathway sense their altered metabolism [Figure 1]. Hence, amino  
9           acid levels in the body are critical for important cellular functions.<sup>5-9</sup>

10          There is a significant difference between the metabolism of normal and malignant  
11          cells.<sup>10</sup> For instance, bio-energetic requirements for homeostasis in normal cells are fulfilled  
12          by catabolic metabolism. On the other hand, the majority of the tumor cells alter their  
13          metabolic program (“*metabolic remodeling*”) and consume additional nutrients in order to  
14          maintain a balance between elevated macromolecular biosynthesis<sup>11</sup> and adequate levels of  
15          ATP for survival.<sup>12,13</sup> However, the endogenous supply of nutrients becomes inadequate  
16          during intense growth. Thus tumor cells depend on exogenous nutrients in their  
17          microenvironment to fulfill the elevated energy requirements *i.e.* they become auxotrophic  
18          for nutrient and energy sources.<sup>14-16</sup> Deprivation of amino acids results in growth inhibition  
19          or death of tumor cells by the modulation of various signaling cascades.<sup>6-9,17,18</sup>

20          Exogenously incorporated enzymes that deprive amino acids could be a novel  
21          strategy for the treatment of auxotrophic tumors. The first FDA approved heterologous  
22          enzyme for the treatment of cancer was *E. coli* L-asparaginase.<sup>19</sup> L-asparaginase exploits the  
23          differences on their dependence of normal and leukemic cells towards L-asparagine.<sup>20</sup> L-  
24          asparaginase has been proven to be a promising agent for the treatment of L-asparagine  
25          auxotrophic T-cell acute lymphoblastic lymphoma (T-ALL). Use of L-asparaginase in T-

1 ALL opened up new windows of ‘amino acid-depriving therapy’. Currently, there is a  
2 resurgence of interest in enzyme-mediated amino acid deprivation as a new therapeutic  
3 approach for cancer treatment.<sup>6,7,21,22</sup> For example, arginine depletion can inhibit tumor cell  
4 proliferation and induce cell death pathways. Here we endeavor to provide a basic  
5 understanding of the roles of arginine in normal and tumor cell with emphasis on current  
6 knowledge and developments in the application of enzyme-mediated arginine depriving  
7 therapy as a potential anticancer approach.

### 8 **Enzyme-mediated arginine deprivation: a potential anti-cancer approach**

9 Arginine is involved in the regulation of various molecular pathways and thus  
10 availability of arginine can modulate key metabolic, immunological, neurological and  
11 signaling pathways of the cells [Figure 2 and 3].<sup>23,24</sup> Auxotrophy towards arginine by certain  
12 tumor cells (particularly that of hepatocellular carcinoma and melanoma) has been well  
13 characterized.<sup>25,26</sup> Normal cells, when deprived of arginine, undergo cell cycle arrest at G<sub>0</sub>/G<sub>1</sub>  
14 phase and become quiescent. If reinstated with arginine, the majority of the normal cells  
15 recover to their normal proliferation status. However, arginine deprivation in tumor cells does  
16 not arrest cell cycle at G<sub>1</sub> phase and continue to be in a cell cycle, leading tumor cells to  
17 undergo unbalanced growth and eventually lead to the activation of apoptotic pathways.<sup>27,28</sup>

18 Owing to the involvement of arginine in a plethora of cellular pathways, arginine  
19 dependence of tumor cells has rapidly emerged as a potential target for cancer.<sup>29</sup> However,  
20 dietary restriction results in the reduction of only 30% of plasma free arginine.<sup>30</sup> Thus,  
21 arginine degrading enzyme-mediated arginine deprivation has been proposed as a potential  
22 anti-cancer therapy by various research groups.<sup>27-35</sup> Enzymes that can be used for arginine  
23 deprivation therapy (ADT) include arginine deiminase (ADI), arginase and arginine  
24 decarboxylase (ADC) as discussed below [Figure 3].

25

## 1           **1. Arginine deiminase**

2           Arginine deiminase (ADI) (E.C.3.5.3.6) is a prokaryotic enzyme originally isolated  
3 from *Mycoplasma*, which catalyzes an irreversible deimination of the guanidine group of L-  
4 arginine to citrulline and ammonium ion.<sup>36</sup> Normal cells are able to convert citrulline into  
5 arginine through argininosuccinate synthetase (ASS) and argininosuccinate lyase (ASL),  
6 expression of which are tightly regulated. However, the expression of ASS/ASL is down-  
7 regulated in certain tumor cells by unknown mechanisms and these cells are unable to  
8 convert citrulline to arginine.<sup>30-33,37</sup> This makes the tumor cells auxotrophic for arginine for  
9 their growth and cellular functioning. ADI-mediated arginine deprivation leads to apoptotic  
10 cell death, selectively of arginine auxotrophic ASS (-) tumor cells sparing the ASS (+) ADI  
11 resistant normal cells<sup>38</sup> [Table 1]. Incidence of ASS deficiency varies depending on the  
12 tumor type and expression level of ASS has been proposed as a biomarker for identification  
13 of ADI sensitive tumors.<sup>24,25,39-42</sup>

14           In 1990, Miyazaki and co-workers<sup>43</sup> were the first to report the growth inhibition of  
15 *Mycoplasma* infected human tumor cells. The cause of growth inhibition of human tumor cell  
16 lines was identified as a ADI produced by *Mycoplasma*. *In vitro* growth-inhibitory dose of  
17 *Mycoplasma* ADI appeared to be 1000 times lower than that of bovine liver arginase.  
18 Subsequently in 1992, growth inhibitory activity of ADI was demonstrated in ASS-  
19 downregulated human melanoma cells.<sup>44</sup> These pioneering studies established ADI as a  
20 potential anti-cancer enzyme [Figure 4].

### 21           **1.1 PEGylated ADI**

22           Being microbial in origin, ADI has serious disadvantages of eliciting strong  
23 antigenicity and rapid plasma clearance (half-life of 4 h). To circumvent these limitations,  
24 several studies have aimed to extend the plasma half-life of ADI and to minimize its  
25 antigenicity. In 1993, Takaku *et al* addressed these problems for the first time by

1 polyethylene glycol (PEG) modification.<sup>45</sup> Remarkably, PEGylation of *Mycoplasma arginini*  
2 ADI enhanced its cytotoxic potential *in vivo* and once a week intravenous injection of PEG-  
3 ADI at a dose of 5 U/mouse (10 mg protein/Kg) depleted plasma arginine to an undetectable  
4 level at least for a week, whereas native enzyme required 10 daily injections to achieve  
5 similar effects. Nevertheless, PEGylation of *Mycoplasma hominis* ADI also resulted in  
6 significant enhancement of arginine lowering potential of native *Mycoplasma hominis*  
7 ADI.<sup>46,47</sup> Recently, PEGylation and pharmacological properties of an engineered ADI  
8 originated from *Pseudomonas plecoglossicida* have been studied. PEGylated *Pseudomonas*  
9 *plecoglossicida* ADI remarkably improves the systemic half-life (by 11-folds) and found to  
10 exhibit superior efficacy than native ADI in depleting plasma arginine.<sup>48</sup>

11 PEG-ADI has also shown promising outcomes for the treatment of human  
12 malignancies. In March 1999, ADI-PEG20, PEGylated recombinant *Mycoplasmal* ADI was  
13 approved as an orphan drug by US-FDA for the treatment of HCC and malignant melanomas.  
14 Subsequently in July 2005, European Agency for the Evaluation of Medicinal Products  
15 (EMA) granted orphan drug status to ADI-PEG20 for the treatment of HCCs.<sup>49</sup>

16 ADI-PEG20 is currently undergoing clinical investigation as a randomized double-  
17 blind Phase III trial in patients with advanced HCC (NCT 01287585), Phase II studies in  
18 patients with ASS-negative metastatic melanoma (NCT 01279967) and Phase II studies in  
19 patients with relapsed small-cell lung cancer (SCLC) (NCT 01266018)<sup>50</sup> [Table 2]. Outcomes  
20 of the previous clinical studies were also encouraging, achieving response rates of 25% and  
21 47% in melanoma and HCC, respectively [Table 2]. Moreover, grade III and IV toxicities  
22 have not been observed in clinical investigations involving ADI-PEG20 in metastatic  
23 melanoma and HCC patients.<sup>51,52</sup> Therefore, clinicians are looking forward to the  
24 establishment of ADI-PEG20 as a potent anti-cancer modality.

25

## 1 1.2 Tumor sensitivity towards ADI

2 The auxotrophicity of tumors towards arginine and their sensitivity towards **it can be**  
3 **attributed** to the lack or reduced expression of ASS in tumors.<sup>25,37-39,53</sup> **Notably**, numerous  
4 tumor cells which are deficient in ASS expression, are sensitive towards ADI treatment  
5 **[Table 1]**. Transfection of an expression plasmid containing human ASS cDNA in HCC and  
6 melanoma cells confers severe resistance to ADI treatment compared to ASS-negative cells.<sup>47</sup>  
7 **Till** date, most promising targets for ASS expression dependent ADT identified are human  
8 melanoma and HCCs. Other promising targets include malignant pleural mesothelioma  
9 (MPM), renal cell carcinoma, prostate cancer, T-ALL and osteosarcoma.<sup>50</sup> However,  
10 molecular mechanisms underlying tumor sensitivity towards ADI treatment, by down-  
11 regulation of ASS expression in tumor cells, are still elusive. Promoter hypermethylation-  
12 dependent silencing of ASS gene is an endorsed mechanism of ASS gene repression.<sup>37,54-56</sup>  
13 Methylation frequency of the ASS promoter upto 50-80% level at the CpG loci is  
14 documented across a broad range of lymphomas. **In contrast**, normal lymphoid samples were  
15 found unmethylated.<sup>26</sup> Treatment of ADI-PEG20 to ASS-methylated lymphoma cell lines  
16 **revealed** dramatic decrease in **the** proliferation rate and viability count, by inducing caspase-  
17 dependent apoptosis, without **affecting** normal lymphoblastoid cell lines. Demethylation-  
18 induced resistance to ADI-PEG20 treatment has also been confirmed in Cutaneous T-cell  
19 Lymphoma (CTCL) cell lines, as their incubation with 5-Aza-dC (demethylating agent) for 8  
20 days **which** resulted in partial demethylation, followed by transcriptional activation and  
21 synthesis of ASS protein.<sup>26</sup>

22 **Recently Rabinovich *et al* have confirmed that proliferation of the osteosarcoma cells**  
23 **is supported by down-regulation of ASS, by facilitating pyrimidine synthesis via activation of**  
24 **CAD (carbamoyl-phosphate synthase 2, aspartate transcarbamylase and dihydroorotase)**  
25 **complex.<sup>57</sup> As cytosolic aspartate serves as a substrate for both ASS and for CAD complex,**



1 ASS down-regulation can enhance aspartate availability for CAD for the synthesis of  
2 pyrimidine nucleotides to promote proliferation. Thus, aspartate transport can be exploited as  
3 an additional therapeutic target in tumors with ASS down-regulation, especially in those ones  
4 which develop resistance to arginine-depriving enzymes.

### 5 **1.3 Tumor resistance towards ADI**

6 ASS-deficient tumors are sensitive to ADI treatment; however, arginine deprivation  
7 eventually up-regulates ASS expression in tumor cells and thereby confers resistance towards  
8 ADI.<sup>25,58</sup> Transcriptional induction of ASS expression and increase in ASS mRNA level is  
9 reported in human embryonic kidney cells and melanoma cells during arginine starvation.<sup>59,60</sup>  
10 Transcription factors such as c-Myc and HIF-1 $\alpha$  are involved in the up-regulation of ASS  
11 expression under arginine depleted conditions.<sup>60</sup> E-box and GC-box are the important  
12 sequences located between -85 and -35 nucleotides in the ASS promoter region that modulate  
13 ASS expression through their interactions with c-Myc and HIF-1 $\alpha$ . Under the normal  
14 concentrations of arginine, HIF-1 $\alpha$  (but not c-Myc) binds to E-box and thus acts as a negative  
15 regulator of ASS expression. Under the conditions of arginine depletion, HIF-1 $\alpha$  is degraded  
16 and replaced by up-regulated c-Myc, which directly binds to E-box; thus, c-Myc acts as a  
17 positive regulator of ASS expression [Fig. 6 of Ref. 60]. Recently reported in melanoma  
18 cells, inhibition of ubiquitin-mediated protein degradation is a molecular mechanism  
19 responsible for the stabilization and accumulation of c-Myc.<sup>61</sup> Furthermore, various cellular  
20 pathways, such as Ras and its downstream ERK/PI3K/AKT kinase cascade are associated  
21 with the post-translational modifications of c-Myc, leading to its phosphorylation and  
22 stabilization during ADI-PEG20-mediated arginine deprivation conditions. Involvement of  
23 Ras/PI3K/ERK signaling pathway in the development of resistance towards ADI treatment  
24 suggests that combination of ADI with Ras/ERK, PI3K/AKT inhibitors is a potential  
25 therapeutic strategy to improve the anti-cancer response.<sup>62,63</sup>

1 Development of anti-drug neutralizing antibodies is another possible mechanism of  
2 resistance towards ADI-PEG20 treatment.<sup>64</sup> Arginine concentrations were recovered up-to  
3 pre-treatment levels in a patient with malignant pleural mesothelioma and in Asian patients  
4 with advanced hepatocellular carcinoma following the ADI-PEG20 treatment. This recovery  
5 in arginine concentration was found concomitant with an increase in anti-ADI-PEG20  
6 antibody titer.<sup>65</sup> These studies suggest the involvement of drug-associated resistance i.e. anti-  
7 drug neutralizing antibodies, rather than tumor-related factors as another possible mechanism  
8 of resistance of some tumor cell types towards ADI-PEG20 treatment.<sup>62,63</sup>

## 10 **1.4 Anti-tumor mechanisms of ADI treatment**

### 11 *1.4.1 Role of autophagy and apoptosis in ADI-mediated arginine deprivation therapy*

12 Due to the involvement of arginine in numerous cellular pathways [Figure 2], the  
13 exact anti-proliferative mechanisms of ADI treatment, besides that of arginine depletion, are  
14 still elusive. One of the potential pathways involved in the cytostatic and cytotoxic potential  
15 of ADI is TRAIL (tumor necrosis factor-related apoptosis-inducing ligand).<sup>66-68</sup> TRAIL plays  
16 an important role in the cleavage of Beclin-1 (Atg6) and Atg5 in arginine deprived melanoma  
17 cells.<sup>69</sup> Beclin-1 and Atg5 are essential for the formation of autophagosomes and thus crucial  
18 for autophagy. Since autophagy serves as a mean to evade apoptosis in arginine depleted  
19 cells, TRAIL induced cleavage of Beclin-1 and Atg5 leads to decreased autophagy, thereby  
20 increasing apoptosis.<sup>69</sup> Additionally, these two drugs (ADI and TRAIL) complement each  
21 other by activating the intrinsic apoptosis pathways. ADI-PEG20 increases cell surface  
22 receptors DR4/5 for TRAIL thereby binding TRAIL to these death receptors. As a result,  
23 caspase-8 or 10 are activated.<sup>66</sup> ADI-PEG20 treatment also modulates different autophagic  
24 pathways involved in the cell survival. AMPK and ERK pathways are activated in ADI-  
25 treated prostate cancer cells; while AKT, mTOR and S6K pathways are attenuated. ADI-

1 PEG20 treatment to CWR22Rv1 prostate cancer cells induced autophagy, as revealed by the  
2 appearance of LC-II only after 30 minutes exposure [continues](#) its persistence after 24 hours  
3 following ADI-PEG20 treatment.<sup>70,71</sup> [Additionally](#), inhibition of autophagy by chloroquine, a  
4 clinically approved anti-malarial agent which inactivates lysosomal functions, accelerates the  
5 ADI-induced apoptotic cell death of prostate cancer<sup>70,71</sup> and SCLCs.<sup>39</sup> Thus autophagy has  
6 been proposed as a pro-survival mechanism of tumor cells during arginine deprivation.<sup>71</sup>

7 ADI-mediated arginine deprivation is also known to induce caspase-dependent  
8 apoptotic pathways in many of the tumor cells types. ADI-PEG20 treatment activates  
9 caspase-3 in ASS-methylated malignant lymphoma cells, whereas ASS-positive normal  
10 lymphoblastoid cells are resistant to it.<sup>26</sup> Similarly, cell death has been attributed to caspases  
11 activation in glioblastoma,<sup>54</sup> melanoma,<sup>38,72</sup> leukemia<sup>73</sup> and pancreatic cancer cells.<sup>74</sup>  
12 Moreover, all these studies indicate that inhibition of autophagy leads to further advancement  
13 in [the](#) ADI-PEG20-mediated demise of tumor cells, suggesting the induction of autophagy as  
14 a mechanism of tumor resistance to ADI-PEG20 treatment.

15 Cumulative [pieces of](#) evidence suggest that [the](#) activation of caspases is not a sole  
16 decisive phenomenon in programmed cell death pathways. Caspase-dependent apoptosis is a  
17 major mode of cell death, but in its absence or failure, there are other pathways which can  
18 also execute cell death.<sup>75-77</sup> ADI-PEG20 treatment to SCLC, leukemia, retinoblastoma and  
19 prostate cancer cells induces apoptotic cell death pathways, however, without activation of  
20 caspases, suggesting the role of caspase-independent apoptosis as a cell death pathway.<sup>33,39,69,</sup>  
21 <sup>70,78</sup> The inter-membrane space of mitochondrion contains proteins such as apoptosis-  
22 inducing factor (AIF) and endonuclease G (EndoG), which can induce apoptotic cell death in  
23 a caspase-independent fashion.<sup>79</sup> EndoG is one of the predominant endonucleases that are  
24 involved in the regulation of cellular functions such as mitochondrial biogenesis, DNA  
25 synthesis and repair. AIF is an FAD-containing flavoprotein which plays an important role in

1 the stability of an electron transport chain.<sup>80</sup> Nutrient deficiency-mediated stress signals  
2 induce mitochondrial outer membrane permeabilization (MOMP), which consequently  
3 releases inter-membrane space proteins such as AIF, EndoG and cytochrome *c*. AIF plays a  
4 role of central mediator in caspase-independent cell death pathway.<sup>81</sup> AIF, once released into  
5 the cytosol, interacts with EndoG and cyclophilin A prior to its translocation into the  
6 nucleus.<sup>82</sup> Subsequently after translocation into the nucleus, it triggers cell death either  
7 directly, through interaction with DNA, or indirectly, through the production of reactive  
8 oxygen species.<sup>73,74,79,80</sup> MOMP promotes both, caspase-dependent and caspase-independent  
9 apoptotic pathways, but with different kinetics.<sup>83</sup> Although, the upstream signaling stimulus  
10 for both, a caspase-dependent and caspase-independent pathway is the same, *i.e.* via  
11 induction of MOMP, their downstream pathways are different. Moreover, nuclear alterations  
12 and the changes occurring in mitochondrial trans-membrane potential during caspase-  
13 independent pathways are different than those observed in a caspase-dependent apoptotic  
14 pathway.<sup>84</sup>

15 To summarize, growing evidence suggests that autophagy is a prevailing cell survival  
16 mechanism in tumor cells undergoing ADI-mediated arginine deprivation. The overall  
17 cellular response to ADI-mediated arginine deprivation in different tumor cells operates  
18 through a complex cascade, initiating with induction of autophagy and followed by the  
19 activation of either caspase-dependent or caspase-independent cell death pathways. It is  
20 worth emphasizing that the discrepancy of cellular responses of tumor cells to ADI-mediated  
21 arginine depletion in activation of either caspases-dependent or caspases-independent cell  
22 death pathways can vary depending on tumor cell type.<sup>38,39,70,71,74</sup> As a result, the precise  
23 mechanisms of tumor cell death- consequential of cellular response to ADI-mediated arginine  
24 depletion- appear to be complex and variable, and need to be further elucidated.

25

### 1 ***1.4.2 Inhibition of de novo protein synthesis by ADI-mediated arginine deprivation***

2 Inhibition of *de novo* protein synthesis is another mechanism which can be attributed  
3 to the anti-tumor potential of ADI. As extracellular arginine pool is responsible for 40% of *de*  
4 *novo* protein synthesis, ADI treatment to human lung carcinoma cells results in an anti-  
5 proliferative effect, mediated by inhibition of protein synthesis.<sup>85</sup> Arginine is present in  
6 various compartments such as extracellular, intracellular and citrulline-arginine regeneration  
7 *i.e.* cytosolic compartment and it is known to regulate various cellular pathways differently.  
8 Protein synthesis mainly utilizes arginine either from the intracellular pool or the citrulline-  
9 arginine regeneration mechanism, while polyamines synthesis largely utilizes arginine pool  
10 from the intracellular origin.<sup>86,87</sup> Polyamines are synthesized through the methionine salvage  
11 pathway via decarboxylation of S-adenosylmethionine (SAM). SAM is a donor metabolite  
12 necessary for the transfer of methyl group to DNA and proteins. Human colon cancer  
13 (HCT116) cells treated with short hairpin CD44 RNA interference showed a decrease in the  
14 total amount of methionine-pool metabolites including polyamines, suggesting the role of  
15 polyamines in cancer proliferation.<sup>88</sup>

16 ADI treatment towards human mammary adenocarcinoma and lung carcinoma cells  
17 differently modulates polyamine synthesis and the global protein synthesis. Interestingly,  
18 inhibition of protein synthesis has been correlated with the ASS-mediated regeneration of  
19 arginine. Cells expressing low levels of ASS (A549) result in decreased protein synthesis  
20 (without affecting polyamine synthesis) and those expressing higher ASS levels (MCF-7) are  
21 resistant to ADI treatment, as the decreased arginine levels can be replaced by citrulline-  
22 arginine regeneration pathway.<sup>85</sup>

### 23 ***1.4.3 Anti-angiogenic effects of ADI-mediated arginine deprivation***

24 As a tumor grows beyond a certain size (2 mm in diameter for most solid tumors),  
25 available vasculature within the tumor becomes inadequate to supply sufficient quantities of

1 essential nutrients for their growth.<sup>89</sup> This results in the generation of hypoxic tumor  
2 microenvironment and leads to the development of new blood vessels (angiogenesis) as a  
3 colossal requisite of the developing tumors.<sup>90</sup> Accordingly, neovascularization can be stated  
4 as one of the decisive phenomena during tumor growth and metastasis.<sup>91</sup> Emerging studies  
5 now indicate that not only molecular signals but also metabolic mechanisms regulate  
6 angiogenesis.<sup>92</sup> Under stress conditions such as hypoxia, tumor cells secrete angiogenic  
7 factors such as vascular endothelial growth factor (VEGF).<sup>93</sup> Increased levels of VEGF  
8 activate VEGF receptor 2 (VEGFR2) signaling in the quiescent endothelial cells which in  
9 turn initiate angiogenesis.<sup>94-96</sup> Endothelial cells produce 85% of their total amount of ATP via  
10 glycolysis. Addition of endothelial cells on anaerobic rather than aerobic pathway enables  
11 them for the formation of vascular sprouts in hypoxic areas.<sup>97,98</sup> Metabolism of tumor  
12 endothelial cells resembles that of highly activated endothelial cells because of the tumor  
13 induced switch from quiescence to proliferation due to metabolically regulated migration  
14 during sprouting.<sup>99,100</sup>

15 Besides ADI's role in modulation of apoptotic pathways, it has an anti-angiogenic  
16 activity that contributes to its anti-tumor potential. The growth, migration and differentiation  
17 of human umbilical vein endothelial cells (HUVECs) are strongly impaired in a medium  
18 containing recombinant ADI.<sup>101</sup> As a consequence; it results in decreased tube formation with  
19 intermittent and incomplete microvascular network. Similarly, Park *et al.* found that *E. coli*  
20 ADI inhibits angiogenesis by inhibiting tube formation of endothelial cells and  
21 neovascularization in Chick Chorioallantoic Membrane (CAM) and Matrigel plug assay.<sup>102</sup>

22 Suppression of nitric oxide (NO) generation is also another possible mechanism for  
23 anti-angiogenic activity of ADI. Since L-arginine is required for nitric oxide synthases  
24 (NOSs) to generate NO, the depletion of arginine by ADI suppresses NO synthesis.<sup>102</sup>  
25 Potential role of ADI-mediated arginine depletion in inhibition of NO synthesis has been

1 reported.<sup>103,104</sup> We and others have previously reported that NO promotes tumor growth  
2 through the stimulation of angiogenesis<sup>105-107</sup> and regulates cellular interaction by controlling  
3 adhesion molecule expression and ultimately cell adhesion.<sup>108,109</sup> NO directly, or indirectly  
4 through NO-mediated reactive nitrogen species (RNS), induces the activation of certain  
5 angiogenic signaling pathways in the endothelial cells.<sup>110</sup> NO acts as an autocrine mediator in  
6 endothelial cell functioning and as a final modulator in VEGF stimulated angiogenesis.<sup>109,111</sup>  
7 NO not only mediates angiogenesis but also subsequent vessel maturation<sup>112,113</sup> Moreover,  
8 NO is known to inhibit angiostatin and thrombospondin-1, two main inhibitors of  
9 angiogenesis.<sup>114</sup> Owing to the important role of NO in angiogenesis, ADI inhibits tumor  
10 growth not only by draining the supply of arginine, but also by its anti-angiogenic activity via  
11 suppression of NO generation.

12 To summarize, certain tumor cell types such as, HCCs and metastatic melanomas are  
13 invariably deficient in ASS expression and can be specifically targeted by ADI-mediated  
14 ADT. It is worth noting that more than one pathway may be attributed to the cytotoxic  
15 potential of ADI-mediated ADT [Figure 5]. The anti-tumor potential of ADI may not only be  
16 simply accredited to its action as arginine degrading enzyme but also to several other  
17 mechanisms important in the cellular functioning of tumor cells. Induction of apoptotic  
18 pathways, inhibition of angiogenesis and inhibition of *de novo* protein synthesis are the  
19 important mechanisms attributed to the cytotoxic potential of ADI. Moreover, studies have  
20 revealed the ADI-mediated modulations in tumor cell-cycle. The fundamental difference of  
21 cell cycle modulations in normal and malignant cells should be exploitable as a means of  
22 selective demise of tumor cells and ADI, in combination with other anti-cancer  
23 chemotherapeutic agents, which can be a potential strategy to improve chemo-sensitization  
24 against tumor cells.<sup>115-118</sup>

25

1

## 2 **2. Arginase**

3 Arginase (E.C.3.5.3.1) is a mammalian enzyme which catalyzes the conversion of  
4 arginine to ornithine and urea. Arginase is considered as an enzyme responsible for the cyclic  
5 nature of urea cycle, since only the organisms containing arginase are able to carry out the  
6 complete urea cycle.<sup>119</sup> Two distinct isoforms of mammalian arginase have been identified  
7 which are encoded by two separate genes.<sup>120</sup> Type I arginase (arginase I) is located in the  
8 cytosol and is mainly expressed in liver. Type II arginase is located in the mitochondrial  
9 matrix and is expressed in extra-hepatic tissues.<sup>121,122</sup> Intracellular regulation of arginase  
10 expression is of immense importance as it has crucial implications for the synthesis of  
11 essential cellular metabolites,<sup>123</sup> For example, cytosolic co-localization of arginase I with  
12 ornithine decarboxylase (ODC) preferentially utilizes ornithine for the biosynthesis of  
13 polyamine. On the other hand, due to its co-localization with ornithine aminotransferase  
14 (OAT) in the mitochondria, arginase II directs ornithine for the production of proline and  
15 glutamine.<sup>124,125</sup>

### 16 **2.1 PEGylated recombinant human arginase I**

17 Elevated requirements of arginine by tumor cells were first identified in 1947 and  
18 preferential utilization of arginine by tumor bearing animals was revealed in 1953.<sup>126,127</sup> The  
19 use of bovine and murine arginase in arginine deprivation therapy was prevailing until the  
20 advent of recombinant DNA technology,<sup>128-130</sup> followed by the pervasive use of recombinant  
21 human arginase in subsequent decades.<sup>131,132</sup> Arginase from bovine and murine sources has  
22 been extensively used for the arginine deprivation therapy *in vitro*. However, limited success  
23 was achieved *in vivo* due to its alkaline optimum pH and very low affinity for the substrate.  
24 Human arginase I also has a serious limitation of very short circulatory half-life (Approx. 30  
25 minutes).



1 To extend plasma half-life of arginase, PEGylation has been **applied successfully**.  
2 PEGylated recombinant human arginase I (rhArg-Peg<sub>5000mw</sub>) had efficient catalytic activity at  
3 physiological pH with improved *in vivo* half-life of 3 days. Furthermore, rhArg-Peg<sub>5000mw</sub>  
4 was found to have significant tumor inhibitory activity in BALB/c nude mice bearing HCC  
5 xenografts.<sup>131</sup> **Notably**, these results were consistent with those demonstrated by Tsui and co-  
6 workers.<sup>133</sup> Recently, a bio-engineered form of human arginase I was developed by the co-  
7 factor replacement, the replacement of two Mn<sup>2+</sup> ions by Co<sup>2+</sup> ions. The modified Co<sup>2+</sup>-  
8 arginase I resulted in 10-fold increase in the catalytic activity and 5-fold greater stability at  
9 the physiological pH. Nevertheless, IC<sub>50</sub> values for killing human HCC and melanoma cell  
10 lines were lowered by 12-15 folds.<sup>134</sup> More recently, modifications in bioengineered Co<sup>2+</sup>-  
11 arginase I were performed by conjugating 5-kDa PEG to enhance plasma half-life. This  
12 modified version of bioengineered arginase I (Co-hArgI-PEG) was proven to be cytotoxic by  
13 significantly increasing the expression of caspases-3 in HCC and pancreatic carcinoma (PC)  
14 tumor xenografts.<sup>135</sup> **Lately**, the cytotoxic potential of Co-hArgI-PEG was identified in acute  
15 myeloid leukemia (AML) and glioblastoma cells. AML cell lines were found sensitive  
16 towards Co-hArgI-PEG-mediated arginine deprivation with very low (58-722 PM) IC<sub>50</sub>  
17 values, suggesting a very high potential of Co-hArgI-PEG-mediated arginine depletion in  
18 AML cells.<sup>136</sup> Moreover, Co-hArgI-PEG-mediated arginine deprivation has been  
19 demonstrated to induce caspase-independent, non-apoptotic cell death in human glioblastoma  
20 cells.<sup>137</sup> **Alternative** method to extend the plasma half-life of recombinant human arginase  
21 **also** has been established. Plasma half-life of a fusion protein form of a recombinant human  
22 arginase (rhArg-Fc, constructed by linking rhArg to the Fc region of human immunoglobulin  
23 IgG1), **was** evidenced to significantly extend up-to approx. 4 days.<sup>138</sup> In addition, rhArg-Fc  
24 was confirmed to conspicuously inhibit the cell growth of human HCC cells *in vitro* and *in*  
25 *vivo*.<sup>138</sup>

1 Last decade has evidenced a prevalent use of recombinant human arginase-mediated  
2 ADT in numerous cancer cell types, mainly metastatic HCC and melanomas.<sup>131,139,140</sup>  
3 Currently, PEGylated derivative of recombinant human arginase I is undergoing clinical trials  
4 for the treatment of human HCC.<sup>141,142</sup> Moreover, initiatives are now being taken to  
5 overcome the possible problem of accumulation of PEGylated products in the liver by  
6 impending approaches such as fusion proteins.<sup>138</sup>

## 7 **2.2 Anti-tumor mechanisms of arginase-mediated arginine deprivation**

8 Selective starvation of L-arginine in tumor cells, which are auxotrophic for L-  
9 arginine, is one of the most important anti-tumor mechanisms of ADT. Arginase can render  
10 its cytostatic effect as a result of modulations in the cell cycle proteins, whereas, cytotoxic  
11 effects rendered by arginase I-mediated arginine deprivation have been proposed as a result  
12 of induction of potential cell death pathways namely apoptosis and probably by ‘autophagic  
13 cell death’. Summarized below are the current understandings of the molecular mechanisms  
14 of cytostatic and cytotoxic effects rendered by arginase-mediated ADT.

### 15 ***2.2.1 Role of autophagy in arginase-mediated arginine deprivation***

16 Autophagy is a key sensing and regulatory mechanism of cells in nutrient deprived  
17 conditions. Under stress conditions, autophagy functions as a bio-energy management  
18 system by recycling cell organelles and damaged and/or long-lived proteins.<sup>143</sup> Although  
19 autophagy seems to be a survival mechanism of the cells, there is a growing evidence of  
20 accumulation of autophagosomes and other autophagic markers in dying cells unable to  
21 process apoptosis, raising the term ‘autophagic cell death’.<sup>144-147</sup> However, the term  
22 ‘autophagic cell death’ is based on morphological features rather than the causative role of  
23 autophagy in cell death. New definition of ‘autophagic cell death’ has been proposed,

1 implying that cell death must occur without the involvement of apoptotic machinery,  
2 (caspase activation) but **with** an increase in autophagic flux.<sup>148,149</sup>

3 Mammalian target of rapamycin (mTOR) is a key regulator **of** coupling cell growth  
4 and nutritional status of the cell.<sup>150,151</sup> Autophagy is induced by the inhibition of mTOR  
5 signaling pathway.<sup>152</sup> During nutrient affluent conditions, mTOR is involved in the negative  
6 regulation of Atg1 (autophagy related gene 1) **which** inhibits autophagy.<sup>153,154</sup> Arginase-  
7 mediated arginine deprivation leads to decreased levels of ATP, which in turn activates the  
8 adenosine 5'-monophosphate-activated protein kinase (AMPK). Activated AMPK  
9 eventually inhibits the mTOR-signaling pathway, manifested by the reduced  
10 phosphorylation of key downstream molecules, such as 4E-BP1 (Eukaryotic translation  
11 initiation factor 4E-binding protein-1). Dephosphorylation of 4E-BP1 is observed in Chinese  
12 hamster ovary (CHO), human melanoma cells and human prostate cancer cells following  
13 their exposure to recombinant human arginase I.<sup>65,155,156</sup> Phagosome/lysosome activity is  
14 also significantly increased following an incubation of human tumor cells in L-arginine  
15 deficient medium.<sup>157</sup> Additionally, studies carried out by Hsueh *et al.*<sup>156</sup> evidenced no  
16 significant induction of apoptotic mechanisms in prostate cells after their exposure to  
17 rhArgI, suggesting the role of autophagic cell death, rather than apoptosis, as an alternative  
18 cell death mechanism. In addition, autophagy has often accompanied damaged mitochondria  
19 and higher levels of reactive oxygen species (ROS).<sup>158,159</sup> Acute generation of ROS has been  
20 attributed to causing severe damages to the cellular macromolecules, which in consequence,  
21 leads to necrosis of the tumor cells.<sup>160,161</sup> Overall, arginase leads to deprivation of arginine,  
22 in consequence, it inhibits mTOR pathway during the deprivation and thus forcing tumor  
23 cells to undergo 'autophagic cell death' pathway.<sup>162</sup>

24 **SLC38A9, a member 9 of the solute carrier family 38, has been recently identified as**  
25 **an integral component of the lysosomal machinery that controls amino acid-induced mTOR**

1 activation.<sup>163,164</sup> Amino acid starvation in human embryonic kidney (HEK293T) cells with  
2 stable expression of SLC38A9 has been shown to activate mTOR in a sustained manner.  
3 Moreover, shRNA-mediated silencing of SLC38A9 results in a reduction of arginine-  
4 induced mTOR activation. Also, depletion of SLC38A9 impaired mTOR activation induced  
5 by cycloheximide (a protein synthesis inhibitor which induces accumulation of intracellular  
6 amino acids), further suggests the role of SLC38A9 in mTOR activation at the lysosomal  
7 rather than at the plasma membrane. These studies have demonstrated that SLC38A9 acts as  
8 an upstream positive regulator for mTOR functioning and thereby modulating autophagy in  
9 arginine-deprived tumor cells.

10 Although some studies have advocated autophagy as a cell death mechanism of  
11 arginase-mediated ADT,<sup>156,157</sup> many groups have explained it as a pro-survival mechanism;  
12 mainly by postponing the activation of apoptosis.<sup>38,161</sup> Thus, understanding the exact role of  
13 autophagy in arginase-mediated cell death pathways is a complicated episode.<sup>162,165</sup>  
14 Therefore, much need to be elucidated about these new findings related to ‘autophagic cell  
15 death’ and caution must be taken to assign autophagy as a cell death pathway in arginase-  
16 mediated ADT.

### 17 ***2.2.2 Role of apoptosis in arginase-mediated arginine deprivation***

18 The role of autophagy, either in cell survival or in cell death, depends on many factors  
19 such as cell type, nature and severity of the stimuli and so on.<sup>166</sup> If the attempt of the cells to  
20 survive through autophagy fails, apoptotic pathways take over and ultimately cause cell  
21 death.<sup>143</sup> Inhibition of autophagy in amino acid deficient conditions induces tumor cell  
22 death, mainly because of further exacerbation of energy dearth.<sup>167,168</sup> Also, longer  
23 persistence of autophagy is proposed to eventually lead the activation of caspase-dependent  
24 cell death pathways, as autophagy and apoptotic cell death pathways are interconnected and

1 also share some common pathways through the induction of the membrane permeability  
2 transitions.<sup>169-171</sup> Induction of apoptotic pathways is another consequence of arginine  
3 depletion and anti-tumor mechanism of arginase I-mediated arginine deprivation.

4 Involvement of apoptosis as a cell death mechanism in arginase-mediated ADT has  
5 been illustrated in various literature reports. Annexin V is known to selectively stain the  
6 cells, which are destined for apoptosis or in the process of apoptosis. 33% of human  
7 melanoma cell population was destined for apoptotic cell death following rhArg  
8 treatment.<sup>139</sup> Arginase I-mediated arginine deprivation led to the transcriptional up-  
9 regulation of *caspase 3*, the intrinsic mitochondrial pathway of apoptosis, which is marked  
10 by the change in mitochondrial membrane potential.<sup>172</sup> Recently, an anti-leukemic potential  
11 of PEGylated-arginase has been attributed to kinases general control nonderepressible 2  
12 (GCN2)-mediated induction of apoptosis in T-ALL cells.<sup>173</sup>

### 13 ***2.2.3 Cell cycle arrest by arginase-mediated arginine deprivation and combination*** 14 ***approaches***

15 rhArg-Peg<sub>5000mw</sub>-mediated arginine deprivation in various HCC cells results in their  
16 cell cycle arrest at G<sub>2</sub>/M phase, by decreased expression levels of cyclin B1 and cdc2, or in S  
17 phase, by a transcriptional up-regulation of cyclin A1 [Ref. 140]. rhArg-Peg<sub>5000mw</sub>-mediated  
18 arginine depletion was witnessed to impair the expression of cyclin D3 in T-ALL cells, which  
19 was followed by an arrest of the cells in the G<sub>0</sub>-G<sub>1</sub> phase of the cell cycle and induction of  
20 apoptosis.<sup>172</sup> Recent investigations of rhArg-Fc-mediated arginine deprivation in human HCC  
21 cells exhibited cell cycle arrest at S phase.<sup>138</sup> The exact mechanisms of these findings are still  
22 elusive, but the possible reasons seem to be the increased expression of cyclin A and declined  
23 transcription levels of p27 and p21 (the key cyclin kinase inhibitors).

1 Owing to the evidence of cell cycle arrest, a combination of arginase and other cell-  
2 cycle specific anti-cancer chemotherapeutics as potential anti-tumor approaches have been  
3 established. Synergistic effects of rhArg-Peg<sub>5000mw</sub> with 5-fluorouracil (5-FU, uracil analog  
4 which interferes with RNA and DNA synthesis) and cytarabine (Ara-C, anti-metabolic  
5 chemotherapeutic agent) have been investigated on the inhibition of proliferation of HCC and  
6 T-ALL cells, respectively.<sup>131,172</sup> Treatment of either rhArg-Peg<sub>5000mw</sub> or Ara-C alone induces  
7 a heterogeneous anti-tumor effect *in vivo*, whereas, combined treatment of rhArg-Peg<sub>5000mw</sub>  
8 and Ara-C induces a homogenous prevention of spleen growth, leading to the prolonged  
9 survival in all of the T-ALL bearing mice.<sup>172</sup> Moreover, combined treatment of PEGylated  
10 recombinant human arginase I and oxaliplatin has been demonstrated to synergize the  
11 inhibiting effect on tumor growth and enhanced overall survival probability as compared to  
12 PEGylated recombinant human arginase I or oxaliplatin treatment alone.<sup>174</sup>

13 Altogether, arginase has an advantage over ADI that it is efficacious in both ASS-  
14 negative and OTC- negative tumors,<sup>59</sup> whereas ADI is efficacious only in ASS-negative  
15 tumors. The tumor cell types expressing ASS are resistant to arginine deprivation treatment  
16 by ADI.<sup>25,26,54,61,131</sup> Even though arginase has been considered as a potential drug candidate  
17 over a period of six decades, low substrate specificity (high  $k_m$  of 2-4 mM), short plasma life  
18 and optimum alkaline pH (pH 9.3) limit *in vivo* applications of arginase.<sup>131,140</sup> In addition,  
19 robust homeostatic mechanisms in the body allow faster restoration of plasma free arginine,  
20 making *in vivo* arginine deprivation by arginase more difficult. Most of the scientific efforts  
21 nowadays pay attention to these limiting characteristics of arginase.<sup>134,175,176</sup>

### 22 3. Arginine decarboxylase

23 Arginine decarboxylase (ADC) (E.C. 4.1.1.19) metabolizes arginine to agmatine, one  
24 of the minor metabolic products of arginine. ADC is mainly found in plants, bacteria and  
25 mammalian liver and brain membranes.<sup>177,178</sup> The mammalian ADC is different from other

1 sources and distinct but related to ODC.<sup>179</sup> Although, arginine decarboxylation by ADC is a  
2 minor metabolic route, its product i.e. agmatine has a significant role in numerous cellular  
3 pathways.<sup>180</sup> Agmatine modulates the polyamine metabolism through its negative interaction  
4 with ODC.<sup>181</sup> Agmatine also confers an inhibitory effect on intracellular polyamine content  
5 by inhibiting polyamine uptake<sup>182</sup> and probably by increased polyamine catabolism.<sup>183</sup>  
6 Mayeur *et al.*,<sup>184</sup> has reported the effect of agmatine accumulation on polyamine metabolism,  
7 cell proliferation and cell cycle distribution in human colon adenocarcinoma epithelial cell  
8 lines. Due to the agmatine-mediated reduction in polyamine synthetic capacity of the cells,  
9 agmatine markedly inhibits the cell proliferation of HT-29 and Caco-2 cells in a dose  
10 dependent manner, without affecting cell membrane integrity. Moreover, agmatine modulates  
11 the cell cycle progression by decreasing ODC activity and expression.<sup>181,185</sup> As ODC plays an  
12 important role in the G<sub>1</sub>/S progression of the cells, agmatine-mediated modulations in ODC  
13 expression lead to modifications in the cell cycle progression.<sup>186</sup> Additionally, agmatine also  
14 has been shown to delay the expression of cyclins in tumor cells, leading to the modifications  
15 in the cell cycle progression.<sup>184</sup>

16 ADC has been investigated for the enzymatic degradation of arginine in normal and  
17 malignant cell cultures.<sup>187</sup> Arginine deprivation in human diploid fibroblasts (normal cells),  
18 achieved using human recombinant ADC, resulted in the cell cycle arrest at G<sub>1</sub>/G<sub>0</sub>. While  
19 treatment of 0.1 unit ml<sup>-1</sup> ADC to HeLa (Human cervical cancer) cells resulted in cell cycle  
20 arrest with an initiation of cell death after 2 days.<sup>187</sup> Similar results were evidenced in the  
21 studies by Wheatley *et al.*,<sup>188</sup> where 5 units ml<sup>-1</sup> ADC was found as effective as arginase in  
22 the inhibition of HeLa cells and cell cycle arrest at G<sub>1</sub> (quiescence) in fibroblasts.

23 Although some research groups have exhibited ADC as a potential anti-tumor  
24 enzyme, only a few reports are available to support this fact [Table 1].<sup>187,188</sup> Even though  
25 ADC possesses low *K<sub>m</sub>* and can degrade arginine very rapidly, the serious problem is related

1 to its product i.e. agmatine. Agmatine is toxic to normal cells when its concentration reaches  
2 a millimolar level, particularly when free arginine levels are low. Additionally, agmatine is  
3 not converted back to arginine under normal physiological conditions, which may lead to its  
4 accumulation and toxicity to normal cells.<sup>189</sup> Though recombinant human ADC expressed in  
5 *E. coli* has been evidenced more active than Sigma enzymes prepared from other sources, its  
6 PEGylation has been shown to result in the loss of its entire activity.<sup>187,189</sup> To consider [the](#)  
7 further rational use of this prospective enzyme as potential anti-cancer modality, it clearly  
8 warrants further evaluation [[Table 3](#)].

### 9 **Concluding remarks**

10 Sufficient evidence has been accumulated indicating that arginine catabolic enzymes-  
11 based approaches may be an effective way to target malignant cells. These enzymes control  
12 tumor cell proliferation as well as make them highly vulnerable to cell-cycle specific  
13 chemotherapeutic agents. This combinatorial approach is one of the potential strategies to  
14 maximize the efficacy to obliterate the tumor cells. Extensive research of the arginine  
15 metabolic pathways led to the establishment of arginine-depriving enzymes as a potential  
16 anti-cancer strategy against arginine auxotrophic tumors. However, many of these enzymes  
17 can be co-expressed in the cells, which results in complex interactions. For example, arginine  
18 is a common substrate for arginase as well as NOS. [The specific role of NO, either in](#)  
19 [inhibition or induction of cell proliferation is dependent on numerous factors like its](#)  
20 [interaction with other free radicals, cellular makeup, tumor milieu, proteins present the](#)  
21 [cellular microenvironment and also upon the chemical and biological heterogeneity of NO.](#)  
22 [NO has been known to demonstrate bipolar cellular effects and often termed as “double-](#)  
23 [edged sword”.](#) Although, NOS remains a viable candidate for cancer treatment, the precise  
24 [role of NO in the tumor microenvironment is extremely complex and conflicting.](#) Also, the  
25 [preferential utilization of arginine by arginase and/or NOS pathway is not fully understood.](#)



1 Thus, many of these pathways warrant further research to understand the arginine metabolism  
2 at cellular and molecular levels involving upstream and downstream pathways of the  
3 enzymes involved.

4 It should be noted that modulation of the immunological responses is one of the major  
5 roles of arginine availability. Arginine metabolism in myeloid-derived suppressor cells via  
6 arginase and/or NOS markedly impairs the T-cell responses that would eradicate and remove  
7 tumor cells.<sup>190</sup> Many excellent articles are available which focus on the role of arginine in  
8 immunological aspects of the tumors.<sup>191-194</sup> It would suffice to say here that the arginine  
9 deprivation therapy may have further anti-tumor effect through restoration of anti-tumor  
10 immunity.

11 Arginine dependence of the tumor cells has been considered as the “*Achilles heel*” of  
12 tumor cells.<sup>195</sup> Inability of tumor cells to proliferate in the absence of arginine can be targeted  
13 for their selective destruction by arginine depriving enzymes. Large numbers of enzyme-  
14 based anti-cancer therapies are currently undergoing clinical evaluation. It is encouraging that  
15 arginase and arginine deiminase already have achieved considerable success, without causing  
16 detrimental side effects and with high tolerability.<sup>51,63,141</sup> The knowledge acquired about the  
17 PEGylation has helped in the generation of adducts of potential value, overcoming the  
18 serious limitations of the anti-cancer enzymes of the non-human origin. The approach of  
19 enzyme-mediated arginine deprivation therapy is highly challenging, however rewarding  
20 upon success due to the provision of overturning the cancer dogma.

## 21 **Acknowledgments**

22 MP gratefully acknowledges Department of Biotechnology (DBT), New Delhi, India  
23 for the award of Senior Research Fellowship. Authors are also thankful to Prof. Rakesh K.  
24 Jain, Department of Radiation Oncology, Massachusetts General Hospital, Harvard Medical

1 School, USA; Dr. Utpal Mohan, Department of Biotechnology, NIPER, Guwahati, India and  
2 Dr. Umesh Patil, School of Chemical Sciences, North Maharashtra University, Maharashtra,  
3 India for the assistance provided during preparation of the manuscript. We apologize to those  
4 authors whose work could not be cited owing to space limitations.

5

## 6 **References**

- 7 1) Wu G. Functional amino acids in nutrition and health. *Amino Acids* 2013; **45**: 407–  
8 411.
- 9 2) F. Loreni, M. Mancino, S. Biffo, Translation factors and ribosomal proteins control  
10 tumor onset and progression: how? *Oncogene* 2014; **33**: 2145-2156.
- 11 3) Proud CG. Control of the translational machinery by amino acids. *Am J Clin Nutr*  
12 2014; **99**: 231s-236s.
- 13 4) Luo J-Q, Chen D-W, Yu B. Upregulation of amino acid transporter expression  
14 induced by l-leucine availability in L6 myotubes is associated with ATF4 signaling  
15 through mTORC1-dependent mechanism. *Nutr* 2013; **29**: 284-90.
- 16 5) Palii SS, Kays CE, Deval C, Bruhat A, Fafournoux P, Kilberg MS. Specificity of  
17 amino acid regulated gene expression: analysis of genes subjected to either complete  
18 or single amino acid deprivation. *Amino Acids* 2009; **37**: 79-88.
- 19 6) Qie S, Liang D, Yin C, Gu W, Meng M, Wang C *et al.* Glutamine depletion and  
20 glucose depletion trigger growth inhibition via distinctive gene expression  
21 reprogramming. *Cell Cycle* 2012; **11**: 3679-3690.
- 22 7) Agrawal V, Alpini SEJ, Stone EM, Frenkel EP, Frankel AE. Targeting methionine  
23 auxotrophy in cancer: discovery & exploration. *Expert Opin Biol Ther* 2012; **12**: 53-  
24 61.

- 1 8) Wu G, Wu Z, Dai Z, Yang Y, Wang W, Liu C *et al.* Dietary requirements of  
2 “nutritionally non-essential amino acids” by animals and humans. *Amino Acids*  
3 2013; **44**: 1107–1113
- 4 9) Fu YM, Yu Z-X, Li Y-Q, Ge X, Sanchez PJ, Fu X *et al.* Specific amino acid  
5 dependency regulates invasiveness and viability of androgen-independent prostate  
6 cancer cells. *Nutr Cancer* 2003; **45**: 60–73.
- 7 10) Icard P, Lincet H. A global view of the biochemical pathways involved in the  
8 regulation of the metabolism of cancer cells. *Biochim Biophys Acta Rev Cancer* 2012;  
9 **1826**: 423-433.
- 10 11) Cantor JR, Sabatini DM. Cancer cell metabolism: One hallmark, many faces. *Cancer*  
11 *Disc* 2012; **2**: 881-898.
- 12 12) Locasale JW, Cantley LC. Metabolic flux and the regulation of mammalian cell  
13 growth, *Cancer Cell* 2011; **14**: 443-451.
- 14 13) Ferreira LMR, Hebrant A, Dumont JE. Metabolic reprogramming of the tumor.  
15 *Oncogene* 2012; **31**: 3999-4011.
- 16 14) Yamamoto T, Takano N, Ishiwata K, Ohmura M, Nagahata Y, Matsuura T, *et al.*  
17 Reduced methylation of PFKFB3 in cancer cells shunts glucose towards the pentose  
18 phosphate pathway. *Nat Commun* 2014; **5**: DOI: 10.1038/ncomms4480.
- 19 15) Matthew G, Heiden V. Targeting cancer metabolism: a therapeutic window opens.  
20 *Nat Rev Drug Disc* 2011; **10**: 671-684.
- 21 16) Ward PS, Thompson CB. Metabolic reprogramming: a cancer hallmark even Warburg  
22 did not anticipate. *Cancer Cell* 2012; **21**: 297-308.
- 23 17) Gelb T, Pshenichkin S, Rodriguez OC, Hathaway HA , Grajkowska E, DiRaddo JO *et*  
24 *al.* Metabotropic glutamate receptor 1 acts as a dependence receptor creating a

- 1 requirement for glutamate to sustain the viability and growth of human melanomas.  
2 *Oncogene* 2015; **34**:2711-2720.
- 3 18) Cetinbas N, Daugaard M, Mullen AR, Hajee S, Rotblat B, Lopez A et al. Loss of the  
4 tumor suppressor Hac1 leads to ROS-dependent glutamine addiction. *Oncogene*  
5 2015; **34**:4005-4010.
- 6 19) Graham ML. Pegaspargase:a review of clinical studies. *Adv Drug Deliv Rev* 2003; **55**:  
7 1293-1302.
- 8 20) Durden DL, Distasio JA. Characterization of the effects of asparaginase from  
9 *Escherichia coli* and a glutaminase-free asparaginase from *Vibrio succinogenes* on  
10 specific cell-mediated cytotoxicity. *Int J cancer* 1981; **27**: 59-65.
- 11 21) Dodd KM, Tee AR. Leucine and mTORC1: a complex relationship. *Am J Physiol*  
12 *Endocrinol Metab* 2012; **302**: E1329–E1342.
- 13 22) Takano N, Sarfraz Y, Gilkes DM, Chaturvedi P, Xiang L, Suematsu M, et al. *Mol*  
14 *Cancer Res* 2014; **12**: 1398-1406.
- 15 23) Wu G, Bazer FW, Davis TA, Kim SW, Li P, Rhoads JM et al. Arginine metabolism  
16 and nutrition in growth, health and disease. *Amino Acids* 2009; **37**: 153–168.
- 17 24) Appleton J. Arginine: Clinical potential of a semi-essential amino acid. *Altern Med*  
18 *Rev* 2007; **7**: 512-522.
- 19 25) Feun LG, Marini A, Walker G, Elgart G, Moffat F, Rodgers SE et al. Negative  
20 argininosuccinate synthetase expression in melanoma tumors may predict clinical  
21 benefit from arginine-depleting therapy with pegylated arginine deiminase. *Br J*  
22 *Cancer* 2012; **106**: 1481-1485.
- 23 26) Delage B, Luong P, Maharaj L, O’Riain C, Syed N, Crook T et al. Promoter  
24 methylation of argininosuccinate synthetase-1 sensitises lymphomas to arginine

- 1 deiminase treatment, autophagy and caspase-dependent apoptosis. *Cell Death Dis*  
2 2012; **3**: e342.
- 3 27) Wheatley DN. Controlling cancer by restricting arginine availability-arginine  
4 catabolizing enzymes as anticancer agents. *Anti-Cancer Drugs* 2004; **15**: 825-833.
- 5 28) García-Navas R, Munder M, Mollinedo F. Depletion of L-arginine induces autophagy  
6 as a cytoprotective response to endoplasmic reticulum stress in human T  
7 lymphocytes. *Autophagy* 2012; **8**: 1557–1576
- 8 29) Synakiewicz A, Stachowicz-Stencel T, Adamkiewicz-Drozynska E. The role of  
9 arginine and the modified arginine deiminase enzyme ADI-PEG 20 in cancer therapy  
10 with special *emphasis* on phase I/II clinical trials. *Expert Opin Inv Drug* 2014; **23**:  
11 1517-1529.
- 12 30) Dillon BJ, Holtsberg FW, Ensor CM, Bomalaski JS, Clark MA. Biochemical  
13 characterization of the arginine degrading enzymes arginase and arginine deiminase  
14 and their effect on nitric oxide production. *Med Sci Monit* 2002; **8**: BR248- BR253.
- 15 31) [Qui F, Huang J, Sui M. Targeting arginine metabolism pathway to treat arginine-](#)  
16 [dependent cancers. \*Cancer Lett\* 2015; \*\*364\*\*: 1-7.](#)
- 17 32) [Phillips MM, Sheaff MT, Szlosarek PW. Targeting arginine-dependent cancers with](#)  
18 [arginine-degrading enzymes: Opportunities and challenges. \*Cancer Res Treat\* 2013;](#)  
19 [45: 251-262.](#)
- 20 33) Noh EJ, Kang SW, Shin YJ, Choi SH, Kim CG, Park IS *et al.* Arginine deiminase  
21 enhances dexamethasone-induced cytotoxicity in human T-lymphoblastic leukemia  
22 CCRF-CEM cells. *Int J Cancer* 2004; **112**: 502-508.
- 23 34) Stasyk OV, Boretsky YR, Gonchar MV, Sibirny AA. Recombinant arginine-  
24 degrading enzymes in metabolic anticancer therapy and bioanalytics. *Cell Biol Int*  
25 2015; **39**:246-252. DOI: 10.1002/cbin.10383

- 1 35) Feun LG, Kuo MT Savaraj N. Arginine deprivation in cancer therapy. *Curr Opin Clin*  
2 *Nutr* 2015; **18**: 78–82. doi: 10.1097/MCO.000000000000122
- 3 36) Shirai H, Blundell TL, Mizuguchi K. A novel superfamily of enzymes that catalyse  
4 the modification of guanidine groups. *Trends Biochem Sci* 2001; **26**: 465-468.
- 5 37) Huang H-Y, Wu H-Y, Wang Y-H, Wang J-W, Fang F-M, Tsai J-W *et al.* ASS1 as a  
6 novel tumor suppressor gene in myxofibrosarcomas: Aberrant loss via epigenetic  
7 DNA methylation confers aggressive phenotypes, negative prognostic impact, and  
8 therapeutic relevance. *Clin Cancer Res* 2013; **19**: 2861–2872.
- 9 38) Savaraj N, You M, Wu C, Wangpaichitr M, Kuo MT, Feun LG. Arginine deprivation,  
10 autophagy, apoptosis (AAA) for the treatment of melanoma. *Curr Mol Med* 2010; **10**:  
11 405-412.
- 12 39) Kelly MP, Jungbluth AA, Wu B-W, Bomalaski J, Old LJ, Ritter G. Arginine  
13 deiminase PEG20 inhibits growth of small cell lung cancers lacking expression of  
14 argininosuccinate synthetase. *Br J Cancer* 2012; **106**: 324-32.
- 15 40) Manca A, Sini MC, Izzo F, Ascierio P, Tatangelo F, Botti G *et al.* Induction of  
16 arginosuccinate synthetase (ASS) expression affects the antiproliferative activity of  
17 arginine deiminase (ADI) in melanoma cells. *Oncol Rep* 2011; **25**: 1495-1502.
- 18 41) Jungbluth AA, Tassello J, Frosina D, Hanson N, Ritter G, Wu B-W *et al.* Expression  
19 pattern of Argininosuccinate-Synthetase (ASS) in normal and tumor tissue as a  
20 marker for susceptibility to Arginine-Deiminase (ADI) therapy. *Mod Pathol* 2010; **23**  
21 (Suppl 1): 387A.
- 22 42) Savaraj N, Wu C, Li YY, Wangpaichitr M, you M, Bomalaski J. Targeting  
23 argininosuccinate synthetase negative melanomas using combination of arginine  
24 degrading enzyme and cisplatin. *Oncotarget* 2015; **6**: 6295-6309.

- 1 43) Miyazaki K, Takaku H, Umeda M, Fujita T, Huang W, Kimura T, et al. Potent growth  
2 inhibition of human tumor cells in culture by arginine deiminase purified from a  
3 culture medium of a *Mycoplasma*-infected cell line. *Cancer Res* 1990; **50**: 4522-4527.
- 4 44) Sugimura K, Ohno T, Kusuyama T, Azuma I. High sensitivity of human melanoma  
5 cell lines to the growth inhibitory activity of mycoplasmal arginine deiminase *in vitro*.  
6 *Melanoma Res* 1992; **2**: 191-196.
- 7 45) Takaku H, Misawa S, Hayashi H, Miyazaki K. Chemical modification by  
8 polyethylene glycol of the anti-tumor enzyme arginine deiminase from *Mycoplasma*  
9 *arginini*. *Jpn J Cancer Res* 1993; **84**: 1195-1200.
- 10 46) Holtsberg FW, Ensor CM, Steiner MR, Bomalaski JS, Clark MA. Poly (ethylene  
11 glycol) (PEG) conjugated arginine deiminase: effects of PEG formulations on its  
12 pharmacological properties. *J Control Release* 2002; **80**: 259-271.
- 13 47) Ensor CM, Holtsberg FW, Bomalaski JS, Clark MA. Pegylated arginine deiminase  
14 (ADI-SS PEG20,000mw) inhibits human melanomas and hepatocellular carcinomas  
15 *in vitro* and *in vivo*. *Cancer Res* 2002; **62**: 5443-5450.
- 16 48) Zhang L, Liu M, Jamil S, Han R, Xu G, Ni Y. PEGylation and pharmacological  
17 characterization of a potential anti-tumor drug, an engineered arginine deiminase  
18 originated from *Pseudomonas plecoglossicida*. *Cancer Lett* 2015; **357**: 346-354.
- 19 49) Ni Y, Schwaneberg U, Sun ZH. Arginine deiminase, a potential anti-tumor drug.  
20 *Cancer Lett* 2008; **261**: 1-11.
- 21 50) Yoon J, Frankel AE, Feun LG, Ekmekcioglu S, Kim KB. Arginine deprivation  
22 therapy for malignant melanoma. *J Clin Pharmacol* 2013; **5**: 11-19.
- 23 51) Glazer ES, Piccirillo M, Albino V, Di Giacomo R, Palaia R, Mastro AA *et al*. Phase  
24 II study of pegylated arginine deiminase for nonresectable and metastatic  
25 hepatocellular carcinoma. *J Clin Oncol* 2010; **28**: 2220-2226.

- 1 52) Ascierto PA, Scala S, Castello G, Daponte A, Simeone E, Ottaiano A *et al.* Pegylated  
2 arginine deiminase treatment of patients with metastatic melanoma: results from  
3 phase I and II studies. *J Clin Oncol* 2005; **23**: 7660–7668.
- 4 53) Kobayashi E, Masuda M, Nakayama R, Ichikawa H, Satow R, Shitashige M *et al.*  
5 Reduced argininosuccinate synthetase is a predictive biomarker for the development  
6 of pulmonary metastasis in patients with osteosarcoma. *Mol Cancer Ther* 2010; **9**:  
7 535-544.
- 8 54) Syed N, Langer J, Janczar K, Singh P, Lo Nigro C, Lattanzio L *et al.* Epigenetic  
9 status of argininosuccinate synthetase and argininosuccinate lyase modulates  
10 autophagy and cell death in glioblastoma. *Cell Death Dis* 2013; **4**: e458,  
11 DOI:10.1038/cddis.2012.197.
- 12 55) Nicholson LJ, Smith PR, Hiller L, Szlosarek PW, Kimberley C, Sehouli J *et al.*  
13 Epigenetic silencing of argininosuccinate synthetase confers resistance to platinum-  
14 induced cell death but collateral sensitivity to arginine auxotrophy in ovarian cancer.  
15 *Int J Cancer* 2009; **125**: 1454-1463.
- 16 56) Szlosarek PW, Klabatsa A, Pallaska A, Sheaff M, Smith P, Crook T *et al.* *In vivo* loss  
17 of expression of argininosuccinate synthetase in malignant pleural mesothelioma is a  
18 biomarker for susceptibility to arginine depletion. *Clin Cancer Res* 2006; **12**: 7126-  
19 7131.
- 20 57) Rabinovich S, Adler L, Yizhak K, Sarver A, Silberman A, Agron S *et al.* Diversion of  
21 aspartate in ASS1-deficient tumours fosters *de novo* pyrimidine synthesis. *Nature*  
22 2015; **527**: 379-383. DOI:10.1038/nature15529.
- 23 58) Feun L, Savaraj N. Pegylated arginine deiminase: A novel anticancer enzyme agent.  
24 *Expert Opin Investig Drugs* 2006; **15**: 815-822.



- 1 59) Bobak YP, Vynnytska BO, Kurlishchuk YV, Sibirny AA, Stasyk OV. Cancer cell  
2 sensitivity to arginine deprivation *in vitro* is not determined by endogenous levels of  
3 arginine metabolic enzymes. *Cell Biol Int* 2010; **34**: 1085-1089.
- 4 60) Tsai WB, Aiba I, Lee SY, Feun L, Savaraj N, Kuo MT. Resistance to arginine  
5 deiminase treatment in melanoma cells is associated with induced argininosuccinate  
6 synthetase expression involving c-Myc/HIF-1 $\alpha$ /Sp4. *Mol Cancer Ther* 2009; **8**: 3223-  
7 3233.
- 8 61) Tsai WB, Aiba I, Long Y, Lin HK, Feun L, Savaraj N *et al.* Activation of  
9 Ras/PI3K/ERK pathway induces c-Myc stabilization to upregulate argininosuccinate  
10 synthetase, leading to arginine deiminase resistance in melanoma cells. *Cancer Res*  
11 2012; **72**: 2622-2633.
- 12 62) Long Y, Tsai W-B, Wangpaichitr M, Tsukamoto T, Savaraj N, Feun LG *et al.*  
13 Arginine deiminase resistance in melanoma cells is associated with metabolic  
14 reprogramming, glucose dependence and glutamine addiction. *Mol Cancer Ther*  
15 2013; **12**: 2581-2590.
- 16 63) Feun L, You M, Wu CJ, Kuo MT, Wangpaichitr M, Spector S *et al.* Arginine  
17 deprivation as a targeted therapy for cancer. *Curr Pharm Des* 2008; **14**: 1049-1057.
- 18 64) Szlosarek PW, Luong P, Phillips MM, Baccharini M, Ellis S, Szyszko T *et al.*  
19 Metabolic response to pegylated arginine deiminase in mesothelioma with promoter  
20 methylation of argininosuccinate synthetase. *J Clin Oncol* 2013; **31**: e111-e113. DOI:  
21 10.1200/JCO.2012.42.1784.
- 22 65) Yang TS, Lu SN, Chao Y, Sheen IS, Lin CC, Wang TE *et al.* A randomised phase II  
23 study of pegylated arginine deiminase (ADI-PEG 20) in Asian advanced  
24 hepatocellular carcinoma patients. *Br J Cancer* 2010; **103**: 954-960.

- 1 66) You M, Savaraj N, Wangpaichitr M, Wu C, Kuo TM, Varona-Santos J *et al.* The  
2 combination of ADI-PEG20 and TRAIL effectively increases cell death in melanoma  
3 cell lines. *Biochem Biophys Res Commun* 2010; **394**: 760-766.
- 4 67) Feun LG, Wu G, Clark M, Bombalaski J, Holtsberg F, Wangpajit M *et al.*  
5 Mechanism of anti-tumor effect of arginine deiminase-polyethylene (ADI-PEG20)  
6 and the possible mechanism of resistance in melanoma. *Proc Am Assoc Cancer Res*  
7 2004; 45: Abstract number 4565.
- 8 68) Wangpaichitr M, Wu C, Bigford G, Theodoropoulos G, You M, Li YY *et al.*  
9 Combination of Arginine Deprivation with TRAIL Treatment as a Targeted-Therapy  
10 for Mesothelioma. *Anticancer Res* 2014; **34**: 6991-7000.
- 11 69) You M, Savaraj N, Kuo MT, Wangpaichitr M, Varona-Santos J, Wu C *et al.* TRAIL  
12 induces autophagic protein cleavage through caspase activation in melanoma cell  
13 lines under arginine deprivation. *Mol Cell Biochem* 2013; **374**: 181–190.
- 14 70) Kim RH, Coates JM, Bowles TL, McNerney GP, Sutcliffe J, Jung JU *et al.* Arginine  
15 deiminase as a novel therapy for prostate cancer induces autophagy and caspase-  
16 independent apoptosis. *Cancer Res* 2009; **69**: 700–708.
- 17 71) Kim RH, Bold RJ, Kung HJ. ADI, autophagy and apoptosis: Metabolic stress as a  
18 therapeutic option for prostate cancer. *Autophagy* 2009; **5**: 567-568.
- 19 72) Savaraj N, Wu C, Kuo MT, You M, Wangpaichitr M, Robles C *et al.* The relationship  
20 of arginine deprivation, argininosuccinate synthetase and cell death in melanoma.  
21 *Drug Target Insights* 2007; **2**: 119–128.
- 22 73) Gong H, Zölzer F, Recklinghausen G, Havers W, Schweigerer L. Arginine deiminase  
23 inhibits proliferation of human leukemia cells more potently than asparaginase by  
24 inducing cell cycle arrest and apoptosis. *Leukemia* 2000; **14**: 826-829.

- 1 74) Bowles TL, Kim R, Galante J, Parsons CM, Virudachalam S, Kung HJ *et al.*  
2 Pancreatic cancer cell lines deficient in argininosuccinate synthetase are sensitive to  
3 arginine deprivation by arginine deiminase. *Int J Cancer* 2008; **123**: 1950–1955.
- 4 75) Surova O, Zhivotovsky B. Various modes of cell death induced by DNA damage.  
5 *Oncogene* 2013; **32**: 3789–3797
- 6 76) Changou CA, Chen Y-R, Xing L, Yen Y, Chuang FYS, Cheng RH *et al.* Arginine  
7 starvation-associated atypical cellular death involves mitochondrial dysfunction,  
8 nuclear DNA leakage, and chromatin autophagy. *Proc Natl Acad Sci USA* 2014; **111**:  
9 14147-14152.
- 10 77) Kung H-J, Changou CA, Li C-F, Ann DK. Chromatophagy: Autophagy goes nuclear  
11 and captures broken chromatin during arginine-starvation. *Autophagy* 2015; **11**: 419-  
12 421.
- 13 78) Gong H, Zölzer F, Recklinghausen GV, Rössler J, Breit S, Havers W *et al.* Arginine  
14 deiminase inhibits cell proliferation by arresting cell cycle and inducing apoptosis.  
15 *Biochem Biophys Res Commun* 1999; **261**: 10-14.
- 16 79) Lorenzo HK, Susin SA. Mitochondrial effectors in caspase-independent cell death.  
17 *FEBS Lett* 2004; **557**: 14-20.
- 18 80) Polster BM. AIF, reactive oxygen species, and neurodegeneration: A “complex”  
19 problem. *Neurochem Int* 2013; **62**: 695-702.
- 20 81) Norberg E, Orrenius S, Zhivotovsky B. Mitochondrial regulation of cell death:  
21 processing of apoptosis-inducing factor (AIF). *Biochem Biophys Res Commun* 2010;  
22 **396**: 95–100.
- 23 82) Zhu C, Wang X, Deinum J, Huang Z, Gao J, Modjtahedi N *et al.* Cyclophilin A  
24 participates in the nuclear translocation of apoptosis-inducing factor in neurons  
25 after cerebral hypoxia-ischemia. *J Exp Med* 2007; **204**: 1741–1748.

- 1 83) Pradelli LA, Bénétteau M, Ricci J-E. Mitochondrial control of caspase-dependent and  
2 -independent cell death. *Cell Mol Life Sci* 2010; **67**: 1589–1597.
- 3 84) Ulukaya E, Acilan C, Yilmaz Y. Apoptosis: why and how does it occur in biology?  
4 *Cell Biochem Funct* 2011; **29**: 468–480.
- 5 85) Shen LJ, Beloussow K, Shen WC. Modulation of arginine metabolic pathways as the  
6 potential anti-tumor mechanism of recombinant arginine deiminase. *Cancer Lett*  
7 2006; **231**: 30-35.
- 8 86) Mandal S, Mandal A, Johansson HE, Orjalo AV, Park MH. Depletion of cellular  
9 polyamines, spermidine and spermine, causes a total arrest in translation and growth  
10 in mammalian cells. *Proc Natl Acad Sci USA* 2013; **110**: 2169-2174.
- 11 87) Gerner EW, Meyskens FL Jr. Polyamines and cancer: old molecules, new  
12 understanding. *Nat Rev Cancer* 2004; **4**: 781-792.
- 13 88) Ohmura M, Hishiki T, Yamamoto T, Nakanishi T, Kubo A, Tsuchihashi K et al.  
14 Impacts of CD44 knockdown in cancer cells on tumor and host metabolic systems  
15 revealed by quantitative imaging mass spectrometry. *Nitric Oxide* 2015; **46**:102–113.
- 16 89) Folkman J. What is the evidence that tumors are angiogenesis dependent? *J Natl*  
17 *Cancer Inst* 1990; **82**: 4-6.
- 18 90) Jain RK. Tumor angiogenesis and accessibility: Role of vascular endothelial growth  
19 factor. *Semin Oncol* 2002; **29**: 3-9.
- 20 91) Cao Z, Shang B, Zhang G, Miele L, Sarkar FH, Wang F *et al.* Tumor cell-mediated  
21 neovascularisation and lymphangiogenesis contrive tumor progression and cancer  
22 metastasis. *Biochim Biophys Acta Rev Cancer* 2013; **1836**: 273-286.
- 23 92) Stapor P, Wang X, Goveia J, Moens S, Carmeliet P. Angiogenesis revisited – role and  
24 therapeutic potential of targeting endothelial metabolism. *J Cell Sci* 2014; **127**: 4331-  
25 4341.

- 1 93) Goel S, Duda DG, Xu L, Munn LL, Boucher Y, Fukumura D *et al.* Normalization of  
2 the vasculature for treatment of cancer and other diseases. *Physiol Rev* 2011; **91**:  
3 1071-1121.
- 4 94) Cantelmo AR, Brajic A, Carmeliet P. Endothelial Metabolism Driving Angiogenesis:  
5 Emerging Concepts and Principles. *Cancer J* 2015; **21**: 244-249.
- 6 95) Chamorro-Jorganes A, Lee MY, Araldi E, Landskroner-Eiger S, Fernández-Fuertes  
7 M, Sahraei M, *et al.* VEGF-induced expression of miR-17~92 cluster in endothelial  
8 cells is mediated by ERK/ELK1 activation and regulates angiogenesis. *Circ Res* 2015;  
9 DOI: 10.1161/CIRCRESAHA.115.307408.
- 10 96) Zhao D, Pan C, Sun J, Gilbert C, Drews-Elger K, Azzam DJ *et al.* VEGF drives  
11 cancer-initiating stem cells through VEGFR-2/Stat3 signaling to upregulate Myc and  
12 Sox2. *Oncogene* 2015; **34**:3107-3119.
- 13 97) Kalucka J, Missiaen R, Georgiadou M, Schoors S, Lange C, De Bock K. Metabolic  
14 control of the cell cycle. *Cell cycle* 2015. DOI: 10.1080/15384101.2015.1090068.
- 15 98) Moens S, Goveia J, Stapor PC, Cantelmo AR, Carmeliet P. The multifaceted activity  
16 of VEGF in angiogenesis-Implications for therapy responses. *Cytokine Growth*  
17 *Factor Rev* 2014; **25**: 473-482.
- 18 99) Eelen G, Zeeuw P, Simons M, Carmeliet P. Endothelial cell metabolism in normal  
19 and diseased vasculature. *Circ Res* 2015; **116**: 1231-1244.
- 20 100) Zecchin A, Stapor PC, Goveia J, Carmeliet P. Metabolic pathway  
21 compartmentalization: an underappreciated opportunity? *Curr Opin Biotech* 2015; **34**:  
22 73-81.
- 23 101) Beloussow K, Wang L, Wu J, Ann D, Shen W-C. Recombinant arginine deiminase as  
24 a potential anti-angiogenic agent. *Cancer Lett* 2002; **183**: 155-162.

- 1 102) Park I-S, Kang S-W, Shin Y-J, Chae K-Y, Park M-O, Kim M-Y *et al.* Arginine  
2 deiminase: a potential inhibitor of angiogenesis and tumor growth. *Br J Cancer* 2003;  
3 **89**: 907-914.
- 4 103) Thomas JB, Holtsberg FW, Ensor CM, Bomalaski JS, Clark MA. Enzymic  
5 degradation of plasma arginine using arginine deiminase inhibits nitric oxide  
6 production and protects mice from the lethal effects of tumor necrosis factor  $\alpha$  and  
7 endotoxin. *Biochem J* 2002; **363**: 581-587.
- 8 104) Noh EJ, Kang SW, Shin YJ, Kim DC, Park I-S, Kim MY *et al.* Characterization of  
9 *Mycoplasma arginini* deiminase expressed in *E. coli* and its inhibitory regulation of  
10 nitric oxide synthesis. *Mol Cells* 2002; **13**: 137-143.
- 11 105) Fraisl P. Crosstalk between oxygen- and nitric oxide-dependent signaling pathways in  
12 angiogenesis. *Exp Cell Res* 2013; **319**: 1331-1339.
- 13 106) Morbidelli L, Donnini S, Ziche M. Role of nitric oxide in tumor angiogenesis. *Cancer*  
14 *Treat Res* 2004; **117**: 155-167.
- 15 107) Jurasz P, Sawicki G, Duszyk M, Sawicka J, Miranda C, Mayers I *et al.* Matrix  
16 metalloproteinase 2 in tumor cell-induced platelet aggregation: Regulation by nitric  
17 oxide. *Cancer Res.* 2001; **61**: 376-382.
- 18 108) Carreau A, Kieda C, Grillon C. Nitric oxide modulates the expression of endothelial  
19 cell adhesion molecules involved in angiogenesis and leukocyte recruitment. *Exp Cell*  
20 *Res* 2011; **317**: 29-41.
- 21 109) Fukumura D, Kashiwagi S, Jain RK. The role of nitric oxide in tumour progression.  
22 *Nat Rev Cancer* 2006; **6**: 521-534.
- 23 110) Lee MY, Luciano AK, Ackah E, Rodriguez-Vita J, Bancroft TA, Eichmann A, *et al.*  
24 Endothelial Akt1 mediates angiogenesis by phosphorylating multiple angiogenic  
25 substrates. *Proc Natl Acad Sci USA* 2014; **111**: 12865-12870.

- 1 111) Fukumura D, Gohongi T, Kadambi A, Izumi Y, Ang J, Yun CO *et al.* Predominant  
2 role of endothelial nitric oxide synthase in vascular endothelial growth factor-induced  
3 angiogenesis and vascular permeability. *Proc Natl Acad Sci USA* 2001; **98**: 2604-  
4 2609.
- 5 112) Kashiwagi S, Izumi Y, Gohongi T, Demou ZN, Xu L, Huang PL *et al.* NO mediates  
6 mural cell recruitment and vessel morphogenesis in murine melanomas and tissue-  
7 engineered blood vessels. *J Clin Invest* 2005; **115**: 1816–1827.
- 8 113) Kashiwagi S, Tsukada K, Xu L, Miyazaki J, Kozin SV, Tyrrell JA *et al.* Perivascular  
9 nitric oxide gradients normalize tumor vasculature. *Nat Med* 2008; **14**: 255-257.
- 10 114) Roberts DD, Isenberg JS, Ridnour LA, Wink DA. Nitric oxide and its gatekeeper  
11 thrombospondin-1 in tumor angiogenesis. *Clin Cancer Res* 2007; **13**: 795-798.
- 12 115) McAlpine JA, Lu H-T, Wu KC, Knowles SK, Thomson JA. Down-regulation of  
13 argininosuccinate synthetase is associated with cisplatin resistance in hepatocellular  
14 carcinoma cell lines: implications for PEGylated arginine deiminase combination  
15 therapy. *BMC Cancer* 2014; **14**: 621.
- 16 116) Liu J, Ma J, Wu Z, Li W, Zhang D, Han L *et al.* Arginine deiminase augments the  
17 chemosensitivity of argininosuccinate synthetase-deficient pancreatic cancer cells to  
18 gemcitabine via inhibition of NF- $\kappa$ B signaling. *BMC Cancer* 2014; **14**: 686.
- 19 117) Allen MD, Luong P, Hudson C, Leyton J, Delage B, Ghazaly E *et al.* Prognostic and  
20 therapeutic impact of argininosuccinate synthetase 1 control in bladder cancer as  
21 monitored longitudinally by PET imaging. *Cancer Res* 2013; **74**: 896–907.
- 22 118) Daylami R, Muilenburg DJ, Virudachalam S, Bold RJ. Pegylated arginine deiminase  
23 synergistically increases the cytotoxicity of gemcitabine in human pancreatic cancer.  
24 *J Exp Clin Cancer Res* 2014; **33**: 102.

- 1 119) Jenkinson CP, Grody WW, Cederbaum SD. Comparative properties of arginases.  
2 *Comp Biochem Physiol* 1996; **114B**: 107-132.
- 3 120) Wu G, Morris SM Jr. Arginine metabolism: nitric oxide and beyond. *Biochem J* 1998;  
4 **336**: 1-17.
- 5 121) Morris SM Jr. Regulation of enzymes of urea and arginine synthesis. *Annu Rev Nutr*  
6 1992; **12**: 81-101.
- 7 122) Gotoh T, Sonoki T, Nagasaki A, Tereda K, Takiguchi M, Mori M. Molecular cloning  
8 of cDNA for nonhepatic mitochondrial arginase (arginase II) and comparison of its  
9 induction with nitric oxide synthase in a murine macrophage-like cell line. *FEBS Lett*  
10 1996; **395**: 119-122.
- 11 123) Elms S, Chen F, Wang Y, Qian J, Askari B, Pandey D *et al*. Insights into the arginine  
12 paradox: Evidence against the importance of subcellular location of arginase and  
13 eNOS. *Am J Physiol Heart Circ Physiol* 2013; **305**: H651-H666.
- 14 124) Morris SM Jr. Arginine metabolism: Boundaries of our Knowledge. *J Nutr* 2007; **137**:  
15 1602S–1609S.
- 16 125) Li H, Meininger CJ, Hawker JR Jr., Haynes TE, Kepka-Lenhart D, Mistry SK *et al*.  
17 Regulatory role of arginase I and II in nitric oxide, polyamine, and proline syntheses  
18 in endothelial cells. *Am J Physiol Endocrinol Metab* 2001; **280**: E75-E82.
- 19 126) Bach SJ, Lasnitzki I. Some aspects of the role of arginine and arginase in mouse  
20 carcinoma. *Enzymologia* 1947; **12**: 198-205.
- 21 127) Bach SJ, Maw GA. Creatine synthesis by tumor-bearing rats. *Biochim Biophys Acta*  
22 1953; **11**: 69-78.
- 23 128) Koji T, Terayama H. Arginase as one of the inhibitory principles in the density-  
24 dependent as well as plasma membrane-mediated inhibition of liver cell growth in  
25 vitro. *Exp Cell Res* 1984; **155**: 359-370.



- 1 129) Terayama H, Koji T, Kontani M, Okumoto T. Arginase is an inhibitory principle in  
2 liver growth of various mammalian cells in vitro Plasma membranes arresting the  
3 growth of various mammalian cells in vitro. *Biochim Biophys Acta* 1982; **720**: 188-  
4 192.
- 5 130) Huang M-H, Yang C-C, Wang C-C. Inhibition of lymphocyte proliferation by liver  
6 arginase. *Life Sci* 1992; **51**: 1725-1730.
- 7 131) Cheng PNM, Lam TL, Lam WM, Tsui SM, Cheng AWM, Lo WH *et al.* Pegylated  
8 recombinant human arginase (rharg-peg 5,000mw) inhibits the *in vitro* and *in vivo*  
9 proliferation of human hepatocellular carcinoma through arginine depletion. *Cancer*  
10 *Res* 2007; **67**: 309-317.
- 11 132) Wheatley DN, Campbell E. Arginine deprivation, growth inhibition and tumor cell  
12 death: 3. Deficient utilisation of citrulline by malignant cells. *Br J Cancer* 2003; **89**:  
13 573-576.
- 14 133) Tsui SM, Lam WM, Lam TL, Chong HC, So PK, Kwok SY *et al.* Pegylated  
15 derivatives of recombinant human arginase (rhArg1) for sustained *in vivo* activity in  
16 cancer therapy: preparation, characterization and analysis of their pharmacodynamics  
17 *in vivo* and *in vitro* and action upon hepatocellular carcinoma cell (HCC). *Cancer Cell*  
18 *Int* 2009; **9**: 9.
- 19 134) Stone EM, Glazer ES, Chantranupong L, Cherukuri P, Breece RM, Tierney DL *et al.*  
20 Replacing Mn<sup>2+</sup> with Co<sup>2+</sup> in human arginase I enhances cytotoxicity toward L-  
21 arginine auxotrophic cancer cell lines. *ACS Chem Biol* 2010; **5**: 333-342.
- 22 135) Glazer ES, Stone EM, Zhu C, Massey KL, Hamir AN, Curley SA. Bioengineered  
23 human arginase I with enhanced activity and stability controls hepatocellular and  
24 pancreatic carcinoma xenografts. *Transl Oncol* 2011; **4**: 138-146.

- 1 136) Tanios R, Bekdash A, Kassab E, Stone E, Georgiou G, Frankel AE *et al.* Human  
2 recombinant arginase I(Co)-PEG5000 [HuArgI(Co)-PEG5000]-induced arginine  
3 depletion is selectively cytotoxic to human acute myeloid leukemia cells. *Leukemia*  
4 *Res* 2013; **37**: 1565-1571.
- 5 137) Khoury O, Ghazale N, Stone E, El-Sibai M, Frankel AE, Abi-Habib RJ. Human  
6 recombinant arginase I (Co)-PEG5000 [HuArgI(Co)-PEG5000]-induced arginine  
7 depletion is selectively cytotoxic to human glioblastoma cells. *J Neuro-Oncol* 2015;  
8 **122**: 75-85. DOI 10.1007/s11060-014-1698-5.
- 9 138) Li L, Wang Y, Chen J, Cheng B, Hu J, Zhou Y *et al.* An engineered arginase FC  
10 protein inhibits tumor growth *In Vitro* and *In Vivo*. *Evid Based Complement Alternat*  
11 *Med* 2013, Article ID 423129. DOI: <http://dx.doi.org/10.1155/2013/423129>.
- 12 139) Lam TL, Wong GKY, Chow HY, Chong HC, Chow TL, Knok SY *et al.* Recombinant  
13 human arginase inhibits the *in vitro* and *in vivo* proliferation of human melanoma by  
14 inducing cell cycle arrest and apoptosis. *Pigment cell melanoma Res* 2010; **24**: 366-  
15 376.
- 16 140) Lam TL, Wong GKY, Chong HC, Cheng PNM, Choi SC, Chow TL *et al.*  
17 Recombinant human arginase inhibits proliferation of human hepatocellular  
18 carcinoma by inducing cell cycle arrest. *Cancer Lett* 2009; **277**: 91-100.
- 19 141) Yau T, Cheng PNM, Chan P, Chan W, Chen L, Yuen J *et al.* A phase 1 dose-  
20 escalating study of pegylated recombinant human arginase 1 (Peg-rhArg1) in patients  
21 with advanced hepatocellular carcinoma. *Invest New Drugs* 2013; **31**: 99–107.
- 22 142) Yau CC, Chan P, Pang R, Chan W, Cheng PNM, Poon R. A phase I study of  
23 recombinant human arginase I (rhArgI) for patients with advanced hepatocellular  
24 carcinoma. *J Clin Oncol* 2010; **28** (ASCO Annual meeting abstracts) e13503.

- 1 143) Ferraro E, Cecconi F. Autophagic and apoptotic response to stress signals in  
2 mammalian cells. *Arch Biochem Biophys* 2007; **462**: 210–219.
- 3 144) Jiang P, Mizushima N. Autophagy and human diseases. *Cell Res* 2014; **24**: 69-79.  
4 DOI:10.1038/cr.2013.161.
- 5 145) Gozuacik D, Kimchi A. Autophagy as a cell death and tumor suppressor mechanism.  
6 *Oncogene* 2004; **23**: 2891–2906.
- 7 146) Macintosh RL, Ryan KM. Autophagy in tumour cell death. *Semin Cancer Biol* 2013;  
8 **23**: 344-351.
- 9 147) Fulda S and Köge D. Cell death by autophagy: emerging molecular mechanisms and  
10 implications for cancer therapy. *Oncogene* 2015; **34**:5105-5113.
- 11 148) Denton D, Nicolson S, Kumar S. Cell death by autophagy: facts and apparent  
12 artefacts. *Cell Death Differ* 2012; **19**: 87–95.
- 13 149) Shen H-M., Codongo P. Autophagic cell death: Loch Ness monster or endangered  
14 species? *Autophagy* 2011; **7**: 457-465.
- 15 150) Efeyan A, Zoncu R, Sabatini DM. Amino acids and mTORC1: from lysosomes to  
16 disease. *Trends Mol Med* 2012; **18**: 524-533.
- 17 151) Beauchamp EM, Platanias LC. The evolution of the TOR pathway and its role in  
18 cancer. *Oncogene* 2013; **32**: 3923–3932
- 19 152) Yan L, Lamb RF. Amino acid sensing and regulation of mTORC1. *Semin Cell Dev*  
20 *Biol* 2012; **23**: 621-625.
- 21 153) Ryter SW, Cloonan SM, Choi AMK. Autophagy: A critical regulator of cellular  
22 metabolism and homeostasis. *Mol Cells* 2013; **36**: 7-16.
- 23 154) Cui J, Gong Z, Shen H-M. The role of autophagy in liver cancer: Molecular  
24 mechanisms and potential therapeutic targets, *Biochim Biophys Acta Rev Cancer*  
25 2013; **1836**: 15-26.

- 1 155) Hara K, Yonezawa K, Weng QP, Kozlowski MT, Belham C, Avruch J. Amino acid  
2 sufficiency and mTOR regulate p70 S6 kinase and eIF-4E BP1 through a common  
3 effector mechanism. *J Biol Chem* 1998; **273**: 14484-14494.
- 4 156) Hsueh EC, Knebel SM, Lo W-H, Leung Y-C, Cheng PNM, Hsueh CT. Deprivation of  
5 arginine by recombinant human arginase in prostate cancer cells. *J Hematol Oncol*  
6 2012; **5**: 17-22.
- 7 157) Scott L, Lamb J, Smith S, Wheatley DN. Single amino acid (arginine) deprivation:  
8 rapid and selective death of cultured transformed and malignant cells. *Br J Cancer*  
9 2000; **83**: 800-810.
- 10 158) Wang Y, Nartiss Y, Steipe B, McQuibban GA, Kim PK. ROS-induced mitochondrial  
11 depolarization initiates PARK2/PARKIN-dependent mitochondrial degradation by  
12 autophagy. *Autophagy* 2012; **8**: 1462–1476.
- 13 159) Li Z-Y, Yang Y, Ming M, Liu B. Mitochondrial ROS generation for regulation of  
14 autophagic pathways in cancer. *Biochem Biophys Res Commun* 2011; **414**: 5-8.
- 15 160) Chen Y, McMillan-Ward E, Kong J, Israels SJ, Gibson SB. Oxidative stress induces  
16 autophagic cell death independent of apoptosis in transformed and cancer cells. *Cell*  
17 *Death Differ* 2008; **15**: 171–182.
- 18 161) Wang Z, Shi X, Li Y, Zeng X, Fan J, Sun Y *et al.* Involvement of autophagy in  
19 recombinant human arginase-induced cell apoptosis and growth inhibition of  
20 malignant melanoma cells. *Appl Microbiol Biotechnol* 2014; **98**: 2485-2494.
- 21 162) Wu WKK, Coffelt SB, Cho CH, Wang XJ, Lee CW, Chan FKL *et al.* The autophagic  
22 paradox in cancer therapy. *Oncogene* 2012; **31**: 939–953.
- 23 163) Rebsamen M, Pochini L, Stasyk T, de Araujo MEG, Galluccio M, Kandasamy RK *et*  
24 *al.* SLC38A9 is a component of the lysosomal amino acid sensing machinery that  
25 controls mTORC1. *Nature* 2015; **519**: 477-481.

- 1 164) Wang S, Tsun Z-Y, Wolfson RL, Shen K, Wyant GA, Plovanich ME et al. Lysosomal  
2 amino acid transporter SLC38A9 signals arginine sufficiency to mTORC1. *Science*  
3 2015; **347**:188-194.
- 4 165) Eng CH, Abraham RT. The autophagy conundrum in cancer: influence of tumorigenic  
5 metabolic reprogramming. *Oncogene* 2011; **30**: 4687–4696.
- 6 166) Pérez E, Das G, Bergmann A, Baehrecke EH. Autophagy regulates tissue overgrowth  
7 in a context-dependent manner. *Oncogene* 2015; **34**:3369-3376.
- 8 167) Tsujimoto Y, Shimizu S. Another way to die: autophagic programmed cell death. *Cell*  
9 *Death Differ* 2005; **12**: 1528–1534.
- 10 168) Wang Z, Shi X, Li Y, Fan J, Zeng X, Xian Z, et al. Blocking autophagy enhanced  
11 cytotoxicity induced by recombinant human arginase in triple-negative breast cancer  
12 cells. *Cell Death Dis* 2014; **5**: e1563. doi:10.1038/cddis.2014.503
- 13 169) Booth LA, Tavallai S, Hamed HA, Cruickshanks N, Dent P. The role of cell  
14 signalling in the crosstalk between autophagy and apoptosis. *Cellular Signal* 2014;  
15 **26**: 549-555.
- 16 170) Fimia GM, Corazzari M, Antonioli M, Piacentini M. Ambra1 at the crossroad  
17 between autophagy and cell death. *Oncogene* 2013; **32**: 3311–3318.
- 18 171) Djavaheri-Mergny M, Maiuri MC, Kroemer G. Cross talk between apoptosis and  
19 autophagy by caspase-mediated cleavage of Beclin 1. *Oncogene* 2010; **29**: 1717–  
20 1719.
- 21 172) Hernandez CP, Morrow K, Lopez-Barcons LA, Zabaleta J, Sierra R, Velasco C *et al.*  
22 Pegylated arginase I: a potential therapeutic approach in T-ALL. *Blood* 2010; **115**:  
23 5214-5221.

- 1 173) Morrow K, Hernandez CP, Raber P, Del Valle L, Wilk AM, Majumdar S *et al.* Anti-  
2 leukemic mechanisms of pegylated arginase I in acute lymphoblastic T-cell leukemia.  
3 *Leukemia* 2013; **27**: 569-577.
- 4 174) Chow AK, Ng L, Sing Li H, Cheng CW, Lam CS, Yau TC, Cheng PN *et al.* Anti-  
5 tumor efficacy of a recombinant human arginase in human hepatocellular carcinoma.  
6 *Curr Cancer Drug Targets* 2012; **12**: 1233–1243.
- 7 175) Marino T, Russo N, Toscano M. What occurs by replacing Mn<sup>2+</sup> with Co<sup>2+</sup> in human  
8 arginase I: First- principles computational analysis. *Inorg Chem* 2013; **52**: 655-659.
- 9 176) Glazer ES, Kaluarachchi WD, Massey KL, Zhu C, Curley SA. Bioengineered  
10 arginase I increases caspase-3 expression of hepatocellular and pancreatic carcinoma  
11 cells despite induction of argininosuccinate synthetase-1. *Surgery* 2010; **148**: 310-  
12 318.
- 13 177) Regunathan S, Reis DJ. Characterization of arginine decarboxylase in rat brain and  
14 liver: distinction from ornithine decarboxylase. *J Neurochem* 2000; **74**: 2201-2208.
- 15 178) Li G, Regunathan S, Barrow CJ, Eshraghi J, Cooper R, Reis DJ. Agmatine: an  
16 endogenous clonidine-displacing substance in the brain. *Science* 1994; **263**: 966-969.
- 17 179) Zhu MY, Iyo A, Piletz JE, Regunathan S. Expression of human arginine  
18 decarboxylase, the biosynthetic enzyme for agmatine. *Biochim Biophys Acta* 2004;  
19 **1670**: 156-164.
- 20 180) Molderings GJ, Haenisch B. Agmatine (decarboxylated l-arginine): Physiological role  
21 and therapeutic potential. *Pharmacol Therapeut* 2012; **133**: 351–365.
- 22 181) Dudkowska M, Lai J, Gardini G, Stachurska A, Grzelakowska-Sztabert B,  
23 Colombatto S *et al.* Agmatine modulates the *in vivo* biosynthesis and interconversion  
24 of polyamines and cell proliferation. *Biochim Biophys Acta* 2003; **1619**: 159-166.

- 1 182) Satriano J, Matsufuji S, Murakami Y, Lortie MJ, Schwartz D, Kelly CJ *et al.*  
2 Agmatine suppresses proliferation by frameshift induction of antizyme and  
3 attenuation of cellular polyamine levels. *J Biol Chem* 1998; **273**: 15313-15316.
- 4 183) Choi YS, Cho YD. Effects of agmatine on polyamine metabolism and the growth of  
5 prostate tumor cells. *J Biochem Mol Biol* 1999; **32**: 173-180.
- 6 184) Mayeur C, Veuillet G, Michaud M, Raul F, Blottière H, Blachier F. Effects of  
7 agmatine accumulation in human colon carcinoma cells on polyamine metabolism,  
8 DNA synthesis and the cell cycle. *Biochim Biophys Acta* 2005; **1745**: 111-123.
- 9 185) Moinard C, Cynober L, Bandt JPD. Polyamines: metabolism and implications in  
10 human diseases. *Clin Nutr* 2005; **24**: 184-197.
- 11 186) Satriano J. Arginine pathways and the inflammatory response: Interregulation of nitric  
12 oxide and polyamines. *Amino Acids* 2004; **26**: 321-329.
- 13 187) Philip R, Campbell E, Wheatley DN. Arginine deprivation, growth inhibition and  
14 tumor cell death: 2. Enzymatic degradation of arginine in normal and malignant cell  
15 cultures. *Br J Cancer* 2003; **88**: 613-623.
- 16 188) Wheatley DN, Scott L, Lamb J, Smith S. Single amino acid (arginine) restriction:  
17 Growth and death of cultured HeLa and human diploid fibroblasts. *Cellular Physiol*  
18 *Biochem* 2000; **10**: 37-55.
- 19 189) Wheatley DN, Campbell E. Arginine catabolism, liver extracts and cancer. *Pathol*  
20 *Oncol Res* 2002; **8**: 18-25.
- 21 190) Bronte V and Zanovello P. Regulation of immune responses by L-arginine  
22 metabolism. *Nat Rev Immunol* 2005; **5**: 641-654.
- 23 191) Sikalidis AK. Amino acids and immune response: A role for cysteine, glutamine,  
24 phenylalanine, tryptophan and arginine in T-cell function and cancer?. *Pathol Oncol*  
25 *Res* 2015; **21**: 9-17.

- 1 192) Raber P, Ochoa AC, Rodriguez PC. Metabolism of L-arginine by myeloid-derived  
2 suppressor cells in cancer: Mechanisms of T-cell suppression and therapeutic  
3 perspectives. *Immunol Invest* 2012; **41**: 614-634.
- 4 193) Peranzoni E, Marigo I, Dolcetti L, Ugel S, Sonda N, Taschin E et al. Role of arginine  
5 metabolism in immunity and immunopathology. *Immunobiol* 2008; **212**: 795-812.
- 6 194) Popovic PJ, Zeh HJ III, Ochoa JB. Arginine and immunity. *J Nutr* 2007; **137**:1681s-  
7 1686s
- 8 195) Wheatley DN, Campbell E, Lai PBS, Cheng PNM. A rational approach to the  
9 systemic treatment of cancer involving medium-term depletion of arginine. *Gene Ther*  
10 *Mol Biol* 2005; **9**: 33-40.
- 11 196) Kilberg MS, Balasubramanian M, Fu L, Shan J. The transcription factor network  
12 associated with the amino acid response in mammalian cells. *Adv Nutr* 2012; **3**: 295–  
13 306.
- 14 197) Jackson RJ, Hellen CU, Pestova TV. The mechanism of eukaryotic translation  
15 initiation and principles of its regulation. *Nat Rev Mol Cell Biol* 2010; **11**: 113-127.
- 16 198) Zang H, Forman HJ. Glutathione synthesis and its role in redox signaling. *Semin Cell*  
17 *Dev Biol* 2012; **23**: 722-728.
- 18 199) Weiger TM, Hermann A. Cell proliferation, potassium channels, polyamines and their  
19 interactions: a mini review. *Amino acids* 2014; **46**: 681-688.
- 20 200) Lind D. Arginine and cancer. *J Nutr* 2004; **134**: 2837s-2841s
- 21 201) Kohler ES, Sankaranarayanan S, Van Ginneken CJ, Van Dijk P, Vermeulen JLM,  
22 Ruijter JM *et al.* The human neonatal small intestine has the potential for arginine  
23 synthesis; developmental changes in the expression of arginine-synthesizing and  
24 catabolizing enzymes. *BMC Dev Biol* 2008; **8**: 107.



- 1 202) Kim JE, Kim SY, Lee KW, Lee HJ. Arginine deiminase originating from  
2 Lactobacillus lactis ssp. lactis American Type Culture Collection (ATCC) 7962  
3 induces G<sub>1</sub>-phase cell-cycle arrest and apoptosis in SNU-1 stomach adenocarcinoma  
4 cells. *Br J Nutr* 2003; **102**: 1469-1476.
- 5 203) Gill P, Pan J. Inhibition of cell division in L5178Y cells by arginine-degrading  
6 mycoplasmas: the role of arginine deiminase. *Can J Microbiol* 1970; **16**: 415-419.
- 7 204) Kim JH, Kim JH, Yu YS, Kim DH, Min BH, Kim KW. Anti-tumor activity of  
8 arginine deiminase via arginine deprivation in retinoblastoma. *Oncol Rep* 2007; **18**:  
9 1373-1377.
- 10 205) Huang CC, Tsai ST, Kuo CC, Chang JS, YT Jin, Chang JY *et al.* Arginine  
11 deprivation as a new treatment strategy for head and neck cancer. *Oral Oncol* 2012;  
12 **48**: 1227-1235.
- 13 206) Tan B, Yin Y, Kong X, Li P, Li X, Gao H *et al.* L-Arginine stimulates proliferation  
14 and prevents endotoxin-induced death of intestinal cells. *Amino Acids* 2010; **38**: 1227-  
15 1235.
- 16 207) Wheatley DN, Philip R, Campbell E. Arginine deprivation and tumor cell death:  
17 Arginase and its inhibition. *Mol Cell Biochem* 2003; **244**: 177-185.
- 18 208) Izzo F, Marra P, Beneduce G, Castello G, Vallone P, De Rosa V *et al.* Pegylated  
19 Arginine Deiminase Treatment of Patients With Unresectable Hepatocellular  
20 Carcinoma: Results From Phase I/II Studies. *J Clin Oncol* 2004; **22**: 1815-1822.
- 21 209) Feun LG, You M, Wu C, Wangpaichitr M, Kuo MT, Marini A *et al.* Final results of  
22 phase II trial of pegylated arginine deiminase (ADI-PEG20) in metastatic melanoma  
23 (MM) [abstract]. *J Clin Oncol* 2010; **28** (Suppl 15): 8528.

- 1 210) Ott PA, Carvajal RD, Pandit-Taskar N, Jungbluth AA, Hoffman EW, Wu BW *et al.*  
2 Phase I/II study of pegylated arginine deiminase (ADI-PEG 20) in patients with  
3 advanced melanoma. *Invest New Drugs* 2013; **31**: 425-434.
- 4 211) Szlosarek PW, Steele J, Sheaff M, Szyszko T, Ellis S, Nolan L. A randomised phase  
5 II trial of pegylated arginine deiminase in patients with malignant pleural  
6 mesothelioma. 2013 World Conference on Lung Cancer; 2013. Abstr no. MO09.02.
- 7 212) Tomlinson BK, Bomalaski JS, Diaz M, Akande T, Mahaffey N, Li T *et al.* Phase I  
8 trial of ADI-peg 20 plus docetaxel (DOC) in patients (pts) with advanced solid  
9 tumors. *J Clin Oncol* (ASCO Meeting Abstracts) 2013; **31** (suppl) 2569.
- 10 213) Tomlinson BK, Thomson JA, Bomalaski JS, Diaz M, Akande T, Mahaffey N *et al.*  
11 Phase I trial of arginine deprivation therapy with ADI-PEG 20 plus Docetaxel in  
12 patients with advanced malignant solid tumors. *Clin Cancer Res* 2015; **21**:2480-2486.  
13 DOI: 10.1158/1078-0432.CCR-14-2610.
- 14 214) Bach SJ, Swaine D. The effect of arginase on the retardation of tumor growth. *Br J*  
15 *Cancer* 1965; **19**: 379-386.
- 16 215) Currie GA. Activated macrophages kill tumor cells by releasing arginase. *Nature*  
17 1978; **273**: 758-759.

18

19

### Figure legends

20

#### 21 **Figure 1: Amino acid response (AAR) pathway**

22 Restriction of essential amino acids activates the general control nondepressible protein 2  
23 (GCN2) kinase by increasing uncharged t-RNA pool.<sup>196</sup> Activated GCN2 kinase  
24 phosphorylates the translation initiation factor eIF2 $\alpha$ . Phosphorylated eIF2 $\alpha$  binds more  
25 tightly to eIF2 $\beta$ , inhibiting the exchange of GDP for GTP. Inhibition of GDP exchange for  
26 GTP further inhibits the binding of eIF2 complex to methionine aminoacyl tRNA, leading to  
27 inhibition of translational initiation.<sup>197</sup> Recently, SLC38A9 has been identified as an

1 upstream positive regulator of the mTOR pathway. Amino acids activate the RAG GTPases,  
2 which then recruit mTOR to the lysosomal surface. Rheb also localizes to lysosomal  
3 membrane. mTOR activation occurs only when both RAG GTPases and Rheb are active.  
4 Upon amino acid deprivation, tuberous sclerosis complex (TSC) translocates to lysosomal  
5 surface and promotes GTP hydrolysis by Rheb and thereby inhibiting mTOR complex.<sup>164</sup>

6

## 7 **Figure 2: Involvement of arginine in human physiology**

8 Arginine is a dibasic, cationic amino acid and is considered as ‘conditionally essential’  
9 amino acid. Arginine plays a crucial role in innate and adaptive immunity. For example,  
10 increased role of arginine in myeloid-derived suppressor cells results in the impairment of T-  
11 cell proliferation and function.<sup>190</sup> Arginine has been identified as the sole physiological  
12 precursor for nitric oxide (NO), a key performer in many cellular regulatory functions.  
13 Arginine also is a precursor of two important amino acids, proline and glutamate.<sup>198</sup> One of  
14 the most important roles of arginine is its implication in the synthesis of polyamines through  
15 the diversion from NO synthesis pathway. Polyamines are known to promote tumor growth,  
16 invasion and metastasis.<sup>199</sup> Arginine also plays a vital role in the synthesis of nucleotides,  
17 creatine, agmatine and hormones such as insulin and prolactin.<sup>200</sup>

## 18 **Figure 3: Arginine synthesis and homeostasis pathways**

19 Arginine is synthesized as an intermediate in the urea cycle. Arginine homeostasis is mainly  
20 achieved by catabolism. In neonates, the gene expression of arginine anabolic enzymes such  
21 as 1-pyrroline-5-carboxylase, argininosuccinate synthetase (ASS) and argininosuccinate lyase  
22 (ASL) is low. Thus, arginine is considered as an essential amino acid in neonates. After birth,  
23 the expression of ASS and ASL increases and expression of arginase is found undetectable at  
24 this stage.<sup>201</sup> Arginine can be degraded by arginase, ADC, ADI and NOSs (Please note that  
25 ADI is not a mammalian enzyme). The products of arginine catabolism play important roles  
26 in tumor cell biology. For example, ornithine, the product of arginase, is diverted to  
27 polyamine synthesis via ornithine decarboxylase. NOSs degrade arginine into citrulline and  
28 NO. Citrulline is recycled to urea cycle, while NO is as a modulator of important metabolic  
29 and signaling cascades. Agmatine is synthesized by decarboxylation of arginine via ADC and  
30 plays an important role in neurotransmission.

31

1 **Figure 4:** Timeline of important advancement in arginine deprivation therapy of cancer

2

3 **Figure 5:** Schematic representation of cytostatic and cytotoxic pathways involved in arginine  
4 deprivation therapy

5 Arginine deprivation therapy (ADT) can potentially modulate numerous cellular and  
6 signaling pathways rendering their cytotoxic and cytostatic pathways. Induction of apoptotic  
7 pathways, inhibition of angiogenesis and inhibition of *de novo* protein synthesis are the  
8 important mechanisms attributed to the cytotoxic potential of ADT. Moreover, ADT-  
9 mediated modulations in tumor cell-cycle can be exploited as a means of tumor growth arrest.

List of Figures

Figure 1:




Figure 2




Figure 3




Figure 4







Figure 5



## **List of tables**

**Table 1: Use of arginine catabolizing enzymes in arginine deprivation therapy (experimental studies)**

**Table 2: Clinical investigations involving arginine depriving enzymes**

**Table 3: Properties of arginine depriving enzymes**

**Table 1:** Use of arginine catabolizing enzymes in ADT (Experimental studies) [ \* indicates Tumor xenograft experiments]

Enzyme used for deprivation	Cell line	Source and Cell type	Studies carried out	Reference
ADI	HSC-3 HSC-4 CaSki C41 A549 SCC T98G  HeLa CHO FF9 HUVEC  SNU-1  L5178Y MCF7 A549  SNUOT-Rb1 Y79  CWR22Rv1* A2058 SK-Mel-2 HUVe SaOS WAC2 Y-79 Meth AC14  SK-LC-13* SW1271	Human tongue squamous carcinoma Human cervix squamous carcinoma Human carcinoma Human cervix squamous epithelium Human colon adenocarcinoma Human glioblastoma  Human cervix Chinese hamster ovary Fetal foreskin fibroblast Human umbilical vein endothelium  Human stomach adenocarcinoma  Mouse lymphoblastic leukemia Human mammary adenocarcinoma Human lung carcinoma  Human retinoblastoma Human retinoblastoma  Human prostate Human melanoma Human melanoma Human umbilical vein endothelium Human osteosarcoma Human neuroblastoma Human retinoblastoma Human sarcoma  Human small cell lung Human small cell lung	Cell growth inhibitory effect of ADI (purified from <i>Mycoplasma</i> infected cell lines) in comparison with arginase  Concentration dependent effect of ADI on cell proliferation Anti-angiogenesis effect of ADI by inhibiting capillary-like tube formation Anti-proliferative effect and ADI induced cell cycle arrest and apoptosis Inhibition of cell division Effect of ADI on the regulation of cellular protein and polyamine synthesis  ASS expression related sensitivity of cells towards ADI  Autophagy and caspase independent apoptosis Combination effect of ADI and TRAIL,  Cell cycle progression and apoptosis Inhibition of NO using Pegylated ADI  Effect of ADI-PEG20-mediated deprivation on the production of NO  ASS expression related sensitivity of cells towards PEG-ADI, Induction of autophagy and caspase-independent	[43]  [102] [202] [203] [85]  [204]  [71] [66] [78]  [103]  [39]

<p>Bovine liver arginase</p> <p>rh-Arginase I</p>	<p>NCI-H82</p> <p>A375</p> <p>SK-mel-2*</p> <p>SK-mel-28*</p> <p>SK-hep 2*</p> <p>SK-hep 3*</p> <p>HEP3B</p> <p>A2058*</p> <p>SK-MEL-2</p> <p>MDA-MB-231</p> <p>Karpas-422</p> <p>MyLa</p> <p>SeaX</p> <p>OEC-M1</p> <p>SCC-15</p> <p>HONE-1</p> <p>A375</p> <p>Sk-Mel2</p> <p>A2058</p> <p>MEL-1220</p> <p>MIA-PaCa-2*</p> <p>PANC-1</p> <p>Capan-1</p> <p>HPAF II</p> <p>L1210</p> <p>HeLa</p> <p>A375</p> <p>MEWO</p> <p>SAos-2</p> <p>IPEC-1</p>	<p>Human small cell lung</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human HCC</p> <p>Human HCC</p> <p>Human HCC</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human breast</p> <p>Human B-cell lymphoma</p> <p>Human T-cell lymphoma</p> <p>Human T-cell lymphoma</p> <p>Human head and neck cancer</p> <p>Human head and neck cancer</p> <p>Human head and neck cancer</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human pancreatic cancer</p> <p>Human pancreatic cancer</p> <p>Human pancreatic cancer</p> <p>Human pancreatic cancer</p> <p>Murine lymphocytic leukemia</p> <p>Human cervical adenocarcinoma</p> <p>Human melanoma</p> <p>Human melanoma</p> <p>Human osteogenic sarcoma</p> <p>Pig intestinal porcine epithelial cells-1</p>	<p>apoptosis</p> <p>Specificity of ADI for degradation of arginine and other amino acids; ASS expression dependent sensitivity of HCC and melanomas towards ADI</p> <p>Involvement of Ras/PI3K/ERK pathway in induction of c-Myc stabilization and up-regulation of ASS</p> <p>Correlation between ASS methylation status and sensitivity of the cells towards ADI</p> <p>Potential clinical correlation between ASS expression and tumor prognosis</p> <p>The role of ASS gene expression in ADI response/resistance</p> <p>The role of ASS gene expression in ADI response/resistance</p> <p>Cell proliferation and non-recoverable cell death of malignant cells on restoration of arginine</p> <p>Cell proliferation and ASS expression dependent recycling of citrulline to arginine</p> <p>LPS- induced cell damage involving mTOR and TLR4 pathways</p>	<p>[47]</p> <p>[61]</p> <p>[26]</p> <p>[205]</p> <p>[72]</p> <p>[74]</p> <p>[187]</p> <p>[132]</p> <p>[206]</p>
---	--	---	--	---

	PC-3 DU-145 LNCaP A375* SK-MEL-2 SK-MEL-28 B16-F0* L1210 HeLa	Human prostate Human prostate Human prostate Human melanoma Human melanoma Human melanoma Mouse melanoma Murine lymphocytic leukemia Human cervical adenocarcinoma	Expression levels of ASS and OCT, rhArginase I- mediated modulations in mTOR signaling pathway  Proliferation and cell cycle progression of melanoma cells, modulations in the cell cycle and apoptosis-related genes  Rescue of the arginase treated cells by norvaline (Arginase inhibitor)  Gene expression profiling of ASS and OTC, Synergistic effect of pegylated rhArginase I with 5-Fluorouracil on cell growth inhibition  Combination effect of pegylated rhArginase I with Cytarabine (Ara-C) on expression of cyclins  Effect of pegylation of rh-arginase I on its anti-tumor efficacy, immunogenicity and circulation half life  Cell cycle progression and transcriptional modulation of cyclins and/or CDKs  Global arrest in protein synthesis; Central role of phospho-eIF2a signaling and the kinases (GCN2 and PERK) in the induction of T-ALL cell apoptosis by rhArginase I-PEG <sub>500mw</sub>	[156]  [139]  [207]  [131]  [172]  [133]  [140]  [173]
rhArginase I-PEG <sub>5000mw</sub>	Hep-3B* SK-HEP-1 Huh7 SK-MEL-28 PLC/PRF/5 CCRF-CEM* Jurkat Molt-3 HepG2* Hep3B*  HepG2* PLC/PRF/5* Hep3B CCRF-CEM* Molt-4 H9 Lousy Jurkat HPB-ALL KOPTK1	Human HCC Human liver adenocarcinoma Human HCC Human melanoma Human primary hepatoma Human T-ALL Human T-ALL Human T-ALL Human HCC Human HCC  Human HCC Human HCC Human HCC Human T-ALL Human T-ALL Human T-ALL Human T-ALL Human T-ALL Human T-ALL Human T-ALL Human T-ALL	Effect of Co <sup>2+</sup> substitution of the Mn <sup>2+</sup> on catalytic activity and stability of human arginase I  Effect of Co <sup>2+</sup> substitution of the Mn <sup>2+</sup> on cytotoxicity	[135]  [134]
Bioengineered human arginase I	HepG2* Panc-1*  Hep3b A375	Human HCC Human pancreatic carcinoma  Human HCC Human melanoma		

**Table 2: Clinical investigations involving arginine depriving enzymes**

Enzyme	Cancer type	Phase of a clinical trial	Number of patients	Clinical outcomes	Common side effects	Post-treatment levels of plasma arginine#	Reference
ADI-PEG20	HCC	II	71	SD:31% (22/71) DCR: 31% (22/71)	Hypersensitivity/skin rash, local tissue reaction at injection site, hyperuricemia, pruritus, fatigue, hyperammonemia, fever, diarrhea	< 2 µM	[63]
	ASS (-) melanoma	I	17	PR: 23.5 % (4/17) SD: 29.4 % (5/17) CBR: 52.9 % (9/17)	Mild/moderate discomfort at the intramuscular injection site, neutropenia and thrombocytopenia, anaemia, fatigue	Undetectable	[25]
	HCC	I/II	19	CR: 11% (2/19) PR: 37 % (7/19) SD: 37% (7/19)	Occasional elevation in serum lipase, bilirubin and amylase levels, hyperuricemia, mild pain at the site of injection, increase in fibrinogen	< 2 µM	[208]
	MM	I/II	24	OR: 25 % (6/24) SD: 25 % (6/24)	Mild pain at the site of injection, hyperuricemia, elevated serum lipase, bilirubin, amylase and LDH, decreased hemoglobin, platelet and WBC count	< 2 µM	[52]
	HCC	II	76	OR: 3% (2/76) SD: 61% (50/76)	Transient and reversible encephalopathy, skin irritation, or discomfort at the site of injection combined with low-grade fever, decreased serum sodium, hemoglobin, albumin, fibrinogen levels, increased Potassium levels, uric acid and lipase	Undetectable	[51]
	MM	II	36	OR+SD: 28 % (10/36)	Discomfort at the injection site		[209]
	Melanoma	I/II	31	SD:31% (9/29) PMR: 27% (8/29)	Pain and rash at injection site, nausea, anorexia, pruritus, arthralgia	Undetectable	[210]

ADl-PEG20 plus Cisplatin	MPM	II	39	PMR: 46% (18/39) SD: 31% (12/39)	Skin injection site reactions, neutropenia, anaphylactoid reactions, serum sickness	2 µM <sup>\$</sup>	[64,211]
ADl-PEG20 plus Cisplatin and Pemetrexed	HCC	III		Ongoing (NCT01287585)			
	Non-Hodgkin's Lymphoma	II		Ongoing (NCT01910025)			
	SLCL	II		Ongoing (NCT01266018)			
	MM	I		Ongoing (NCT01665183)			
	Arginine auxotrophic tumors such as MPM and NSCLC	I		Ongoing (NCT02029690)			
ADl-PEG20 plus Docetaxel	Solid Prostate and NSCLC tumors	I	18	PR: 6% (1/18) SD: 33% (6/18)		Undetectable	[212,213]
ADl-PEG20 Plus Doxorubicin	HER2 (-) Breast Cancer	I		Ongoing (NCT01948843)			
Peg-rhArgI	HCC	I	15	SD:26.7% (4/15)	Abdominal pain, diarrhea, nausea, elevated ALT, AST, GGT & bilirubin	< 8 µM	[141]
Peg-rhArgI plus Oxaliplatin and Capecitabine	HCC	II		Ongoing (NCT02089633)			

Peg-rhArgI (the second-line therapy after sorafenib)	HCC	II	Ongoing (NCT02089763)		
---	-----	----	--------------------------	--	--

# Basal (Pre-treatment) level of arginine was ~ 130 µM

\$ Basal (Pre-treatment) level of arginine was ~ 63 µM

DCR- Disease-control rate (complete/partial response + stable disease)

SD- Stable disease

OS- Overall survival

PR- Partial response

CBR- Clinical benefit rate

MM- Metastatic melanoma

OR-Overall response (Complete + partial response)

CR- Complete response

PMR- partial metabolic response

MPM- Malignant Pleural Mesothelioma

Peg-rhArgI - Pegylated recombinant human arginase 1

ALT - Alanine Transaminase

AST - Aspartate Transaminase

GGT - Gamma-glutamyl transferase

SLCL- Small Cell Lung Cancer

NSCLC- Non-Small Cell Lung Cancer

HER2- Human epidermal growth factor receptor 2



**Table 3:** Properties of arginine depriving enzymes

<b>Arginine deiminase (E.C. 3.5.3.6)</b>	<b>Arginase (E.C.3.5.3.1)</b>	<b>Arginine decarboxylase (E.C.4.1.1.19)</b>
Main products are citrulline and NH <sub>3</sub>	Main products are ornithine and urea	Main products are agmatine and CO <sub>2</sub>
At physiological pH, Mycoplasma ADI is 300x more effective than arginase at depleting arginine	Very high alkaline pH optimum (pH 9.3) and has little enzymic activity at physiological pH	Mammalian ADC has a basic pH optimum (pH 8.23)
Circulatory half-life of ~ 4 h	Very short circulatory half-life (Approx. 30 minutes)	Not reported
Very high affinity for arginine (K <sub>m</sub> of 0.1-1 mM)	Low affinity for arginine (K <sub>m</sub> of 2-4 mM)	High affinity for arginine (K <sub>m</sub> of ~ 1mM)
Most normal cells and tissues are able to take up citrulline from the circulation	Ornithine can only be reconverted back into arginine in the liver and can cause toxicity to extra-hepatic tissues by inhibiting protein synthesis	Agmatine is not converted back to arginine under normal physiological conditions, may lead to its accumulation and toxicity to normal cells
Only found in microorganisms and is strongly antigenic in mammals	Human enzyme, non-immunogenic	Found in plants, microbes and human brain
Tumor sensitivity to ADI is dependent on ASS expression	The sensitivity of tumors to rhArg is independent of ASS expression	Studied only in human cervical cancer ( HeLa) cell lines
Efficacious only in ASS-negative tumors	Efficacious in both ASS-negative and OTC-negative tumors	
No cofactor requirement	Mn <sup>2+</sup> is essential for catalytic activity	Pyridoxal phosphate is a cofactor
Pegylation improves catalytic activity at physiological pH	Pegylation improves catalytic activity at physiological pH	PEGylation results in the total loss of catalytic activity