New Procedure to Generate Multipliers in Complex Neumann Problem and Effective Kohn Algorithm

Citation

Siu, Yum-Tong. 2017. New Procedure to Generate Multipliers in Complex Neumann Problem and Effective Kohn Algorithm. Science China Mathematics 60, no. 6: 1101–1128.

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:41467542>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

New Procedure to Generate Multipliers in Complex Neumann Problem and Effective Kohn Algorithm

Dedicated to the Memory of Professor Lu Qikeng

Yum-Tong Siu

The purpose of this note is threefold. (i) To explain the effective Kohn algorithm for multipliers in the complex Neumann problem and its difference with the full-real-radical Kohn algorithm, especially in the context of an example of Catlin-D'Angelo concerning the ineffectiveness of the latter. (ii) To extend the techniques of multiplier ideal sheaves for the complex Neumann problem to general systems of partial differential equations. (iii) To present a new procedure of generation of multipliers in the complex Neumann problem as a special case of the multiplier ideal sheaves techniques for general systems of partial differential equation.

For *a priori* estimates in the theory of partial differential equations some of the standard techniques are the following. (i) Using integration by parts to get L^2 estimates of derivatives in certain directions, for example, L^2 estimates of all first-order partial derivatives of a function with compact support from applying integration by parts to its inner product with its Laplacian. (ii) Using Lie bracket of two vector fields to conclude, from given derivative estimates of fractional orders along each of the two vectors, the derivative estimates of lower fractional order along their Lie bracket, for example, Hörmander's work on the sum of squares of vector fields [Hörmander1967].

The technique of multiplier ideal sheaves introduced by Kohn for the complex Neumann problem is a new method to conclude, from L^2 estimates of derivatives along certain *complex-valued* vector fields, the derivative estimates of lower fractional orders in all directions by introducing the notion of multipliers. In a system of partial differential equations where the estimate is for a vector-valued test function with components ψ_ν , for given complex-valued vector fields Y_j , when the estimates for several linear combinations $\sum_{j,\nu} \rho_{j,\nu} Y_j \psi_\nu$ with some given smooth functions $\rho_{j,\nu}$ are known, the multipliers are the smooth coefficients a_ν of a linear combination $\sum_\nu a_\nu \psi_\nu$ such that there is an estimate of the Sobolev L^2 norm of $\sum_\nu a_\nu \varphi_\nu$ of some positive fractional order. The goal is to derive differential relations among the multipliers to obtain some geometric condition to solve the regularity problem for the given system of partial differential equations, by the method of

using the differential relations and some initial multipliers to conclude that the function which is identically 1 is a scalar multiplier.

For Kohn's original setting of a bounded weakly pseudoconvex domain Ω in $\mathbb{C}^n$ the vector fields Y_j are the vector fields $L_1, \dots, L_{n-1}$ of type $(1, 0)$ tangential to the boundary $\partial\Omega$ of Ω together with their complex conjugates $\bar{L}_1, \dots, \bar{L}_{n-1}$. The test function is a $(0, 1)$ -form in the domains of $\bar{\partial}$ and $\bar{\partial}^*$ whose $n - 1$ tangential components are $\varphi_1, \dots, \varphi_{n-1}$. The given linear combinations $\sum_{j,\nu} \rho_{j,\nu} Y_j \psi_\nu$ are the $(n - 1)^2 + 1$ linear combinations $\bar{L}_j \varphi_\nu$ (for $1 \leq j, \nu \leq n - 1$) and $\sum_{\nu=1}^{n-1} L_\nu \varphi_\nu$.

The structure of this note is as follows. We start out with the background and motivation for the technique of multiplier ideal sheaves. We then discuss the two Kohn algorithm of generating multipliers. One is the full-real-radical Kohn algorithm. The other is the effective Kohn algorithm. We explain the algebraic geometric techniques in the effective Kohn algorithm, with special attention paid to the effectiveness of the orders of subellipticity in each step. The effective Kohn algorithm is then applied to Catlin-D'Angelo's example to highlight the difference in effectiveness between the full-real-radical Kohn algorithm and the effective Kohn algorithm. We present the generalization of the technique of multipliers to general systems of partial differential equations. Finally we apply the generalized techniques to the complex Neumann problem to obtain a new procedure to generate vector multipliers from matrix multipliers for special domains.

The notations $\mathbb{N}$, $\mathbb{R}$, and $\mathbb{C}$ mean respectively all positive integers, all real numbers, and all complex numbers. The notations $\mathcal{O}_{\mathbb{C}^n, P}$ and $\mathfrak{m}_{\mathbb{C}^n, P}$ mean respectively all holomorphic germs on $\mathbb{C}^n$ at P and the maximum ideal of the local ring $\mathcal{O}_{\mathbb{C}^n, P}$. Unless specified otherwise, $\|\cdot\|$ denotes the L^2 norm and $(\cdot, \cdot)$ denotes the L^2 inner product. The notation $\|\cdot\|_{L^2(U)}$ is also used to more clearly specify that it is the L^2 norm over U . The notation $(\cdot, \cdot)_{L^2(U)}$ is also used to more clearly specify that it is the L^2 inner product over U . The notation L_k^2 means the Sobolev norm defined by using the L^2 norm of derivatives up to order k . The notations ∂_j and $\bar{\partial}_j$ mean respectively $\frac{\partial}{\partial z_j}$ and $\frac{\partial}{\partial \bar{z}_j}$, where $z_1, \dots, z_n$ are the coordinates of $\mathbb{C}^n$.

TABLE OF CONTENTS

§1. Background and Motivation for Multiplier Ideal Sheaves

- §2. Generation of Multipliers in Kohn's Algorithm
- §3. Orders of Subellipticity in Algebraic Geometric Techniques for 3-Dimensional Special Domain
- §4. Effective Kohn Algorithm Applied to Catlin-D'Angelo's Example
- §5. Multipliers in More General Setting
- §6. New Procedure to Generate Vector Multiplier from Matrix Multiplier in Complex Neumann Problem of Special Domain

§1. BACKGROUND AND MOTIVATION FOR MULTIPLIER IDEAL SHEAVES

In regularity problems of local or global systems of partial differential equations, multiplier ideal sheaves describe the location and the extent of the failure of *a priori* estimates. There are two ways to introduce such multiplier ideal sheaves. The first way is from the continuity method of solving partial differential equations (which are usually nonlinear partial differential equations defined on compact Riemannian manifolds) where a multiplier ideal sheaf arises as the limit of rescaling of local coordinate charts to make possible the use of Ascoli-Arzelà techniques for convergence. The second way is just to directly introduce multiplier ideal sheaves as factors required in the integral norms to make *a priori* estimates hold for the regularity problem. We very briefly describe both.

(1.1) *Multiplier Ideal Sheaves from Limit of Rescaling of Local Coordinate Charts.* One method to solve nonlinear partial differential equations is to use the *continuity method* which uses a family of partial differential equations parametrized by $t \in [0, 1]$ so that

- (i) $t = 1$ is the given partial differential equations,
- (ii) $t = 0$ can be solved (which means solvability of partial differential equation at initial parameter), and
- (iii) from the solution for $t = t_0 < 1$ it is possible to solve for $t < t_0 + \varepsilon$ for some $\varepsilon > 0$ (which means that the *openness* property is assumed).

The difficulty is to prove the *closedness* property that for $0 < t^* \leq 1$ solutions s_ν for the parameter value $t = t_\nu$ for $t_\nu \nearrow t^*$ can be used to construct a solution s for the parameter value $t = t^*$.

The natural approach is to obtain s by taking the limit of s_{ν_k} for some subsequence $\{\nu_k\}$ of the sequence $\{\nu\}$. Usually the boundedness in some weak norm for s_ν , for example, L^2 , can be derived from the setup of the given partial differential equation. The method of Ascoli-Arzelà calls for boundedness in some stronger norm for s_ν , for example, L^2_1 defined by the first-order derivatives being L^2 .

In the setting of a manifold the derivative involved in L^2_1 depends on the choice of local coordinates so that L^2_1 is determined up to equivalence (of sandwiching between its products with two constants). One can always cheat by using different rescaling of local coordinates for each s_ν , but one has to pay the price that at the end the variable rescaling of local coordinates results in a factor (or *multiplier*) which is the limit of the Jacobian determinant in coordinate change in the integral for L^2 .

The limit s of s_{ν_k} from the Ascoli-Arzelà argument is L^2 only after the insertion of the multiplier in its L^2 integral. This can be interpreted as transforming the requirement in the the method of Ascoli-Arzelà for two norms L^2 and L^2_1 (with difference in orders of differentiation in their definitions) to two norms which are the L^2 norm and the multiplier-modified L^2_1 norm. Multipliers form an ideal sheaf, called the *multiplier ideal sheaf*. Global conditions, such as topological conditions, can be used to conclude that the multiplier ideal sheaf must be the full structure sheaf, giving the solvability of the partial differential equation from global conditions. Examples are (i) the existence of Hermitian-Einstein metrics for stable holomorphic vector bundles over compact algebraic (or Kähler) manifolds, where the method of limit of rescaling of local coordinates was first applied by Donaldson in [Donaldson1985] for the surface case and (ii) the existence of Kähler-Einstein metrics for certain Fano manifolds, where the method of multiplier ideal sheaves as defined by taking limit of metrics of the anti-canonical line bundle was first applied by Nadel in [Nadel1990].

We now look at the second way of introducing multiplier ideal sheaves, which is best described by using, as an example, the problem of the regularity of the Kohn solution for the complex Neumann problem on a bounded weakly pseudoconvex domain in $\mathbb{C}^n$ with smooth boundary.

(1.2) *Regularity Problem of Kohn Solution for Complex Neumann Problem on Weakly Pseudoconvex Domain.* Let Ω be a bounded weakly pseudoconvex

domain in $\mathbb{C}^n$ with smooth boundary $\partial\Omega$. It means that there is a smooth function r defined on some open neighborhood W of $\partial\Omega$ in $\mathbb{C}^n$ such that

- (i) $\Omega \cap W = \{ z \in W \mid r < 0 \}$,
- (ii) dr is nowhere zero on $\partial\Omega$, and
- (iii) the $(1,1)$ -form $\sqrt{-1} \partial \bar{\partial} r$ on W assumes nonnegative value when evaluated at $(\xi, \bar{\xi})$ for $\xi \in T_{\partial\Omega}^{(1,0)}$, where $T_{\partial\Omega}^{(1,0)}$ is the bundle of all tangent vectors ξ of $\mathbb{C}^n$ of type $(1,0)$ at points of $\partial\Omega$ which are tangential to $\partial\Omega$ (in the sense that $\xi(r) = 0$).

For notational simplicity we will consider only the complex Neumann problem in the case of $(0,1)$ -forms (instead of $(0,p)$ -form for $1 \leq p \leq n$). For a $\bar{\partial}$ -closed $(0,1)$ -form f on Ω which are smooth on the closure $\bar{\Omega}$ of Ω , the *Kohn solution* u for the $\bar{\partial}$ -equation $\bar{\partial}u = f$ on Ω means the unique smooth function u on Ω such that $\bar{\partial}u = f$ on Ω and u is perpendicular to all L^2 holomorphic functions on Ω with respect to the usual Euclidean volume form of $\mathbb{C}^n$.

The regularity problem of the Kohn solution for the complex Neumann problem on a bounded weakly pseudoconvex domain with smooth boundary is to study under what additional assumption on Ω the Kohn solution u is always smooth on $\bar{\Omega}$ when the given $(0,1)$ -form f is smooth on $\bar{\Omega}$.

(1.3) *Regularity from Subelliptic Estimate.* The *subelliptic estimate* of order $\varepsilon > 0$ is said to hold at a point P of $\partial\Omega$ if there exist an open neighborhood U of P in $\mathbb{C}^n$ and positive numbers ε and C satisfying

$$|||\varphi|||_\varepsilon^2 \leq C (\|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 + \|\varphi\|^2)$$

for all smooth $(0,1)$ -forms φ on $U \cap \bar{\Omega}$ with compact support which belong to the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$ (with respect to the usual L^2 inner product), where

- (i) $|||\cdot|||_\varepsilon$ is the Sobolev L^2 norm on Ω involving derivatives up to order ε in the boundary tangential directions of Ω , and
- (ii) $\|\cdot\|$ is the usual L^2 norm on Ω without involving any derivatives.

See [Kohn1979, p.92, (3.4)] for a detailed definition of the Sobolev norm $|||\cdot|||_\varepsilon$. In this note we will also use the notation Λ^s introduced in [Kohn1979,

p.92, (3.3)] which is the pseudo-differential operator corresponding to the $(\frac{s}{2})$ -th power of 1 plus the Laplacian in tangential coordinates of the boundary of Ω .

We would like to comment on the reason for the use of the Sobolev norm $||| \cdot |||_\varepsilon$ in whose definition only derivatives along the tangent directions of $\partial\Omega$ is used. The condition for a smooth $(0,1)$ -form g on $\bar{\Omega}$ to belong to the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$ (instead of the *formal* adjoint of $\bar{\partial}$) is that the components of g normal to $\partial\Omega$ vanish at all points of $\partial\Omega$ (which, in other words, means that the pointwise inner product of g and $\bar{\partial}r$ vanishes at every point of $\partial\Omega$). When g is in the domain of $\bar{\partial}^*$, the derivative of g along tangent directions of $\partial\Omega$ still belongs to the domain of $\bar{\partial}^*$, but if g is differentiated in the normal direction of $\partial\Omega$, the result in general will no longer belong to the domain of $\bar{\partial}^*$. That is the reason why we would like to avoid using differentiation along the normal directions of $\bar{\partial}$ in the Sobolev norm of order ε adopted in the definition of the subelliptic estimate.

Sobolev norms involving derivatives are used to enable us to conclude the order of differentiability of the solution u of the $\bar{\partial}$ -equation $\bar{\partial}u = f$ from the order of differentiability of the right-hand side f of the equation. Though we do not include the differentiation in the normal direction of $\partial\Omega$ in the Sobolev norm used, from the order of differentiability of the solution u along the *tangent directions* of $\partial\Omega$ we can still conclude the order of differentiability of u along the normal direction of $\partial\Omega$ because the equation $\bar{\partial}u = f$ itself provides us directly the differentiability of u along the *real normal* direction of $\partial\Omega$ from the differentiability of u along the $(0,1)$ component of the *complex normal* direction of $\partial\Omega$.

It was proved by Kohn-Nirenberg in 1965 [Kohn-Nirenberg1965, p.458, Theorem 4] that if for some $\varepsilon > 0$ the subelliptic estimate of order ε holds at every point of $\partial\Omega$, then the smoothness on $\bar{\Omega}$ of the Kohn solution u of $\bar{\partial}u = f$ follows from the smoothness of the $\bar{\partial}$ -closed $(0,1)$ -form f on $\bar{\Omega}$.

(1.4) *Multipliers to Measure Location and Extent of Failure of A Priori Estimates.* A smooth function germ F at a point P of $\partial\Omega$ (defined on some open neighborhood U_F of P in $\mathbb{C}^n$) is called a *scalar multiplier* if for some positive number $\varepsilon_F > 0$ and some positive constant C_F the subelliptic estimate

$$(1.4.1) \quad |||F\varphi|||_{\varepsilon_F}^2 \leq C_F (\|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 + \|\varphi\|^2)$$

of order ε_F , *modified* by the factor F , holds for all smooth $(0, 1)$ -forms φ on $U_F \cap \bar{\Omega}$ with compact support which belong to the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$. We say that the *order of subellipticity* for the scalar multiplier F is $\geq \varepsilon$. We are only interested in an effective lower bound for the order of subellipticity ε_F and will not study the supremum of all possible such ε_F .

The collection of smooth function germs at P which are scalar multipliers at P forms an ideal. This ideal is called the *multiplier ideal* at P and is denoted by I_P . The precise definition of the multiplier ideal makes precise the intuitive motivation that the direction and the order of vanishing of the multiplier ideal measures the location and extent of the failure of the *a priori* subelliptic estimate.

The test function φ which is multiplied by the scalar multiplier F to yield the modified subelliptic estimate (1.4.1) is not a scalar and is a $(0, 1)$ -form with n components (of which the normal component is 0). It is possible to get more information by using *vector multipliers* instead of just scalar multipliers. A smooth germ of $(1, 0)$ -form θ at a point of $\partial\Omega$ (defined on some open neighborhood U_θ of P in $\mathbb{C}^n$) is called a *vector multiplier* if for some positive number $\varepsilon_\theta > 0$ and some positive constant C_θ the subelliptic estimate

$$\|\theta \cdot \varphi\|_{\varepsilon_\theta}^2 \leq C_\theta (\|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 + \|\varphi\|^2)$$

of order ε_θ , *modified* by the dot product with θ , holds for all smooth $(0, 1)$ -forms φ on $U_\theta \cap \bar{\Omega}$ with compact support which belong to the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$. Here the dot product $\theta \cdot \varphi$ means the pointwise inner product $\langle g, \bar{\theta} \rangle$ of the two $(0, 1)$ -forms $\bar{\theta}$ and g with respect to the usual Euclidean Hermitian inner product of $\mathbb{C}^n$. In other words, if $\theta = \sum_{j=1}^n \theta_j dz_j$ and $\varphi = \sum_{j=1}^n \varphi_{\bar{j}} d\bar{z}_j$ (where $z_1, \dots, z_n$ are the global coordinates of $\mathbb{C}^n$), then $\theta \cdot \varphi = \sum_{j=1}^n \theta_j \varphi_{\bar{j}}$. The convention to introduce a vector multiplier θ as a $(1, 0)$ -form and the dot product $\theta \cdot \varphi$, instead of introducing a $(0, 1)$ -form ψ and the pointwise inner product $\langle \varphi, \psi \rangle$, is chosen so that, in the case of a special domain described in (2.8.1) below, one needs only consider scalar multipliers and vector multipliers which are holomorphic (see (2.9.3) below).

We say that the *order of subellipticity* for the vector multiplier θ is $\geq \varepsilon_\theta$. Again we are only interested in an effective lower bound for the order of subellipticity ε_θ and will not study the supremum of all possible such ε_θ .

The collection of smooth germs of $(1, 0)$ -forms at P which are vector multipliers at P forms a module over the algebra of all smooth function

germs at P . This module is called the *module of vector multipliers* at P and is denoted by A_P .

§2. GENERATION OF MULTIPLIERS IN KOHN'S ALGORITHM

It is easy to define multipliers to describe the location and the extent of failure of subelliptic estimates, but it is difficult to use multipliers to find easily verifiable conditions on the weakly pseudoconvex domain Ω to obtain subelliptic estimates and thereby the smoothness of the Kohn solution u on $\bar{\Omega}$ from the smoothness of the right-hand side f on $\bar{\Omega}$. The most important part of the theory of multipliers for the complex Neumann problem is the generation of scalar and vector multipliers by Kohn's algorithm which makes it possible to study subelliptic estimates from the geometric condition (known as *finite type*) of the finiteness of the maximum normalized order of contact between the boundary $\partial\Omega$ of Ω and a local holomorphic curve $f : \Delta \rightarrow \mathbb{C}^n$ (where Δ is the open unit disk in $\mathbb{C}$). Here the normalized order of contact means that the vanishing order of the pullback by f of the defining function r of $\partial\Omega$ divided by the vanishing order of f . The precise definition of finite type condition and its history will be given later in (2.3) below.

(2.1) *Kohn's Algorithm to Generate Scalar and Vector Multipliers.* The following three procedures constitute the Kohn algorithm of generating scalar and vector multipliers at a boundary point P of $\partial\Omega$.

(A) INITIAL MULTIPLIERS

- (i) The function r belongs to the ideal of multipliers I_P at P . Its order of subellipticity for the scalar multiplier r is ≥ 1 .
- (ii) For any germ of smooth vector field $\xi = \sum_{k=1}^n a_k \frac{\partial}{\partial z_k}$ at P of type $(1, 0)$ which is tangential to $\partial\Omega$, the $(1, 0)$ -form

$$(\partial\bar{\partial}r) \lrcorner \bar{\xi} = \sum_{j,k=1}^n \bar{\xi}_k \frac{\partial^2 r}{\partial z_j \partial \bar{z}_k} dz_j$$

(which is the interior product of the $(1, 1)$ -form $\partial\bar{\partial}r$ and the $(0, 1)$ -vector field $\bar{\xi}$) belongs to the module A_P of vector multipliers at P . The order of subellipticity for the vector multiplier $\partial\bar{\partial}_j r$ is $\geq \frac{1}{2}$ for $1 \leq j \leq n-1$ at points P where ∂r is normalized to be dz_n . (See [Kohn1979, p.97, (4.29)]).

(B) GENERATION OF NEW MULTIPLIERS

- (i) If $f \in I_P$, then $\partial f \in A_P$. The order $\varepsilon_{\partial f}$ of subellipticity for the vector multiplier ∂f is $\geq \frac{\varepsilon_f}{2}$ if the order of subellipticity of f is $\geq \varepsilon_f$. (See [Kohn1979, p.99, (4.42)].)
- (ii) If $\theta_1, \dots, \theta_{n-1} \in A_P$ and

$$\theta_1 \wedge \dots \wedge \theta_{n-1} \wedge \partial r = f_{\theta_1, \dots, \theta_{n-1}} dz_1 \wedge \dots \wedge dz_n,$$

then $f_{\theta_1, \dots, \theta_{n-1}} \in I_P$. In other words, for vector multipliers $\theta_1, \dots, \theta_{n-1}$ the coefficient of $dz_1 \wedge \dots \wedge dz_n$ in $\theta_1 \wedge \dots \wedge \theta_{n-1} \wedge \partial r$ is a scalar multiplier. Moreover, if the order of subellipticity of each θ_j (for $1 \leq j \leq n-1$) is $\geq \varepsilon$, then the order of subellipticity of $f_{\theta_1, \dots, \theta_{n-1}}$ is $\geq \varepsilon$. We denote the scalar multiplier $f_{\theta_1, \dots, \theta_{n-1}}$ by $\det_{n-1}(\theta_1, \dots, \theta_{n-1})$.

(C) REAL RADICAL PROPERTY

If $g \in I_P$ and $|f|^m \leq |g|$ for some $m \in \mathbb{N}$, then $f \in I_P$. The order of subellipticity for the scalar multiplier f is $\geq \frac{\varepsilon}{m}$ if the order of subellipticity of g is $\geq \varepsilon$. (See [Kohn1979, p.98, Lemma 4.34].)

For the purpose of discussing the effectiveness of Kohn's algorithm later, we introduce now two terms concerning the radicals of ideals. For an ideal J of smooth function germs at P , we call the ideal of all smooth functions f such that $|f|^m \leq |g|$ for some $m \in \mathbb{N}$ and for some $g \in J$ the *full real radical* of J . If $q \in \mathbb{N}$ is given, we call the ideal of all smooth functions f such that $|f|^q \leq |g|$ for some $g \in J$ the *real radical of root order q* .

(2.2) *Key Features of Kohn's Algorithm.* The subelliptic estimate holds at a point P of $\partial\Omega$ when there is a scalar multiplier which is nonzero at P . Kohn's algorithm seeks to reduce the vanishing order of scalar multipliers by differentiation and by root-taking. The procedure of differentiation described in (B) in (2.1) above would only allow certain differential operators to lower the vanishing order of multipliers, namely only $(1, 0)$ -differentiation is allowed and only the determinants of coefficients of $(1, 0)$ -differentials of scalar multipliers (from Cramer's rule) can be used to produce new scalar multipliers. The procedure of root-taking described in (C) in (2.1) above identifies a smooth function germ as a scalar multiplier when a positive integral power of its absolute-value is dominated by the absolute-value of some known scalar multiplier.

(2.3) *Condition of Finite Type.* The *type* m at a point P of the boundary of weakly pseudoconvex Ω is the supremum of the normalized touching order

$$\frac{\text{ord}_0(r \circ f)}{\text{ord}_0 f},$$

to $\partial\Omega$, of all local holomorphic curves $f : \Delta \rightarrow \mathbb{C}^n$ with $\varphi(0) = P$, where Δ is the open unit 1-disk and ord_0 is the vanishing order at the origin 0. The notion of finite type was first introduced by Kohn in 1972 [Kohn1972, p.525, Def.2.3] for the case of $n = 2$ where the formulation is in terms of the nonvanishing of ∂r on the iterated Lie brackets of tangential vector fields to $\partial\Omega$ of types $(1, 0)$ and $(0, 1)$. It was extended to the case of a general n by D'Angelo in [D'Angelo1979, p.59] in terms of finite algebraic obstructions to the existence of a nontrivial holomorphic complex curve germ in $\mathbb{C}^n$ tangential to $\partial\Omega$ and then in the formulation in terms of normalized touching order in [D'Angelo1982, p.625, Definition 2.16 and Definition 2.18].

(2.4) *Kohn's Conjecture.* The goal of the theory of multipliers for the complex Neumann problem is to prove by using the procedures of generating new multipliers to prove that the function identically 1 is generated as a multiplier so that for a bounded weakly pseudoconvex domain Ω with finite type m the subelliptic estimate for some positive order $\varepsilon > 0$ holds and as a consequence the Kohn solution u of the $\bar{\partial}$ -equation $\bar{\partial}u = f$ on Ω with the right-hand side f smooth on $\bar{\Omega}$ is also smooth on $\bar{\Omega}$. Moreover, there is effectiveness in the use of the procedures to generate new multipliers so that ε is some explicit function of the type m of the domain Ω and its complex dimension n .

(2.5) *Full-Real-Radical Kohn Algorithm.* If the question of effectiveness is to be set aside, the following algorithm can be used for the generation of new multipliers in I_P .

(i) We start with the initial member r of I_P and denote by $I_P^{(0)}$ the ideal generated by this initial scalar multiplier r . Likewise, we start out with the initial members $(\partial\bar{\partial}r) \lrcorner \bar{\xi}$ of A_P for all choices of smooth $(1, 0)$ -vector fields ξ tangential to $\partial\Omega$ and denote by $A_P^{(0)}$ the module generated by these initial vector multipliers.

(ii) We use induction on the nonnegative integer ν to define $I_P^{(\nu)}$ and $A_P^{(\nu)}$ as follows. We add to $I_P^{(\nu)}$ all the multipliers $\det_{n-1}(\theta_1, \dots, \theta_{n-1})$ for $\theta_1, \dots, \theta_{n-1}$ in $A_P^{(\nu)}$ to form an ideal which we denote by $\hat{I}_P^{(\nu)}$. For the induction step from

ν to $\nu + 1$ we define $I_P^{(\nu+1)}$ as the *full real radical* of the ideal $\hat{I}_P^{(\nu)}$. We add to $A_P^{(\nu)}$ all the vector multipliers ∂F for $F \in I_P^{(\nu+1)}$ to form $A_P^{(\nu+1)}$.

We then hope that if the pseudoconvex domain Ω is of finite type m , the above construction of $I_P^{(\nu)}$ and $A_P^{(\nu)}$ by induction on ν will result in $I_P^{(\nu_m)}$ containing the function 1 for some ν_m which effectively depends on m and n . In other words, the algorithm *terminates*. In particular, the subelliptic estimate of some positive order $\varepsilon > 0$ holds. However, since there is no control on the order of the root-taking used in going from the ideal $\hat{I}_P^{(\nu)}$ to its *full real radical* $I_P^{(\nu+1)}$ there is no way for us to conclude that the order ε of the subellipticity proved depends effectively on m and n . For the purpose of our discussion we call this algorithm of using induction on ν to construct $I_P^{(\nu)}$ with the goal of ending up with $1 \in I_P^{(\nu_m)}$ the *full-real-radical Kohn algorithm*.

(2.6) *Effective Kohn Algorithm*. In order to end up with an effective order ε of subellipticity, it will turn out that one needs to modify the full-real-radical Kohn algorithm to use the following *effective Kohn algorithm*.

(i) The starting point for the effective Kohn algorithm is the same as for the full-real-radical Kohn algorithm. Again we start with the initial member r of I_P and denote by $I_P^{(0)}$ the ideal generated by this initial scalar multiplier r . Likewise, we start out with the initial members $(\partial\bar{\partial}r) \lrcorner \bar{\xi}$ of A_P for all choices of smooth $(1, 0)$ -vector fields ξ tangential to $\partial\Omega$ and denote by $A_P^{(0)}$ the module generated by these initial vector multipliers.

(ii) What is different with the effective Kohn algorithm is that we introduce a positive integer q_ν for every nonnegative integer index ν so that we take the real radical of root order q_ν instead of the full real radical in the step of going from $\nu - 1$ to ν . First we set $q_0 = 1$ and set $I_P^{(0,1)} = I_P^{(0)}$. This is motivated by the fact that the full real radical of $I_P^{(0)}$ is $I_P^{(0)}$ again because dr is nowhere zero on $\partial\Omega$. We use induction on the nonnegative integer ν to define a sequence of positive integers q_ν and then $I_P^{(\nu, q_\nu)}$ and $A_P^{(\nu, q_\nu)}$ as follows. We add to $I_P^{(\nu, q_\nu)}$ all the multipliers $\det_{n-1}(\theta_1, \dots, \theta_{n-1})$ for $\theta_1, \dots, \theta_{n-1}$ in $A_P^{(\nu, q_\nu)}$ and then use the resulting collection of multipliers to form an ideal which we denote by $\hat{I}_P^{(\nu, q_\nu)}$. For the induction step from ν to $\nu + 1$ we define $I_P^{(\nu+1, q_{\nu+1})}$ as the *real radical* of root order $q_{\nu+1}$ of the ideal $\hat{I}_P^{(\nu, q_\nu)}$. We add to $A_P^{(\nu, q_\nu)}$ all the vector multipliers ∂F for $F \in I_P^{(\nu+1, q_{\nu+1})}$ and then use the resulting collection of vector multipliers to form a module which is $A_P^{(\nu+1, q_{\nu+1})}$.

for the next step of going from ν to $\nu + 1$ in the construction by induction on ν .

We then hope that if the pseudoconvex domain Ω is of finite type m , the above construction of $I_P^{(\nu, q_\nu)}$ and $A_P^{(\nu, q_\nu)}$ by induction on ν will result in $I_P^{(\nu_m, q_{\nu_m})}$ containing the function 1 for some ν_m which effectively depends on m and n . Moreover, we hope that each q_ν for $0 \leq \nu \leq \nu_m$ depends also effectively on m and n so that the subelliptic estimate of some positive order $\varepsilon > 0$ holds with ε depending effectively on m and n .

Note that $I_P^{(\nu, \mu)}$ and $A_P^{(\nu, \mu)}$ are defined only when $\mu = q_\nu$. Instead of using the notations $I_P^{(\nu, q_\nu)}$ and $A_P^{(\nu, q_\nu)}$ we could have used the notations which depend only on ν , for example, $\tilde{I}_P^{(\nu)}$ and $\tilde{A}_P^{(\nu)}$. We prefer the clumsier notations $I_P^{(\nu, q_\nu)}$ and $A_P^{(\nu, q_\nu)}$ to highlight the effective choice of q_ν .

In the effective Kohn algorithm described above, no procedure is given to determine the sequence q_ν . The sequence q_ν is obtained by a rather complicated algebraic geometric argument. Since the purpose is to achieve $1 \in I_P^{(\nu_m, q_{\nu_m})}$, it suffices to describe explicitly how to use algebraic geometric techniques to determine the procedures of differentiation and root-taking of order $\leq q_\nu$ to construct scalar multipliers and vector multipliers from some initial scalar multipliers and vector multipliers to achieve $1 \in I_P^{(\nu_m, q_{\nu_m})}$.

Kohn's papers [Kohn1977, Kohn1979] discussed relations between subelliptic estimates, finite type property, and the termination of the full-real-radical Kohn algorithm for smooth weakly pseudoconvex domains. The relations can be summarized in the following Kohn conjecture, formulated separately for the full-real-radical Kohn algorithm and the effective Kohn algorithm.

(2.7) *Conjecture on Full-Real-Radical Kohn Algorithm.* For a bounded weakly pseudoconvex domain Ω in $\mathbb{C}^n$ with smooth boundary and of finite type m and for a point P of the boundary of Ω , in the ascending chain of multiplier ideals $I_P^{(\nu)}$ for $\nu \in \mathbb{N} \cup \{0\}$ there exists some $\nu^* \in \mathbb{N}$ such that $I_P^{(\nu^*)}$ contains the constant function 1. In other words, the chain of multiplier ideals $I_P^{(\nu)}$ for $\nu \in \mathbb{N} \cup \{0\}$ terminates at $\nu = \nu^*$.

If the positive integer ν^* depends effectively on m and n , then we say that the full-real-radical Kohn algorithm terminates effectively.

(2.8) *Conjecture on Effective Kohn Algorithm.* For a bounded weakly pseudoconvex domain Ω in $\mathbb{C}^n$ with smooth boundary and of finite type m and for a point P of the boundary of Ω , there exist $\tilde{\nu} \in \mathbb{N}$ and a sequence of positive numbers $q_1, \dots, q_{\tilde{\nu}}$ such that in the ascending chain of multiplier ideals $I_P^{(\nu, q_\nu)}$ for $0 \leq \nu \leq \tilde{\nu}$, the multiplier ideal $I_P^{(\tilde{\nu}, q_{\tilde{\nu}})}$ contains the constant function 1. Moreover, the positive integer $\tilde{\nu}$ and the sequence of positive integers $q_1, \dots, q_{\tilde{\nu}}$ depend effectively on m and n .

For the conjecture on the full-real-radical Kohn algorithm, when the boundary $\partial\Omega$ of the bounded weakly pseudoconvex domain Ω is assumed real-analytic, the finite type condition becomes a conclusion instead of an assumption. Kohn first showed that if the chain of multiplier ideals $I_P^{(\nu)}$ for $\nu \in \mathbb{N} \cup \{0\}$ does not terminate, then the boundary $\partial\Omega$ contains a local real-analytic subvariety of holomorphic dimension ≥ 1 ([Kohn1977, p.2215, Lemma 20] and [Kohn1979, p.113, Proposition 6.20]). Then Diederich-Fornaess proved that the real-analytic boundary of a bounded weakly pseudoconvex domain cannot contain a local real-analytic subvariety of holomorphic dimension ≥ 1 ([Diederich-Fornaess1978, p.373, Lemma 2] and [Diederich-Fornaess1978, p.374, Theorem 3]). This result of Kohn and Diederich-Fornaess holds not only for the case of $(0, 1)$ -forms but for the general case of $(0, q)$ -forms for $1 \leq q \leq n$. Since their method of proof is by contradiction, there is no effectiveness in the termination of the chain of multiplier ideals $I_P^{(\nu)}$.

(2.8.1) For the conjecture on Kohn's effective algorithm, the paper [Siu2010] introduces algebraic geometric techniques to study the problem by looking first at the special case of Ω being a special domain. For notational convenience we now consider a domain Ω in $\mathbb{C}^{n+1}$ instead of $\mathbb{C}^n$. A *special domain* Ω in $\mathbb{C}^{n+1}$ (with coordinates $w, z_1, \dots, z_n$) is a bounded domain given by

$$(2.8.1.1) \quad \operatorname{Re} w + \sum_{j=1}^N |F_j(z_1, \dots, z_n)|^2 < 0,$$

where $F_j(z_1, \dots, z_n)$ which is defined on some open neighborhood of $\bar{\Omega}$ in $\mathbb{C}^{n+1}$ depends only on the variables $z_1, \dots, z_n$ and is holomorphic in $z_1, \dots, z_n$ for each $1 \leq j \leq N$. In [Siu2010] the verification of the conjecture on Kohn's effective algorithm for the case of $n = 2$ (which means for special domains of complex dimension 3) was given in detail, with only indications for the case of special domains of general complex dimension. A rough outline was given there for the extension of the method of algebraic geometric

techniques first to the general real-analytic case and then the general smooth case. We carry out here the effective Kohn algorithm, which was introduced in [Siu2010], in a way which keeps track of the order of subellipticity in each step, in the context of comparing the full-real-radical Kohn algorithm and the effective Kohn algorithm.

In the chain of multiplier ideals $I_P^{(\nu)}$ in the conjecture for the full-real-radical Kohn algorithm, if for every ν there is an effective positive integer p_ν (i.e., dependent only on m and n) such that $\left(I_P^{(\nu+1)}\right)^{p_\nu} \subset I_P^{(\nu)}$, then we can use $p_\nu = q_\nu$ and the proof of the conjecture on Kohn's effective algorithm is simply reduced to the conjecture on the full-real-radical Kohn algorithm with effective termination. Unfortunately, there are simple examples, even for special domains of complex dimension 3, with no effective p_ν . This means that the conjecture for effective Kohn algorithm is different from the conjecture for the full-real-radical Kohn algorithm with effective termination. A simple example of this kind was given by Catlin and D'Angelo in [Catlin-D'Angelo2010], which we will discuss in §4 below.

(2.9) *Algebraic Geometric Techniques in Effective Kohn Algorithm.* We now explain how the algebraic geometric techniques precisely work to provide naturally the positive integer q_ν to make the process effective. The reason for considering special domains in $\mathbb{C}^{n+1}$ is that instead of ideals of smooth function germs on $\mathbb{C}^{n+1}$ we need only consider ideals of holomorphic function germs on $\mathbb{C}^n$. That is the reason why for notational convenience we suddenly consider domains in $\mathbb{C}^{n+1}$ instead of domains in $\mathbb{C}^n$. We focus on the case where each F_j vanish at the origin of $\mathbb{C}^n$ (for $1 \leq j \leq N$) so that the origin belongs to the boundary of Ω . We are concerned only with the problem of the subelliptic estimate at the origin.

The new notion of pre-multipliers needs to be introduced. A holomorphic function germ $f(z_1, \dots, z_n)$ on $\mathbb{C}^n$ at the origin is a *pre-multiplier* if its differential df is a vector multiplier at the origin. We now verify that the holomorphic function germs $F_1, \dots, F_N$ are pre-multipliers and the order of subellipticity of each dF_j is $\geq \frac{1}{4}$.

(2.9.1) *Levi Form and Initial Vector Multiplier for Special Domain.* First, for a special domain we write down explicit expressions for a tangent vector, its Levi form, and a smooth test $(1, 0)$ -form in the domain of the actual adjoint

$\bar{\partial}^*$ of $\bar{\partial}$. Since the defining function for the special domain Ω is

$$r = \operatorname{Re} w + \sum_{j=1}^N |F_j(z_1, \dots, z_n)|^2,$$

it follows that

$$\partial r = \frac{1}{2} dw + \sum_{j=1}^N \bar{F}_j dF_j$$

and all the $(1, 0)$ -vector tangential to $\partial\Omega$ at the origin are of the form

$$b \frac{\partial}{\partial w} + \sum_{k=1}^n a_k \frac{\partial}{\partial z_k}$$

for $b, a_1, \dots, a_n \in \mathbb{C}$ with

$$b = -2 \sum_{1 \leq j \leq N, 1 \leq k \leq n} a_k \bar{F}_j \frac{\partial F_j}{\partial z_k}.$$

The value of the Levi-form $\partial \bar{\partial} r = \sum_{j=1}^N dF_j \wedge \bar{dF}_j$ at $\eta \wedge \bar{\eta}$ for the element

$$\eta = -2 \left(\sum_{1 \leq j \leq N, 1 \leq k \leq n} a_k \bar{F}_j \frac{\partial F_j}{\partial z_k} \right) \frac{\partial}{\partial w} + \sum_{k=1}^n a_k \frac{\partial}{\partial z_k}$$

of $T_{\partial\Omega}^{(1,0)}$ is equal to

$$\sum_{1 \leq j \leq N, 1 \leq k \leq n} \left| a_k \frac{\partial F_j}{\partial z_k} \right|^2,$$

which means that a special domain is always weakly pseudoconvex and is strict pseudoconvex at a boundary point if and only if $dF_{j_1}, \dots, dF_{j_n}$ are $\mathbb{C}$ -linearly independent at that point for some $1 \leq j_1 < \dots < j_n \leq N$. For the element

$$\xi = -2 \left(\sum_{j=1}^N \bar{F}_j \frac{\partial F_j}{\partial z_\mu} \right) \frac{\partial}{\partial w} + \frac{\partial}{\partial z_\mu}$$

of $T_{\partial\Omega}^{(1,0)}$ (for some fixed $1 \leq \mu \leq n$) the $(1, 0)$ -form

$$(\partial \bar{\partial} r) \lrcorner \bar{\xi} = \sum_{\nu=1}^n \left(\sum_{j=1}^N (\partial_\nu F_j) \overline{(\partial_\mu F_j)} \right) dz_\nu$$

is a vector multiplier.

For an open neighborhood U of a point of $\partial\Omega$ in $\mathbb{C}^n$, if $\varphi = \sum_{\nu=1}^n \varphi_{\bar{\nu}} d\bar{z}_{\nu} + \hat{\varphi} d\bar{w}$ is a smooth test $(0, 1)$ -form on $U \cap \bar{\Omega}$ with compact support which belongs to the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$, then the normal component of φ vanishes on $U \cap \partial\Omega$, which means that

$$\begin{aligned} (\partial r) \cdot \varphi &= \left(\frac{1}{2} dw + \sum_{j=1}^N \overline{F_j} dF_j \right) \cdot \left(\sum_{\nu=1}^n \varphi_{\bar{\nu}} d\bar{z}_{\nu} + \hat{\varphi} d\bar{w} \right) \\ &= \frac{1}{2} \hat{\varphi} + \sum_{j=1}^N \sum_{\nu=1}^n \overline{F_j} (\partial_{\nu} F_j) \varphi_{\bar{\nu}} \end{aligned}$$

vanishes on $U \cap \partial\Omega$, or

$$\hat{\varphi} = -2 \sum_{j=1}^N \sum_{\nu=1}^n \overline{F_j} (\partial_{\nu} F_j) \varphi_{\bar{\nu}}$$

on $U \cap \partial\Omega$.

It will turn out that in the case of a special domain the use of multipliers and vector multipliers can be limited to those which are holomorphic in $z_1, \dots, z_n$ and are independent of w and as a consequence in the study of the subelliptic estimate for a special domain the component $\hat{\varphi}$ of a test $(0, 1)$ -form φ actually plays no role. (See (2.9.3) below.)

(2.9.2) *Initial Pre-Multiplier for Special Domain.* To verify that dF_j is a vector multiplier with order of subellipticity $\geq \frac{1}{4}$, we take an open neighborhood U of 0 on $\mathbb{C}^{n+1}$ on which the holomorphic functions $F_1, \dots, F_N$ of $(z_1, \dots, z_n) \in \mathbb{C}^n$ are defined. For $0 < \varepsilon \leq \frac{1}{2}$ and any smooth $(0, 1)$ -form $\varphi = \sum_{\nu=1}^n \varphi_{\bar{\nu}} d\bar{z}_{\nu} + \hat{\varphi} d\bar{w}$ on $U \cap \bar{\Omega}$ with compact support which belongs to

the domain of the *actual* adjoint $\bar{\partial}^*$ of $\bar{\partial}$, we have

$$\begin{aligned}
& \sum_{j=1}^N \left\| \left\| \sum_{\nu=1}^n (\partial_\nu F_j) \varphi_{\bar{\nu}} \right\| \right\|_\varepsilon^2 \\
&= \sum_{j=1}^N \sum_{\nu, \mu=1}^n (\Lambda^\varepsilon((\partial_\nu F_j) \varphi_{\bar{\nu}}), \Lambda^\varepsilon((\partial_\mu F_j) \varphi_{\bar{\mu}})) \\
&= \sum_{j=1}^N \sum_{\nu, \mu=1}^n \left(\overline{(\partial_\mu F_j)} \Lambda^{2\varepsilon}((\partial_\nu F_j) \varphi_{\bar{\nu}}), \varphi_{\bar{\mu}} \right) \\
&= \sum_{j=1}^N \sum_{\nu, \mu=1}^n \left(\Lambda^{2\varepsilon} \left(\overline{(\partial_\mu F_j)} (\partial_\nu F_j) \varphi_{\bar{\nu}} \right), \varphi_{\bar{\mu}} \right) + \sum_{j=1}^N \sum_{\nu, \mu=1}^n \left(\left[\overline{(\partial_\mu F_j)}, \Lambda^{2\varepsilon} \right] ((\partial_\nu F_j) \varphi_{\bar{\nu}}), \varphi_{\bar{\mu}} \right) \\
&= \sum_{\mu=1}^n \left(\Lambda^{2\varepsilon} \left(\sum_{\nu=1}^n \left(\left(\sum_{j=1}^N \overline{(\partial_\mu F_j)} (\partial_\nu F_j) \right) \varphi_{\bar{\nu}} \right) \right), \varphi_{\bar{\mu}} \right) \\
&\quad + \sum_{j=1}^N \sum_{\nu, \mu=1}^n \left(\left[\overline{(\partial_\mu F_j)}, \Lambda^{2\varepsilon} \right] ((\partial_\nu F_j) \varphi_{\bar{\nu}}), \varphi_{\bar{\mu}} \right) \\
&\leq \left(\sum_{\mu=1}^n \left\| \left\| \sum_{\nu=1}^n \left(\sum_{j=1}^N (\partial_\nu F_j) \overline{(\partial_\mu F_j)} \right) \varphi_{\bar{\nu}} \right\| \right\|_{2\varepsilon}^2 \right)^{\frac{1}{2}} \|\varphi\| + C_1 \|\varphi\|^2 \\
&\leq C_2 (\|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 + \|\varphi\|^2)
\end{aligned}$$

where C_1 and C_2 are positive constants independent of φ (but depend on U and ε), because

$$\sum_{\nu=1}^n \left(\sum_{j=1}^N (\partial_\nu F_j) \overline{(\partial_\mu F_j)} \right) dz_\nu$$

is a vector multiplier whose order of subellipticity is $\geq \frac{1}{2}$ for $1 \leq \nu \leq n$. This finishes the verification that each $dF_j = \sum_{\nu=1}^n (\partial_\nu F_j) dz_\nu$ is a vector multiplier whose order of subellipticity is $\geq \frac{1}{4}$.

(2.9.3) *Holomorphic Multipliers for Special Domain.* We start out with pre-multipliers $F_1, \dots, F_N$ and the vector multipliers $dF_1, \dots, dF_N$ which are holomorphic 1-forms in the variables $z_1, \dots, z_n$ obtained from them. For the special domain Ω we will only work with vector multipliers which are

holomorphic 1-forms in the variables $z_1, \dots, z_n$. When we have n such vector multipliers $G^{(j)} = \sum_{\nu=1}^n G^{(j)}(z_1, \dots, z_n) dz_j$ of order of subellipticity $\geq \varepsilon_\nu$ for $1 \leq \nu \leq n$, when we apply the procedure (B)(ii) in (2.1) above to generate new multipliers, from

$$\partial r = \frac{1}{2} dw + \sum_{j=1}^N \overline{F_j} dF_j$$

we get

$$G^{(1)} \wedge \dots \wedge G^{(n)} \wedge \partial r = \frac{1}{2} \det \left(G_j^{(\nu)} \right)_{1 \leq \nu, j \leq n} dz_1 \wedge \dots \wedge dz_n \wedge dw$$

to conclude that the holomorphic function

$$\det \left(G_j^{(\nu)} \right)_{1 \leq \nu, j \leq n}$$

of the variables $z_1, \dots, z_n$ is a scalar multiplier whose order of subellipticity is $\geq \min(\varepsilon_1, \dots, \varepsilon_n)$. So for the special domain Ω , when we start out only with vector multipliers $dF_1, \dots, dF_N$ (whose orders of subellipticity are all $\geq \frac{1}{4}$) and use only the procedures in B(i)(ii) and C described in (2.1) above to generate new scalar and vector multipliers, we need only work with scalar and vector multipliers which are holomorphic functions or holomorphic 1-forms in the variables $z_1, \dots, z_n$.

We now translate the algorithm from the language of analysis to the language of algebraic geometry. The procedures in the algorithm now read as follows.

(2.9.4) *Algebraic Geometric Formulation of Kohn Algorithm for Special Domain.* The multiplicity q of the ideal generated by $F_1, \dots, F_N$ given by

$$\dim_{\mathbb{C}} \left(\mathcal{O}_{\mathbb{C}^n, 0} \left/ \sum_{j=1}^N \mathcal{O}_{\mathbb{C}^n, 0} F_j \right. \right)$$

is related to the type m of the special domain Ω at the origin by $2m \leq q \leq (n+2)2m$. See [Siu2010, Lemma(I.3) and Lemma(I.4)]. In particular, the special domain is of finite type at 0 if and only if 0 is an isolated point of the common zero-set of $F_1, \dots, F_N$. Assume that the multiplicity q of the ideal generated by $F_1, \dots, F_N$ is finite. The N pre-multipliers $F_1, \dots, F_N$ (with order of subellipticity of each dF_j at least $\frac{1}{4}$) are all that we start out with. For a special domain of complex dimension $n+1$ there are only the following two procedures from Kohn's algorithm.

(i) If holomorphic function germs $g_1, \dots, g_n$ at 0 on $\mathbb{C}^n$ are pre-multipliers (which automatically include all multipliers), then the coefficient of $dz_1 \wedge \dots \wedge dz_n$ in $dg_1 \wedge \dots \wedge dg_n$ is a multiplier. In other words, the Jacobian determinant

$$\frac{\partial(g_1, \dots, g_n)}{\partial(z_1, \dots, z_n)}$$

of the holomorphic functions $g_1, \dots, g_n$ with respect to the variables $z_1, \dots, z_n$ is a multiplier. If the order of subellipticity of each $g_1, \dots, g_n$ is $\geq \eta$, then the order of subellipticity of their Jacobian determinant is $\geq \frac{\eta}{2}$.

(ii) If g and f are holomorphic function germs at 0 on $\mathbb{C}^n$ with $f^m = g$ for some positive integer m and if g is a multiplier whose order of subellipticity is $\geq \eta$, then f is also a multiplier whose order of subellipticity is $\geq \frac{\eta}{m}$.

Note that the set of all multipliers forms an ideal in the ring of all holomorphic function germs and the set of all vector-multipliers form a module over the ring of all holomorphic function germs, but in general the set of all pre-multipliers does *not* form a module over the ring of all holomorphic functions, because though the differential dF of a pre-multiplier F is a multiplier, yet for any holomorphic function germ g the differential $d(gF)$ is equal to $gdF + Fdg$ and the term Fdg , unlike the term gdF , is in general *not* a vector-multiplier.

We now discuss the algebraic geometric formulations of the steps in both the full-real-radical Kohn algorithm and the effective Kohn algorithm. We will describe first the steps in the full-real-radical Kohn algorithm. Then we will describe the effective Kohn algorithm but only in the case when the special domain is of complex dimension 3.

(2.10) *Steps in Full-Real-Radical Kohn Algorithm.* We start out with the $\mathbb{C}$ -vector space V_0 of initial pre-multipliers generated by $F_1, \dots, F_N$. Let J_0 be the ideal generated by all the Jacobian determinants of any n elements $g_1, \dots, g_n$ of V_0 . Let I_0 be the radical of J_0 .

Inductively we construct the $\mathbb{C}$ -vector space V_ν , the ideal J_ν , and the ideal I_ν for any nonnegative integer ν as follows. For the step of going from ν to $\nu + 1$, we let $V_{\nu+1}$ be the $\mathbb{C}$ -vector space generated by elements of V_ν and all elements of the ideal I_ν . Let $J_{\nu+1}$ be the ideal generated by all the Jacobian determinants of any n elements of $V_{\nu+1}$. Let $I_{\nu+1}$ be the radical of the ideal of $J_{\nu+1}$. This finishes the construction by induction.

Let p_ν be the smallest positive integer such that $I_\nu^{p_\nu}$ is contained in J_ν . Note that each element of I_ν is a multiplier, but each element of V_ν is only a pre-multiplier.

The full-real-radical Kohn algorithm for $F_1, \dots, F_N$ *terminates* if there exists some nonnegative $\tilde{\nu}$ such that $I_{\tilde{\nu}}$ is the unit ideal, which means the entire ring of all holomorphic function germs. In that case we choose $\tilde{\nu}$ to be the smallest such nonnegative integer.

We say that the full-real-radical Kohn algorithm for $F_1, \dots, F_N$ *terminates effectively* if $\tilde{\nu}$ and each p_ν for $0 \leq \nu \leq \tilde{\nu}$ are bounded by explicit functions of n and q .

The order of subellipticity for the multiplier 1 is at least

$$\frac{1}{2^{\tilde{\nu}+2} \prod_{\nu=0}^{\tilde{\nu}} p_\nu}.$$

This order of subellipticity from the termination of the original Kohn algorithm is effective only when $\tilde{\nu}$ is effective and each p_ν is effective for each $0 \leq \nu \leq \tilde{\nu}$.

(2.11) *Ideal Containing an Effective Power of its Radical in Effective Kohn Algorithm.* The key difference between the full-real-radical Kohn algorithm (which in general is not effective) by the effective Kohn algorithm is to replace the taking of the radical I_ν of the ideal J_ν by an appropriately chosen sub-ideal $\tilde{I}_\nu$ of I_ν with the property that an effective power $(\tilde{I}_\nu)^{s_\nu}$ of $\tilde{I}_\nu$ is contained in J_ν . The choice of the sub-ideal $\tilde{I}_\nu$ of I_ν and the positive integer s_ν involves rather complicated algebraic geometric techniques. In order to facilitate the explicit comparison of the full-real-radical Kohn algorithm with the effective Kohn algorithm in a concrete example (such as the example of Catlin-D'Angelo [Catlin-D'Angelo2010]), in the description of the steps in the effective Kohn algorithm for special domains we will confine ourselves to special domains of complex dimension 3, which means the case of $n = 2$.

§3. ORDERS OF SUBELLIPTICITY IN ALGEBRAIC GEOMETRIC TECHNIQUES FOR 3-DIMENSIONAL SPECIAL DOMAIN

We now describe the steps in effective Kohn algorithm for special domains of complex dimension 3. Again we start with holomorphic function germs $F_1, \dots, F_N$ at 0 on $\mathbb{C}^2$ (which define the special domain in $\mathbb{C}^3$ and which generates an ideal in $\mathcal{O}_{\mathbb{C}^2,0}$ of multiplicity $\leq q$).

(3.1) *Step One.* We take two generic $\mathbb{C}$ -linear combinations $\hat{F}_1, \hat{F}_2$ of $F_1, \dots, F_N$ of vanishing order $\leq q$ at 0 such that the Jacobian determinant

$$h_2^* := \frac{\partial(\hat{F}_1, \hat{F}_2)}{\partial(z_1, z_2)}$$

has vanishing order $\leq q$ at 0 as a holomorphic function germ at 0 on $\mathbb{C}^2$. An order of subellipticity of h_2^* as a multiplier at the origin is at least $\frac{1}{4}$. Let

$$h_2^* = (h_{2,1}^*)^{k_1} \cdots (h_{2,\ell_2}^*)^{k_{\ell_2}}$$

be the factorization into irreducible holomorphic function germs $h_{2,1}^*, \dots, h_{2,\ell_2}^*$ with $k_1 \geq k_2 \geq \dots \geq k_{\ell_2} \geq 1$. Since the vanishing order of h_2^* is $\leq q$, we have $k_1 + \dots + k_{\ell_2} \leq q$ and, in particular, $k_1 \leq q$. The holomorphic function germ

$$(h_{2,1}^* \cdots h_{2,\ell_2}^*)^{k_1}$$

contains h_2^* as a factor and is therefore a multiplier whose order of subellipticity is $\geq \frac{1}{4}$. Let

$$\hat{h}_2 = h_{2,1}^* \cdots h_{2,\ell_2}^*.$$

Since the k_1 -th power of $\hat{h}_2$ is a multiplier, it follows from $k_1 \leq q$ that $\hat{h}_2$ is a multiplier whose order of subellipticity is $\geq \frac{1}{4q}$. The construction of $\hat{h}_2$ from h_2^* is to make sure that the divisor of $\hat{h}_2$ is reduced (which means that its multiplicity at any of its regular points is 1, though it is possibly reducible with many branches). Up to this point the only goal accomplished is to produce a *reduced* holomorphic function germ $\hat{h}_2$ at 0, with vanishing order $\leq q$ at 0, which is a multiplier whose order of subellipticity is $\geq \frac{1}{4q}$. This step of construction of $\hat{h}_2$ from h_2^* is included only for the sake of convenience and is not actually absolutely necessary.

(3.2) *Step Two.* Take a generic $\mathbb{C}$ -linear combination h_1 of $F_1, \dots, F_N$ and a generic $\mathbb{C}$ -linear coordinate system w_1, w_2 of $\mathbb{C}^2$ such that the following three conditions (i), (ii) and (iii) are satisfied. Here generic means that $h_1 = \sum_{j=1}^N a_j F_j$ and $w_k = \sum_{\ell=1}^2 g_{k\ell} z_\ell$ with the element $(a_j, g_{k\ell})_{1 \leq j \leq N, 1 \leq k, \ell \leq 2}$ of $\mathbb{C}^{N+1}$ chosen outside some proper subvariety $\mathcal{Z}$ of $\mathbb{C}^{N+4}$. The proper subvariety $\mathcal{Z}$ of $\mathbb{C}^{N+4}$ can be obtained as the union of three proper subvarieties of $\mathbb{C}^{N+4}$, one for each of the three conditions (i), (ii) and (iii).

- (i) The origin 0 of $\mathbb{C}^2$ is an isolated zero of w_1 and the holomorphic function germs h_1 .
- (ii) The multiplicity at 0 of the ideal generated by h_1 and $\hat{h}_2$ is $\leq q^2$.
- (iii) The multiplicity at 0 of the ideal generated by $\hat{h}_2$ and $\left(\frac{\partial h_1}{\partial w_1}\right)_{w_2=\text{const}}$ is $\leq 3q^2$.

The reason why a generic $\mathbb{C}$ -linear combination h_1 of $F_1, \dots, F_N$ satisfies Condition (ii) is that the multiplicity of the ideal generated by $F_1, \dots, F_N$ at the origin is $\leq q$. The number q^2 in Condition (iii) is the product of the multiplicity q of $\hat{h}_1$ at 0 and the multiplicity q of the ideal generated by $F_1, \dots, F_N$ at 0.

A choice of a generic $\mathbb{C}$ -linear combination h_1 of $F_1, \dots, F_N$ and a generic $\mathbb{C}$ -linear coordinate system w_1, w_2 of $\mathbb{C}^2$ satisfying Condition (iii) is obtained from the following statement.

(3.2.1) For each holomorphic function germ f on $\mathbb{C}^n$ at 0 which vanishes at 0, the function germ f^{n+1} on $\mathbb{C}^n$ at 0 belongs to the ideal generated by $\frac{\partial f}{\partial z_j}$ for $1 \leq j \leq n$ (where $z_1, \dots, z_n$ are the coordinates of $\mathbb{C}^n$).

The statement (3.2.1) is a consequence of Skoda's result on ideal generation [Skoda1972] (see [Siu2010, p.1232, Propositioni(A.2)]) and can be considered as a generalization, to general holomorphic function germs, of Euler's formula expressing a homogeneous polynomials in terms of its first-order partial derivatives. We apply the statement to $f = h_1$ to conclude that $(h_1)^3$ belongs to the ideal generated by $\frac{\partial h_1}{\partial z_1}, \frac{\partial h_1}{\partial z_2}$. Condition (ii) implies that a generic $\mathbb{C}$ -linear combination $H = \sum_{k=1}^2 c_k \frac{\partial h_1}{\partial z_k}$ satisfies the condition that the multiplicity at 0 of the ideal generated by H and $\hat{h}_2$ is $\leq 3q^2$. We can choose a generic $\mathbb{C}$ -linear coordinate system w_1, w_2 of $\mathbb{C}^2$ such that $H = \left(\frac{\partial h_1}{\partial w_1}\right)_{w_2=\text{const}}$.

We are now ready to construct more multipliers from the choice of h_1 and w_1, w_2 . Since h_1 vanishes at 0, it follows Condition (iii) that there exist holomorphic function germs α, β at 0 on $\mathbb{C}^2$ such that

$$(h_1)^{3q^2} = \alpha \hat{h}_2 + \beta \left(\frac{\partial h_1}{\partial w_1}\right)_{w_2=\text{const}}.$$

Though h_1 which is a $\mathbb{C}$ -linear combination $F_1, \dots, F_N$ of pre-multipliers $F_1, \dots, F_N$ may not be a multiplier, the long key argument given below is

to show that actually h_1 is a multiplier. We need the following statement concerning Weierstrass polynomials.

(3.2.2) *Weierstrass Polynomial for Image Curve Under Branched-Cover Map.* Let ζ_1 and ζ_2 be holomorphic function germs at 0 on $\mathbb{C}^2$ vanishing at 0 such that the origin 0 of $\mathbb{C}^2$ is an isolated point of the common zero-set of ζ_1 and ζ_2 . Let H be a holomorphic function germ at 0 on $\mathbb{C}^2$ vanishing at 0 such that the origin 0 of $\mathbb{C}^2$ is an isolated point of the common zero-set of H and ζ_1 . Let ℓ be a positive integer. If ζ_2^ℓ belongs to the ideal generated by H and ζ_1 , then some holomorphic function germ on $\mathbb{C}^2$ at 0 of the form of a Weierstrass polynomial

$$\zeta_2^\ell + \sum_{j=0}^{\ell-1} \theta_j(\zeta_1) \zeta_2^j$$

(with ζ_1, ζ_2 as variables) contains H as a factor, where θ_j is a holomorphic function germ on $\mathbb{C}$ at 0 which vanishes at 0 for $0 \leq j \leq \ell - 1$.

For the proof of the statement (3.2.2) on Weierstrass polynomials, first we observe that for the special case where ζ_1, ζ_2 are the coordinate functions z_1, z_2 of $\mathbb{C}^2$ and the restriction of H to $\{z_1 = 0\}$ is equal to z_2^ℓ as holomorphic function germ on $\mathbb{C}$ with z_2 as coordinate, the statement is simply the usual factorization of a holomorphic function germ H as a product of a nowhere zero holomorphic function germ and a Weierstrass polynomial of degree ℓ in the variable z_2 . For the proof of the general case, we consider the germ at 0 of the holomorphic map $\pi : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ defined by $(z_1, z_2) \rightarrow (\zeta_1, \zeta_2)$. Since the origin 0 of $\mathbb{C}^2$ is an isolated point of the common zero-set of ζ_1 and ζ_2 , the map π is an analytic branched cover (as the germ of a holomorphic map). Let C be the divisor of H and $\tilde{C}$ be the image of C under π (with multiplicities counted) and let $\tilde{H}$ be a holomorphic function germ on $\mathbb{C}^2$ at 0 whose divisor is $\tilde{C}$. Since the origin is an isolated point of the common zero-set of H and ζ_1 and since ζ_2^ℓ belongs to the ideal generated by ζ_1 and H , it follows that the restriction of $\tilde{H}$ to $\{\zeta_1 = 0\}$ is equal to $\zeta_2^{\tilde{\ell}}$ as holomorphic function germ on $\mathbb{C}$ with ζ_2 as coordinate for some positive integer $\tilde{\ell} \leq \ell$. The general case now follows from applying the special case when H is replaced by $\zeta_2^{\ell-\tilde{\ell}} \tilde{H}$ and the coordinates z_1, z_2 are replaced by ζ_1, ζ_2 .

(3.3) *Step Three.* Because of Condition (ii) in (3.2), we can now apply the second part of the statement (3.2.2) on Weierstrass polynomials to the case

of $H = \hat{h}_2$ and $\zeta_1 = h_1$ and $\zeta_2 = w_2$ to get a holomorphic function germ h_2 of the form

$$w_2^{q^2} + \sum_{j=0}^{q^2-1} \theta_j(h_1) w_2^j$$

which contains $\hat{h}_2$ as a factor. (The property of $\hat{h}_2$ being a reduced holomorphic function germ means that in applying the above statement on Weierstrass polynomial, the divisor C in the proof of the statement (3.2.2) on Weierstrass polynomials is reduced and we do not have to worry about multiplicities of its branches, but this point, though offering some convenience, is not essential.) Since h_2 contains as a factor the multiplier $\hat{h}_2$ whose order of multiplicity is $\geq \frac{1}{4q}$, it follows h_2 is a multiplier whose order of subellipticity is $\geq \frac{1}{4q}$. Let $h_{2,0} = h_2$ and for $1 \leq \nu \leq q^2$ let

$$h_{2,\nu} = q^2(q^2 - 1) \cdot (q^2 - \nu + 1) w_2^{q^2-\nu} + \sum_{j=0}^{q^2-1} j(j-1) \cdots (j-\nu+1) \theta_j(h_1) w_2^{j-\nu},$$

which is obtained by differentiating ν -times the function h_2 with respect to w_2 with h_1 fixed when h_2 is regarded as a function of h_1 and w_2 . Then

$$dh_{2,\nu} = \eta_\nu dh_1 + h_{2,\nu+1} dw_2 \quad \text{for } 0 \leq \nu \leq q^2 - 1$$

for a holomorphic function germ η_ν which is the partial derivative of $h_{2,\nu}$ with respect to h_1 with w_2 fixed when $h_{2,\nu}$ is regarded as a function of h_1 and w_2 . Let

$$\tilde{h}_{2,1} = \left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} h_{2,1}.$$

From

$$\begin{aligned} dh_1 \wedge dh_2 &= dh_1 \wedge (\eta_0 dh_1 + h_{2,1} dw_2) \\ &= dh_1 \wedge h_{2,1} dw_2 \\ &= \left(\left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} dw_1 + \left(\frac{\partial h_1}{\partial w_2} \right)_{w_1=\text{const}} dw_2 \right) \wedge h_{2,1} dw_2 \\ &= \tilde{h}_{2,1} dw_1 \wedge dw_2, \end{aligned}$$

it follows that $\tilde{h}_{2,1}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{8q}$.

Since $\hat{h}_2 h_{2,1}$, being a multiple of the multiplier $\hat{h}_2$, is itself a multiplier, it follows that the linear combination

$$\alpha \hat{h}_2 h_{2,1} + \beta \tilde{h}_{2,1} = \alpha \hat{h}_2 h_{2,1} + \beta \left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} h_{2,1} = (h_1)^{3q^2} h_{2,1}$$

of the two multipliers $\hat{h}_2 h_{2,1}$ and $\tilde{h}_{2,1}$ with coefficients, which are holomorphic function germs at 0 on $\mathbb{C}^2$, is a multiplier whose order of subellipticity is $\geq \frac{1}{8q}$.

(3.4) *Recursive Argument in Step Three.* Now we repeat the above argument with $h_{2,\nu}$ replacing $\hat{h}_2$ in the following way to conclude by induction on ν that $(h_1)^{3q^2\nu} h_{2,\nu}$ is a multiplier with order of subellipticity $\geq \frac{1}{2\nu+2q}$ for $1 \leq \nu \leq q^2$.

The case of $\nu = 1$ was just proved. Suppose we have proved the step up to some $\nu < q^2$ and we would like to prove the next step of $\nu + 1$. From

$$\begin{aligned} & dh_1 \wedge d \left((h_1)^{3q^2\nu} h_{2,\nu} \right) \\ &= dh_1 \wedge \left(\left((3q^2\nu - 1)(h_1)^{3q^2\nu-1} h_{2,\nu} + (h_1)^{3q^2\nu} \eta_\nu \right) dh_1 + (h_1)^{3q^2\nu} h_{2,\nu+1} dw_2 \right) \\ &= dh_1 \wedge (h_1)^{3q^2\nu} h_{2,\nu+1} dw_2 \\ &= \left(\left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} dw_1 + \left(\frac{\partial h_1}{\partial w_2} \right)_{w_1=\text{const}} dw_2 \right) \wedge (h_1)^{3q^2\nu} h_{2,\nu+1} dw_2 \\ &= \left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} (h_1)^{3q^2\nu} h_{2,\nu+1} dw_1 \wedge dw_2 \end{aligned}$$

it follows that $\left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} (h_1)^{3q^2\nu} h_{2,\nu+1}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{2\nu+3q}$.

Since $\hat{h}_2 (h_1)^{3q^2\nu} h_{2,\nu+1}$, being a multiple of the multiplier $\hat{h}_2$, is itself a multiplier, it follows that the linear combination

$$\begin{aligned} & \alpha \hat{h}_2 (h_1)^{3q^2\nu} h_{2,\nu+1} + \beta \left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} (h_1)^{3q^2\nu} h_{2,\nu+1} \\ &= \left(\alpha \hat{h}_2 + \beta \left(\frac{\partial h_1}{\partial w_1} \right)_{w_2=\text{const}} \right) (h_1)^{3q^2\nu} h_{2,\nu+1} \\ &= (h_1)^{3q^2(\nu+1)} h_{2,\nu+1} \end{aligned}$$

of the two multipliers $\hat{h}_2(h_1)^{3q^2\nu}h_{2,\nu+1}$ and $\beta\left(\frac{\partial h_1}{\partial w_1}\right)_{w_2=\text{const}}(h_1)^{3q^2\nu}h_{2,\nu+1}$ with coefficients, which are holomorphic function germs at 0 on $\mathbb{C}^2$, is a multiplier with order of subellipticity $\geq \frac{1}{2\nu+3q}$. This finishes the proof by induction on ν that $(h_1)^{3q^2\nu}h_{2,\nu}$ is a multiplier with order of subellipticity $\geq \frac{1}{2\nu+2q}$ for $1 \leq \nu \leq q^2$. Since h_{2,q^2} is equal to $(q^2)!$, it follows that $(h_1)^{3q^4}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{2q^2+2q}$. By the real radical property of multipliers we conclude that h_1 is a multiplier whose order of subellipticity is $\geq \frac{1}{3q^52q^2+2}$.

Since by Condition (ii) of (3.2) the ideal generated by h_1 and $\hat{h}_2$ contains the q^2 -th power of the maximum ideal $\mathfrak{m}_{\mathbb{C}^2,0}$ of $\mathbb{C}^2$ at 0, it follows that w_1 and w_2 are multipliers whose order of subellipticity is $\geq \frac{1}{3q^72q^2+2}$. By taking the Jacobian determinant of w_1 and w_2 , we end up with 1 being a multiplier whose order of subellipticity is $\frac{1}{3q^72q^2+3}$.

(3.5) *Remark.* For use in (4.3) below we would like to remark that the above arguments work in the same way when the holomorphic function germ h_2 is chosen to be

$$(h_1)^r \left(w_2^{q^2} + \sum_{j=0}^{q^2-1} \theta_j(h_1)w_2^j \right)$$

instead of

$$w_2^{q^2} + \sum_{j=0}^{q^2-1} \theta_j(h_1)w_2^j$$

if r is effective in the sense that r is bounded by an explicit function of q . Of course, the effective lower bound of the order of subellipticity of the constant function 1 as a multiplier needs to be correspondingly modified to be $\frac{1}{3q^72q^2+r+3}$.

§4. EFFECTIVE KOHN ALGORITHM APPLIED TO CATLIN-D'ANGELO'S EXAMPLE

We now apply the algebraic geometric techniques in the effective Kohn algorithm to the example of Catlin-D'Angelo given in [Catlin-D'Angelo2010] for which the full-real-radical Kohn algorithm is ineffective.

(4.1) *Catlin-D'Angelo's Example of Ineffectiveness of Full-Real-Radical Kohn Algorithm.* Let $K > M \geq 2$ and $N \geq 3$. The special domain Ω

$$\operatorname{Re} w + |F_1(z_1, z_2)|^2 + |F_2(z_1, z_2)|^2 < 0$$

in $\mathbb{C}^3$ is defined by the two holomorphic functions $F_1(z_1, z_2) = z_1^M$ and $F_2 = z_2^N + z_2 z_1^K$ on $\mathbb{C}^2$. The origin of $\mathbb{C}^3$ is the boundary point of Ω whose scalar and vector multipliers we consider. The following is reproduced from pp.81-82 of [Catlin-D'Angelo2010] in the notations and terminology used in this note. By Weierstrass division (applied to the Weierstrass polynomial which is the product F_2 and a nowhere holomorphic function germ), modulo F_2 every element of $\mathcal{O}_{\mathbb{C}^2,0}$ is equal to

$$\sum_{j=0}^{N-1} a_j(z_1) z_2^j$$

for some holomorphic function germs $a_0, \dots, a_{N-1}$ on $\mathbb{C}^2$ at 0. Modulo F_1 we can write

$$a_j(z_1) = \sum_{k=0}^{M-1} b_{jk} z_1^k$$

for $0 \leq j \leq N-1$, where $b_{jk} \in \mathbb{C}$ for $0 \leq j \leq N-1$, $0 \leq k \leq M-1$. Hence the multiplicity q given by

$$\dim_{\mathbb{C}} \left(\mathcal{O}_{\mathbb{C}^2,0} \left/ \sum_{j=1}^2 \mathcal{O}_{\mathbb{C}^2,0} F_j \right. \right)$$

is $\leq MN$.

The full-real-radical Kohn algorithm proceeds as follows in this example. Let g be the Jacobian determinant

$$\frac{\partial(F_1, F_2)}{\partial(z_1, z_2)} = \det \begin{pmatrix} z_1^{M-1} & 0 \\ K z_2 z_1^{K-1} & N z_2^{N-1} + z_1^K \end{pmatrix} = N z_1^{M-1} z_2^{N-1} + z_1^{K+M-1}$$

of F_1, F_2 with respect to z_1, z_2 . We use the notations in (2.10). The ideal J_0 is the principal ideal with the irreducible function germ g as the generator and its radical I_0 is the same as J_0 . The ideal J_1 is

$$\mathcal{O}_{\mathbb{C}^2,0} g + \sum_{j=1}^2 \mathcal{O}_{\mathbb{C}^2,0} \frac{\partial(F_j, g)}{\partial(z_1, z_2)},$$

where

$$\frac{\partial(F_1, g)}{\partial(z_1, z_2)} = \det \begin{pmatrix} z_1^{M-1} & 0 \\ \frac{\partial g}{\partial z_1} & N(N-1)z_1^{M-1}z_2^{N-2} \end{pmatrix} = N(N-1)z_1^{2M-2}z_2^{N-2}.$$

Since g^2 modulo $\frac{\partial(F_1, g)}{\partial(z_1, z_2)}$ is equal to $z_1^{2(K+M-1)}$, it follows from $K+M-1 \geq 1$ that the holomorphic function germ z_1 at 0 belongs to the radical I_1 of J_1 . From $N \geq 3$ we conclude that modulo $(z_2)^2$ the three holomorphic function germs g, F_1, F_2 become respectively $z_1^{K+M-1}, z_1^M, z_2 z_1^K$. Hence the ideal J_1 generated by the three Jacobian determinants formed from pairs out of g, F_1, F_2 is contained in the ideal generated by z_1^{M+K-2} and z_2 . This means that z_1^m cannot be in J_1 for $m < M+K-2$, otherwise z_1^m belongs to the ideal generated by z_1^{M+K-2} and z_2 , which is a contradiction. Since the holomorphic function germ z_1 belongs to I_1 , this means that the smallest positive integer p_1 satisfying $(I_1)^{p_1} \subset J_1$ must be $\geq M+K-2 \geq K$. Thus, the full-real-radical Kohn algorithm is not effective, because K is arbitrary and there cannot be any function of NM which bounds K .

(4.2) *Remark.* In (4.1) when we carry out the full-real-radical Kohn algorithm for Catlin-D'Angelo's example, we stopped after showing the algorithm to be ineffective. For later comparison, we now carry out the remaining steps of the algorithm until we produce the constant function 1 as a multiplier. We have seen that z_1 belongs to I_1 . Since all three holomorphic function germs $g, \frac{\partial g}{\partial z_1}$ and $\frac{\partial g}{\partial z_2}$ contain z_1 as a factor, it follows that J_1 is contained in the principal ideal generated by z_1 and I_1 must be equal to the principal ideal generated by z_1 . The function germ $z_2^N = F_2 - z_2 z_1^K$ is a pre-multiplier in V_2 . The Jacobian determinant $\frac{\partial(z_1, z_2^N)}{\partial(z_1, z_2)} = N z_2^{N-1}$ belongs to J_2 . Hence z_2 belongs to I_2 and we can conclude that I_2 is the maximum ideal $\mathfrak{m}_{\mathbb{C}^2, 0}$ of $\mathbb{C}^2$ at 0. By taking the Jacobian determinant of the elements z_1, z_2 of I_2 , we conclude that 1 is a multiplier. To get to the multiplier 1 from F_1, F_2 , we have to perform differentiation 4 times in the construction of Jacobian determinants.

(4.3) *Effective Kohn Algorithm for Catlin-D'Angelo's Example.* We now carry out concretely the steps in the effective Kohn algorithm for Catlin-D'Angelo's example to illustrate the difference between the full-real-radical Kohn algorithm and the effective Kohn algorithm.

The key point in the effective Kohn algorithm is to construct a Weierstrass polynomial h_2 in one coordinate w_2 such that (i) h_2 contains as a

factor a multiplier $\hat{h}_2$ which is obtained in a procedure involving the Jacobian determinant of two $\mathbb{C}$ -linear combinations of the defining holomorphic functions $F_1, \dots, F_N$ of the special domain and (ii) the coefficients of h_2 are holomorphic function germs of some $\mathbb{C}$ -linear combination h_1 of $F_1, \dots, F_N$. Then by using induction on ν we show, with effectiveness, that the Jacobian determinant of h_1 and the function $(h_1)^{m_\nu} \left(\frac{\partial^\nu h_2}{\partial w_2^\nu} \right)_{h_1=\text{constant}}$ is a multiplier (for some effective positive integer m_ν), resulting finally in the conclusion that h_1 is a multiplier. In the key argument the Weierstrass polynomial h_2 can be replaced by the product of an effective power of h_1 and a Weierstrass polynomial (see Remark (3.4)).

In the example of Catlin-D'Angelo where the defining functions for the special domain in $\mathbb{C}^3$ are the two holomorphic functions $F_1(z_1, z_2) = z_1^M$ and $F_2 = z_2^N + z_2 z_1^K$ on $\mathbb{C}^2$, we can use $h_1 = F_1$ and use

$$\frac{z_1}{N} \frac{\partial(F_1, F_2)}{\partial(z_1, z_2)} = z_1^M z_2^{N-1} + \frac{1}{N} z_1^{K+M}$$

as h_2 which is the product of h_1 and the Weierstrass polynomial

$$z_2^{N-1} + \frac{1}{N} z_1^K = z_2^{N-1} + \frac{1}{N} (h_1)^m$$

in the variable z_2 , where $m = \frac{K}{M}$ which we assume for the time being to be a positive integer. It turns out that the argument used in the effective Kohn algorithm works in the same way without the assumption that $m = \frac{K}{M}$ is a positive integer. This assumption used in the setup merely motivates the steps of the argument.

For Catlin-D'Angelo's example, the induction on ν to show, with effectiveness, that the Jacobian determinant of h_1 and the function $(h_1)^{m_\nu} \left(\frac{\partial^\nu h_2}{\partial w_2^\nu} \right)_{h_1=\text{constant}}$ is a multiplier (for some effective positive integer m_ν) is translated (after obvious modifications) to verifying by induction on j that each H_j defined by $H_j = z_1^{(j+1)(M-1)} z_2^{N-j}$ is a multiplier for $1 \leq j \leq N$, because the ν -th derivative of

$$z_2^{N-1} + \frac{1}{N} z_1^K = z_2^{N-1} + \frac{1}{N} (h_1)^m$$

with respect to z_2 with h_1 being kept constant is $(N-1) \cdots (N-\nu) z_2^{N-1-\nu}$ for $1 \leq \nu \leq N-1$.

At this point we can forget that the use of $H_j = z_1^{(j+1)(M-1)} z_2^{N-j}$ for $1 \leq j \leq N$ is motivated by the steps of the effective Kohn algorithm. We now simply carry out the induction on j to verify that $H_j = z_1^{(j+1)(M-1)} z_2^{N-j}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{j+2}}$ for $1 \leq j \leq N$.

Since the Jacobian determinant

$$g = \frac{\partial(F_1, F_2)}{\partial(z_1, z_2)} = N z_1^{M-1} z_2^{N-1} + z_1^{K+M-1}$$

is multiplier whose order of subellipticity is $\geq \frac{1}{4}$, the Jacobian determinant

$$\frac{\partial(F_1, g)}{\partial(z_1, z_2)} = N(N-1) z_1^{2M-2} z_2^{N-1}$$

is a multiplier whose order of subellipticity is $\geq \frac{1}{8}$, which means that H_1 is a multiplier whose order of subellipticity is $\geq \frac{1}{8}$. Suppose H_j has been verified to be a multiplier whose order of subellipticity is $\geq \frac{1}{2^{j+2}}$ for some $1 \leq j < N$. Then the Jacobian determinant

$$\begin{aligned} \frac{\partial(F_1, H_j)}{\partial(z_1, z_2)} &= \det \begin{pmatrix} M z_1^{M-1} & 0 \\ \frac{\partial H_j}{\partial z_1} & (N-j) z_1^{(j+1)(M-1)} z_2^{N-j-1} \end{pmatrix} \\ &= M(N-j) z_1^{(j+2)(M-1)} z_2^{N-j-1} \end{aligned}$$

is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{j+3}}$, which means that H_{j+1} is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{j+3}}$. This finishes the induction argument and we know that $H_N = z_1^{(N+1)(M-1)}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+2}}$.

By taking the effective $(N+1)(M-1)$ -th root, we conclude that the holomorphic function germ z_1 is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+2}(N+1)(M-1)}$ and the holomorphic function germ $z_2 z_1^K$ which contains z_1 as a factor is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+2}(N+1)(M-1)}$. Then

$$z_2^N = (z_2^N + z_2 z_1^K) - z_2 z_1^K = F_2 - z_2 z_1^K$$

is a pre-multiplier whose differential has order of subellipticity $\geq \frac{1}{2^{N+3}(N+1)(M-1)}$. Since both z_1 and z_2^N are pre-multipliers whose differentials have order of subellipticity $\geq \frac{1}{2^{N+3}(N+1)(M-1)}$, the Jacobian determinant

$$\frac{\partial(z_1, z_2^N)}{\partial(z_1, z_2)} = \det \begin{pmatrix} 1 & 0 \\ 0 & N z_2^{N-1} \end{pmatrix} = N z_2^{N-1}$$

is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+3}(N+1)(M-1)}$, which means that z_2^{N-1} is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+3}(N+1)(M-1)}$. By taking the $(N-1)$ -th root of z_2^{N-1} , we conclude that z_2 is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+3}(N+1)(N-1)(M-1)}$. Finally 1 which is the Jacobian determinant $\frac{\partial(z_1, z_2)}{\partial(z_1, z_2)}$ is a multiplier whose order of subellipticity is $\geq \frac{1}{2^{N+4}(N+1)(N-1)(M-1)}$. This algorithm is effective, because $\frac{1}{2^{N+4}(N+1)(N-1)(M-1)}$ is bounded from below by the explicit function $\frac{1}{2^{q+4}(q+1)(q-1)^2}$ of $q = NM$.

(4.4) *Remark.* In carrying out above the effective Kohn algorithm for Catlin-D'Angelo's example, to get to the multiplier 1 from F_1, F_2 , we have to perform differentiation $N+2$ times in the construction of Jacobian determinants. Compared to the full-real-radical Kohn algorithm which requires only 4 differentiation to terminate, to avoid ineffectiveness in the taking of roots in the effective Kohn algorithm we choose the option of performing more, but still an effective number of, differentiations.

(4.5) *Geometric Reason for Ineffectiveness of Full-Real-Radical Kohn Algorithm for Catlin-D'Angelo's Example.* The above discussion shows by computation why in Catlin-D'Angelo's example the full-real-radical Kohn algorithm is ineffective while the effective Kohn algorithm gives effectiveness. Now we would like to analyze geometrically why such a phenomenon occurs. When $F_2 = z_2^N + z_2 z_1^K$ is regarded as a polynomial in z_2 , its degree N is effective (in the sense of being bounded by an explicit function of $q = MN$) but its discriminant obtained by eliminating z_2 from F_2 and $\frac{\partial F_2}{\partial z_2}$, as a function germ in z_1 vanishes to an order at $z_1 = 0$ which is a function of K and is not effective. In other words, the N roots (in z_2) of $F_2 = 0$ as N functions of z_1 are close together near $z_1 = 0$ to an order which is a function of K and is not effective. The discriminant of F_2 and the closeness of the N roots of $F_2 = 0$ enter the picture, because $F_1 = z_1^M$ depends only on z_1 and the Jacobian determinant of F_1, F_2 is the first multiplier in the algorithm. Because of the ineffectiveness of the vanishing order of the discriminant of F_2 at a function in z_1 at $z_1 = 0$, the step of root-taking is ineffective. On the other hand, the effective Kohn algorithm replaces ineffective root-taking of the discriminant of a Weierstrass polynomial by differentiating the Weierstrass polynomial with respect to its variable as many times as its degree to avoid the ineffective root-taking.

§5. MULTIPLIERS IN MORE GENERAL SETTING

We now discuss the generalization of Kohn's technique of multipliers to more general systems of partial differential equations.

(5.1) *Generalization of Kohn's Technique of Multiplier Ideal Sheaves to More General Setting.* The generalization of Kohn's technique of multiplier ideal sheaves to a more general setting comes from looking at Kohn's technique for the complex Neumann problem from the following perspective. The subelliptic estimate for a bounded smooth weakly pseudoconvex domain Ω in $\mathbb{C}^n$ at its boundary point P seeks to estimate

$$\|\varphi\|_\varepsilon^2 = \|\Lambda^\varepsilon \varphi\|^2$$

by a constant C_ε times

$$Q(\varphi, \varphi) = \|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 + \|\varphi\|^2,$$

for some $\varepsilon > 0$, for all smooth test $(0,1)$ -form φ on $U \cap \bar{\Omega}$ with compact support which belongs to the domain of the actual adjoint $\bar{\partial}^*$ of $\bar{\partial}$ (where U is an open neighborhood of P in $\mathbb{C}^n$). For the convenience of discussion, we simply say that $\Lambda^\varepsilon \varphi$ is *estimable* on U when

$$\|\Lambda^\varepsilon \varphi\|^2 \leq C_\varepsilon Q(\varphi, \varphi).$$

In general, we say that some expression ψ defined from φ is *estimable* on U (or simply *estimable*) if

$$\|\psi\|^2 \leq C Q(\varphi, \varphi)$$

for some constant C independent of φ (which is smooth on $U \cap \partial\Omega$ with compact support). The starting point is the basic identity

$$\|\bar{\partial}\varphi\|^2 + \|\bar{\partial}^*\varphi\|^2 = \|\bar{\nabla}\varphi\|^2 + \int_{\partial\Omega} \langle \text{Levi}_{\partial\Omega}, \bar{\varphi} \wedge \varepsilon \rangle,$$

where $\bar{\nabla}$ is the (covariant) differentiation of φ in the $(0,1)$ -direction and $\text{Levi}_{\partial\Omega}$ is the Levi form $\partial\bar{\partial}r$ of the boundary $\partial\Omega$ of Ω when Ω is locally defined by $r < 0$ with $dr \equiv 1$ on $\partial\Omega$. In particular,

$$\|\bar{\nabla}\varphi\|^2 \leq Q(\varphi, \varphi) \quad \text{and} \quad \|\bar{\partial}^*\varphi\|^2 \leq Q(\varphi, \varphi).$$

Together with $\|\varphi\|^2 \leq Q(x, y)$, this means that both $\bar{\nabla}\varphi$ and $\bar{\partial}^*\varphi$, as well as φ , are estimable. The expressions

$$\bar{\nabla}\varphi = (\bar{\partial}_j \varphi_{\bar{k}})_{1 \leq j, k \leq n} \quad \text{and} \quad \bar{\partial}^*\varphi = - \sum_{j=1}^n \partial_j \varphi_{\bar{j}}$$

are linear combinations of first-order partial derivatives of the components of φ . A multiplier F means the estimability of $\Lambda^\varepsilon(F\varphi)$ and a vector multiplier θ means the estimability of $\Lambda^\varepsilon(\theta \cdot \varphi)$. Kohn's technique is to use the estimability of $\bar{\nabla}\varphi$, $\bar{\partial}^*\varphi$ and φ and apply algebraic manipulations and integration by parts to construct from the estimability of $\Lambda^\varepsilon(F\varphi)$ and $\Lambda^\varepsilon(\theta \cdot \varphi)$ other F' and θ' with estimable $\Lambda^\varepsilon(F'\varphi)$ and $\Lambda^\varepsilon(\theta' \cdot \varphi)$. For such manipulations it does not matter what the meaning of $Q(\varphi, \varphi)$ is. Moreover, the operations of integration by parts are along the tangent directions of the boundary, because Λ^ε is the pseudo-differential operator corresponding to the $(\frac{\varepsilon}{2})$ -th power of 1 plus the Laplacian in tangential coordinates of the boundary.

(5.1.1) For our generalization of the technique of multiplier ideal sheaves, we use the following simple setting which highlights the core argument of the technique. Fix an integer $q \geq 2$. Let Ω be an open neighborhood of 0 in $\mathbb{R}^m$ and $Y_{j\nu}$ be complex-valued smooth differential operators on Ω for $1 \leq j \leq N$ and $1 \leq \nu \leq q$. For any q -tuple $\varphi = (\varphi_1, \dots, \varphi_q)$ of smooth complex-valued functions with compact support on Ω , let

$$Q(\varphi, \varphi) = \|\varphi\|^2 + \sum_{1 \leq j \leq N, 1 \leq \nu \leq q} \|Y_{j\nu}\varphi_\nu\|^2,$$

where $\|\cdot\|$ means the L^2 norm on Ω . An expression ψ of φ of the form $\sum_{\nu=1}^q Z_\nu \varphi_\nu$ (where each Z_ν is a pseudo-differential operator on Ω) is said to be *estimable* on an open neighborhood U of 0 in Ω (or simply *estimable*) if there is a positive constant C such that

$$\|\psi\|^2 \leq CQ(\varphi, \varphi)$$

for all q -tuple $\varphi = (\varphi_1, \dots, \varphi_q)$ of smooth complex-valued test functions with compact support on U . When ψ is vector-valued instead of scalar-valued, the estimability of ψ on U means the estimability of each of its components on U . When we have two such expressions ψ and $\tilde{\psi}$, we say that the inner product $(\psi, \tilde{\psi})$ is *estimable* on U if

$$(\psi, \tilde{\psi}) \leq CQ(\varphi, \varphi)$$

all q -tuple $\varphi = (\varphi_1, \dots, \varphi_q)$ of smooth complex-valued test functions with compact support on U . We refer to C as the constant of estimability of ψ or $(\psi, \tilde{\psi})$.

(5.1.2) Let Λ^ε be the pseudo-differential operator which is the $(\frac{\varepsilon}{2})$ -th power of 1 plus the Laplacian in the coordinates of $\mathbb{R}^m$. We introduce three kinds of multipliers: (i) scalar multiplier, (ii) vector multiplier, and (iii) matrix multiplier.

The germ at 0 of a smooth function α is a *scalar multiplier* at 0 with *order of subellipticity* $\geq \varepsilon$ if $\Lambda^\varepsilon(\alpha\varphi_\nu)$ (for $1 \leq \nu \leq q$) is estimable on U for some $\varepsilon > 0$ and some open neighborhood U of 0 in Ω on which α is defined. Clearly the product of a scalar multiplier with any smooth function is again a scalar multiplier with no change in the order of subellipticity. By considering the commutator $[\Lambda^\varepsilon, \alpha]$ of pseudo-differential operators, we conclude that for $0 < \varepsilon \leq 1$ the estimability of $\Lambda^\varepsilon(\alpha\varphi_\nu)$ on U is equivalent to the estimability of $\alpha(\Lambda^\varepsilon\varphi_\nu)$ on U , because $\|\varphi\|^2$ is estimable on U .

The germ at 0 of a smooth q -tuple of smooth complex-valued functions $\vec{a} = (a_1, \dots, a_q)$ is called a *vector multiplier* at 0 with *order of subellipticity* $\geq \varepsilon$ if $\Lambda^\varepsilon(\sum_{\nu=1}^q a_\nu\varphi_\nu)$ is estimable on U for some $\varepsilon > 0$ and some open neighborhood U of 0 in Ω on which $\vec{a}$ is defined. Clearly the product of a vector multiplier with any smooth function is again a vector multiplier with no change in the order of subellipticity. Again, for $0 < \varepsilon \leq 1$ the estimability of $\Lambda^\varepsilon(\sum_{\nu=1}^q a_\nu\varphi_\nu)$ on U is equivalent to the estimability of $\sum_{\nu=1}^q a_\nu(\Lambda^\varepsilon\varphi_\nu)$ on U .

An $q \times q$ matrix $\mathbf{a} = (a_{jk})_{1 \leq j, \ell \leq q}$ is called a *matrix multiplier* at 0 with *order of subellipticity* $\geq \varepsilon$ if every one of its rows $\vec{a}_j = (a_{j1}, \dots, a_{jq})$ (for $1 \leq r \leq q$) is a vector multiplier at 0 with order of subellipticity $\geq \varepsilon$. Clearly a matrix multiplier multiplied on the left by a $q \times q$ matrix with smooth functions as entries yields a matrix multiplier with no change in the order of subellipticity.

Some simple relations among scalar multipliers, vector multipliers, and matrix multipliers are as follows. The product of a scalar multiplier with any row q -vector with smooth functions as components is a vector multiplier. The product of a scalar multiplier with any $q \times q$ matrix with smooth functions as entries is a matrix multiplier. Any vector multiplier (as a row vector) multiplied on the left by a column q -vector with smooth functions as components yields a matrix multiplier. By Cramer's rule the determinant of a matrix multiplier is a scalar multiplier. Any matrix multiplier multiplied on the left by a row q -vector with smooth functions as components yields a vector multiplier.

(5.1.3) Just like the real radical property (C) of Kohn's multipliers in (2.1), scalar multipliers here enjoy the same real radical property that if α is a scalar multiplier at 0 with order of subellipticity $\geq \varepsilon$ (for some $0 < \varepsilon \leq 1$) and β is a smooth complex-valued function germ at 0 such that $|\beta|^\sigma \leq |\alpha|$ for some $\sigma \in \mathbb{N}$, then β is a scalar multiplier with order of subellipticity $\geq \frac{\varepsilon}{\sigma}$. The proof is completely analogous to the proof of [Kohn1979, p.98, Lemma 4.3.4] and is as follows. Let U be an open neighborhood of 0 in Ω such that both α and β are represented by smooth functions on U and φ is a test q -tuple of smooth functions on U with compact support. Let $\eta = \frac{\varepsilon}{\sigma}$. Since

$$\begin{aligned} \|\Lambda^{\sigma\eta}(\beta^\sigma \varphi)\|_{L^2(U)} &\leq C_1^* \|\beta^\sigma \Lambda^{\sigma\eta} \varphi\|_{L^2(U)} + C_1^{**} \|\varphi\|_{L^2(U)} \\ &\leq C_2^* \|\alpha \Lambda^{\sigma\eta} \varphi\|_{L^2(U)} + C_2^{**} \|\varphi\|_{L^2(U)} \\ &\leq C_3^* \|\Lambda^{\sigma\eta}(\alpha) \varphi\|_{L^2(U)} + C_3^{**} \|\varphi\|_{L^2(U)}, \end{aligned}$$

it suffices to prove the statement that

$$(5.1.3.1)_\tau \quad \|\Lambda^{\tau\eta}(\beta^\tau \varphi)\|_{L^2(U)} \leq C_\tau \|\Lambda^{\sigma\eta}(\beta^{\sigma\eta} \varphi)\|_{L^2(U)} + C'_\tau \|\varphi\|_{L^2(U)}$$

for $1 \leq \tau \leq \sigma$. The statement $(5.1.3.1)_\tau$ follows from descending induction on τ for $1 \leq \tau \leq \sigma$, because

$$\begin{aligned} \|\Lambda^{\tau\eta}(\beta^\tau \varphi)\|_{L^2(U)}^2 &= (\Lambda^{(\tau+1)\eta}(\beta^{\tau+1} \varphi), \Lambda^{(\tau-1)\eta}(\beta^{\tau-1} \varphi))_{L^2(U)} + \hat{C} \|\varphi\|^2 \\ &\leq \|\Lambda^{(\tau+1)\eta}(\beta^{\tau+1} \varphi)\|_{L^2(U)} \|\Lambda^{(\tau-1)\eta}(\beta^{\tau-1} \varphi)\|_{L^2(U)} + \tilde{C} \|\varphi\|^2 \end{aligned}$$

for $1 \leq \tau < \sigma$, where $C_j^*, C_j^{**}, C_\tau, C'_\tau, \hat{C}, \tilde{C}$ are constants independent of φ .

In particular, if α is a scalar multiplier at 0 with order of subellipticity $\geq \varepsilon$ (for some $0 < \varepsilon \leq 1$), then its complex-conjugate $\bar{\alpha}$ is also scalar multiplier at 0 with order of subellipticity $\geq \varepsilon$.

A very important part of the technique of multiply ideal sheaves is the differential relations among the scalar multipliers, vector multipliers, and matrix multipliers. This is represented by two procedures involving differentiation. The first procedure produces a new vector multiplier from a matrix multiplier. The second procedure produces a new vector multiplier from a scalar multiplier. The second procedure is similar to the procedure (B)(i) for Kohn's multipliers for the complex Neumann problem in (2.1). The first procedure is a new one, even in the special case of Kohn's multipliers for the complex Neumann problem. The following theorem presents a unified version of both procedures, which yields both procedures as special cases.

(5.2) *Theorem (Generation of Vector Multiplier from Matrix Multiplier or Scalar Multiplier).* Let $X_1, \dots, X_q$ be complex-valued smooth first-order differential operators on Ω whose adjoint operators are $X_1^*, \dots, X_q^*$ with respect to the L^2 inner product on Ω such that each $X_j^* \varphi$ is estimable on Ω for $1 \leq j \leq q$. Let $\Gamma_{k\ell}$ be a smooth complex-valued function on Ω for $1 \leq k, \ell \leq q$ such that $\sum_{1 \leq k, \ell \leq q} \Gamma_{k\ell} X_k \varphi_\ell$ is estimable on Ω . Let $\varepsilon_1, \varepsilon_2$ be positive numbers ≤ 1 . Let $\mathbf{a} = (a_{jk})_{1 \leq j, k \leq q}$ be a matrix of multipliers at 0 so that each of its rows $\vec{a}_j = (a_{jk})_{1 \leq k \leq q}$ is a vector multiplier at 0 with order of subellipticity $\geq \varepsilon_1$ for $1 \leq j \leq q$. Let α be a scalar multiplier at 0 with order of subellipticity $\geq \varepsilon_2$. Let $(A_{jk})_{1 \leq j, k \leq q}$ be a matrix of smooth complex-valued function germs at 0 such that $\sum_{\ell=1}^q A_{j\ell} a_{\ell k}$ equals to the Kronecker delta δ_{jk} times α for $1 \leq j, k \leq q$. Let

$$b_j = \sum_{1 \leq p, \ell, k \leq q} \Gamma_{pk} A_{k\ell} (X_p a_{\ell j})$$

and $\vec{b} = (b_j)_{1 \leq j \leq q}$. Then $\vec{b}$ is a vector multiplier at 0 whose order of subellipticity is $\geq \frac{1}{2} \min(\varepsilon_1, \varepsilon_2)$. In particular, the following two special cases hold.

(i) For $1 \leq j \leq q$ let

$$c_j = \sum_{1 \leq p, \ell \leq q} \text{adj}(\mathbf{a})_{p\ell} (X_p a_{\ell j})$$

and $\vec{c} = (c_j)_{1 \leq j \leq q}$, where $\text{adj}(\mathbf{a})$ is the adjoint matrix of $\mathbf{a}$ (so that the matrix product of $\text{adj}(\mathbf{a})$ and $\mathbf{a}$ is equal to $\det(\mathbf{a})$ times the identity matrix of order q). Then $\vec{c}$ is a vector multiplier at 0 whose order of subellipticity is $\geq \frac{\varepsilon_1}{2}$.

(ii) Let $d_j = \sum_{k=1}^q \Gamma_{kj} X_k \alpha$ for $1 \leq j \leq q$ and $\vec{d} = (d_j)_{1 \leq j \leq q}$. Then $\vec{d}$ is a vector multiplier at 0 whose order of subellipticity is $\geq \frac{\varepsilon_2}{2}$.

(5.2.1) *Proof of Theorem (5.2).* Let $\varepsilon = \min(\varepsilon_1, \varepsilon_2)$. Let U be an open neighborhood of 0 in Ω such that α and the vector multiplier $\vec{a}_j = (a_{jk})_{1 \leq k \leq q}$ (for $1 \leq j \leq q$) are defined and smooth on U and $\Lambda^\varepsilon(\sum_{\nu=1}^q a_{j\nu} \varphi_\nu)$ is estimable on U for smooth test functions $\varphi = (\varphi_1, \dots, \varphi_q)$ on U with compact support. Let ψ be a linear combination of φ_j (for $1 \leq j \leq q$) with smooth functions on U as coefficients and which we will specify more precisely later. By the Cauchy-Schwarz inequality, for $1 \leq p, \ell \leq q$ the inner product

$$\left(\Lambda^\varepsilon \left(\sum_{j=1}^q a_{\ell j} \varphi_j \right), X_p^* \psi \right)$$

is estimable on U , because $\vec{a}_\ell$ is a vector-multiplier for $1 \leq \ell \leq q$ and $X_p^* \psi$ is estimable (from the estimability of $X_p^* \varphi$) for $1 \leq p \leq q$. Note that the constant of estimability depends on the smooth coefficient functions in the linear combination ψ of $\varphi_1, \dots, \varphi_q$ which are yet to be specified. Integration by parts applied to X_p (by switching X_p over to X_p^* in the inner product) yields the estimability of

$$(5.2.1.1) \quad \left(\Lambda^\varepsilon \left(\sum_{j=1}^q (X_p a_{\ell j}) \varphi_j \right), \psi \right) + \left(\Lambda^\varepsilon \left(\sum_{j=1}^q a_{\ell j} (X_p \varphi_j) \right), \psi \right)$$

on U for $1 \leq p, \ell \leq q$ after we take care of the error terms from the commutator of pseudodifferential operators in the standard way.

Now we apply $\sum_{\ell=1}^q A_{k\ell}$ to (5.2.1.1) to get the estimability of

$$\left(\Lambda^\varepsilon \left(\sum_{\ell,j=1}^q A_{k\ell} (X_p a_{\ell j}) \varphi_j \right), \psi \right) + \left(\Lambda^\varepsilon \left(\sum_{\ell,j=1}^q A_{k\ell} a_{\ell j} (X_p \varphi_j) \right), \psi \right)$$

on U for $1 \leq k \leq q$, which is the same as

$$(5.2.1.2) \quad \left(\Lambda^\varepsilon \left(\sum_{\ell,j=1}^q A_{k\ell} (X_p a_{\ell j}) \varphi_j \right), \psi \right) + \left(\Lambda^\varepsilon (\alpha(X_p \varphi_k)), \psi \right),$$

because $\sum_{j=1}^q A_{ij} a_{j\ell} = \alpha \delta_{i\ell}$ for $1 \leq j, \ell \leq q$.

We apply $\sum_{1 \leq p, k \leq q} \Gamma_{pk}$ to (5.2.1.2) to get the estimability of

$$\left(\Lambda^\varepsilon \left(\sum_{1 \leq p, \ell, j, k \leq q} \Gamma_{pk} A_{k\ell} (X_p a_{\ell j}) \varphi_j \right), \psi \right) + \left(\Lambda^\varepsilon \left(\alpha \sum_{1 \leq p, k \leq q} \Gamma_{pk} (X_p \varphi_k) \right), \psi \right)$$

on U , which is the same as

$$\left(\Lambda^\varepsilon \left(\sum_{1 \leq p, \ell, j, k \leq q} \Gamma_{pq} A_{q\ell} (X_p a_{\ell j}) \varphi_j \right), \psi \right) + \left(\sum_{1 \leq p, k \leq q} \Gamma_{pq} (X_p \varphi_q), \Lambda^\varepsilon (\bar{\alpha} \psi) \right)$$

up to estimable error terms from the commutators of pseudodifferential operators.

Since $\bar{\alpha}$ is a scalar-multiplier at 0 (on account of α being a scalar multiplier at 0), from the estimability of $\sum_{1 \leq p, k \leq q} \Gamma_{pk}(X_p \varphi_k)$ on U and the Cauchy-Schwarz inequality we conclude that

$$\left(\sum_{1 \leq p \leq r, 1 \leq k \leq q} \Gamma_{pk}(X_p \varphi_k), \Lambda^\varepsilon(\bar{\alpha} \psi) \right)$$

is estimable on U . Hence

$$\left(\Lambda^\varepsilon \left(\sum_{1 \leq p \leq r, 1 \leq \ell, j, k \leq q} \Gamma_{pk} A_{k\ell}(X_p a_{\ell j}) \varphi_j \right), \psi \right)$$

is estimable on U .

We can now choose

$$\psi = \sum_{1 \leq p, \ell, j, k \leq q} \Gamma_{pk} A_{k\ell}(X_p a_{\ell j}) \varphi_j$$

so that

$$\left(\Lambda^\varepsilon \left(\sum_{1 \leq p \leq r, 1 \leq \ell, j, k \leq q} \Gamma_{pk} A_{k\ell}(X_p a_{\ell j}) \varphi_j \right), \psi \right)$$

is equal to

$$\left\| \Lambda^{\frac{\varepsilon}{2}} \sum_{j=1}^q b_j \varphi_j \right\|^2$$

This means that $\vec{b}$ is a vector multiplier at 0 whose order of subellipticity is $\geq \frac{\varepsilon}{2}$.

We now look at the two special cases. The special case (i) follows from setting $(A_{jk})_{1 \leq j, k \leq q}$ to be the adjoint matrix $\text{adj}(\mathbf{a})$ of the matrix $\mathbf{a}$ and setting α to be $\det(\mathbf{a})$ with $\varepsilon_2 = \varepsilon_1$. The special case (ii) follows from setting A_{jk} to be the Kronecker delta δ_{jk} for $1 \leq j, k \leq q$ and setting a_{jk} to be $\alpha \delta_{jk}$ for $1 \leq j, k \leq q$ with $\varepsilon_1 = \varepsilon_2$ Q.E.D.

(5.3) *Remark.* Though Theorem (5.2) is presented as involving interior estimates, the same argument works also in boundary situations like the complex Neumann problem where, for the argument, integration by parts is needed only for the boundary tangential directions which do not affect the condition of the test forms to be in the domain of the actual adjoint $\bar{\partial}^*$ of $\bar{\partial}$. The special case (ii) of Theorem (5.2), after modification for application to the situation of the complex Neumann

problem for special domains, gives a procedure to generate a vector multiplier from a matrix multiplier. In (6.5) below, computations of examples are given to show that this procedure is a new procedure of generating vector multipliers for special domains in $\mathbb{C}^n$ with $n \geq 4$.

(5.4) *Estimable Linear Combinations and Initial Multipliers.* The goal of the technique of multiplier ideal sheaves is to use the differential relations among the multipliers and some initial multipliers to conclude, under some geometric conditions, that the function which is identically 1 is a scalar multiplier. An increase in the differential relations among the multipliers facilitates the achievement of the goal. Theorem (5.2) uses the collection $(\Gamma_{k\ell})_{1 \leq k, \ell \leq q}$ of smooth functions on Ω to construct a new vector multiplier from a matrix multiplier. The condition on the collection $(\Gamma_{k\ell})_{1 \leq k, \ell \leq q}$ of smooth functions on Ω is that $\sum_{1 \leq k, \ell \leq q} \Gamma_{k\ell} X_k \varphi_\ell$ is estimable on Ω . For that reason we refer to the collection $(\Gamma_{k\ell})_{1 \leq k, \ell \leq q}$ of smooth functions on Ω an *estimable linear combination*. To facilitate the construction of new vector multipliers, we can use a family of such estimable linear combinations $(\Gamma_{k\ell}^{(\lambda)})_{1 \leq k, \ell \leq q}$ indexed by $1 \leq \lambda \leq \lambda^*$ instead of a single one. There remains the crucial question of geometric conditions to guarantee solution of the regularity problem. This condition (which is similar to the condition of finite type for the complex Neumann problem) should be a condition on the family of estimable linear combinations $(\Gamma_{k\ell}^{(\lambda)})_{1 \leq k, \ell \leq q}$ for $1 \leq \lambda \leq \lambda^*$ and the choice of initial scalar multipliers $\alpha^{(\sigma)}$ (for $1 \leq \sigma \leq \sigma^*$) and initial vector multipliers $\vec{a}^{(\tau)}$ (for $1 \leq \tau \leq \tau^*$). This question has not yet been satisfactorily answered.

§6. NEW PROCEDURE TO GENERATE VECTOR MULTIPLIER FROM MATRIX MULTIPLIER IN COMPLEX NEUMANN PROBLEM OF SPECIAL DOMAIN

We now modify the argument in the special case (ii) of Theorem (5.2) to apply to the complex Neumann problem to obtain a new procedure of generating a vector multiplier from a matrix multiplier. This new procedure works for any bounded weakly pseudoconvex domain with smooth boundary, but we will carry out the modification only for a special domain, because the notations for a special domain have already been introduced here to make the argument for a special domain easier to present. Then we show by explicit computation for some special domains in $\mathbb{C}^4$ that this new way of generating a vector multiplier cannot be derived from Kohn's procedures in (2.1).

(6.1) *New Procedure to Generate Vector Multiplier from Matrix Multiplier for the Complex Neumann Problem.* Let Ω be a special domain in $\mathbb{C}^{n+1}$ (with coordinates $w, z_1, \dots, z_n$) defined by holomorphic functions $F_j(z_1, \dots, z_n)$ on some open

neighborhood of $\bar{\Omega}$, as described in (2.8.1). For the complex Neumann problem for the special domain Ω in $\mathbb{C}^{n+1}$, the roles of the vector fields $X_1, \dots, X_q$ are played by $\partial_j = \frac{\partial}{\partial z_j}$ for $1 \leq j \leq n$ and the roles of $X_1^*, \dots, X_q^*$ are played by $\bar{\partial}_{\bar{j}} = \frac{\partial}{\partial \bar{z}_j}$ for $1 \leq j \leq n$ and the role of Γ_{jk} for $1 \leq j, k \leq q$ is played by the Kronecker delta δ_{jk} for $1 \leq j, k \leq n$. Let $\mathbf{a} = (a_{jk})_{1 \leq j, k \leq n}$ be a matrix whose entry a_{jk} is a holomorphic function of $z_1, \dots, z_n$ defined on an open neighborhood of 0 in $\mathbb{C}^n$ for $1 \leq j, k \leq n$ such that each row vector $\vec{a}_j = (a_{jk})_{1 \leq k \leq n}$ is a vector multiplier at 0 with order of subellipticity $\geq \eta$ (for some $0 < \eta \leq 1$). Let U be an open neighborhood of 0 in $\mathbb{C}^{n+1}$ such that each a_{jk} , as a holomorphic function in $z_1, \dots, z_n, w$ but independent of w , is defined on U .

Let $\varphi = \sum_{j=1}^n \varphi_{\bar{j}} d\bar{z}_j + \hat{\varphi} d\bar{w}$ be a smooth test $(1, 0)$ -form on $\bar{\Omega} \cap U$ with compact support which is in the domain of the actual adjoint $\bar{\partial}^*$ of $\bar{\partial}$. Let ψ be a scalar function which is a linear combination of $\varphi_{\bar{j}}$ with smooth functions as coefficients and which we will specify more precisely later. Let $0 < \varepsilon < \frac{\eta}{2}$. The L^2 inner product

$$\left(\Lambda^{2\varepsilon} \left(\sum_{j=1}^n a_{\ell j} \varphi_{\bar{j}} \right), \bar{\partial}_p \psi \right)$$

is estimable on U by the Cauchy-Schwarz inequality from the assumption that $\vec{a}_\ell = \sum_{j=1}^n a_{\ell j} dz_j$ is a vector-multiplier at 0 with order of subellipticity $\geq \eta \geq 2\varepsilon$ for $1 \leq \ell \leq n$ and the assumption that the L^2 norm of $\bar{\partial}_p \psi$ is estimable on U for $1 \leq p \leq n$ (from the estimability of $\bar{\partial}_p \varphi_{\bar{j}}$ on U for any $1 \leq p, j \leq n$). Integration by parts applied to $\bar{\partial}_p$ yields the estimability of

$$(6.1.1) \quad \left(\Lambda^{2\varepsilon} \left(\sum_{j=1}^n (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right) + \left(\Lambda^{2\varepsilon} \left(\sum_{j=1}^n a_{\ell j} (\partial_p \varphi_{\bar{j}}) \right), \psi \right)$$

on U after we take care of the error terms from the commutators of operators in the standard way.

Let $(A_{q\ell})_{1 \leq q, \ell \leq n}$ be the adjoint matrix of $\mathbf{a}$ so that $\sum_{j=1}^n A_{ij} a_{jk} = (\det \mathbf{a}) \delta_{ik}$ for $1 \leq i, k \leq n$ (where δ_{jk} is the Kronecker delta). Now for $1 \leq p \leq n$ we apply $\sum_{\ell=1}^n A_{p\ell}$ to (6.1.1) to get the estimability of

$$\left(\Lambda^{2\varepsilon} \left(\sum_{\ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right) + \left(\Lambda^{2\varepsilon} \left(\sum_{\ell, j=1}^n A_{p\ell} a_{\ell j} (\partial_p \varphi_{\bar{j}}) \right), \psi \right)$$

on U , which is the same as

$$(6.1.2) \quad \left(\Lambda^{2\varepsilon} \left(\sum_{\ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right) + \left(\Lambda^{2\varepsilon} ((\det \mathbf{a})(\partial_p \varphi_{\bar{p}})), \psi \right),$$

because $\sum_{j=1}^n A_{ij} a_{jk} = (\det \mathbf{a}) \delta_{ik}$ for $1 \leq i, k \leq n$.

We now sum (6.1.2) over $1 \leq p \leq n$ to get the estimability of

$$\left(\Lambda^{2\varepsilon} \left(\sum_{p, \ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right) + \left(\Lambda^{2\varepsilon} \left((\det \mathbf{a}) \left(\sum_{p=1}^n \partial_p \varphi_{\bar{p}} \right) \right), \psi \right)$$

on U , which is the same as

$$\left(\Lambda^{2\varepsilon} \left(\sum_{p, \ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right) + \left(\sum_{p=1}^n \partial_p \varphi_{\bar{p}}, \Lambda^{2\varepsilon} (\overline{(\det \mathbf{a})} \psi) \right).$$

As a determinant whose rows are vector-multipliers with order of subellipticity $\geq \eta \geq 2\varepsilon$, the determinant $\det(\mathbf{a})$ (as well as its complex-conjugate) is a scalar-multiplier with order of subellipticity $\geq \eta \geq 2\varepsilon$. The function $\Lambda^{2\varepsilon} (\overline{(\det \mathbf{a})} \psi)$ is estimable on U , because ψ is a linear combination of $\varphi_{\bar{j}}$ for $1 \leq j \leq n$ with smooth functions as coefficients. From the estimability of $\sum_{p=1}^n \partial_p \varphi_{\bar{p}}$ on U and the Cauchy-Schwarz inequality we conclude that

$$\left(\Lambda^{2\varepsilon} \left(\sum_{p, \ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right)$$

is estimable on U . We now choose

$$\psi = \sum_{p, \ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}}$$

so that

$$\left(\Lambda^{2\varepsilon} \left(\sum_{p, \ell, j=1}^n A_{p\ell} (\partial_p a_{\ell j}) \varphi_{\bar{j}} \right), \psi \right)_{L^2(\Omega)}$$

is equal to $\|\Lambda^\varepsilon \psi\|_{L^2(\Omega)}^2$. This means that the $(1, 0)$ -form

$$\sum_{j=1}^n \left(\sum_{1 \leq p, \ell \leq n} A_{p\ell} (\partial_p a_{\ell j}) \right) dz_j$$

is a vector-multiplier with order of subellipticity $\geq \frac{\eta}{2}$. This is a new process of producing vector-multipliers from a matrix of vector-multipliers. We now summarize the result in the following theorem.

(6.2) *Theorem.* Let Ω be a special domain in $\mathbb{C}^{n+1}$ (with coordinates $w, z_1, \dots, z_n$) defined by (2.8.1.1). Assume that 0 is a boundary point of Ω . Let $\mathbf{a} = (a_{jk})_{1 \leq j, k \leq n}$ be a matrix whose entry a_{jk} is a holomorphic function of $z_1, \dots, z_n$ defined on an open neighborhood of 0 in $\mathbb{C}^n$ for $1 \leq j, k \leq n$ such that each row vector $\vec{a}_j = (a_{jk})_{1 \leq k \leq n}$ is a vector multiplier at 0 with order of subellipticity $\geq \eta$ (for some $0 < \eta \leq 1$). Let $\text{adj}(\mathbf{a})$ be the adjoint matrix of $\mathbf{a}$. Then the holomorphic $(1, 0)$ -form

$$(6.2.1) \quad \sum_{j=1}^n \left(\sum_{1 \leq p, \ell \leq n} (\text{adj}(\mathbf{a}))_{p\ell} (\partial_p a_{\ell j}) \right) dz_j$$

is a vector multiplier at 0 with order of subellipticity $\geq \frac{\eta}{2}$, where $(\text{adj}(\mathbf{a}))_{p\ell}$ is the entry of $\text{adj}(\mathbf{a})$ in the p -th row and the ℓ -th column.

(6.3) *Comparison with Known Procedure of Generating Vector Multiplier from Matrix Multiplier.* In the case of a special domain, the known procedure (B)(i)(ii) in (2.1) to generate a vector multiplier from a given matrix multiplier $\mathbf{a}$ is to first use (B)(ii) in (2.1) to get the determinant $\det(\mathbf{a})$ of $\mathbf{a}$ as a scalar multiplier and then use (B)(i) in (2.1) to get the $(1, 0)$ -form $\partial(\det(\mathbf{a}))$ as a vector multiplier. Here we use the same notations as in (6.2).

We would like to compare $\partial(\det(\mathbf{a}))$ with the vector multiplier from Theorem (6.2). Since $\sum_{j=1}^n (\text{adj}(\mathbf{a}))_{kj} a_{jk} = \det(\mathbf{a})$ for any $1 \leq k \leq n$, we have

$$(6.3.1) \quad \partial(\det \mathbf{a}) = \sum_{j=1}^n (\partial(\text{adj}(\mathbf{a}))_{kj}) a_{jk} + \sum_{j=1}^n (\text{adj}(\mathbf{a}))_{kj} (\partial a_{jk})$$

which is different from the vector multiplier (6.2.1) where the index j of $a_{\ell j}$ is used as the index for the component of the vector multiplier instead of the subscript j of ∂_j which is the index for the component of the vector multiplier (6.3.1).

As shown in the computations given below in (6.4) and (6.5), it turns out that in the case of $n = 2$ the old procedure (B)(i)(ii) in (2.1) produces the same result as the new procedure given in Theorem (6.2), but in the case of $n \geq 3$ the new procedure indeed gives some new vector multipliers different from those produced by the procedures B(i)(ii).

(6.4) *New Procedure Gives No New Vector Multipliers for 3-Dimensional Special Domain.* We explicitly compute (6.2.1) and (6.3.1) in the case of a special domain Ω in $\mathbb{C}^3$ to determine whether the result (6.2.1) from the new procedure is different from the result (6.3.1) from the old procedure. We need only consider holomorphic functions and holomorphic 1-forms on $\mathbb{C}^2$ as scalar and vector multipliers. Let $a_{j1}dz_1 + a_{j2}dz_2$ as two holomorphic 1-forms which are vector multipliers. For the matrix multiplier $\mathbf{a} = (a_{jk})_{1 \leq j,k \leq 2}$, the adjoint matrix $\text{adj}(\mathbf{a})$ is

$$(A_{jk})_{1 \leq j,k \leq 2} = \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix}.$$

The vector multiplier generated by the new procedure is $b_1dz_1 + b_2dz_2$ with

$$\begin{aligned} b_j &= A_{11}\partial_1 a_{1j} + A_{12}\partial_1 a_{2j} + A_{21}\partial_2 a_{1j} + A_{22}\partial_2 a_{2j} \\ &= a_{22}\partial_1 a_{1j} - a_{12}\partial_1 a_{2j} - a_{21}\partial_2 a_{1j} + a_{11}\partial_2 a_{2j}. \end{aligned}$$

On the other hand,

$$\partial_j(\det \mathbf{a}) = a_{11}\partial_j a_{22} + a_{22}\partial_j a_{11} - a_{12}\partial_j a_{21} - a_{21}\partial_j a_{12}$$

so that the difference of their dot products with a test $(0,1)$ -form φ is

$$\begin{aligned} &(b_1dz_1 + b_2dz_2) \cdot \varphi - (\partial(\det \mathbf{a})) \cdot \varphi \\ &= \left(a_{22}(\partial_1 a_{11})\varphi_{\bar{1}} - a_{12}(\partial_1 a_{21})\varphi_{\bar{1}} - a_{21}(\partial_2 a_{11})\varphi_{\bar{1}} + a_{11}(\partial_2 a_{21})\varphi_{\bar{1}} \right. \\ &\quad \left. + a_{22}(\partial_1 a_{12})\varphi_{\bar{2}} - a_{12}(\partial_1 a_{22})\varphi_{\bar{2}} - a_{21}(\partial_2 a_{12})\varphi_{\bar{2}} + a_{11}(\partial_2 a_{22})\varphi_{\bar{2}} \right) \\ &\quad - \left(a_{11}(\partial_1 a_{22})\varphi_{\bar{1}} + a_{22}(\partial_1 a_{11})\varphi_{\bar{1}} - a_{12}(\partial_1 a_{21})\varphi_{\bar{1}} - a_{21}(\partial_1 a_{12})\varphi_{\bar{1}} \right. \\ &\quad \left. + a_{11}(\partial_2 a_{22})\varphi_{\bar{2}} + a_{22}(\partial_2 a_{11})\varphi_{\bar{2}} - a_{12}(\partial_2 a_{21})\varphi_{\bar{2}} - a_{21}(\partial_2 a_{12})\varphi_{\bar{2}} \right) \\ &= (\partial_2 a_{21} - \partial_1 a_{12})(a_{11}\varphi_{\bar{1}} + a_{12}\varphi_{\bar{2}}) + (\partial_1 a_{12} - \partial_2 a_{21})(a_{21}\varphi_{\bar{1}} + a_{22}\varphi_{\bar{2}}) \end{aligned}$$

with the cancellation of the four pairs of terms $((1,1),(3,2))$, $((1,2),(3,3))$, $((2,3),(4,4))$, $((2,4),(4,1))$ where (j,k) refers to the term on the j -th row in the k -th position in the array of terms in four rows of four terms each. The difference is a linear combination of $a_{j1}\varphi_{\bar{1}} + a_{j2}\varphi_{\bar{2}}$ for $j = 1, 2$ with smooth coefficients and is known to be estimable, because $a_{j1}dz_1 + a_{j2}dz_2$ is a vector multiplier for $j = 1, 2$. It means that in the case of a special domain in $\mathbb{C}^3$ the new procedure does not give any new multipliers.

(6.5) *New Procedure Gives More Vector Multipliers for 4-Dimensional Special Domain.* The new procedure of generating vector multipliers already gives vector

multiplies different from those generated by the procedure B(i)(ii) in (2.1) in the case of special domain in $\mathbb{C}^4$, as shown in the following computation. Consider the upper triangular matrix

$$\mathbf{a} = \begin{pmatrix} a_{11} & \xi & 0 \\ 0 & a_{22} & \eta \\ 0 & 0 & a_{33} \end{pmatrix}$$

whose three row vectors are vector multipliers which are holomorphic in the variables z_1, z_2, z_3 . Its adjoint matrix $\text{adj}(\mathbf{a})$, as the inverse of $\mathbf{a}$ times its determinant, is

$$(A_{jk})_{1 \leq j, k \leq 3} = \begin{pmatrix} a_{22}a_{33} & -a_{33}\xi & \xi\eta \\ 0 & a_{11}a_{33} & -a_{11}\eta \\ 0 & 0 & a_{11}a_{22} \end{pmatrix}.$$

We now compare the vector multiplier $\partial(\det \mathbf{a})$ with the vector multiplier $\sum_{j=1}^3 b_j dz_j$ generated by the new procedure by taking the difference of their dot products with a test function $\varphi = \sum_{j=1}^3 \varphi_{\bar{j}} d\bar{z}_j + \hat{\varphi} dw$ to get

$$\begin{aligned} & \left(\sum_{j=1}^3 b_j dz_j \right) \cdot \varphi - \partial(\det \mathbf{a}) \cdot \varphi \\ &= \sum_{1 \leq j < k \leq 3} A_{jk} (\partial_{\nu} a_{jk}) \varphi_{\bar{\nu}} \\ &= \sum_{\nu=1}^3 (-a_{33}\xi \partial_{\nu} \xi - a_{11}\eta \partial_{\nu} \eta) \varphi_{\bar{\nu}} \\ &= -\frac{1}{2} \sum_{\nu=1}^3 (a_{33} \partial_{\nu} (\xi^2) - a_{11} \partial_{\nu} (\eta^2)) \varphi_{\bar{\nu}}. \end{aligned}$$

The assumption that $\mathbf{a}$ is a matrix multiplier gives us only the estimability of

$$a_{11}\varphi_{\bar{1}} + \xi\varphi_{\bar{2}}, \quad a_{22}\varphi_{\bar{2}} + \eta\varphi_{\bar{3}}, \quad a_{33}\varphi_{\bar{3}},$$

from which we cannot conclude the estimability of

$$-\frac{1}{2} \sum_{\nu=1}^3 (a_{33} \partial_{\nu} (\xi^2) - a_{11} \partial_{\nu} (\eta^2)) \varphi_{\bar{\nu}}$$

for arbitrary functions ξ and η , because even in the special case of $\varphi_{\bar{1}} = \varphi_{\bar{2}} = 0$, the estimability of

$$-\frac{1}{2} \sum_{\nu=1}^3 (a_{33} \partial_{\nu} (\xi^2) - a_{11} \partial_{\nu} (\eta^2)) \varphi_{\bar{\nu}}$$

would mean the estimability of $a_{11}\partial_3(\eta^2)\varphi_{\bar{3}}$ for arbitrary a_{11} and η , which cannot be derived from the estimability of $a_{33}\varphi_{\bar{3}}$. This shows that the new procedure gives more vector multipliers.

REFERENCES

- [Catlin-D’Angelo2010] David Catlin and John P. D’Angelo, Subelliptic estimates. In: *Several Complex Variables and Connections with PDE Theory and Geometry (Proc. 2008 Fridtjof Conference for Linda Rothschild)*. ed. Ebenfelt *et al.* Trends in Mathematics. Birkhäuser/Springer Basel AG, Basel, 2010, pp.75 – 94.
- [D’Angelo1979] John P. D’Angelo, Finite type conditions for real hypersurfaces. *J. Differential Geom.* **14** (1979), 59 – 66.
- [D’Angelo1982] John P. D’Angelo, Real hypersurfaces, orders of contact, and applications. *Ann. of Math.* **115** (1982), 615 - 637.
- [Diederich-Fornaess1978] Klas Diederich and John-Erik Fornaess, Pseudoconvex domains with real-analytic boundary. *Ann. of Math.* **107** (1978), 371 – 384.
- [Donaldson1985] Simon Donaldson, Anti self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles. *Proc. London Math. Soc.* **50** (1985), 1 - 26.
- [Hörmander1967] Lars Hörmander, Hypoelliptic second order differential equations. *Acta Math.* **119** (1967), 147 - 171.
- [Kohn1972] Joseph J. Kohn, Boundary behavior of $\bar{\partial}$ on weakly pseudo-convex manifolds of dimension two. Collection of articles dedicated to S. S. Chern and D. C. Spencer on their sixtieth birthdays. *J. Differential Geometry* **6** (1972), 523 - 542.
- [Kohn1977] Joseph J. Kohn, Sufficient conditions for subellipticity on weakly pseudo-convex domains. *Proc. Nat. Acad. Sci. U.S.A.* **74** (1977), 2214 - 2216.
- [Kohn1979] Joseph J. Kohn, Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudo-convex domains: sufficient conditions. *Acta Math.* **142** (1979), 79 – 122.
- [Kohn-Nirenberg1965] Joseph J. Kohn and Louis Nirenberg, Non-coercive boundary value problems. *Comm. Pure Appl. Math.* **18** (1965), 443 - 492.
- [Nadel1990] Alan Michael Nadel, Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature. *Ann. of Math.* **132** (1990), 549 - 596.
- [Siu2010] Yum-Tong Siu, Effective Termination of Kohn’s Algorithm for Subelliptic Multipliers, *Pure and Applied Mathematics Quarterly* **6** (2010) (Special Issue in honor of Joseph J. Kohn), 1169 – 1241. (arXiv:0706.4113)

[Skoda1972] Henri Skoda, Application des techniques L^2 à la théorie des idéaux d'une algèbre de fonctions holomorphes avec poids. *Ann. Sci. École Norm. Sup.* **5** (1972), 545 – 579.

Author's address: Department of Mathematics, Harvard University, Cambridge, MA 02238, U.S.A.