New Small Mammals From the Hasnot-Tatrot Area of the Potwar Plateau, Northern Pakistan

Citation

Khan, M. Akbar, Lawrence J. Flynn. 2017. New Small Mammals From the Hasnot-Tatrot Area of the Potwar Plateau, Northern Pakistan. *Paläontologische Zeitschrift* 91, no. 4: 589-599.

Published Version

10.1007/s12542-017-0375-3

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:42656566>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

[Click here to view linked References](#)

NEW SMALL MAMMALS FROM THE HASNOT-TATROT AREA OF THE POTWAR PLATEAU, NORTHERN PAKISTAN

Muhammad Akbar Khan¹ and Lawrence J. Flynn²

1. Department of Zoology, University of the Punjab, Lahore, Punjab, Pakistan (38000)
2. Department of Human Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA

Key Words Rhizomyinae, Hystrix, Alilepus, Siwalik, Pakistan, late Miocene, Pliocene

Abstract

New rodent and lagomorph fossils from the Hasnot-Tatrot area of northern Pakistan are presented here to complement knowledge of stratigraphic ranges and morphology of key late Neogene Siwalik taxa. Most of the material is from two sites near the village of Bhandar in strata of the late Miocene age Dhok Pathan Formation; one specimen comes from the Pliocene Tatrot beds. We apply previously established magnetostratigraphy to date the fossils, the Bhandar sites dating to 6.6 to 6.7 Ma. In describing the fossils, we emphasize new morphological information represented by the material. As surface finds, these fossils represent relatively large body size species, three bamboo rat relatives, a porcupine, and a rabbit. The bamboo rats (Rhizomyinae) are an endemic group, and both the porcupine (*Hystrix*) and the rabbit (*Alilepus*) represent late Miocene immigrants into the Indian subcontinent.

Introduction

One of the classical areas in the Siwalik Group deposits to produce late Miocene and Pliocene vertebrate fossils over many years is the Hasnot-Tatrot area of northern Pakistan, particularly the terrestrial deposits near the village of Bhandar. Barnum Brown (Brown et al. 1924) and G.E. Lewis (1937) provided provenance for their collections from the area early last century. The Yale-Geological Survey of Pakistan (Y-GSP) project developed refined geographic and biostratigraphic controls for new fossil finds in the region, especially during the 1990s (Barry et al. 2002). In recent years, field teams from the University of the Punjab, Lahore, have continued work in the area, collecting important new fossils and documenting locality provenance.

Among the fossil materials recovered by University of the Punjab teams are mammalian groups not well represented among surface finds. Here we report on small mammal fossils from the area, which are significant from two aspects. They increase the number of biostratigraphic records with good provenance for taxa that were poorly documented previously, and they provide new morphological data for incompletely known species. The new fossils come, primarily, from two localities very near the village of Bhandar (Fig. 1). One other specimen comes from the well-known Pliocene Tatrot horizon, a fossil-rich surface developed on a sandstone dip slope that crops out prominently at Tatrot Village, 10 km. to the northeast.

The Bhandar fossil localities are at the same stratigraphic level in exposures immediately north of the village Bhandar. These sites H16 and H18 lie just below a distinctive sand layer in the upper part of the Dhok Pathan Formation that the Y-GSP project referred to as the Pillar Sandstone. The Pillar Sandstone can be traced laterally for several kilometers and is associated with fossils above and below it. The new sites are subjacent to the sandstone and occur in the vicinity of fossil

localities worked by Lewis (his site L97) and by Brown (his site B135). Y-GSP teams worked in this area and documented another locality (Y453) about 1 km to the north. Y453 lies in the Y-GSP magnetostratigraphic section at the top of Chron C3An.2n, at 6.5 Ma. The older sites H16 and H18 fall lower in the magnetozone, where they would date to between 6.6 and 6.7 Ma. Several of the fossils are mandibles representing bamboo rats (*Rhizomyinae*), and other specimens represent a porcupine and a rabbit, both late Miocene immigrants to northern Pakistan.

Methods. Eight new University of the Punjab fossils, all surface finds, are described and discussed here in the following systematic account. They are designated with the acronym PUPC, and are housed in the permanent collections of the Department of Zoology of the university. Specimens were measured in millimeters (mm) by a Nikon microscope fitted with a reticle, and large dimensions were taken by calipers. Specimens were photographed initially with a Canon EOS 350D 8.0 megapixel DSLR camera, lens 18-55mm; close-ups were also made with a Dino-Lite digital microscope.

Abbreviations. L, W, for length and width of teeth; A-P for the antero-posterior dimension of the incisor (right angle to its width). Standard tooth terminology is used: upper case letters plus a numeral for upper teeth (e.g., M3 for the upper third molar) and lower case plus numeral for lower teeth (e.g., p4 for lower fourth premolar). “Y” is as prefix for Y-GSP localities.

Systematic palaeontology

Order **Rodentia** Bowdich, 1821

Family **Spalacidae** Gray, 1821

Subfamily **Rhizomyinae** Winge, 1887

Genus ***Miorhizomys*** Flynn, 2009

Type species. *M. nagrii* (Hinton, 1933), from “Nagri zone” near Haritalyangar, northern India.

Diagnosis. Member of Tribe Rhizomyini with moderately deep dentary and a masseteric crest lacking a strong anterior extension, lower crest not inflated; lower incisor with flattened enamel and no median ridge; molars relatively small, of moderate crown height, with unilateral hypsodonty; M1 rounded anteriorly, without anterolingual flexus; m2 with three roots and strong mesolophid; longitudinal connection (mure) on m2 and m3 with small posterolingual enamel lake and often a deep buccal reentrant; incomplete closure of ventral slit of the infraorbital foramen; pronounced fossorial adaptations, low elevation of the skull.

Discussion. *Miorhizomys* is related to living bamboo rats, and therefore exhibits the fossorial characteristics of crown group Tribe Rhizomyini, namely large size, robust jaws and skulls, digging adaptations of the postcrania, chisel-like incisors, and deep dentaries. The lower incisor is particularly derived (Fig. 2). Unlike the round incisor with a single median ridge in the rhizomyines of Tribe Tachyoryctini, the chisel-like lower incisor of *Miorhizomys* and later Rhizomyini is deep anteroposteriorly, shows flattening of the enamel, and is smooth, without a median ridge.

Miorhizomys choristos (Flynn, 1982)

(Fig. 3a-c)

Type specimen. YGSP 4053, partial skull and skeleton from Y172, Dhok Pathan Formation near Mahluwala, Khaur area, Potwar Plateau, 8.3 Ma.

Studied material. PUPC 00/37, partial left dentary with incisor and m2-3; locality H16, 6.6 to 6.7 Ma.

Description. The left dentary represents a large bamboo rat, and as a member of the Rhizomyini clade, it shows burrowing features. It is robust and deep, and despite loss of bone along the ventral border of the incisor, jaw depth below m2 is 14.9 mm. The dentary is broken 6 mm anterior to the alveolus of m1 and 8.5 mm posterior to m3. The masseteric crest is prominent, but the lower portion is not greatly inflated, lacks an anterior extension, and intersects the upper portion as a rounded junction below m1, above the midline of the dentary. Anterior to this junction, at about the midline, is the mental foramen. The first lower molar is absent, but would have been small, relative to m2-3 as in the type specimen of *M. choristos*. The alveolus for m1 narrows anteriorly, given the small, round anterior root and the broad, oblique posterior root. The large posterior root of the last molar makes the alveolus of m3 extend beyond its posterior wall. Although m1 is missing, the alveolar length of the molar row (13.6 mm, Table 1) would exceed somewhat the actual length of m1-3.

Molars are lophodont and slightly higher crowned buccally than lingually. The protoconid of m2 is the largest cusp, and the metaconid is the highest component of the occlusal surface. The m2 is tetralophodont, with the metaconid and entoconid being large, transverse cusps incorporated into the metalophid and hypolophid, respectively. The connections of the metaconid to the protoconid and of the entoconid to the hypoconid are narrow, and these cusps would be isolated in early wear. Similarly, the central connection between protoconid and hypoconid is weak at this stage of wear. The posterolophid continues lingually from the hypoconid, nearly to the margin of the tooth, but the mesolophid is a short extension of the protoconid, terminating lingual to the midline of m2. Dimensions of m2 are 4.35 mm long by 5.05 mm wide.

1
2
3
4 The third molar (5.8 x 4.4 mm) is longer but narrower than m2, and less worn (a rhizomyine
5
6 characteristic is a strong wear gradient, increasing anteriorly). Its protoconid and metaconid are
7
8 smaller but similar to those of m2; the entoconid is much smaller than on m2. The mesolophid is
9
10 quite short and abuts tightly the entoconid, with which it will unite after further wear. The major
11
12 buccal reentrant is deep, extending directly to the entoconid, such that the hypolophid is
13
14 reduced to a thin posteriorly-directed connection that joins the posterolophid slightly lingual to
15
16 the midline of m3. The posterolophid is less broad than that of m2. As a consequence of the
17
18 transverse reduction of the tooth, the posterior lake of m3 is small and positioned lingually.
19
20 The incisor is stout (4.3 mm transverse, 6.1 mm antero-posterior), triangular in cross section,
21
22 with flattened anterior surface. The anteromedial corner is slightly more acute than a right
23
24 angle; the anterolateral corner is rounded, with the enamel wrapping onto about 30% of the
25
26 lateral side of the incisor. The enamel bears fine rugosities, but is otherwise smooth.
27
28
29
30
31
32
33
34
35
36
37

38 *Genus **Rhizomyides** Bohlin, 1946*

39
40 *Type species. R. sivalensis* Bohlin, 1946, from late Miocene Haritalyangar, northern India.

41
42 *Diagnosis (emended).* Medium size to large tachyoryctine with moderate hypsodonty, high
43
44 crowned in early wear; lower masseteric crest inclined, extending anteriorly beyond upper
45
46 masseteric crest; lower molar mesolophids well developed, large posterior enamel lake on m3;
47
48 dentary not deepened and incisor has rounded enamel.
49
50
51
52

53 *Discussion.* *Rhizomyides* is a long-ranging (temporally) tachyoryctine, and not a member of the
54
55 crown bamboo rats. The genus does not display the array of strongly fossorial traits evident in
56
57 *Miorhizomys* (Flynn, 2009). For example (Fig. 3b, e), the dentary is much shallower than in the
58
59
60

rhizomyine *Miorhizomys*. Its masseteric crest differs greatly from that of *Miorhizomys*: the lower part is inflated and extends anteriorly beyond the upper crest. The genus also retains primitive incisor morphology (Fig. 2): smaller incisor with rounded enamel bearing a longitudinal ridge.

***Rhizomyides sivalensis* (Lydekker, 1884)**

(Fig. 3d-f; Fig. 4a-f)

Studied material. PUPC 09/30, right dentary fragment with incisor, m2-3 from H16. From H18, PUPC 09/31 and 09/32, right and left halves of mandible of one individual with all teeth. All specimens 6.6 to 6.7 Ma.

Description. This large member of the genus has a dentary that is shallow relative to that of Rhizomyini (Fig. 3e, depth below m2 = 9.9 mm on PUPC 09/30; 9.8 mm on PUPC 09/32). Although incomplete, PUPC 09/30 preserves much of the dentary morphology; its 31.4 mm length includes a high condyle and a strong coronoid process (both broken). Alveolar length of the molar row is 11.6 mm. On all specimens, the masseteric crest is prominent, the lower portion particularly inflated and inclined. It joins the upper part of the crest below the posterior root of m1, and continues anteriorly with an extension below the anterior root of m1, at the midlevel of the dentary. Anterior to m1 at the midlevel of the jaw is the mental foramen.

Molars are slightly higher crowned buccally than lingually. Wear transforms the lophodont pattern into a flattened occlusal surface, yet it should be noted that, for both *Rhizomyides* and *Miorhizomys*, the surface of the molar row is not a plane. This is most evident for the mandible PUPC 09/31-32. The wear surface is a slightly concave, helical groove (due to wear by the upper dentition), deep at the lingual side of m3, running anteriorly and slightly diagonally across m2 toward its anterobuccal side, and continuing on the buccal side of m1. The result is greater wear

on the buccal sides of, especially, m2 and m1. The molar row is preserved in PUPC 09/31 and 09/32, with length of 13.7 and 13.2 mm, respectively.

Unworn lower molars are mesodont and inclined anteriorly. The first molar is narrowest, but elongated. The large hypoconid continues lingually with a sweeping posterolophid. It has a narrow connection to the posterior arm of the protoconid near the midline of the tooth. The posterior arm of the protoconid continues lingually and bifurcates as an entoconid and a short, narrow mesolophid, both lingual in position. The entoconid is large, with accessory spurs, and upon wear joins the posterolophid lingually. Opposite the protoconid is a broad lingual reentrant whose anterior wall is the metaconid. The metaconid joins the protoconid not directly, but at the base of its anterior arm, which continues as the anterolophid. The anterolophid is a D-shaped anterior complex that joins the protoconid buccal to the midline of the tooth and the metaconid at the lingual side. There is a broad transverse sinus between hypoconid and protoconid, and a smaller sinus between protoconid and anterior complex.

The m2 and m3 are represented by specimens in early and late wear. The shape of the tooth causes apparent change in the dimensions of the occlusal surface: the longer than wide tooth in early wear seems to become wider than long in late wear. This accounts for the considerable variation in measurements of Table 1. Also, the entire molar row (m1-m3) wears such that old individuals appear to have short tooth rows. The dominant features of m2 are a triangular protoconid and hypoconid with deep, posteriorly-directed buccal reentrant, and four transverse crests, the metalophid, mesolophid, hypolophid, and posterolophid. Because the buccal reentrant or sinus is deep, crossing half of the tooth, the hypolophid and mesolophid opposite the sinus are short. The metalophid and hypolophid widen at the lingual wall, representing an

1
2
3
4 expanded metaconid and entoconid, respectively. The mesolophid between is narrow and nearly
5
6 reaches the lingual wall (later wear unites the mesolophid with the wall). The sweeping
7
8 posterolophid is the distal wall of the tooth. These structures yield three lingual enamel lakes for
9
10 m2, the smallest between mesolophid and hypolophid. An interesting variation is observed in
11
12 slightly worn m2 (not m3) of PUPC 09/31 and 09/32: the straight anterior wall of the
13
14 anterolophid includes a short buccal arm that makes an ephemeral notch in the occlusal trace of
15
16 the enamel. This feature with other molar and incisor morphology is inherited from *Kanisamys*-
17
18 like structures of earlier Rhizomyinae.
19
20
21
22
23

24
25 The last lower molar is narrower than m2 (see Table 1), but its base expands distally so that a
26
27 length measurement including the base of the tooth is greater than at the occlusal surface. Its
28
29 anterior width is less than that of m2, and m3 narrows posteriorly. The structure of the
30
31 metalophid is much like that of m2, but the mesolophid is somewhat shorter. In lightly worn
32
33 PUPC 09/31 and 09/32, the lingual end of the metalophid preserves a tiny enamel lake that
34
35 recalls the free lingual end of the anterolophid of early Rhizomyinae. The narrow posterior end of
36
37 m3 has its hypoconid slightly displaced lingually, and its entoconid greatly displaced buccally and
38
39 reduced, relative to m2. The posterolophid is greatly shortened relative to that of m2, and the
40
41 posterior enamel lake is correspondingly small. In slightly worn PUPC 09/31 and 09/32 the
42
43 posterior lake is even smaller than the middle lake between hypolophid and mesolophid, but
44
45 those proportions reverse with wear. In later wear, the shallower middle lake disappears, but the
46
47 posterior lake persists, and is relatively bigger than that of the m3 of *Miorhizomys*. Intra-
48
49 individual variation is evident in comparing left and right m3; PUPC 90/32 has a third molar that
50
51
52
53
54
55
56
57
58
59
60
61

1
2
3
4 is more vertically disposed, with a less posteriorly projecting distal extremity – thus its length
5
6 measurement is less than that of PUPC 90/31.
7

8
9 The lower incisor of *Rhizomyides* contrasts greatly with that of *Miorhizomys*. The incisor has a
10 rounded enamel face, sharply rounded medially and broadly rounded laterally, with enamel
11 wrapping half way around its lateral side. There is a strong, persistent ridge of raised enamel
12 running medial to the midline of the shaft. This contrasts sharply with the smooth, flat enamel of
13 the stout incisor of *Miorhizomys* (Fig. 2). Measurements (Table 1) vary depending on incisor
14 maturity (individual age, or where measured in the jaw).
15
16
17
18
19
20
21
22
23
24
25
26

27 Genus ***Protachyoryctes*** Hinton, 1933
28

29
30 *Type species.* *P. tatroti* Hinton, 1933, from late Miocene deposits below Tatrot Village, northern
31 Pakistan, only known species.
32
33

34
35 *Diagnosis (emended).* Medium size and hypsodont; lower masseteric crest nearly horizontal, not
36 strongly inclined; lower molars with deep buccal reentrants, reduced mesolophids, and strong
37 posterolophids; lower incisor relatively small, with strongly rounded enamel.
38
39
40
41

42
43 *Discussion.* This monotypic genus is known only from the late Miocene of the Potwar Plateau,
44 northern Pakistan. A member of Tachyoryctini, it is not strongly fossorial, but rather has molars
45 of crown height great enough to be considered hypsodont. Its masseteric crest is similar to, but
46 not as inclined as in *Rhizomyides*, and its incisor is relatively smaller and strongly rounded.
47
48
49
50
51

52
53 ***Protachyoryctes tatroti*** Hinton, 1933
54

55
56 (Fig. 4g, h)
57
58
59
60
61

1
2
3
4 *Studied material.* PUPC 90/34, damaged right dentary with broken incisor and m2-3, from H18,
5
6
7 6.6 to 6.7 Ma in age.
8

9
10 *Description:* Alveolar length of tooth row, on poorly preserved PUPC 90/34 about 10.4 mm;
11
12 mandibular depth below m2 is 7.7 mm. Tooth dimensions (Table 1) are less than corresponding
13
14 measurements for *Rhizomyides sivalensis*. The bone surface is weathered, but the lower
15
16 masseteric crest is nearly horizontal, not inclined. Molars are still high crowned despite
17
18 significant wear. The m2 is partly missing, but a thin mesolophid is evident and the middle
19
20 enamel lake is small. Buccal sinuses of m2 and m3 are deep. The posterior lake of m3 is large.
21
22
23 The incisor is strongly rounded in cross section, the enamel bearing a single ridge.
24
25
26

27 *Discussion.* The damaged PUPC 90/34 resembles *Rhizomyides* specimens, but preserves few
28
29 distinctive features. The partially-preserved heavy masseteric crest is nearly horizontal, the
30
31 relatively small incisor is rounded with a ridge, and the molars in their worn state show complete
32
33 transverse lophs and a large posterior lake. Tooth dimensions (Table 1) are too small for
34
35 *Rhizomyides sivalensis*, but consistent with *Protachyoryctes tatroti*, previously known from
36
37 nearby sites in the Dhok Pathan Formation. This specimen is younger than previously known
38
39 fossils, which date to 7.8 to 7.1 Ma.
40
41
42
43
44
45
46
47

48 Family **Hystriidae** Fischer, 1817
49

50 Genus **Hystrix** Linnaeus, 1758
51

52 *Type species.* *H. cristata* Linnaeus, 1758, extant in Europe.
53
54

55 **Hystrix** sp. (Fig. 5c-e)
56
57
58
59
60
61

1
2
3
4 *Studied material.* PUPC 15/444, upper left incisor tip from H18, 6.6 to 6.7 Ma; PUPC 09/35, right
5
6
7 M3 from Tatrot Village, ~3.4 Ma (estimate in Barry et al. 2002).
8

9
10 *Description.* The strongly curved incisor represents an upper tooth, with tip intact. It is stout,
11
12 transverse dimension = 5 mm, antero-posterior depth = 8 mm, and in transverse section nearly
13
14 flattened anteriorly, more rounded on its lateral side. The enamel band wraps onto 1/3 of the
15
16 lateral side, and bears no grooves or ridges, but is pigmented orange-red.
17
18

19
20 The molar from Tatrot is somewhat polished by abrasion and the enamel is pitted by exposure. It
21
22 is considered to represent a right M3 due to its asymmetrical shape, and flattened anterior wall,
23
24 although the single interdental appression facet is nearly unrecognizable. Its numerous
25
26 transverse enamel lakes represent the traces of major transverse lophs, with spurs between
27
28 them, subdividing the lakes. Atypical for a right M3, as here interpreted, a lingual lake is
29
30 elongated posterobuccally, rather than anterobuccally. Remnants of two small buccal roots are
31
32 evident; a large lingual root extends posteriorly to support the medial side and back of the tooth.
33
34
35 Dimensions are 10.2 mm long, 10.1 mm wide.
36
37
38

39
40 *Discussion.* These fossils do not include material definitive at the species level, and identification
41
42 as *Hystrix* is based on the multiple enamel loops in the molar and on large size similar to
43
44 porcupines previously found in the late Miocene of Pakistan. Careful analysis by Van Weers and
45
46 Rook (2003) showed that known Siwalik fossils of Miocene age are not distinguishable from
47
48 *Hystrix primigenia*. The incisor from Bhandar likely represents this species, but the Pliocene
49
50 molar is considerably younger and conceivably represents a distinct species. Its dimensions are
51
52 close to but above the size range cited by Van Weers and Rook (2003) for *Hystrix primigenia*.
53
54
55
56
57
58
59
60
61

Order **Lagomorpha** Brandt, 1855

Family **Leporidae** Fischer, 1817

Genus ***Alilepus*** Dice, 1931

Type species. *A. annectens* (Schlosser, 1924) from the late Miocene locality, Ertemte, Inner Mongolia.

Alilepus elongatus Winkler et al., 2011

(Fig. 5a, b)

Studied material. PUPC 09/33, right dentary fragment with alveolus for p3, damaged p4 and m1, and alveolus for anterior root of m2 from H18, 6.6 to 6.7 Ma.

Description. The jaw fragment represents a large rabbit, the size of described *Alilepus elongatus*. Preserved dentary depth below p4 is 11 mm. Although not well preserved, it has matrix filling the alveolus for p3, which indicates a length of ~4.4 mm and width of 3.7 mm for the anterior premolar. Dimensions of p4 are 2.8 mm long by 3.3 mm maximum width; and of m1, 2.7 by 3.45 mm. Enamel of the cheek teeth is heaviest buccally and on the distal wall of the trigonid, and in occlusal view the enamel trace undulates only slightly.

Discussion. The rabbit is identified based on occurrence in the area of a single kind of leporid in strata of similar age. The cheek tooth dimensions for PUPC 09/33 are consistent with the species as published by Winkler et al. (2011), except that the p3 seems large, due likely to an overestimate, since length and width are estimated only from the alveolus. Although poorly preserved, the specimen provides an approximate jaw depth, not previously known. It complements the fossils reported by Winkler et al. (2011), which date to about 7 to 7.4 Ma;

1
2
3
4 isolated teeth in that study date to about 6.5 Ma. *Alilepus* is a late Miocene immigrant into the
5
6
7 Indian subcontinent.
8
9

10 11 12 **Discussion** 13

14 The fossils reported here are important because they add size and morphological data to assess
15
16 variation in poorly known species, and because they extend or fill in the known time ranges for
17
18 these species. The bamboo rat represented by PUPC 00/37 is comparable morphologically to the
19
20 holotype of *Miorhizomys choristos* (Flynn 1982), but is a bit larger. Its deep dentary and chisel-
21
22 like incisor are typical of fossorial crown bamboo rats, Tribe Rhizomyini. Its distinctive molar
23
24 morphology with isolated entoconids, weak longitudinal connections, and diminutive posterior
25
26 enamel lake on m3 match the holotype and other specimens attributed to the species, including
27
28 material from Haritalyangar, India (Flynn et al. 1990; Table 1 here). Several previously known *M.*
29
30 *choristos* jaws from the Potwar Plateau date to 8.4-8.3 Ma, and the Haritalyangar material is
31
32 perhaps somewhat older (Pillans et al. 2005). One fragmentary specimen found in the Hasnot
33
34 area (YGSP 37068) in recent years dates to about 7 Ma. The new Bhandar specimen, in good
35
36 condition, is the youngest known *M. choristos* (Fig. 6).
37
38
39
40
41
42
43
44

45 The remaining rhizomyine dentaries share features that are primitive with respect to Rhizomyini
46
47 and are classified as Tachyoryctinae. Most striking are the shallower dentary with relatively
48
49 heavier muscle scar of the lower masseteric crest. The lower crest meets the upper crest slightly
50
51 more posteriorly than in *Miorhizomys*, but then continues with an anterior extension below m1.
52
53 The incisor is quite different: round with median longitudinal ridge. The incisor does not show
54
55 the heavy, chisel morphology of tooth-diggers. The molars are less derived than those of
56
57
58
59
60

Rhizomyini; they are more elongated with complete transverse lophs, longitudinal crests, and less reduced m3.

New hemi-mandibles with dentition nicely complement what is known of *Rhizomyides sivalensis*.

The new fossils, representing young and old individuals complement the large dental variation and size range (Table 1) for *Rhizomyides sivalensis*. The dentary is thickened laterally at the lower masseteric crest, and the lower incisor invariably is strongly rounded with a single persistent median ridge on the enamel. Molars appear to change in dimensions with wear, and the ontogeny is captured by comparing specimens in early and late wear. The worn (occlusal) surface increases in width with wear, and decreases in length, although length shortening is countered in m3 by the fact that its distal wall extends posteriorly toward the root. *Rhizomyides* has stronger, more independent mesolophids than *Miorhizomys*, and less inflated cusps. The entoconids are smaller and merged into hypolophids, and longitudinal crests continue into late wear. Left and right dentitions of the same individual differ somewhat.

Early workers recorded *Rhizomyides sivalensis* at Haritalyangar, India, and the Hasnot region, Pakistan (Lydekker 1878, Black 1972). Jacobs (1978) recovered specimens in the Hasnot region which proved to be assignable to this species. *Rhizomyides sivalensis* occurs in the Dhok Pathan Formation of the Potwar Plateau, Pakistan, in rocks spanning 7.3 to 6.5 Ma. The new Bhandar specimens fall at the young end of that time range. The specimen of *Protachyoryctes tatroti* reported here comes from near the type area for the species and is younger than specimens with good provenance (Fig. 6).

Hystrix has been known from the late Miocene Siwaliks for well over a century, beginning when Lydekker (1878, 1884) named and described a jaw from the Hasnot region *Hystrix sivalensis*.

1
2
3
4 Currently, based on the comprehensive work of van Weers (e.g., van Weers and Rook, 2003), the
5
6 Miocene Siwalik porcupine is considered indistinguishable from the western contemporaneous
7
8 member of the Pikermian chronofauna, *Hystrix primigenia*. Barry et al. (2002) note the late
9
10 Miocene introduction of this large rodent into the Siwalik fauna at about 8 Ma. *Hystrix* appears
11
12 to have remained an element of the South Asian mammal fauna to the present. We document
13
14 the genus in the late Pliocene at the classical Tatrot site, overlying the Dhok Pathan Formation.
15
16 Being large and significantly younger than known late Miocene porcupines, it possibly represents
17
18 another species of the genus.
19
20

21
22 Every specimen of *Alilepus elongatus* is important as it is a poorly-represented immigrant to the
23
24 Indian subcontinent in the late Miocene, arriving before 7.4 Ma (Barry et al. 2002). The new
25
26 Bhandar specimen is the only known partial mandible. This record is younger than the type
27
28 material of the species.
29
30
31
32
33
34
35
36
37

38 Conclusion

39
40 New surface finds of fossil jaws and teeth from the Hasnot region of the Potwar Plateau,
41
42 Pakistan, add morphological and biostratigraphic data for known late Miocene Siwalik small
43
44 mammal species ranges (Fig. 6). The several Miocene age rhizomyines include a member of the
45
46 crown bamboo rats, Tribe Rhizomyini (*Miorhizomys choristos*), and two tachyoryctin genera,
47
48 *Rhizomyides* and *Protachyoryctes*. In addition to the rhizomyine species, the new material
49
50 represents porcupines (*Hystrix*) and the leporid *Alilepus elongatus*, both late Miocene immigrants
51
52 to the Indian Subcontinent. The new occurrences are plotted with previously well-documented
53
54 occurrences to update the biostratigraphic ranges of these species. The newly recovered
55
56
57
58
59
60

specimens point to the importance of continued collecting in the Siwalik deposits of the Indian Subcontinent combined with careful preservation of geological provenance data.

Acknowledgments We thank the government of Pakistan and University of the Punjab authorities for supporting this research. Prof. Dr. Muhammad Akhtar and Syed Ghyour Abbas are thanked for their help in the field. Mr. Adeeb Babar prepared Figure 1. Dr. John C. Barry kindly advised us on age assessment of the strata of the Hasnot area. Irina Ruf made many helpful suggestions, and we thank Lutz Maul (and anonymous) for careful reading that led to an improved manuscript.

References

- Barry, J.C., M.E. Morgan, L.J. Flynn, D. Pilbeam, A.K. Behrensmeyer, S. Mahmood Raza, I.A. Khan, C. Badgley, J. Hicks, and J. Kelley. 2002. Faunal and environmental change in the Late Miocene Siwaliks of Northern Pakistan. *Paleobiology Memoir* 3: 1-71.
- Black, C.C. 1972. Review of fossil rodents from the Neogene Siwalik Beds of India and Pakistan. *Palaeontology* 15(2): 238-266.
- Bohlin, B. 1946. The fossil mammals from the Tertiary deposit of Taben-buluk, western Kansu. Part II: Simplicidentata, Carnivora, Artiodactyla, Perissodactyla, and Primates. *Sino-Swedish Expedition Publication* 28, 6 (4): 1–259, 9 pls.
- Bowdich, T.E. 1821. *An analysis of the natural classifications of Mammalia, for the use of students and travelers*. Paris: J. Smith.

Brandt, J.F. 1855. Beitrage zur nähern Kenntniss der Säugethiere Russland's. *Kaiserlichen Akademie der Wissenschaften, Saint Petersburg, Mémoires Mathématiques, Physiques et Naturelles* 7: 1-365.

Brown, B., W.K. Gregory, and M. Hellman. 1924. On three incomplete anthropoid jaws from the Siwaliks, India. *American Museum Novitates* 130: 1-9.

Dice, L.R. 1931. *Alilepus*, a new name to replace *Allolagus* Dice, preoccupied, and notes on several species of fossil hares. *Journal of Mammalogy*, 12:159-160.

Fischer von Waldheim, G. 1817. Adversaria zoologica. *Mémoires de la Société Impériale des Naturalistes de Moscou*. 5 : 368-428.

Flynn, L.J. 1982. Systematic revision of Siwalik Rhizomyidae (Rodentia). *Géobios* 15: 327-389.

Flynn, L.J. 2009. The antiquity of *Rhizomys* and independent acquisition of fossorial traits in subterranean muroids. In *Systematic Mammalogy: Contributions in Honor of Guy G. Musser*, eds. R.S. Voss and M.D. Carleton, 128-156. New York: Bulletin of the American Museum of Natural History 331.

Flynn, L.J., A. Sahni, J.J. Jaeger, B. Singh, S.B. Bhatia. 1990. Additional fossil rodents from the Siwalik Beds of India. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* B 93 (1): 7-20.

Gray, J.E. 1821. On the natural arrangement of vertebrate animals. *London Medical Repository* 15: 296-310.

Hinton, M.A.C. 1933. Diagnoses of new genera and species of rodents from Indian Tertiary deposits. *Annals and Magazine of Natural History Series* 10, 12: 620–622.

Jacobs, L.L. 1978. Fossil rodents (Rhizomyidae & Muridae) from Neogene Siwalik deposits, Pakistan. *Museum of Northern Arizona Press Bulletin* 52: 1-103.

Lewis, G. E. 1937. A new Siwalik correlation. *American Journal of Science* (Series 5) 33: 191-204.

Linnaeus, C. 1758. *Systema naturae per regna tria naturae, Secundum Classes, Ordines, Genera, Species cum characteribus, differentiis, synonymis, locis. Vol. 1, Regnuma Animale, 10th ed.* Stockholm: Laurentii Salvi.

Lydekker, R. 1878. Notices of Siwalik mammals. *Records of the Geological Survey of India* 11: 64-104.

Lydekker, R. 1884. Indian Tertiary and post-Tertiary Vertebrata. Part 3. Rodents and new ruminants from the Siwaliks and synopsis of Mammalia. *Memoirs of the Geological Survey of India* 10(3): 105-134.

Pillans, B., M. Williams, D. Cameron, R. Patnaik, J. Hogarth, A. Sahni, J.C. Sharma, F. Williams, and R.L. Bernor. 2005. Revised correlation of the Haritalyangar magnetostratigraphy, Indian Siwaliks: implications for the age of the Miocene hominids *Indopithecus* and *Sivapithecus*, with a note on a new hominid tooth. *Journal of Human Evolution* 48: 507-515.

Van Weers, D.J., and L. Rook. 2003. Turolian and Ruscinian porcupines (genus *Hystrix*, Rodentia) from Europe, Asia, and North Africa. *Paläontologische Zeitschrift* 77(1): 95-113.

Winge, H. 1887. Jordfundne og nulevende Gnavere (Rodentia) fra Lagoa Santa, Minas Geraes, Brasilien. *E Museo Lundii, University of Copenhagen* 1: 1-178.

Winkler, A.J., L.J. Flynn, and Y. Tomida. 2011. Fossil lagomorphs from the Potwar Plateau, northern Pakistan. *Palaeontologia Electronica* 14.3.38A, 16 p., <http://palaeo-electronica.org/toc.htm>.

FIGURE CAPTIONS

Figure 1. Map showing overview of much of the Indian Subcontinent and outcrop area of the Siwalik Group formations, inset showing the Punjab province, which includes the Potwar Plateau, and close-up of the Hasnot area. The principal fossil localities are H16 and H18, very near Bhandar Village.

Figure 2. Cross sections of lower incisors of Rhizomyinae. Left, the tachyoryctin *Rhizomyides sivalensis*, showing the less derived rounded enamel with characteristic ridge (arrow), and the bamboo rat *Miorhizomys choristos* (Tribe Rhizomyini, right), showing the features of large size, long A-P dimension, and flattened enamel surface. Anterior views of PUPC 09/30, *R. sivalensis* right incisor, and PUPC 0-0/37, *M. choristos* left incisor drawn to the same scale.

Figure 3. PUPC 00/37, *Miorhizomys choristos* left jaw (a-c), and PUPC 09/30, adult *Rhizomyides sivalensis* right jaw with worn m2-3 (d-f). Lateral views of the dentaries (b, e) show the difference in jaw depth, and the views of *R. sivalensis* (d, e) show the inflated lower masseteric crest.

Figure 4. PUPC 09/31 and 09/32, left (a, d, f) and right (b, e) dentaries of the same individual of *Rhizomyides sivalensis*; these are paired in anatomical position (c) with 5 mm scale. Occlusal, lateral and medial views of the left dentary are stacked for viewing and posed next to occlusal and lateral views of the right dentary. Lateral and occlusal views (g, h) of the smaller PUPC 90/34, right dentary fragment of *Protachyoryctes tatroti*, 5 mm scale. *P. tatroti* has a shallow dentary as does *R. sivalensis*.

Figure 5. *Alilepus elongatus*, PUPC 09/33, right jaw fragment with p4m1 in occlusal and lateral view (a, b), anterior to right. *Hystrix* sp. (c-e): occlusal and lingual views of PUPC 09/35, right M3 (c, d); (e) is an isolated left upper incisor in apical, medial, lateral, labial views (left to right).

Figure 6. Biostratigraphic ranges in the Potwar Plateau of taxa studied here. On the left is the time scale based on magnetostratigraphy updated from Barry et al. (2002), with ages in millions of years. To the right of the time line are locality numbers. Vertical lines show the ranges of five species, circles show dated localities: filled circles are Y-GSP localities and open circles represent the new Bhandar sites. We interpret oldest sites as first occurrences of taxa. The genus *Hystrix* is known to persist into the Pliocene, but there is no indication that any of the species became extinct by the end of the Miocene.

Figure

Table 1. Dimensions (molar rows and teeth) for *Miorhizomys choristos* and *Protachyoryctes tatroti* specimens compared with *Rhizomyides sivalensis*. *: row length based on alveoli

<i>Miorhizomys choristos</i>					<i>Protachyoryctes tatroti</i>				
	PUPC 00 37	YGSP 4053	YGSP (old) 16200	VPL/BS/ PU 102	PUPC 09 34	GSI D272	YGSP 8360	YGSP 15336	YGSP 50949
Molar Row	13.6*	12.5*	12.4*	12.1*	10.4*	9.9	11.2*	11	11
L m1	X	X	X	X	X	3.56	X	3.88	3.8
W m1	X	3.1	X	3	X	3.06	3	3.16	2.75
L m2	5.05	5.55	3.9	4.2	3.35	3.56	3.63	3.59	3.6
W m2	4.35	4.38	4.6	4.1	3.5	3.75	3.94	3.52	3.35
L m3	5.8	5.22	4.8	5.2	4.05	4	4.28	3.9	3.1
W m3	4.4	4.25	4.1	4.2	3.4	3.06	3.38	3.28	3.9
A-P incisor	6.1	5.2	6.2	5	2.9	X	2.6	2.8	X
W incisor	4.3	4	3.6	3.2	2.3	X	2.2	1.9	X

<i>Rhizomyides sivalensis</i>										
	PUPC 09 30	PUPC 09 31	PUPC 09 32	GSI D97	GSI D275	GSI D276	GSI D277	YGSP 49876	YGSP 15319	YGSP 50115
Molar Row	11.7*	13.7	13.2	14.4*	13.3*	15.3	14.9	14.2*	14.4*	15.5*
L m1	X	4.65	4.6	X	X	5.15	4.6	X	X	X
W m1	X	3.5	3.45	X	X	3.45	4.1	X	3.7	X
L m2	3.75	4.3	4.35	4.55	4.55	5.25	4.05	4.3	4	4.7
W m2	4.3	4.5	4.25	4.8	4.2	4.31	4.5	4.2	4.94	4.6
L m3	4.45	4.85	4.6	5.9	5.8	5.44	5.5	4.75	5.38	X
W m3	3.85	4.25	4.2	4.2	4.2	4.31	3.6	3.8	4.63	X
A-P incisor	3.85	3.25	3.2	3.95	X	X	3.55	3.45	3.94	3.9
W incisor	2.85	2.5	2.55	3.05	X	X	3.15	2.7	3.44	2.95