



Multi-Ethnic Analysis of Lipid-Associated Loci: The NHLBI CARE Project

Citation

Musunuru, Kiran, Simon P. R. Romaine, Guillaume Lettre, James G. Wilson, Kelly A. Volcik, Michael Y. Tsai, Herman A. Taylor, et al. 2012. Multi-ethnic analysis of lipid-associated loci: The NHLBI CARE project. PLoS ONE 7(5): e36473.

Published Version

doi:10.1371/journal.pone.0036473

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10399871>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Multi-Ethnic Analysis of Lipid-Associated Loci: The NHLBI CARE Project

Kiran Musunuru^{1,2,3}, Simon P. R. Romaine⁴, Guillaume Lettre⁵, James G. Wilson⁶, Kelly A. Volcik⁷, Michael Y. Tsai⁸, Herman A. Taylor Jr.^{6,9}, Pamela J. Schreiner⁸, Jerome I. Rotter¹⁰, Stephen S. Rich¹¹, Susan Redline¹², Bruce M. Psaty^{13,14}, George J. Papanicolaou¹⁵, Jose M. Ordovas¹⁶, Kiang Liu¹⁷, Ronald M. Krauss¹⁸, Nicole L. Glazer¹⁹, Stacey B. Gabriel¹, Myriam Fornage⁷, L. Adrienne Cupples^{19,20}, Sarah G. Buxbaum⁹, Eric Boerwinkle⁷, Christie M. Ballantyne^{21*}, Sekar Kathiresan^{1,2,3*}, Daniel J. Rader^{22*}

1 Broad Institute, Cambridge, Massachusetts, United States of America, **2** Massachusetts General Hospital, Boston, Massachusetts, United States of America, **3** Harvard Medical School, Boston, Massachusetts, United States of America, **4** University of Leeds, Leeds, United Kingdom, **5** Montreal Heart Institute, Université de Montréal, Montreal, Canada, **6** University of Mississippi Medical Center, Jackson, Mississippi, United States of America, **7** The University of Texas Health Science Center at Houston, Houston, Texas, United States of America, **8** School of Public Health, University of Minnesota, Minneapolis, Minnesota, United States of America, **9** Jackson State University, Jackson, Mississippi, United States of America, **10** Medical Genetics Institute, Cedars-Sinai Medical Center, Los Angeles, California, United States of America, **11** University of Virginia School of Medicine, Charlottesville, Virginia, United States of America, **12** University Hospitals, Case Medical Center and Case Western Reserve University, Cleveland, Ohio, United States of America, **13** University of Washington, Seattle, Washington, United States of America, **14** Group Health Research Institute, Group Health Cooperative, Seattle, Washington, United States of America, **15** National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, Maryland, United States of America, **16** JM-United States Department of Agriculture Human Nutrition Research Center on Aging at Tufts University, Boston, Massachusetts, United States of America, **17** Northwestern University Feinberg School of Medicine, Chicago, Illinois, United States of America, **18** Children's Hospital Oakland Research Institute, Oakland, California, United States of America, **19** Boston University, Boston, Massachusetts, United States of America, **20** The National Heart, Lung, Blood Institute's Framingham Heart Study, Framingham, Massachusetts, United States of America, **21** Baylor College of Medicine, Houston, Texas, United States of America, **22** University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania, United States of America

Abstract

Background: Whereas it is well established that plasma lipid levels have substantial heritability within populations, it remains unclear how many of the genetic determinants reported in previous studies (largely performed in European American cohorts) are relevant in different ethnicities.

Methodology/Principal Findings: We tested a set of ~50,000 polymorphisms from ~2,000 candidate genes and genetic loci from genome-wide association studies (GWAS) for association with low-density lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C), and triglycerides (TG) in 25,000 European Americans and 9,000 African Americans in the National Heart, Lung, and Blood Institute (NHLBI) Candidate Gene Association Resource (CARE). We replicated associations for a number of genes in one or both ethnicities and identified a novel lipid-associated variant in a locus harboring *ICAM1*. We compared the architecture of genetic loci associated with lipids in both African Americans and European Americans and found that the same genes were relevant across ethnic groups but the specific associated variants at each gene often differed.

Conclusions/Significance: We identify or provide further evidence for a number of genetic determinants of plasma lipid levels through population association studies. In many loci the determinants appear to differ substantially between African Americans and European Americans.

Citation: Musunuru K, Romaine SPR, Lettre G, Wilson JG, Volcik KA, et al. (2012) Multi-Ethnic Analysis of Lipid-Associated Loci: The NHLBI CARE Project. PLoS ONE 7(5): e36473. doi:10.1371/journal.pone.0036473

Editor: Massimo Federici, University of Tor Vergata, Italy

Received: April 6, 2011; **Accepted:** April 6, 2012; **Published:** May 21, 2012

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: The following nine parent studies have contributed parent study data, ancillary study data, and DNA samples through the Broad Institute of Massachusetts Institute of Technology and Harvard (N01-HC-65226): Atherosclerosis Risk in Communities: University of North Carolina at Chapel Hill (N01-HC-55015, N01-HC-55018), Baylor Medical College (N01-HC-55016), University of Mississippi Medical Center (N01-HC-55021), University of Minnesota (N01-HC-55019), Johns Hopkins University (N01-HC-55020), University of Texas, Houston (N01-HC-55022); Cardiovascular Health Study: University of Washington (N01-HC-85079, N01-HC-55222, U01-HL-080295), Wake Forest University (N01-HC-85080), Johns Hopkins University (N01-HC-85081, N01-HC-15103), University of Pittsburgh (N01-HC-85082), University of California, Davis (N01-HC-85083), University of California, Irvine (N01-HC-85084), New England Medical Center (N01-HC-85085), University of Vermont (N01-HC-85086), Georgetown University (N01-HC-35129), University of Wisconsin (N01-HC-75150); Cleveland Family Study: Case Western Reserve University (R01-HL-46380, M01-RR-00080); Cooperative Study of Sickle Cell Disease: University of Illinois (N01-HB-72982, N01-HB-97062), Howard University (N01-HB-72991, N01-HB-97061), University of Miami (N01-HB-72992, N01-HB-97064), Duke University (N01-HB-72993), George Washington University (N01-HB-72994), University of Tennessee (N01-HB-72995, N01-HB-97070), Yale University (N01-HB-72996, N01-HB-97072), Children's Hospital-Philadelphia (N01-HB-72997, N01-HB-97056), University of Chicago (N01-HB-72998, N01-HB-97053), Medical College of Georgia (N01-HB-73000, N01-HB-97060), Washington University (N01-HB-73001, N01-HB-97071), Jewish Hospital and Medical Center of Brooklyn (N01-HB-73002), Trustees of Health and Hospitals of the City of Boston, Inc., (N01-HB-73003), Children's Hospital-Oakland (N01-HB-73004, N01-HB-97054), University of Mississippi (N01-HB-73005), St. Luke's Hospital-New York (N01-HB-73006), Alta Bates-Herrick Hospital (N01-HB-97051), Columbia University (N01-HB-97058), St. Jude's Children's Research Hospital (N01-HB-97066), Research Foundation, State University of New York-Albany (N01-HB-97068, N01-HB-97069), New England Research Institute (N01-HB-97073), Interfaith Medical Center-Brooklyn (N01-HB-97085); Coronary Artery Risk in Young Adults: University of Alabama at Birmingham (N01-HC-48047, N01-HC-95095), University of Minnesota (N01-HC-48048), Northwestern University (N01-HC-48049), Kaiser Foundation Research Institute (N01-HC-48050), Tufts-New England Medical Center (N01-HC-45204), Wake Forest University (N01-HC-45205), Harbor-University of California Los Angeles Research and Education Institute (N01-HC-05187), University of California, Irvine (N01-HC-45134, N01-HC-95100)

;Framingham Heart Study: Boston University (N01-HC-25195, R01-HL-092577, R01-HL-076784, R01-AG-028321); Jackson Heart Study: Jackson State University (N01-HC-95170), University of Mississippi (N01-HC-95171), Tougaloo College (N01-HC-95172); Multi-Ethnic Study of Atherosclerosis: University of Washington (N01-HC-95159), University of California, Los Angeles (N01-HC-95160), Columbia University (N01-HC-95161), Johns Hopkins University (N01-HC-95162, N01-HC-95168), University of Minnesota (N01-HC-95163), Northwestern University (N01-HC-95164), Wake Forest University (N01-HC-95165), University of Vermont (N01-HC-95166), New England Medical Center (N01-HC-95167), Harbor-University of California Los Angeles Research and Education Institute (N01-HC-95169), Cedars-Sinai Medical Center (R01-HL-071205), University of Virginia (subcontract to R01-HL-071205); Sleep Heart Health Study: Johns Hopkins University (U01-HL-064360), Case Western University (U01-HL-063463), University of California, Davis (U01-HL-053916), University of Arizona (U01-HL-053938, U01-HL-053934), University of Pittsburgh (U01-HL-077813), Boston University (U01-HL-053941), MedStar Research Institute (U01-HL-063429), Johns Hopkins University (U01-HL-053937). This work was also supported in part by a T32 grant in Cell and Molecular Training for Cardiovascular Biology and K99-HL098364 from the National Heart, Lung, and Blood Institute (KM). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: cmb@bcm.edu (CMB); sekar@broadinstitute.org (SK); rader@mail.med.upenn.edu (DJR)

Introduction

Plasma concentrations of lipids and lipoproteins [low-density lipoprotein cholesterol (LDL), high-density lipoprotein cholesterol (HDL-C), triglycerides (TG)] are heritable risk factors for cardiovascular disease [1,2]. Recently, genome-wide association mapping of common variants (>5% minor allele frequency) in individuals of European ancestry has proven useful in identifying genetic loci contributing to plasma lipids [3]. Roughly one third of these loci harbor genes with previously recognized involvement in lipid metabolism, many by virtue of having rare variants that result in Mendelian disorders. The other loci are suspected to harbor novel lipid regulators, the study of which has the potential of providing important new insights into lipid biology.

As promising as these observations may be, several critical questions remain – How many of the previously reported loci are genuine causal determinants of plasma lipid levels? Are there more loci associated with plasma lipids that are discoverable with techniques besides genome-wide association mapping? Do any of these loci confer effects on lipids in an ethnic-specific manner? To address these questions, we utilized the National Heart, Lung, and Blood Institute (NHLBI) Candidate Gene Association Resource (CARE) comprising more than 40,000 individuals from nine prospective cohorts with measured lipid phenotypes and genotype information obtained using the “ITMAT-Broad-CARE” array, or IBC array, with ~50,000 polymorphisms from ~2,000 candidate genes/loci [4,5]. Most of the polymorphisms on this array were obtained from a systematic literature search for candidate genes implicated in cardiovascular diseases; these were supplemented with polymorphisms from loci identified in early genome-wide association studies (GWAS) for lipids [5].

We initiated this study with specific hypotheses: (1) Common DNA sequence variants in previously reported genes and loci, as well as additional novel loci, are associated with plasma lipids; (2) At some of these loci, the specific variants related to plasma lipids differ between ethnic groups. To test these hypotheses, we performed association analyses of the polymorphisms on the IBC array in 25,000 European Americans and 9,000 African Americans in CARE.

Materials and Methods

Ethics Statement

All participants in each of the CARE cohorts gave informed written consent. The Institutional Review Boards (IRBs) of each CARE cohort (i.e., the IRBs for each cohort's field centers, coordinating center, and laboratory center) have reviewed and approved the cohort's interaction with CARE. The study described in this manuscript was approved by the Committee on the Use of Humans as Experimental Subjects (COUHES) of the Massachusetts Institute of Technology.

CARE Study Design and Quality Control

A full description of the CARE Study is found elsewhere [4]. In brief, DNA samples and phenotypic information from nine NHLBI prospective cohorts [the Atherosclerosis Risk In Communities (ARIC) study, the Coronary Artery Risk Development In Young Adults (CARDIA) study, the Cleveland Family Study (CFS), the Cardiovascular Health Study (CHS), the Cooperative Study of Sickle Cell Disease (CSSCD), the Framingham Heart Study (FHS), the Jackson Heart Study (JHS), the Multi-Ethnic Study of Atherosclerosis (MESA), and the Sleep Heart Health Study (SHHS)] were assembled at the Broad Institute of MIT and Harvard. We used the following strata of cohort and ethnicity (EA = European American, AA = African American): ARIC-AA, CARDIA-AA, CFS-AA, CHS-AA, JHS, MESA-AA, ARIC-EA, CARDIA-EA, CFS-EA, CHS-EA, FHS, MESA-EA. Plasma lipid levels—LDL-C, HDL-C, TG—from the baseline examinations of cohort participants were used (Table S1). All phenotypes, including plasma lipid levels, had been standardized according to the Clinical Data Interchange Standard Consortium Study Data Tabulation Model (CDISC SDTM).

All DNA samples passing initial quality checks were interrogated with the IBC v2 chip for genotyping of 49,320 total single nucleotide polymorphisms (SNPs). Samples with overall genotyping success rate <95%, individual SNPs with genotyping success rate <95% within a cohort, monomorphic SNPs, and SNPs mapping to multiple genomic loci were removed. An inbreeding coefficient was calculated for each sample as a measure of heterozygosity, with those samples exceeding 4 standard deviations from the mean (suggesting poor DNA quality if too low, or sample contamination if too high) being removed. In cases of identical DNA samples, the sample with the lowest genotyping success rate was removed. Samples that shared 5% or more of their genome with many other samples were also removed. Additional outlier samples as determined by multidimensional scaling were also removed. SNPs for which genotype missingness could be predicted by surrounding haplotypes were removed, as were any SNPs found to be associated with chemistry plates. Pedigrees and SNPs not conforming to Mendelian expectations were identified and, where appropriate, samples were removed. These analyses were all performed with PLINK [6].

Because different ethnic groups were represented, with the expectation of differing genotype frequencies and admixture, no filters were applied for minor allele frequency or Hardy-Weinberg *P* values. The Hardy-Weinberg *P* value for each index SNP in each cohort is listed in Table S2.

About 2,000 ancestry-informative markers were included in the IBC v2 chip [4]. ANCESTRYMAP [7] was used to generate estimates for global ancestry in African American samples. EIGENSTRAT [8] was used for self-reported African Americans and European Americans separately to generate ten principal components each accounting for population stratification. We

found that the first principal component for African Americans was highly correlated with the global ancestry estimates ($r^2 > 0.98$).

Phenotype Modeling

We modeled the lipid phenotypes in the following ways. LDL-C was calculated according to Friedewald's formula: $\text{LDL-C} = \text{total cholesterol} - \text{HDL-C} - (\text{TG} \div 5)$. If a TG value was >400 mg/dL, LDL-C was treated as a missing value. For individuals on lipid-lowering therapy, the LDL-C value was multiplied by 1.42 to model a 30% reduction in LDL-C on therapy. This represents the average expected reduction in LDL-C with a first-generation statin, the most commonly used lipid-lowering medication during the study periods of most of the cohorts [9]. TG values were \log_{10} -transformed. Sex-specific phenotype residuals were constructed within strata of cohort and ethnicity after accounting for age and age². Each set of residuals was standardized to a mean of zero and a standard deviation of one. The standardized residual served as the phenotype in genotype-phenotype association analyses. Generation of residuals was performed with the R statistical package (The R Foundation for Statistical Computing, Vienna, Austria).

Association Testing

For each of the three traits, we used linear regression to test SNP-phenotype associations in each stratum of cohort and ethnicity assuming an additive genetic model, using the ten calculated principal components as covariates: phenotype \sim genotype + PC1+ PC2+ PC3+ PC4+ PC5+ PC6+ PC7+ PC8+ PC9+ PC10. These association analyses were performed in PLINK. Genotype-phenotype associations within each ethnic group were assessed by weighted z-score-based fixed-effects meta-analysis and effect size estimates (β values) generated by inverse-variance weighted meta-analysis, both using METAL (G. Abecasis, University of Michigan). Genomic control correction was applied to each cohort individually prior to meta-analysis, and a final genomic control correction was applied to each ethnicity's meta-analysis dataset. For all cross-ethnic comparisons, only SNPs available in all cohorts (12 total) were considered. We considered $P = 1 \times 10^{-6}$ to be the threshold for statistical significance, applying a Bonferroni correction for the number of SNPs tested ($0.05 \div 50,000$).

We performed sensitivity analyses for two cohorts for which there were significant numbers of related individuals—CFS and FHS—to account for family relationships, using a linear mixed effects (LME) model to analyze the lipid residuals, with the SNP genotype treated as a fixed effect, and a random effect according to the degree of relatedness within a family [10]. Upon substitution of these data into the meta-analyses for each lipid trait, there were essentially no differences in the SNPs found to be statistically significant, and these data were not considered further.

To identify SNPs exerting sex-specific effects, we added sex as a covariate in the above linear regression model and tested for a formal SNP \times sex interaction; the results were meta-analyzed as described above. Similarly, to test for SNP \times SNP interactions among the most strongly associated SNPs at each of the identified loci, we added the minor allele count for each SNP as a covariate, in turn, into the above model (with sex removed), testing for a formal interaction between each SNP pair combination.

For those loci containing IBC-significant SNPs in both ethnicities, we performed SNP conditional analyses. These were conducted by adding SNP allele counts as extra covariates to the initial linear regression model described above. The conditional analyses were performed iteratively, with additional independent SNPs identified at each step in each locus added to the model.

Adjusted R^2 values for each SNP included in the conditional analyses, as well as for the combination of independently significant SNPs at each locus, were calculated by replicating the relevant linear regression model in the largest cohort (ARIC for both ethnicities) using Statistics Package for Social Sciences (SPSS) software (version 16.0; SPSS Inc., Chicago, IL), with principal components excluded.

Results

Replication of Reported Lipid-associated Loci and Identification of New Loci

Nineteen lipid-associated loci that were reported in an early set of lipid GWAS studies of individuals of European ancestry were included in the IBC v2 array—these loci harbor the genes *SORT1*, *CILP2/PBX4*, *GALNT2*, *MLXIPL*, *TRIB1*, *ANGPTL3*, *APOB*, *APOE*, *HMGCR*, *LDLR*, *PCSK9*, *ABCA1*, *APOA1/APOA5*, *CETP*, *LIPC*, *LIPG*, *LPL*, *GCKR*, and *MVK-MMAB* [11–14]. We found that SNPs at all 19 loci replicated in the CARE European American cohorts (Table 1) at the pre-specified IBC significance level ($P < 1 \times 10^{-6}$). Importantly, there was no overlap between the GWAS cohorts and the CARE cohorts. We also found that SNPs at 10 loci replicated in the CARE African American cohorts—*SORT1* (sortilin), *APOB* (apolipoprotein B), *APOE* (apolipoprotein E), *LDLR* (LDL receptor), *PCSK9* (proprotein convertase subtilisin/kexin type 9), *ABCA1* [ATP-binding cassette, sub-family A (ABC1), member 1], *APOA1* (apolipoprotein A-I), *CETP*, (cholesteryl ester transfer protein), *LIPC* (hepatic lipase), and *LPL* (lipoprotein lipase) (Table 1). All of these genes, with the exception of *SORT1*, have well-described functions in lipoprotein metabolism.

We also found significant associations for a number of IBC SNPs chosen for their proximity to candidate genes selected from the literature. Of these, eight loci—*FADS1* (fatty acid desaturase 1), *PLTP* (phospholipid transfer protein), *LCAT* (lecithin-cholesterol acyltransferase), *ANGPTL4* (angiopoietin-like 4), *ABCG5/ABCG8* (ATP-binding cassette sub-family G members 5 and 8; sterolin-1 and -2), *LPA* [lipoprotein(a)], *NPC1L1* [NPC1 (Niemann-Pick disease, type C1, gene)-like 1], *HPR* (haptoglobin-related protein)—were shown to have genome-wide-significant lipid-associated SNP variants in GWAS reported subsequent to the fabrication of the IBC array (Table 1) [3]. All of these genes, with the exception of *FADS1* and *HPR*, have well-described functions in lipoprotein metabolism. The *LCAT* locus was notable for being IBC-significant for HDL-C in both CARE European Americans and African Americans. An additional significant locus in African Americans harboring *CD36* (cluster of differentiation 36; thrombospondin receptor) had previously been reported in a candidate gene study; indeed, the same SNP that was IBC-significant (rs3211938) was shown in that prior study to be a nonsense coding variant that resulted in CD36 deficiency in a homozygous individual and was associated with increased HDL-C levels ($P = 0.00018$) and decreased TG levels ($P = 0.0059$) in an African American cohort [15].

Finally, one locus harbors an IBC-significant variant that has not previously been reported to be associated with lipid traits—*ICAMI* (intercellular adhesion molecule 1) in CARE African Americans (Table 1). Of note, the P value of association (1.24×10^{-8}) for the *ICAMI* SNP rs5030359 is not only IBC-significant but also genome-wide-significant. This variant is of low frequency in African Americans (just under 1%) and is virtually absent in European Americans. Interestingly, this variant demonstrated the largest effect size for any trait + ethnicity combination ($\beta = -0.52$).

Table 1. SNP variants associated with lipid traits in CARE European Americans and African Americans.

Gene(s) of interest in associated interval	Trait	SNP	Chr	Position	Alleles	European American		African American		European American P	African American P	European American β	African American β	Direction (EA, AA)
						MAF	MAF							
PCSK9	LDL	rs11591147	1	55278235	T, G	0.016	0.0033	1.74×10^{-28}	0.00011	5.05×10^{-11}	-0.44	-0.51	--	
		rs11806638	1	55290748	A, C	0.056	0.33	0.012	5.05×10^{-11}	-0.054	-0.11	-0.11	--	
ANGPTL3	TG	rs1748197	1	62828700	A, G	0.34	0.34	6.23×10^{-7}	0.012	6.23×10^{-7}	-0.053	-0.042	--	
	LDL	rs7528419	1	109618715	A, G	0.22	0.28	1.69×10^{-51}	1.34×10^{-16}	0.18	0.15	0.15	++	
SORT1		rs12740374	1	109619113	T, G	0.22	0.26	2.90×10^{-51}	9.33×10^{-20}	-0.18	-0.17	-0.17	--	
	HDL	rs4846918	1	228367209	T, C	0.15	0.14	4.22×10^{-7}	0.45	0.069	0.016	0.016	++	
GALNT2	LDL	rs934197	2	21120966	A, G	0.32	0.12	2.88×10^{-34}	0.0081	0.13	0.065	0.065	++	
		rs562338	2	21141826	A, G	0.18	0.39	7.60×10^{-34}	2.66×10^{-8}	-0.15	-0.092	-0.092	--	
HDL		rs673548	2	21091049	A, G	0.21	0.22	6.45×10^{-9}	0.044	0.068	0.037	0.037	++	
	TG	rs1042034	2	21078786	T, C	0.21	0.16	2.23×10^{-10}	0.00031	0.068	0.079	0.079	++	
GCCR	TG	rs1260326	2	27584444	T, C	0.41	0.14	6.79×10^{-35}	8.23×10^{-5}	0.037	0.088	0.088	++	
	LDL	rs4953023	2	43927504	A, G	0.070	0.078	2.41×10^{-8}	0.0011	-0.11	-0.10	-0.10	--	
ABCG5-ABCG8	LDL	rs12916	5	74692295	T, C	0.41	0.25	5.50×10^{-13}	0.099	-0.075	-0.031	-0.031	--	
	LDL	rs10455872	6	160930108	A, G	0.067	0.010	2.81×10^{-12}	0.24	-0.14	-0.081	-0.081	--	
NPC1L1	LDL	rs17725246	7	44548511	T, C	0.19	0.25	5.56×10^{-7}	0.025	-0.061	-0.041	-0.041	--	
	TG	rs17145750	7	72664314	T, C	0.15	0.11	4.22×10^{-25}	0.12	-0.0017	-0.040	-0.040	--	
CD36	HDL	rs3211938	7	80138385	T, G	0.00010	0.083	0.030	2.60×10^{-12}	-1.31	-0.1931	-0.1931	--	
	HDL	rs3916027	8	19869148	A, G	0.26	0.41	1.01×10^{-31}	1.93×10^{-7}	0.13	0.082	0.082	++	
LPL		rs13702	8	19868772	T, C	0.29	0.48	2.14×10^{-28}	3.12×10^{-11}	-0.12	-0.10	-0.10	--	
	TG	rs3916027	8	19869148	A, G	0.26	0.41	1.21×10^{-40}	2.65×10^{-8}	-0.072	-0.089	-0.089	--	
TRIB1	LDL	rs327	8	19863816	T, G	0.27	0.41	2.72×10^{-36}	1.03×10^{-9}	0.0026	0.097	0.097	++	
	HDL	rs6982636	8	126548497	A, G	0.47	0.36	4.29×10^{-9}	1.0	-0.059	-0.00	-0.00	--	
APOA1	HDL	rs2980880	8	126550154	A, G	0.31	0.32	5.85×10^{-7}	0.46	0.052	0.012	0.012	++	
	TG	rs2980875	8	126550929	A, G	0.47	0.37	1.15×10^{-15}	0.19	0.057	0.023	0.023	++	
APOA1-C3-A4-A5	HDL	rs1883025	9	106704122	T, C	0.26	0.34	6.25×10^{-7}	0.031	-0.055	-0.036	-0.036	--	
	HDL	rs2515629	9	106634185	A, G	0.17	0.17	0.53	2.04×10^{-7}	0.0079	0.11	0.11	++	
FADS1-FADS2-FADS3	HDL	rs1535	11	61354548	A, G	0.33	0.15	5.29×10^{-9}	0.00055	0.059	0.074	0.074	++	
	HDL	rs10750097	11	116169250	A, G	0.21	0.43	2.76×10^{-18}	0.55	0.10	-0.0094	-0.0094	+-	
MMAB-MVK	TG	rs2075290	11	116158506	T, C	0.070	0.055	8.13×10^{-51}	0.00044	-0.064	-0.12	-0.12	--	
	HDL	rs9804646	11	116170289	T, C	0.079	0.36	0.011	4.81×10^{-7}	-0.039	-0.083	-0.083	--	
LIPC	HDL	rs2075440	12	108343483	A, G	0.45	0.35	6.78×10^{-7}	0.45	-0.048	-0.013	-0.013	--	
	HDL	rs2070895	15	56511231	A, G	0.21	0.47	3.90×10^{-24}	3.93×10^{-10}	0.12	0.097	0.097	++	
CEFTP	HDL	rs17231506	16	55552029	T, C	0.32	0.14	1.50×10^{-120}	1.54×10^{-8}	0.24	0.12	0.12	++	
		rs17231520	16	55553328	A, G	-	0.070	-	7.65×10^{-58}	-	0.50	0.50	0+	

Table 1. Cont.

Gene(s) of interest in associated interval	Trait	SNP	Chr	Position	Alleles	European American		African American		European American <i>P</i>	African American <i>P</i>	European American β	African American β	Direction (EA, AA)
						MAF	MAF							
LCAT	HDL	rs2107369	16	66673519	T, G	0.14	0.26	9.23×10^{-8}	0.58	1.22×10^{-8}	-0.076	-0.010	--	
		rs255052	16	66582496	A, G	0.15	0.22	6.63×10^{-7}			0.069	0.11	++	
HPR	LDL	rs2000999	16	70665594	A, G	0.20	-	2.39×10^{-8}	-		0.069	-	+0	
LIPG	HDL	rs1943981	18	45423813	A, T	0.16	0.05	1.83×10^{-13}	0.41		-0.095	-0.027	--	
ANGPTL4	HDL	rs2278236	19	8337581	A, G	0.48	0.47	6.46×10^{-7}	0.96		0.048	-0.00060	+ -	
ICAM1	LDL	rs5030359	19	10249462	A, G	-	0.0086	-	1.24×10^{-8}		-	-0.52	0-	
LDLR	LDL	rs6511720	19	11063306	T, G	0.12	0.14	1.38×10^{-49}	1.75×10^{-18}		-0.23	-0.20	--	
CSPG3-CILP2-PBX4	TG	rs3794991	19	19471596	T, C	0.090	0.066	4.83×10^{-10}	0.98		-0.084	0.0012	- +	
APOE	LDL	rs12721046	19	50113094	A, G	0.15	0.030	2.24×10^{-29}	0.00098		0.16	0.16	+ +	
		rs389261	19	50112183	A, G	-	0.25	-	1.10×10^{-14}		-	0.15	0+	
	HDL	rs12721046	19	50113094	A, G	0.15	0.030	1.67×10^{-7}	0.056		-0.075	-0.087	--	
	TG	rs439401	19	50106291	T, C	0.37	0.14	1.85×10^{-15}	0.88		-0.022	-0.0046	--	
		rs12721054	19	50114427	A, G	0.00047	0.12	0.60	5.55×10^{-26}		0.33	0.26	+ +	
PLTP	HDL	rs4810479	20	43978455	T, C	0.25	0.40	1.90×10^{-9}	0.00013		0.067	0.062	+ +	

AA = African American; EA = European American; MAF = major allele frequency. Minor allele frequencies all derived from ARIC (largest cohort for both African Americans and European Americans). Bold indicates the most highly associated SNP (lowest *P* value) at IBC significance ($P < 1 \times 10^{-6}$) for locus + trait + ethnicity. Direction is modeled on the first allele listed in Alleles column. doi:10.1371/journal.pone.0036473.t001

Architecture of Lipid Loci in African Americans Compared to European Americans

Comparing each index SNP in each of the lipid loci in European Americans to African Americans, we noted quite varied patterns of association. In some cases, the effect size estimates and minor allele frequencies (MAFs) are quite similar in two ethnicities. An example is SNP rs6511720 in the *LDLR* locus, with $\beta = -0.23$ (in standard deviation units) and $\text{MAF} = 0.12$ in European Americans, $\beta = -0.20$ and $\text{MAF} = 0.14$ in African Americans for LDL-C (Table 1). The relative P values ($P = 1.38 \times 10^{-49}$ vs. $P = 1.75 \times 10^{-18}$) can be attributed solely to the different sample sizes (25,000 European Americans vs. 9,000 African Americans). Another example is rs1748197 in the *ANGPTL3* (angiopoietin-like 3) locus, which has $\beta = -0.053$ and $\text{MAF} = 0.34$ in European Americans, $\beta = -0.042$ and $\text{MAF} = 0.34$ in African Americans for TG ($P = 6.23 \times 10^{-7}$ vs. $P = 0.012$).

Other index SNPs diverged considerably in the two ethnicities. Some of the SNPs displayed considerable differences in their estimated effect sizes despite similar MAFs. Notable examples include rs2278236 in the *ANGPTL4* locus, with $\beta = 0.048$ ($\text{MAF} = 0.48$; $P = 6.46 \times 10^{-7}$) in European Americans and $\beta = -0.00060$ ($\text{MAF} = 0.48$; $P = 0.96$) in African Americans; rs4846918 in the *GALNT2* (UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 2) locus, with $\beta = 0.069$ ($\text{MAF} = 0.15$; $P = 4.22 \times 10^{-7}$) in European Americans and $\beta = 0.016$ ($\text{MAF} = 0.14$; $P = 0.45$) in African Americans for HDL-C; and rs2515629 in the *ABCA1* locus, with $\beta = 0.0079$ ($\text{MAF} = 0.17$; $P = 0.53$) in European Americans and $\beta = 0.11$ ($\text{MAF} = 0.17$; $P = 2.04 \times 10^{-7}$) in African Americans for HDL-C. These findings indicate considerable heterogeneity in the architecture of lipid-related loci in the two ethnicities.

We took advantage of the dense genotyping in each locus on the IBC v2 array to perform more detailed comparisons of the 11 loci that harbored IBC-significant SNPs for both ethnicities (Table 2). In only two of the loci (*LIPC* for HDL-C, *LDLR* for LDL-C) was the same SNP the most highly associated for the trait in each ethnicity. At an additional four loci (*SORT1* with LDL-C, *APOB* with LDL-C, *LPL* with HDL-C and TG, *LCAT* with HDL), the most highly associated SNP in one ethnicity was among the most highly associated SNPs in the other ethnicities. In the remaining five loci (*PCSK9* for LDL-C, *ABCA1* for HDL-C, *APOA1-C3-A4-A5* for TG, *CETP* for HDL, *APOE* for LDL-C and TG), the most highly associated SNPs in the ethnicities did not overlap.

For each ethnicity we performed conditional analyses with the most highly associated SNPs in each of the 11 loci to uncover any additional, independently associated SNPs (at IBC significance) in the same gene regions (Table 2). We performed this iteratively until no association signals remained in each locus. By this criterion, we identified up to four independent SNP variants (in the case of *CETP* with HDL-C in European Americans) per locus. Remarkably, only one locus (*SORT1*) harbored a single independent SNP in each ethnic group.

For many gene regions, the independent SNPs did not colocalize in the two ethnicities. For example, in the *PCSK9* locus, the most highly associated SNPs (rs11591147 in European Americans and rs11806638 in African Americans) are more than 10 kb apart and show much weaker associations (and are much less common) in the other ethnicity (Table 2). Similarly, the second independent SNPs (rs499883 in European Americans and rs505151 in African Americans) are more than 10 kb apart and are each absent in the other ethnicity. Similar patterns are seen with *APOB*, *APOA1-C3-A4-A5*, and *LCAT*. More complex patterns are seen in loci such as *LPL*, *CETP*, and *APOE*, where some

independent SNPs colocalize (within a few kb of each other) and others are distant.

Notably, for two loci/trait combinations—*LCAT* with HDL, *APOE* with TG—the strength of association of SNPs is greater in African Americans than in European Americans (Table 2). For *APOE*, this is due to the rarity of the index SNP (rs1721054) in European Americans ($\text{MAF} = 0.00047$ vs. $\text{MAF} = 0.12$), since the effect size estimates are similar in the two ethnic groups ($\beta = 0.33$ in European Americans vs. $\beta = 0.26$ in African Americans). For *LCAT*, rarity accounts for the second independent SNP in African Americans (rs35673026, absent in European Americans) but not the most highly associated SNP (rs255052, similarly common in European Americans, $\text{MAF} = 0.15$ vs. $\text{MAF} = 0.22$, with modestly different effect size estimates, $\beta = 0.069$ vs. $\beta = 0.11$). No second independent SNP for *LCAT* was identified in European Americans.

Table 2 also reports the proportion of the variance (adjusted R^2) in the respective traits explained by each SNP and the combination of the independently significant SNPs at each locus. The greatest proportion of variance explained by a single SNP in African Americans was 3.7% (rs17231520 for HDL-C); for European Americans it was 2.5% (rs17231506 for HDL-C). These SNPs are both located in the *CETP* locus on chromosome 16. Unsurprisingly, the *CETP* locus explained the greatest proportion of variance, in any trait, for both African Americans and European Americans (explaining 5.0% and 3.6% of the variance in HDL-C, respectively). For African Americans, the combination of all independently significant SNPs at the 11 loci that harbored IBC-significant SNPs for both ethnicities explained 4.5%, 7.7%, and 1.9% of the variance in LDL-C, HDL-C, and TG, respectively. For European Americans, 6.2%, 6.0%, and 4.1% of the variance in LDL-C, HDL-C, and TG were explained, respectively.

SNP×SNP Interactions and SNPs Exerting Sex-specific Effects

We investigated whether any SNP×SNP interactions existed between any of the most significant SNPs identified at each of the 19 above-mentioned loci for their respective traits; none were identified (Tables S3, S4, S5, S6, S7, S8). The lowest P value (0.002) was generated for the interaction between SNPs rs10455872 (*LPA*) and rs934197 (*APOB*) for LDL-C in African Americans, but was far from IBC-significant and withstands a Bonferroni correction of only 25 tests (there were 120 SNP×SNP interactions tested for LDL-C among African Americans).

Similarly, no IBC-significant SNP×sex interactions were observed among the SNPs identified in Table 1 (Tables S9, S10, S11). Of note, the SNP rs4810479 (*PLTP*) generated a P value of 1.51×10^{-4} for HDL-C among European Americans and was highly significant among women ($P = 7.11 \times 10^{-12}$) but not men ($P = 0.12$). The P value of this interaction was two orders of magnitude smaller than for any of the other SNPs for any lipid trait, with the exception of rs439401 ($P = 1.30 \times 10^{-3}$ for TG among European Americans). Expanding the search to all SNPs on the IBC chip failed to identify any further IBC-significant SNP×sex interactions (Tables S12, S13).

Discussion

In this work, we identify or provide further evidence for a number of genetic determinants of plasma lipid levels through population association studies of two ethnicities. Specifically, the results of these studies support each of our hypotheses.

Table 2. Independent SNP variants associated with lipid traits within single gene regions.

Gene(s) of interest in associated interval	Trait	SNP	Chr	Position	Alleles	European		African		European American Direction (EA, AA)	African American Direction (EA, AA)	African American P	European American P	European American SNP R^2	African American SNP R^2	European American Locus R^2	African American Locus R^2
						MAF	MAF	MAF	MAF								
PCSK9	LDL	rs11591147	1	55278235	T, G	0.016	0.0033	1.73 × 10 ⁻²⁸ *	0.00025	--	--	0.00025	0.007	0.005	0.011	0.003	
		rs11806638	1	55290748	A, C	0.056	0.33	0.012	4.65 × 10 ⁻¹¹ *	--	--	0.000	0.000	0.002	0.002	0.002	
		rs499883	1	55291762	A, G	0.40	-	1.45 × 10 ⁻¹¹ **	-	+0	+0	0.004	-	-	-	-	
		rs505151	1	55301775	A, G	-	0.2457	-	9.88 × 10 ⁻¹⁰ **	0-	0-	-	-	-	0.002	-	
SORT1	LDL	rs7528419	1	109618715	A, G	0.22	0.28	1.69 × 10 ⁻⁵¹ *	4.98 × 10 ⁻¹⁷	++	++	0.013	0.013	0.005	0.013	0.006	
		rs12740374	1	109619113	T, G	0.22	0.26	2.89 × 10 ⁻⁵¹	2.32 × 10 ⁻²⁰ *	--	--	0.013	0.013	0.006	0.013	0.006	
		rs5742904	2	21082665	T, C	0.00068	-	7.53 × 10 ⁻¹⁴ ***	-	+0	+0	0.004	0.004	-	0.016	0.004	
APOB	LDL	rs934197	2	21120966	A, G	0.32	0.12	2.88 × 10 ⁻³⁴ *	0.0076	++	++	0.007	0.007	0.001	0.007	0.001	
		rs562338	2	21141826	A, G	0.18	0.39	7.60 × 10 ⁻³⁴ **	1.08 × 10 ⁻⁸ *	--	--	0.008	0.008	0.004	0.008	0.004	
		rs268	8	19857809	A, G	0.020	0.0031	4.55 × 10 ⁻¹¹ **	0.040	++	++	0.003	0.003	0.000	0.003	0.000	
		rs13702	8	19868772	T, C	0.29	0.48	2.14 × 10 ⁻²⁸	1.43 × 10 ⁻¹⁰ *	--	--	0.005	0.005	0.006	0.005	0.006	
LPL	HDL	rs3916027	8	19869148	A, G	0.26	0.41	1.01 × 10 ⁻³¹ *	7.99 × 10 ⁻⁷	++	++	0.006	0.006	0.003	0.006	0.003	
		rs10503669	8	19891970	A, C	0.098	0.057	3.77 × 10 ⁻²⁸ ***	0.00084	++	++	0.005	0.005	0.001	0.005	0.001	
		rs268	8	19857809	A, G	0.020	0.0031	9.49 × 10 ⁻¹³ **	0.5053	--	--	0.003	0.003	0.000	0.003	0.000	
		rs327	8	19863816	T, G	0.27	0.41	2.72 × 10 ⁻³⁶	1.03 × 10 ⁻⁹ *	++	++	0.008	0.008	0.004	0.008	0.004	
ABCA1	HDL	rs12679834	8	19864713	T, C	0.10	0.082	1.29 × 10 ⁻³² ***	3.65 × 10 ⁻⁸	++	++	0.006	0.006	0.004	0.006	0.004	
		rs3916027	8	19869148	A, G	0.26	0.41	1.20 × 10 ⁻⁴⁰ *	2.65 × 10 ⁻⁸	--	--	0.009	0.009	0.004	0.009	0.004	
		rs2515629	9	106634185	A, G	0.17	0.17	0.53	2.98 × 10 ⁻⁷ *	++	++	0.000	0.000	0.004	0.000	0.004	
		rs2472449	9	106644018	T, G	0.34	0.22	1.40 × 10 ⁻⁶ **	9.50 × 10 ⁻⁵	++	++	0.001	0.001	0.002	0.001	0.002	
APOA1-C3-A4-A5	TG	rs1883025	9	106704122	T, C	0.26	0.34	6.25 × 10 ⁻⁷ *	0.031	--	--	0.001	0.001	0.000	0.001	0.000	
		rs180327	11	116128869	T, C	0.37	0.49	3.44 × 10 ⁻²³	1.79 × 10 ⁻⁵ **	--	--	0.008	0.008	0.000	0.008	0.000	
		rs2075290	11	116158506	T, C	0.070	0.055	8.14 × 10 ⁻⁵¹ *	0.00044	--	--	0.013	0.013	0.000	0.013	0.000	
		rs12287066	11	116167541	T, G	0.068	0.17	3.54 × 10 ⁻³⁴ **	0.031	++	++	0.010	0.010	0.000	0.010	0.000	
LIPC	HDL	rs9804646	11	116170289	T, C	0.079	0.36	0.011	4.81 × 10 ⁻⁷ *	--	--	0.001	0.001	0.003	0.001	0.003	
		rs4775041	15	56461987	C, G	0.29	0.13	2.69 × 10 ⁻²⁰ **	0.21	++	++	0.004	0.004	0.000	0.004	0.000	
		rs2070895	15	56511231	A, G	0.21	0.47	3.91 × 10 ⁻²⁴ *	1.35 × 10 ⁻¹⁰ *	++	++	0.007	0.007	0.005	0.007	0.005	
		rs17231506	16	55520209	T, C	0.32	0.14	1.50 × 10 ⁻¹²⁰ *	2.41 × 10 ⁻⁸	++	++	0.025	0.025	0.005	0.025	0.005	
CE1P	HDL	rs4783961	16	5552395	A, G	0.499	0.43	3.58 × 10 ⁻³¹	1.17 × 10 ⁻⁴³ **	++	++	0.006	0.006	0.026	0.006	0.026	
		rs17231520	16	5553328	A, G	-	0.070	-	2.46 × 10 ⁻⁵⁶ *	0+	0+	-	-	0.037	-	0.037	
		rs7499892	16	55564091	T, C	0.17	0.39	4.20 × 10 ⁻⁷⁸	1.04 × 10 ⁻²⁶ ***	--	--	0.017	0.017	0.012	0.017	0.012	
		rs5883	16	55564854	T, C	0.058	0.11	2.66 × 10 ⁻⁸ ***	1.72 × 10 ⁻¹²	++	++	0.000	0.000	0.006	0.000	0.006	
LCAT	HDL	rs11076176	16	55564947	T, G	0.17	-	8.88 × 10 ⁻⁶⁷ **	-	+0	+0	0.013	0.013	-	-	-	
		rs5880	16	55572592	C, G	0.050	0.0096	7.28 × 10 ⁻³³ ****	0.0044	--	--	0.007	0.007	0.001	0.007	0.001	
		rs35673026	16	66534352	T, C	-	0.0038	-	1.54 × 10 ⁻⁶ **	0+	0+	-	-	0.004	-	0.004	

Table 2. Cont.

Gene(s) of interest in associated interval	Trait	SNP	Chr	Position	Alleles	European American MAF	African MAF	European American P	African American P	African American Direction (EA, AA)	European American SNP R ²	African American SNP R ²	European American Locus R ²	African American Locus R ²
		rs255052	16	66582496	A, G	0.15	0.22	6.64×10 ⁻⁷	5.02×10 ⁻⁹ *	++	0.002	0.007		
		rs2107369	16	66673519	T, G	0.14	0.26	9.23×10 ⁻⁸ *	0.70	--	0.002	0.000		
LDLR	LDL	rs6511720	19	11063306	T, G	0.12	0.14	1.39×10 ⁻⁴⁹ *	2.29×10 ⁻¹⁸ *	--	0.008	0.009	0.010	0.012
		rs17242787	19	11063460	A, T	-	0.019	-	2.19×10 ⁻⁹ **	0-	-	0.003		
		rs688	19	11088602	T, C	0.43	0.10	1.66×10 ⁻¹¹ **	0.0014	++	0.002	0.003		
APOE	LDL	rs8106922	19	50093506	A, G	0.39	0.24	7.25×10 ⁻⁵	5.75×10 ⁻⁶ ***	--	0.000	0.003	0.010	0.019
		rs405509	19	50100676	T, G	0.49	0.26	1.93×10 ⁻¹¹	1.89×10 ⁻¹² **	++	0.002	0.005		
		rs769450	19	50102284	A, G	0.40	0.37	6.93×10 ⁻⁷ ***	0.00015	++	0.001	0.003		
		rs389261	19	50112183	A, G	-	0.25	-	2.72×10 ⁻¹⁴ *	0+	-	0.003		
		rs12721046	19	50113094	A, G	0.15	0.030	2.23×10 ⁻²⁹ *	0.0017	++	0.006	0.003		
		rs12721109	19	50139061	A, G	0.022	0.0028	7.72×10 ⁻²⁷ **	0.023	--	0.004	0.000		
APOE	TG	rs439401	19	50106291	T, C	0.37	0.14	1.85×10 ⁻¹⁵ *	0.88	--	0.003	0.000	0.003	0.012
		rs12721054	19	50114427	A, G	0.00047	0.12	0.60	3.73×10 ⁻²⁵ *	++	0.000	0.012		

AA = African American; EA = European American; MAF = major allele frequency. Minor allele frequencies and R² values all derived from ARIC (largest cohort for both African Americans and European Americans). Direction is modeled on the first allele listed in "Alleles" column. Independent SNPs (when they exist) are indicated as: * = lowest P value for locus + trait + ethnicity; ** = lowest P value at or near IBC significance ($P < 1 \times 10^{-6}$) after conditioning on first SNP; *** = lowest P value at or near IBC significance ($P < 1 \times 10^{-6}$) after conditioning on first two independent SNPs; etc. Locus R² values calculated from all independently significant SNPs at the corresponding locus.
doi:10.1371/journal.pone.0036473.t002

Common DNA Sequence Variants in Previously Reported Genes and Loci, as well as Additional Novel Loci, are Associated with Plasma Lipids

We found that all 19 lipid-associated loci identified in early GWAS studies replicated in the combined CARE European American cohorts (~25,000 individuals), and 10 of the loci replicated in the combined CARE African American cohorts (~9,000 individuals). It is likely that with genotyping in additional African American individuals, increased power will ultimately allow replication of many of the remaining nine loci. We identified an additional 10 loci associated with one or more lipid traits at Bonferroni-corrected statistical significance. Eight of these loci were mapped in GWAS studies subsequent to the design of the IBC array, and one locus had been identified in a prior non-GWAS study. The remaining locus, *ICAMI*, is novel. The identification of *ICAMI* is of particular interest because its association with LDL-C was exclusive to African Americans.

These findings give a high degree of confidence that most, if not all, of the reported GWAS lipid loci harbor authentic determinants of plasma lipid levels deserving of further functional investigation, and that many of the loci are relevant not just in European American populations but more generally in global populations.

At Some Lipid-associated Loci, the Specific Variants Related to Plasma Lipids Differ between Ethnic Groups

We were able to use the dense SNP genotyping in loci available via the IBC array to analyze lipid-associated loci in an unprecedented level of detail, particularly in African Americans. Our analyses demonstrate that at many loci there are major differences in genetic architecture between European Americans and African Americans. These differences manifest in at least three ways.

First, some of the most highly associated SNPs in one ethnicity are rare or absent in the other ethnicity. This is a well-established phenomenon; for example, truncation mutations in *PCSK9* that are of low frequency in African Americans, but absent in European Americans, have been shown to result in a robust reduction in LDL-C levels and coronary heart disease risk [16,17]. Many of the discrepancies in lipid associations of SNPs between the two ethnicities can be attributed primarily to differences in allele frequency. For example, rs3211938 in *CD36* is much more highly associated with HDL-C in African Americans ($P=2.60 \times 10^{-12}$) than in European Americans ($P=0.030$), with a large discrepancy in MAFs (8.3% vs. 0.01%) (Table 1). Similarly, rs12721054 in *APOE* is both more highly associated with TG and more frequent in African Americans ($P=5.55 \times 10^{-25}$, MAF=12%) than in European Americans ($P=0.60$, MAF=0.047%). These variants may represent causal variants, as is the case with the nonsense coding variant rs3211938 in *CD36* [15].

Second, some of the most highly associated SNPs in one ethnicity are poorly associated in the other ethnicity despite similar MAFs in both groups. An example is rs2515629 in the *ABCA1* locus ($P=2.04 \times 10^{-7}$, MAF=0.17, $\beta=0.11$ in African Americans; $P=0.53$, MAF=0.17, $\beta=0.0079$ in European Americans) (Table 2). In cases like this, it seems unlikely that the SNP represents a causal variant that acts exclusively in one ethnic group. Instead, rs2515629 is likely to be in strong linkage disequilibrium with an unrecognized (not genotyped on the IBC array) causal variant in African Americans; this causal variant may not be present in European Americans, or it may be present but in poor linkage disequilibrium with rs2515629 in European Americans due to differences in correlation structure.

Third, in some gene regions there are differences in the distributions of independent lipid-associated SNPs. Presumably these independent SNPs reflect the presence of independent causal variants. For example, in the *CETP* locus we identified at least 3 independently associated SNPs (with HDL) in African Americans and 4 independently associated SNPs in European Americans (Table 2), scattered over a range of 20 kb, with none clearly marking the same causal variants. Similar complexity is seen in the *PCSK9*, *LPL*, and *APOE* loci and, to a lesser degree, in most of the other 11 loci for which we performed conditional analyses (Figures S1 and S2). Thus, significant variability in the full spectrum of causal DNA variants across gene regions in different ethnic groups may well be the rule rather than the exception.

A previous analysis of lipid-associated loci in the Jackson Heart Study identified several loci—*LPL*, *APOB*, and *GCKR*—that appeared to account for some part of the inter-ethnic variation in lipid profiles, and found that SNPs in the *LPL* locus displayed different effect sizes depending on whether they existed on a background of European vs. African ancestry [18]. Our study found evidence of a similar phenomenon for a larger number of loci. In the *LPL* locus, for example, the most highly associated SNPs for TG in either European Americans and African Americans both have larger estimated effect sizes in the latter than the former (rs3916027, $\beta=-0.089$ vs. $\beta=-0.0072$; rs327, $\beta=0.097$ vs. $\beta=0.0026$), consistent with what was observed in the previous study. Other loci in which specific variants appear to have larger effect sizes in African Americans than in European Americans include *PCSK9* for LDL-C (rs11806638, $\beta=-0.11$ vs. $\beta=-0.054$), *GCKR* for TG (rs1260326, $\beta=0.088$ vs. $\beta=0.037$), *APOA1-C3-A4-A5* for TG (rs2075290, $\beta=-0.12$ vs. $\beta=-0.064$; rs9804646, $\beta=-0.083$ vs. $\beta=-0.039$), and *LCAT* for HDL-C ($\beta=0.11$ vs. $\beta=0.069$), among others. Similarly, there are a number of loci in which specific variants appear to have larger effect sizes in European Americans than in African Americans (Table 1). Thus, genetic differences in lipid determinants between the two ethnicities appear to be widespread across many loci.

Inter-ethnic Comparisons of SNP-phenotype Relationships may Pinpoint Causal DNA Variants

Finally, while it has been assumed that differing patterns of linkage disequilibrium in European Americans and African Americans could be helpful in localizing shared, causal DNA variants, for the reasons stated above this may not often be the case. Of the 29 loci we studied, only in 4 cases did association signals in the two ethnicities appear to unambiguously converge (Figures S1 and S2).

In the *SORT1* locus, the most highly associated SNPs in the two ethnicities were not identical but were in close proximity (within a few kb) and in high linkage disequilibrium, suggesting that they are marking the same causal variant (Figure S1A). In European Americans, we found six SNPs that together bear the strongest association with LDL-C (P values ranging from 1.69×10^{-51} to 1.33×10^{-49}) and are effectively in perfect linkage disequilibrium. In contrast, for the same 6 SNPs in African Americans, they are in varying degrees of linkage disequilibrium, with rs12740374 standing out as having the strongest association ($P=9.33 \times 10^{-20}$ in African Americans; $P=2.90 \times 10^{-51}$ in European Americans). Thus, this analysis nominates rs12740374 as a strong candidate for the causal variant in the locus. Of note, a recent study identified rs12740374 as a causal variant through functional experimentation in cell-based reporter assays [19], confirming the potential value of inter-ethnic comparisons of SNP-phenotype relationships in identifying causal DNA variants.

In the *LDLR* and *LIPC* loci, respectively, rs6511720 and rs2070895 were the most highly associated SNPs in both ethnicities, suggesting that each is either a causal SNP or tightly linked to a causal SNP (Figure S1B, S1C). In the *APOB* locus, the most highly LDL-C-associated SNP in European Americans, rs934197, is not associated with LDL-C in African Americans despite being common in the second ethnic group (Figure S1D). However, the second independent SNP in the locus found in European Americans, rs562338, is the most highly LDL-C-associated SNP in African Americans, suggesting that it may be a causal variant in the *APOB* locus, albeit not the only one detected in European Americans.

Rs6511720 is located in intron 1 of the *LDLR* gene, suggesting that it might affect either transcription or splicing of the gene transcript. Rs2070895 is in the promoter region of *LIPC*, just 300 bp upstream of the start codon, suggesting a role in gene transcription. Finally, rs562338 is located 20 kb upstream of the *APOB* gene, which would suggest some long-range regulatory role. Further experimentation aimed at evaluating whether rs6511720, rs2070895, and rs562338 are indeed causal DNA variants and by what mechanisms they might act in their respective loci is warranted.

Supporting Information

Figure S1 Graphical plots of SNPs in four lipid-associated loci in which the best independently associated SNPs for European Americans and African Americans coincide. The x-axis indicates the basepair position on the indicated chromosome. The y-axis indicates the negative log of the *P* value of association with the indicated lipid trait. Red stars indicate the *P* values of association of the SNPs in European Americans, black circles in African Americans. Each of the best independently associated SNPs in each locus is labeled, and all of the SNPs in strong linkage disequilibrium with one of the best SNPs are enclosed in the same oval.
(DOC)

Figure S2 Graphical plots of SNPs in each lipid-associated locus. The x-axis indicates the basepair position on the indicated chromosome. The y-axis indicates the negative log of the *P* value of association with the indicated lipid trait. Red stars indicate the *P* values of association of the SNPs in European Americans, black circles in African Americans. Each of the best independently associated SNPs in each locus is circled and labeled.
(DOC)

Table S1 CARE participant characteristics.
(DOC)

Table S2 Hardy-Weinberg *P* values for lipid-associated SNP variants.
(DOC)

Table S3 SNP×SNP interactions between the most significant SNPs at each LDL-C-related locus among European Americans.

References

1. Nambudiri KK, Kaplan EB, Heuch I, Elston RC, Green PP, et al. (1985) The Collaborative Lipid Research Clinics Family Study: biological and cultural determinants of familial resemblance for plasma lipids and lipoproteins. *Genet Epidemiol* 2: 227–254.
2. Kannel WB, Dawber TR, Kagan A, Revotskie N, Stokes J 3rd (1961) Factors of risk in the development of coronary heart disease—six year follow-up experience. The Framingham Study. *Ann Intern Med* 55: 33–50.

(DOC)

Table S4 SNP×SNP interactions between the most significant SNPs at each LDL-C-related locus among African Americans.
(DOC)

Table S5 SNP×SNP interactions between the most significant SNPs at each HDL-C-related locus among European Americans.
(DOC)

Table S6 SNP×SNP interactions between the most significant SNPs at each HDL-C-related locus among African Americans.
(DOC)

Table S7 SNP×SNP interactions between the most significant SNPs at each triglyceride-related locus among European Americans.
(DOC)

Table S8 SNP×SNP interactions between the most significant SNPs at each triglyceride-related locus among African Americans.
(DOC)

Table S9 SNP×sex interaction tests for the most significant SNPs at each LDL-C-related locus.
(DOC)

Table S10 SNP×sex interaction tests for the most significant SNPs at each HDL-C-related locus.
(DOC)

Table S11 SNP×sex interaction tests for the most significant SNPs at each triglyceride-related locus.
(DOC)

Table S12 Top ten significant SNP×sex interactions for each lipid trait among European Americans.
(DOC)

Table S13 Top ten significant SNP×sex interactions for each lipid trait among African Americans.
(DOC)

Acknowledgments

The authors wish to acknowledge the support of the National Heart, Lung, and Blood Institute (NHLBI) and the contributions of the research institutions, study investigators, and field staff.

Author Contributions

Conceived and designed the experiments: KM CMB SK DJR. Performed the experiments: KM SPRR. Analyzed the data: KM SPRR GL. Contributed reagents/materials/analysis tools: JGW KAV MYT HAT PJS JIR SSR SR BMP GJP JMO KL RMK NLG SBG MF LAC SGB EB. Wrote the paper: KM.

3. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, et al. (2010) Biological, clinical, and population relevance of 95 loci for blood lipids. *Nature* 466: 707–713.
4. Keating BJ, Tischfield S, Murray SS, Bhargava T, Price TS, et al. (2008) Concept, design and implementation of a cardiovascular gene-centric 50 k SNP array for large-scale genomic association studies. *PLoS ONE* 3: e3583.
5. Musunuru K, Lettre G, Young T, Farlow DN, Pirruccello JP, et al. (2010) Candidate Gene Association Resource (CARE): design, methods, and proof of concept. *Circ Cardiovasc Genet* 3: 267–275.

6. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, et al. (2007) PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 81: 559–575.
7. Patterson N, Hattangadi N, Lane B, Lohmueller KE, Hafler DA, et al. (2004) Methods for high-density admixture mapping of disease genes. *Am J Hum Genet* 74: 979–1000.
8. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, et al. (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* 38: 904–909.
9. Kapur NK, Musunuru K (2008) Clinical efficacy and safety of statins in managing cardiovascular risk. *Vasc Health Risk Manag* 4: 341–353.
10. Chen MH, Yang Q (2010) GWAF: an R package for genome-wide association analyses with family data. *Bioinformatics* 26: 580–581.
11. Willer CJ, Sanna S, Jackson AU, Scuteri A, Bonnycastle LL, et al. (2008) Newly identified loci that influence lipid concentrations and risk of coronary artery disease. *Nat Genet* 40: 161–169.
12. Kathiresan S, Melander O, Guiducci C, Surti A, Burt NP, et al. (2008) Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. *Nat Genet* 40: 189–197.
13. Kooner JS, Chambers JC, Aguilar-Salinas CA, Hinds DA, Hyde CL, et al. (2008) Genome-wide scan identifies variation in MLXIPL associated with plasma triglycerides. *Nat Genet* 40: 149–151.
14. Wallace C, Newhouse SJ, Braund P, Zhang F, Tobin M, et al. (2008) Genome-wide association study identifies genes for biomarkers of cardiovascular disease: serum urate and dyslipidemia. *Am J Hum Genet* 82: 139–149.
15. Love-Gregory L, Sherva R, Sun L, Wasson J, Schappe T, et al. (2008) Variants in the CD36 gene associate with the metabolic syndrome and high-density lipoprotein cholesterol. *Hum Mol Genet* 17: 1695–1704.
16. Cohen J, Pertsemlidis A, Kotowski IK, Graham R, Garcia CK, et al. (2005) Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. *Nat Genet* 37: 161–165.
17. Cohen JC, Boerwinkle E, Mosley TH Jr., Hobbs HH (2006) Sequence variations in PCSK9, low LDL, and protection against coronary heart disease. *N Engl J Med* 354: 1264–1272.
18. Deo RC, Reich D, Tandon A, Akylbekova E, Patterson N, et al. (2009) Genetic differences between the determinants of lipid profile phenotypes in African and European Americans: the Jackson Heart Study. *PLoS Genet* 5: e1000342.
19. Musunuru K, Strong A, Frank-Kamenetsky M, Lee NE, Ahfeldt T, et al. (2010) From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature* 466: 714–719.