



The histone deacetylase SIRT6, a critical modulator of metabolism and tumorigenesis

Citation

Sebastian, Carlos, Lei Zhong, Magali Silberman, Deborah Toiber, Barbara Martinez, Jean-Pierre Etchegaray, Claudia Cosentino, Sofia Giacosa, and Raul Mostoslavsky. 2012. The histone deacetylase sirt6, a critical modulator of metabolism and tumorigenesis. BMC Proceedings 6(Suppl 3): 017.

Published Version

doi:10.1186/1753-6561-6-S3-017

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10436250>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

ORAL PRESENTATION

Open Access

The histone deacetylase SIRT6, a critical modulator of metabolism and tumorigenesis

Carlos Sebastian, Lei Zhong, Magali Silberman, Deborah Toiber, Barbara Martinez, Jean-Pierre Etchegaray, Claudia Cosentino, Sofia Giacosa, Raul Mostoslavsky*

From Metabolism, diet and disease
Washington, DC, USA. 29-31 May 2012

Efficient glucose metabolism is critical for maintaining cellular viability. Under normal nutrient and oxygen conditions, glucose is converted to pyruvate, entering the mitochondria for oxidative phosphorylation and ATP production. Under hypoxia or nutrient stress, metabolism is switched to glycolysis, increasing lactate production and reducing mitochondrial respiration, a switch known to play an important role in cancer cells, as defined by Otto Warburg decades ago. Little is known whether chromatin plays a role in carbohydrate flux. The yeast Sir2 protein is an NAD-dependent histone deacetylase that senses the metabolic status of the cell and functions as a chromatin silencer to promote lifespan and genomic stability. Recently, we discovered that the mammalian SIRT6 is a chromatin factor that influences glucose metabolism and DNA repair. In mice, SIRT6-deficiency provokes a profound and lethal hypoglycemia which culminates in accelerated death. At the cellular level, SIRT6 inactivation leads to increased cellular glucose uptake, higher lactate production and decreased mitochondrial activity. Our results indicate that SIRT6 directly regulates expression of several key glycolytic genes. In this context, SIRT6 co-represses Hif1 α , acting as a histone H3 lysine9 (H3K9) deacetylase to inhibit expression of glycolytic genes (Zhong et al, 2010). Strikingly, our new studies indicate that the “glycolytic switch” observed in the absence of SIRT6 provides a unique growth advantage in the context of tumorigenesis. Indeed, we find that SIRT6 deficiency causes transformation even in the absence of oncogenic signaling, and SIRT6 levels determine progression and disease-free survival in various human cancers. Our results suggest that SIRT6 might play a critical role in modulating the Warburg effect.

Published: 1 June 2012

Reference

1. Zhong L, D'Urso A, Toiber D, Sebastian C, Henry RE, Vadysirisack DD, Guimaraes A, Marinelli B, Wikstrom JD, Nir T, Clish CB, Vaitheeswaran , Iliopoulos OB, Kurland I, Dor Y, Weissleder R, Shirihai OS, Ellisen L, Espinosa JM, Mostoslavsky R: **The histone deacetylase SIRT6 regulates glucose homeostasis via Hif1 α** . *Cell* 2010, **140**:280-293.

doi:10.1186/1753-6561-6-S3-O17

Cite this article as: Sebastian et al.: The histone deacetylase SIRT6, a critical modulator of metabolism and tumorigenesis. *BMC Proceedings* 2012 **6**(Suppl 3):O17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



The Massachusetts General Hospital Cancer Center, Harvard Medical School, Boston, MA USA