



The role of SIRT3 in regulating cancer cell metabolism

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Gonzalez Herrera, Karina N, Lydia W Finley, and Marcia C Haigis. 2012. The role of sirt3 in regulating cancer cell metabolism. BMC Proceedings 6(Suppl 3): P18.
Published Version	doi:10.1186/1753-6561-6-S3-P18
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10436254
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

POSTER PRESENTATION

Open Access

The role of SIRT3 in regulating cancer cell metabolism

Karina N Gonzalez Herrera^{1*}, Lydia W Finley³, Marcia C Haigis^{1,2}

From Metabolism, diet and disease
Washington, DC, USA. 29-31 May 2012

Background

Sirtuins are a family of NAD⁺-dependent deacetylase, deacylase, and/or mono-ADP ribosyltransferase enzymes involved in regulation of many biological processes. Mammals contain seven sirtuins, three of which are localized to the mitochondria (SIRT3-5). SIRT3 has been shown to be the major mitochondrial deacetylase that regulates metabolic enzymes and promotes oxidative metabolism and energy production. Loss of one copy of the SIRT3 gene is observed in various human cancers. Thus, we examined the role of SIRT3 in regulating metabolism in cancer cells.

Materials and methods

We have examined glucose uptake and lactate production in a variety of human cancer cell lines, as well as wild-type and SIRT3 null mouse embryonic fibroblasts (MEFs). In addition, we have utilized steady-state metabolomics to determine the metabolic profile of wild-type and SIRT3 null MEFs. Lastly, we utilized a combination of techniques, including quantitative RT-PCR and Western blotting, to examine the mechanism by which SIRT3 regulates cancer cell metabolism.

Results

Our data show that loss of SIRT3 increases glucose uptake and lactate production. Based on these and previous results from our laboratory, SIRT3 functions as a tumor suppressor to repress the Warburg effect by decreasing reactive oxygen species and destabilizing HIF1 α [1]. We are currently examining other metabolic pathways important to cancer that may be regulated by SIRT3.

Conclusions

In conclusion, we show that loss of SIRT3 results in increased glycolysis, which is the metabolic reprogramming observed in some cancer cells and is also known as the Warburg effect.

Author details

¹Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA. ²The Paul F. Glenn Labs for the Biological Mechanisms of Aging, Harvard Medical School, Boston, MA 02115, USA. ³Cancer Biology and Genetics Program, Memorial Sloan-Kettering Cancer Center, New York, NY 10065, USA.

Published: 1 June 2012

Reference

1. Finley LWS, Carracedo A, Lee J, Souza A, Egia A, Zhang J, Teryua-Feldstein J, Moreira PI, Cardoso SM, Clish CB, Pandolfi PP, Haigis MC: **SIRT3 opposes reprogramming of cancer cell metabolism through HIF1 α destabilization.** *Cancer Cell* 2011, **19**:416-428.

doi:10.1186/1753-6561-6-S3-P18

Cite this article as: Gonzalez Herrera et al.: The role of SIRT3 in regulating cancer cell metabolism. *BMC Proceedings* 2012 **6**(Suppl 3):P18.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Cell Biology, Harvard Medical School, Boston, MA 02115, USA

Full list of author information is available at the end of the article