



Effects of Genetic Variants Previously Associated with Fasting Glucose and Insulin in the Diabetes Prevention Program

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Florez, Jose C., Kathleen A. Jablonski, Jarred B. McAteer, Paul W. Franks, Clinton C. Mason, Kieren Mather, Edward Horton et al. 2012. Effects of genetic variants previously associated with fasting glucose and insulin in the Diabetes Prevention Program. PLoS ONE 7(9): e44424.
Published Version	doi:10.1371/journal.pone.0044424
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:10463938
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

Effects of Genetic Variants Previously Associated with Fasting Glucose and Insulin in the Diabetes Prevention Program

Jose C. Florez^{1,2,3*}, Kathleen A. Jablonski⁴, Jarred B. McAteer^{1,2}, Paul W. Franks^{5,6}, Clinton C. Mason⁷, Kieren Mather⁸, Edward Horton^{3,9}, Ronald Goldberg¹⁰, Dana Dabelea¹¹, Steven E. Kahn¹², Richard F. Arakaki¹³, Alan R. Shuldiner¹⁴, William C. Knowler⁷, for the Diabetes Prevention Program Research Group^{*¶}

1 Center for Human Genetic Research and Diabetes Research Center (Diabetes Unit), Massachusetts General Hospital, Boston, Massachusetts, United States of America, **2** Program in Medical and Population Genetics, Broad Institute, Cambridge, Massachusetts, United States of America, **3** Department of Medicine, Harvard Medical School, Boston, Massachusetts, United States of America, **4** The Biostatistics Center, George Washington University, Rockville, Maryland, United States of America, **5** Lund University Diabetes Center, Department of Clinical Sciences, Lund University, Malmö, Sweden, **6** Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts, United States of America, **7** Diabetes Epidemiology and Clinical Research Section, National Institute of Diabetes and Digestive and Kidney Diseases, Phoenix, Arizona, United States of America, **8** Division of Endocrinology, Indiana University School of Medicine, Indianapolis, Indiana, United States of America, **9** Joslin Diabetes Center, Boston, Massachusetts, United States of America, **10** Lipid Disorders Clinic, Division of Endocrinology, Diabetes, and Metabolism, and the Diabetes Research Institute, Leonard M. Miller School of Medicine, University of Miami, Miami, Florida, United States of America, **11** Department of Preventive Medicine and Biometrics, University of Colorado at Denver and Health Sciences Center, Denver, Colorado, United States of America, **12** Division of Metabolism, Endocrinology and Nutrition, Department of Medicine, VA Puget Sound Health Care System and University of Washington, Seattle, Washington, United States of America, **13** Department of Medicine Clinical Research, University of Hawaii, Honolulu, Hawaii, United States of America, **14** Division of Endocrinology, Diabetes and Nutrition, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland, United States of America

Abstract

Common genetic variants have been recently associated with fasting glucose and insulin levels in white populations. Whether these associations replicate in pre-diabetes is not known. We extended these findings to the Diabetes Prevention Program, a clinical trial in which participants at high risk for diabetes were randomized to placebo, lifestyle modification or metformin for diabetes prevention. We genotyped previously reported polymorphisms (or their proxies) in/near *G6PC2*, *MTNR1B*, *GCK*, *DGKB*, *GCKR*, *ADCY5*, *MADD*, *CRY2*, *ADRA2A*, *FADS1*, *PROX1*, *SLC2A2*, *GLIS3*, *C2CD4B*, *IGF1*, and *IRS1* in 3,548 Diabetes Prevention Program participants. We analyzed variants for association with baseline glycemic traits, incident diabetes and their interaction with response to metformin or lifestyle intervention. We replicated associations with fasting glucose at *MTNR1B* ($P < 0.001$), *G6PC2* ($P = 0.002$) and *GCKR* ($P = 0.001$). We noted impaired β -cell function in carriers of glucose-raising alleles at *MTNR1B* ($P < 0.001$), and an increase in the insulinogenic index for the glucose-raising allele at *G6PC2* ($P < 0.001$). The association of *MTNR1B* with fasting glucose and impaired β -cell function persisted at 1 year despite adjustment for the baseline trait, indicating a sustained deleterious effect at this locus. We also replicated the association of *MADD* with fasting proinsulin levels ($P < 0.001$). We detected no significant impact of these variants on diabetes incidence or interaction with preventive interventions. The association of several polymorphisms with quantitative glycemic traits is replicated in a cohort of high-risk persons. These variants do not have a detectable impact on diabetes incidence or response to metformin or lifestyle modification in the Diabetes Prevention Program.

Citation: Florez JC, Jablonski KA, McAteer JB, Franks PW, Mason CC, et al. (2012) Effects of Genetic Variants Previously Associated with Fasting Glucose and Insulin in the Diabetes Prevention Program. PLoS ONE 7(9): e44424. doi:10.1371/journal.pone.0044424

Editor: Giorgio Sesti, Università Magna-Graecia di Catanzaro, Italy

Received: February 1, 2012; **Accepted:** August 3, 2012; **Published:** September 11, 2012

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) of the National Institutes of Health provided funding to the clinical centers and the Coordinating Center for the design and conduct of the study, and the collection, management, analysis, and interpretation of the data, through Award Number U01DK048489. The Southwestern American Indian Centers were supported directly by the NIDDK and the Indian Health Service. The General Clinical Research Center Program, National Center for Research Resources, and the Department of Veterans Affairs supported data collection at many of the clinical centers. Funding for data collection and participant support was also provided by the Office of Research on Minority Health, the National Institute of Child Health and Human Development, the National Institute on Aging, the Office of Research on Women's Health, the Centers for Disease Control and Prevention, and the American Diabetes Association. This research was also supported, in part, by the intramural research program of the NIDDK, and by R01 DK072041 to JCF, KAJ and ARS. PWF was supported by the Swedish Research Council, Novo Nordisk, Swedish Diabetes Association, and the Swedish Heart-Lung Foundation. The Investigators gratefully acknowledge the commitment and dedication of the participants of the DPP. The opinions expressed are those of the investigators and do not necessarily reflect the official views of the funding agencies. A complete list of Centers, investigators, and staff can be found in the Appendix. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: JCF has received consulting honoraria from Novartis, Pfizer and Eli Lilly. Bristol-Myers Squibb and Parke-Davis provided medication. LifeScan Inc., Health O Meter, Hoechst Marion Roussel, Inc., Merck-Medco Managed Care, Inc., Merck and Co., Nike Sports Marketing, Slim Fast Foods Co., and Quaker Oats Co. donated materials, equipment, or medicines for concomitant conditions. McKesson BioServices Corp., Matthews Media Group, Inc., and the Henry M. Jackson Foundation provided support services under subcontract with the Coordinating Center. The authors also received funding from the commercial source Novo Nordisk. There are no patents, products in development or marketed products to declare. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials, as detailed online in the guide for authors.

* E-mail: dppmail@biostat.bsc.gwu.edu (DPPRG); jcflorez@partners.org (JCF)

¶ A list of Diabetes Prevention Program Research Group investigators is provided in the Appendix S1.

Introduction

Glucose homeostasis is tightly regulated. Control of its variation in non-diabetic individuals is influenced by familial factors, many of which are presumed to be heritable [1,2]. In searching for genetic determinants of quantitative glycemic traits, candidate gene and genome-wide association studies (GWAS) conducted in populations of European descent have identified associations of fasting glucose with genetic variants in or near the genes that encode glucokinase (*GCK*; [3]), the glucose-6-phosphatase catalytic subunit (*G6PC2*; [4,5]) and the melatonin receptor 1b (*MTNR1B*; [6,7]). The Meta-Analysis of Glucose and Insulin-related traits Consortium (MAGIC) recently performed a global meta-analysis of 21 GWAS cohorts followed by replication in 26 studies, totaling >122,000 non-diabetic individuals for fasting glucose and >98,000 non-diabetic individuals for fasting insulin [8]. These efforts confirmed the *GCK*, *G6PC2* and *MTNR1B* associations, and uncovered associations of fasting glucose with single nucleotide polymorphisms (SNPs) in or near *DGKB*, *GCKR*, *ADCY5*, *MADD*, *CRY2*, *ADRA2A*, *FADS1*, *PROX1*, *SLC2A2*, *GLIS3*, *C2CD4B* and the type 2 diabetes genes *TCF7L2* and *SLC30A8*. In addition, SNPs in or near *IGF1*, *GCKR* and perhaps *IRS1* have been found to influence fasting insulin concentrations, a surrogate for insulin resistance. Of these loci, only *GCK*, *MTNR1B*, *DGKB*, *GCKR*, *ADCY5* and *PROX1* (besides *TCF7L2* and *SLC30A8*) were associated with type 2 diabetes at genome-wide significance levels, with several others (but not all) showing a consistent trend but not meeting the same stringent statistical threshold. This work has illustrated that genetic associations with quantitative intermediate traits may lead to the discovery of type 2 diabetes loci, but also that not all genetic loci that influence fasting glucose levels in healthy individuals necessarily contribute to type 2 diabetes pathogenesis.

The MAGIC investigators have also performed more detailed characterization of the mechanisms of glucose regulation influenced by these loci in white individuals [9]. In the Third National Health and Nutrition Examination Survey (NHANES III), a genetic risk score constructed with the glucose-raising alleles was shown to have consistent effects in other ethnic groups representative of the US population [10]. The Gene \times Lifestyle interactions And Complex traits Involved in Elevated disease Risk (GLACIER) investigators showed that several of these loci associate with impaired fasting glucose (IFG) cross-sectionally and prospectively, and some have a progressively deleterious effect on fasting glucose [11]. Shortly thereafter, the Whitehall II investigators reported that a genetic risk score constructed with these variants was strongly associated with fasting glucose and remained stable over time [12]. Finally, we have recently shown that different genetic variants influence type 2 diabetes risk at distinct stages of the normoglycemia to IFG to type 2 diabetes progression, with *MTNR1B* and *GCK* exerting their effects preferentially in the normoglycemia to IFG transition [13].

To understand why some loci raise fasting glucose but do not increase type 2 diabetes risk, it is critical to establish whether their glucose-raising effects remain evident in the setting of impaired glucose tolerance (IGT), as glycemic context may modulate the strength of the genetic effect [13]. Furthermore, the impact of these loci on the prospective development of diabetes has not yet been reported. Finally, establishing whether and how distinct preventive interventions modulate these effects may facilitate the clinical translation of these findings and illuminate the specific genes and mechanisms by which these loci affect glycemic homeostasis. We concentrated on SNPs associated with fasting glucose, rather than those associated with 2-hour glucose [14],

because 1) the two 2-hour glucose SNPs that are not already captured by fasting glucose-associated variants (*GIPR* and *VPSI3C*) have no detectable impact on type 2 diabetes [15], 2) the ascertainment of DPP participants by the strict IGT definition is likely to bias the distribution of 2-hour glucose alleles, 3) longitudinal changes in 2-hour glucose among carriers of the 2-hour glucose-raising alleles have already been reported in a better suited population cohort [16], and 4) evidence obtained by the MAGIC investigators argues against an interaction of known 2-hour glucose loci with physical activity or body mass index (BMI) (Robert Scott, personal communication). We therefore genotyped the fasting glucose-associated SNPs in the multi-ethnic cohort of the Diabetes Prevention Program (DPP), and analyzed their relationships with glycemic measures at baseline and one year, the development of diabetes, and their potential interaction with preventive interventions on diabetes incidence.

Methods

The Diabetes Prevention Program

The DPP study design and baseline characteristics of the participants have been described previously [17,18]. Briefly, the DPP was designed to test whether intensive lifestyle modification or pharmacologic interventions with metformin or troglitazone prevent or delay the onset of diabetes in individuals at high risk. The trial, conducted from 1996 to 2001 in 27 US-based medical centers, included 3,234 participants randomized to intensive lifestyle modification (goal >7% weight loss and >150 min/week of physical activity), metformin (850 mg twice daily), or placebo; the fourth arm, comprising 585 additional participants randomized to troglitazone, was terminated early because of concerns with hepatotoxicity. For enrollment, participants had to have a fasting glucose between 95–125 mg/dL and IGT (2-hour glucose between 140–199 mg/dL after a 75-gram oral glucose tolerance test [OGTT]). Of the total 3,819 DPP participants, 3,548 had DNA and consented to genetic investigation: 56.4% were of European descent, 20.2% African American, 16.8% Hispanic, 4.3% Asian and 2.4% American Indian by self-report. Their mean age was 51 years and mean BMI was 34.0 kg/m². The primary endpoint (diabetes incidence, ascertained biannually and confirmed on a second occasion) was reached in nearly 38% of participants randomized to the placebo arm after a mean of 3.2 years of follow-up; there was a 58% reduction of diabetes incidence in the lifestyle intervention group and a 31% reduction in the metformin group compared to placebo [19]. For the purposes of this study, participants randomized to troglitazone were excluded, leaving a total of 2,890 individuals with valid genotypes for analysis. Institutional Review Board approval was obtained by each participating site, and all participants included in this report provided written informed consent for the main study and for subsequent genetic investigations.

Quantitative Glycemic Traits

We calculated the insulin sensitivity index (ISI) as $22.5 / [(fasting\ insulin \times fasting\ glucose) / 18.01]$; the ISI is the reciprocal of insulin resistance calculated by homeostasis model assessment (HOMA-IR) [20]. We estimated insulin secretion by the insulinogenic index using the formula $[(insulin\ at\ 30\ min) - (insulin\ at\ 0\ min)] / [(glucose\ at\ 30\ min) - (glucose\ at\ 0\ min)]$. The oral disposition index (DIo) was calculated as $1 / (fasting\ insulin \times insulinogenic\ index)$ [21]. We studied genetic associations with these measures at baseline and at 1 year: we chose one year because changes in weight were most

Table 1. SNPs genotyped and their allele frequencies by ethnic group.

SNP	Chromosome	Position (NCBI 36)	Nearest gene	Alleles (effect/other)	Allele frequencies (%)				
					White (n = 1,617)	African-American (n = 592)	Hispanic (n = 475)	Asian (n = 125)	American Indian (n = 81)
Fasting glucose									
rs340874	1	184833918	<i>PROX1*</i>	C/T	55.9	19.8	41.2	42.3	35.4
rs573225	2	161653734	<i>G6PC2</i>	A/G	71.7	91.8	85.4	90.3	92.0
rs11708067	3	120438894	<i>ADCY5*</i>	A/G	79.4	85.9	77.3	91.2	70.0
rs11920090	3	168087406	<i>SLC2A2</i>	T/A	87.1	67.3	86.6	91.1	94.4
rs2191349	7	14947780	<i>DGKB*</i>	T/G	55.7	57.9	48.1	64.0	24.1
rs917793	7	44131132	<i>GCK*</i>	T/A	19.5	23.7	32.3	21.6	48.1
rs7034200	9	4244098	<i>GLIS3</i>	A/C	48.9	64.2	57.3	46.0	65.6
rs10885122	10	106670840	<i>ADRA2A</i>	G/T	88.0	35.6	84.3	86.8	88.9
rs11605924	11	45579933	<i>CRY2</i>	A/C	49.3	87.0	47.7	68.0	51.9
rs7944584	11	47035421	<i>MADD</i>	A/T	71.7	95.0	83.7	90.4	98.1
rs174550	11	57899714	<i>FADS1</i>	T/C	68.0	91.4	43.5	55.2	11.1
rs10830963	11	88799685	<i>MTNR1B*</i>	G/C	28.8	9.1	22.7	41.2	24.1
rs11071657	15	39256547	<i>C2CD4B</i>	A/G	64.4	86.9	53.7	70.0	37.0
Fasting insulin									
rs4675095	2	219495543	<i>IRS1</i>	A/T	93.3	98.5	84.8	85.6	69.1
rs855228	12	99957291	<i>IGF1</i>	T/C	84.3	40.9	76.1	65.4	79.0
Fasting glucose and insulin									
rs780094	2	27483120	<i>GCKR*</i>	C/T	59.6	81.7	62.2	66.8	88.9

*Loci previously associated with type 2 diabetes at genome-wide levels of statistical significance. The allele previously associated with higher levels of the trait (effect allele) is shown first; allele frequencies correspond to the effect allele. Gene names: *PROX1*, prospero homeobox 1; *G6PC2*, glucose-6-phosphatase, catalytic, 2; *ADCY5*, adenylate cyclase 5; *SLC2A2*, solute carrier family 2, member 2; *DGKB*, diacylglycerol kinase, beta 90 kDa; *GCK*, glucokinase; *GLIS3*, GLIS family zinc finger 3; *ADRA2A*, adrenergic, alpha-2A-, receptor; *CRY2*, cryptochrome 2; *MADD*, MAP-kinase activating death domain; *FADS1*, fatty acid desaturase 1; *MTNR1B*, melatonin receptor 1B; *C2CD4B*, C2 calcium-dependent domain containing 4B; *IRS1*, insulin receptor substrate 1; *IGF1*, insulin-like growth factor 1; *GCKR*, glucokinase regulator. doi:10.1371/journal.pone.0044424.t001

pronounced at that time point, and it contained the highest number of participants with available measures.

SNP Selection and Genotyping

We genotyped the index SNPs associated with fasting glycemic traits reported by the MAGIC investigators [8]. Where assay design failed we selected proxies based on linkage disequilibrium in the HapMap CEU population: rs573225 for rs560887 in *G6PC2*, $r^2 = 0.961$; rs917793 for rs4607517 in *GCK*, $r^2 = 1.0$; and rs855228 for rs35767 in *IGF1*, $r^2 = 0.915$. DNA was extracted from peripheral blood leukocytes and quantitated as previously described [22]. Genotyping was carried out by allele-specific primer extension of multiplex amplified products and detection using matrix-assisted laser desorption ionization time-of-flight mass spectrometry on a Sequenom iPLEX platform [23]. Genotyping success rate was $\geq 98.5\%$. Because results for the two previously known type 2 diabetes genes *TCF7L2* and *SLC30A8* have been reported elsewhere [22,24,25], they are not presented here.

Statistical Analyses

We used Cox proportional hazards regression models with genotype, intervention and their interactions as the independent variables predicting time to diabetes over mean 3.2 years follow-up. We adjusted for gender, age at enrollment, ethnicity, treatment

arm, and baseline BMI. For the quantitative glycemic traits, we employed generalized mixed models to test additive effects of genotype on baseline log-transformed quantitative traits, and on the same traits after one year of intervention adjusted for the baseline value, age, sex, self-reported ethnicity, BMI and treatment arm. We note that these SNPs have been associated with glycemic traits at genome-wide levels of significance, and therefore their prior probability of true effects is many orders of magnitude higher than the genome average. As our analyses represent further characterization of each of these established loci, we selected a *P* value threshold of 0.05. Finally, we also tested for any evidence of epistatic interactions between the *MTNR1B* SNP rs10830963 and the *G6PC2* SNP rs573225, both of which have significant effects on fasting glucose in the DPP, by including appropriate interaction terms at baseline and one year.

Results

Baseline Associations

The SNPs genotyped, their chromosomal location, the nearest gene and their allele frequencies in the five DPP ethnic groups are shown in Table 1. Allele frequencies were comparable to those previously reported by MAGIC in Europeans [8] and NHANES III in non-Hispanic whites, African Americans and US Hispanics [10].

Table 2. Nominal genotypic associations with quantitative traits at baseline.

SNP	Nearest gene	Alleles (effect/other)	Trait	LS Means (95% CI)	Additive P value	Pairwise P values
rs573225	G6PC2	A/G	FG (mg/dL)	AA 106.7 (106.2–107.2)	0.002	AA vs AG 0.002
				AG 105.6 (104.9–106.3)		AA vs GG 0.31
				GG 105.8 (104.5–107.1)		AG vs GG 0.73
			Fins (μU/mL)	AA 24.44 (23.65–25.26)	0.006	AA vs AG 0.31
				AG 24.97 (23.87–26.11)		AA vs GG 0.005
				GG 27.71 (25.56–30.05)		AG vs GG 0.02
			Ins Index	AA 1.25 (1.20–1.31)	0.002	AA vs AG 0.16
				AG 1.20 (1.13–1.28)		AA vs GG 0.003
				GG 1.04 (0.92–1.17)		AG vs GG 0.03
			ISI	AA 0.155 (0.15–0.161)	0.03	AA vs AG 0.62
				AG 0.154 (0.147–0.161)		AA vs GG 0.01
				GG 0.138 (0.127–0.15)		AG vs GG 0.03
Dlo	AA 0.049 (0.047–0.051)	<0.001	AA vs AG 0.03			
	AG 0.046 (0.043–0.049)		AA vs GG <0.001			
	GG 0.037 (0.033–0.042)		AG vs GG <0.001			
rs11708067	ADCY5	A/G	Fins (μU/mL)	AA 24.05 (23.24–24.88)	0.001	AA vs AG 0.001
				AG 25.85 (24.79–26.95)		AA vs GG 0.53
				GG 25.29 (23.12–27.67)		AG vs GG 0.64
			ISI	AA 0.158 (0.153–0.164)	0.004	AA vs AG 0.003
				AG 0.148 (0.141–0.154)		AA vs GG 0.72
				GG 0.151 (0.138–0.166)		AG vs GG 0.72
rs11920090	SLC2A2	T/A	Dlo	AA 0.042 (0.037–0.049)	0.006	AA vs AT 0.27
				AT 0.046 (0.043–0.049)		AA vs TT 0.04
				TT 0.049 (0.047–0.051)		AT vs TT 0.03
rs7944584	MADD	A/T	Proins (pmol/L)	AA 16.4 (15.9–16.92)	<0.001	AA vs AT <0.001
				AT 14.98 (14.34–15.65)		AA vs TT <0.001
				TT 13.53 (12.46–14.68)		AT vs TT 0.01
rs174550	FADS1	T/C	Fins (μU/mL)	TT 23.78 (22.83–24.78)	0.008	TT vs CT 0.06
				CT 24.97 (23.96–26.03)		TT vs CC 0.06
				CC 25.52 (24.25–26.86)		CT vs CC 0.47
			ISI	CC 0.149 (0.141–0.157)	0.01	TT vs CT 0.09
				CT 0.153 (0.146–0.159)		TT vs CC 0.09
				TT 0.160 (0.153–0.167)		CT vs CC 0.46
rs10830963	MTNR1B	G/C	FG (mg/dL)	GG 108.7 (107.6–109.9)	<0.001	GG vs CG 0.02
				CG 107.3 (106.7–108.0)		GG vs CC <0.001
				CC 105.6 (105.1–106.2)		CG vs CC <0.001
			Proins (pmol/L)	GG 15.88 (14.80–17.03)	0.009	GG vs CG 0.66
				CG 15.43 (14.85–16.04)		GG vs CC 0.66
				CC 16.44 (15.90–17.00)		CG vs CC 0.003
			Ins Index	GG 1.17 (1.05–1.29)	0.01	GG vs CG 0.74
				CG 1.19 (1.12–1.25)		GG vs CC 0.21
				CC 1.27 (1.21–1.33)		CG vs CC 0.05
rs855228	IGF1	T/C	FG (mg/dL)	TT 106.1 (105.5–106.7)	0.01	TT vs CT 0.37
				CT 106.4 (105.8–107.0)		TT vs CC 0.02
				CC 107.7 (106.6–108.7)		CT vs CC 0.43
rs780094	GCKR	C/T	FG (mg/dL)	CC 106.8 (106.2–107.3)	0.001	CC vs CT 0.12
				CT 106.3 (105.6–106.9)		CC vs TT 0.003
				TT 105.2 (104.3–106.1)		CT vs TT 0.04

FG, fasting glucose; Fins, fasting insulin; Ins Index, insulinogenic index; ISI, insulin sensitivity index; Dlo, oral disposition index; Proins, fasting proinsulin adjusted for fasting insulin. To convert glucose mg/dL to mmol/L, divide by 18.01. To convert insulin μU/ml to pmol/L to, multiply by 6.0.
doi:10.1371/journal.pone.0044424.t002

Table 3. Associations with quantitative traits at one year.

SNP	Nearest gene	Alleles (effect/other)	FG		Fins		Proins		Ins Index		ISI		Dlo	
			<i>P</i> int	<i>P</i> assoc	<i>P</i> int	<i>P</i> assoc	<i>P</i> int	<i>P</i> assoc	<i>P</i> int	<i>P</i> assoc	<i>P</i> int	<i>P</i> assoc	<i>P</i> int	<i>P</i> assoc
rs340874	<i>PROX1</i>	C/T	0.81	0.47	0.90	0.15	0.99	0.08	0.42	0.99	0.89	0.14	0.63	0.22
rs573225	<i>G6PC2</i>	A/G	0.96	0.17	0.08	0.88	0.34	0.91	0.77	0.20	0.11	0.66	0.81	0.08
rs11708067	<i>ADCY5</i>	A/G	0.80	0.27	0.22	0.98	0.31	0.85	0.46	0.32	0.20	0.80	0.86	0.52
rs11920090	<i>SLC2A2</i>	T/A	0.88	0.49	0.24	0.53	0.79	0.38	0.59	0.98	0.25	0.49	0.69	0.89
rs2191349	<i>DGKB</i>	T/G	0.79	0.41	0.04	–	0.09	0.80	0.50	0.55	0.07	0.84	0.84	0.99
rs917793	<i>GCK</i>	T/A	0.07	0.12	0.39	0.12	0.08	0.31	0.86	0.23	0.24	0.08	0.42	0.07
rs7034200	<i>GLIS3</i>	A/C	0.72	0.98	0.02	–	0.25	0.56	0.94	0.94	0.03	–	0.13	0.64
rs10885122	<i>ADRA2A</i>	G/T	0.82	0.14	0.20	0.60	0.16	0.46	0.29	0.18	0.25	0.44	0.31	0.03
rs11605924	<i>CRY2</i>	A/C	0.20	0.56	0.13	0.46	0.76	0.80	0.40	0.28	0.13	0.40	0.21	0.58
rs7944584	<i>MADD</i>	A/T	0.04	–	0.73	0.80	0.63	0.30	0.11	0.10	0.63	0.83	0.36	0.29
rs174550	<i>FADS1</i>	T/C	0.58	0.20	0.87	0.09	0.73	0.23	0.76	0.97	0.86	0.07	0.64	0.65
rs10830963	<i>MTNR1B</i>	G/C	0.68	0.003	0.17	0.37	0.27	0.41	0.69	0.002	0.16	0.90	0.96	0.08
rs11071657	<i>C2CD4B</i>	A/G	0.04	–	0.96	0.42	0.55	0.97	0.38	0.09	0.98	0.54	0.35	0.048
rs4675095	<i>IRS1</i>	A/T	0.21	0.99	0.67	0.26	0.68	0.59	0.53	0.55	0.71	0.30	0.62	0.62
rs855228	<i>IGF1</i>	T/C	0.39	0.60	0.15	0.27	0.52	0.87	0.36	0.72	0.15	0.24	0.63	0.75
rs780094	<i>GCKR</i>	C/T	0.57	0.76	0.07	0.22	0.25	0.38	0.17	0.92	0.09	0.26	0.43	0.28

FG, fasting glucose; Fins, fasting insulin; Ins Index, insulinogenic index; ISI, insulin sensitivity index; Dlo, oral disposition index; Proins, fasting proinsulin adjusted for fasting insulin. *P* int denotes the *P* value for the genotype × intervention interaction test; *P* assoc denotes the *P* value for the main effect association in the full cohort when *P* int >0.05.

doi:10.1371/journal.pone.0044424.t003

Table 4. Levels of quantitative glycemic traits at one year by genotype and treatment arm at loci with a nominally significant interaction.

SNP	gene	Alleles (effect/other)	Trait	Placebo		Metformin		Lifestyle	
				LS Means (95% CI)	<i>P</i> values	LS Means (95% CI)	<i>P</i> values	LS Means (95% CI)	<i>P</i> values
rs2191349		T/G	Fins	GG 24.71 (22.96–26.59)	GG/GT: 0.99	GG 22.62 (21.02–24.34)	GG/GT: 0.21	GG 18.43 (17.03–19.94)	GG/GT: 0.99
	<i>DGKB</i>		(μU/mL)	GT 24.99 (23.59–26.48)	GG/TT: 0.99	GT 21.30 (20.06–22.62)	GG/TT: 0.04	GT 19.01 (17.82–20.28)	GG/TT: 0.99
				TT 25.71 (24.00–27.54)	GT/TT: 0.99	TT 20.41 (19.03–21.88)	GT/TT: 0.21	TT 19.05 (17.71–20.51)	GT/TT: 0.99
rs7034200		A/C	Fins	AA 24.86 (23.27–26.55)	AA/AC: 0.99	AA 22.54 (21.06–24.12)	AA/AC: 0.12	AA 18.01 (16.76–19.36)	AA/AC: 0.28
	<i>GLIS3</i>		(μU/mL)	AC 25.23 (23.83–26.73)	AA/CC: 0.99	AC 21.14 (19.87–22.50)	AA/CC: 0.05	AC 19.20 (18.02–20.45)	AA/CC: 0.28
				CC 25.07 (23.18–27.12)	AC/CC: 0.99	CC 20.45 (19.00–22.01)	AC/CC: 0.37	CC 19.47 (17.96–21.11)	AC/CC: 0.74
			ISI	AA 0.153 (0.142–0.164)	AA/AC: 0.99	AA 0.175 (0.163–0.189)	AA/AC: 0.14	AA 0.222 (0.205–0.240)	AA/AC: 0.28
				AC 0.152 (0.143–0.162)	AA/CC: 0.99	AC 0.187 (0.175–0.200)	AA/CC: 0.06	AC 0.207 (0.193–0.221)	AA/CC: 0.28
				CC 0.151 (0.139–0.165)	AC/CC: 0.99	CC 0.194 (0.179–0.210)	AC/CC: 0.36	CC 0.204 (0.186–0.222)	AC/CC: 0.74
rs7944584		A/T	FG	AA 106.8 (105.5–108.1)	AA/AT: 0.008	AA 102.4 (101.3–103.5)	AA/AT: 0.99	AA 102.1 (101.0–103.2)	AA/AT: 0.99
	<i>MADD</i>		(mg/dL)	AT 104.3 (102.6–106.1)	AA/TT: 0.50	AT 102.7 (101.1–104.2)	AA/TT: 0.99	AT 101.5 (99.98–103.1)	AA/TT: 0.99
				TT 104.8 (101.4–108.3)	AT/TT: 0.78	TT 101.3 (98.45–104.3)	AT/TT: 0.99	TT 101.4 (98.65–104.2)	AT/TT: 0.99
rs11071657		A/G	FG	AA 107.1 (105.6–108.7)	AA/AG: 0.41	AA 102.3 (101.0–103.6)	AA/AG: 0.99	AA 101.9 (100.6–103.2)	AA/AG: 0.96
	<i>C2CD4B</i>		(mg/dL)	AG 105.9 (104.5–107.4)	AA/GG: 0.41	AG 102.7 (101.4–104.0)	AA/GG: 0.99	AG 101.7 (100.4–103.0)	AA/GG: 0.96
				GG 105.3 (103.1–107.6)	AG/GG: 0.63	GG 102.0 (100.2–103.9)	AG/GG: 0.99	GG 102.7 (100.8–104.7)	AG/GG: 0.96

P values for pairwise comparisons between genotypic groups are shown, with groups separated by a "/". Fins, fasting insulin (μU/mL); ISI, insulin sensitivity index; FG, fasting glucose (mg/dL). To convert glucose mg/dL to mmol/L, divide by 18.01. To convert insulin μU/ml to pmol/L to, multiply by 6.0.

doi:10.1371/journal.pone.0044424.t004




Figure 1. Effect of genotype at *MTNR1B* rs10830963 on glycemic traits at baseline and one year. Fasting glucose is shown in panel (a) and the insulinogenic index is shown in panel (b). Because no significant SNP \times intervention interaction was found, the full cohort was analyzed in aggregate. Fasting glucose is higher ($P=0.003$) and the insulinogenic index is lower ($P=0.002$) in carriers of the G risk allele after one year, even after adjustment for the corresponding baseline levels. Least-square means ($\pm 95\%$ CI) are shown. To convert glucose mg/dL to mmol/L, divide by 18.01. doi:10.1371/journal.pone.0044424.g001

We tested associations of these SNPs with baseline fasting glucose, fasting insulin, fasting proinsulin adjusted for fasting insulin, the insulinogenic index, the ISI and the DIo in this multiethnic cohort of individuals with IGT. We replicated associations with fasting glucose at *G6PC2* ($P=0.002$), *MTNR1B* ($P<0.001$) and *GCKR* ($P=0.001$). We also replicated associations of the glucose-raising allele with *reduced* insulino-

genic index at *MTNR1B* and *increased* insulinogenic and disposition indices at *G6PC2*. We again noted a strong association of *MADD* with fasting proinsulin levels, adjusted for concomitant insulin ($P<0.001$). All nominally significant ($P<0.05$) associations and corresponding trait distributions are shown in Table 2.

Table 5. Diabetes incidence by genotype at each locus, in the overall cohort and stratified by treatment arm.

SNP	Nearest gene	Alleles	SNP * Tx	Treatment adjusted HR (95% CI)	P-value	PLACEBO HR (95% CI)	P-value	METFORMIN HR (95% CI)	P-value	LIFESTYLE HR (95% CI)	P-value
rs340874	<i>PROX1*</i>	C (vs T)	N	0.88 (0.78–0.98)	0.02	0.85 (0.71–1.01)	0.06	0.92 (0.75–1.12)	0.39	0.86 (0.68–1.08)	0.20
rs573225	<i>G6PC2</i>	A (vs G)	N	1.11 (0.97–1.27)	0.14	0.96 (0.77–1.19)	0.70	1.18 (0.94–1.47)	0.15	1.27 (0.98–1.64)	0.07
rs11708067	<i>ADCY5*</i>	A (vs G)	N	1.06 (0.92–1.23)	0.38	1.04 (0.84–1.28)	0.73	1.08 (0.84–1.35)	0.60	1.10 (0.82–1.47)	0.51
rs11920090	<i>SLC2A2</i>	T (vs A)	N	1.02 (0.88–1.19)	0.75	1.08 (0.86–1.33)	0.56	1.00 (0.77–1.30)	0.99	0.99 (0.71–1.37)	0.93
rs2191349	<i>DGKB*</i>	T (vs G)	N	1.06 (0.94–1.18)	0.34	1.05 (0.88–1.27)	0.56	1.10 (0.90–1.33)	0.34	1.01 (0.80–1.27)	0.96
rs917793	<i>GCK*</i>	T (vs A)	N	0.96 (0.84–1.10)	0.59	0.87 (0.70–1.07)	0.20	1.14 (0.90–1.44)	0.29	0.92 (0.69–1.22)	0.56
rs7034200	<i>GLIS3</i>	A (vs C)	N	1.00 (0.89–1.12)	1.00	0.90 (0.75–1.08)	0.22	1.04 (0.85–1.27)	0.68	1.15 (0.91–1.47)	0.25
rs10885122	<i>ADRA2A</i>	G (vs T)	N	1.03 (0.91–1.16)	0.63	1.01 (0.84–1.22)	0.89	1.03 (0.84–1.28)	0.76	1.08 (0.84–1.39)	0.56
rs11605924	<i>CRY2</i>	A (vs C)	N	1.01 (0.90–1.12)	0.91	0.93 (0.78–1.10)	0.40	1.06 (0.88–1.28)	0.56	1.09 (0.87–1.37)	0.47
rs7944584	<i>MADD</i>	A (vs T)	N	0.93 (0.80–1.08)	0.29	0.86 (0.69–1.08)	0.20	0.89 (0.69–1.14)	0.35	1.11 (0.84–1.47)	0.47
rs174550	<i>FADS1</i>	T (vs C)	N	0.94 (0.83–1.05)	0.26	0.95 (0.80–1.14)	0.57	0.98 (0.80–1.19)	0.81	0.86 (0.67–1.09)	0.21
rs10830963	<i>MTNR1B*</i>	G (vs C)	N	1.07 (0.94–1.22)	0.29	1.20 (0.98–1.47)	0.07	1.01 (0.80–1.26)	0.95	0.95 (0.73–1.24)	0.69
rs11071657	<i>C2CD4B</i>	A (vs G)	N	0.93 (0.83–1.05)	0.26	0.93 (0.77–1.12)	0.43	0.92 (0.75–1.14)	0.42	0.96 (0.75–1.22)	0.72
rs4675095	<i>IRS1</i>	A (vs T)	N	0.96 (0.78–1.18)	0.68	0.93 (0.67–1.30)	0.69	1.16 (0.84–1.59)	0.37	0.71 (0.44–1.15)	0.17
rs855228	<i>IGF1</i>	T (vs C)	N	1.09 (0.97–1.23)	0.14	1.04 (0.87–1.25)	0.66	1.16 (0.95–1.43)	0.15	1.12 (0.88–1.43)	0.38
rs780094	<i>GCKR*</i>	C (vs T)	N	0.96 (0.85–1.08)	0.48	0.91 (0.75–1.10)	0.33	1.01 (0.82–1.23)	0.93	0.96 (0.75–1.22)	0.72

*Loci previously associated with type 2 diabetes. Effect allele denotes the allele associated with higher glucose or insulin levels in MAGIC. There are no significant SNP \times treatment interactions. One nominally significant P value for association with diabetes incidence is not consistent with the expected direction of effect. doi:10.1371/journal.pone.0044424.t005

Associations at One Year

We tested whether the metformin or lifestyle preventive interventions interacted with each SNP to modulate quantitative glycemic traits at one year. We adjusted one-year traits for the corresponding baseline trait, to indicate change in each variable during active treatment. Where no nominally significant interaction with treatment was found, SNP main effects on the one-year trait were tested in the whole cohort with an adjustment for treatment arm; if an interaction was detected at $P < 0.05$, analyses were stratified by treatment arm (Table 3). Nominally significant interactions were found for *DGKB* and fasting insulin, *GLIS3* and both fasting insulin and ISI, and both *MADD* and *C2CD4B* and fasting glucose. Least-square means for each genotype group and the corresponding pairwise comparisons are shown in Table 4.

At *MTNR1B*, the glucose-raising allele continued to have a significant main effect on raising fasting glucose and lowering the insulinogenic index at one year (Figure 1). Because one-year traits are adjusted for the baseline level, this effect is indicative of a worsening deleterious effect of this locus on β -cell function. We further explored the concordant effects of SNPs at *MTNR1B* and *G6PC2* on fasting glucose but discordant effects for insulinogenic index by testing for epistatic interactions between the two on fasting glucose at baseline and one year: the interaction terms were not statistically significant.

Diabetes Incidence

We tested whether the metformin or lifestyle preventive interventions interact with each SNP on the risk of developing diabetes during 3.2 years of mean follow-up. As no nominal interactions were found, the effects of each SNP on diabetes incidence were evaluated in the full cohort while adjusting for treatment arm; stratified analyses are also shown (Table 5). The only nominal association with diabetes incidence was found for the glucose-lowering allele at *PROX1* ($P = 0.02$), in a direction opposite to that reported in case-control analyses in MAGIC, where the C allele increased type 2 diabetes risk (odds ratio 1.07 [95% CI 1.05–1.09], $P = 7.2 \times 10^{-10}$) [8].

Discussion

The MAGIC investigators reported a number of loci that influence fasting glucose and fasting insulin levels in non-diabetic populations of European descent; only a few of the loci were also associated with type 2 diabetes at genome-wide levels of significance [8]. The authors speculated that it is not the mere elevation in fasting glucose, but how fasting glucose is raised, that determines overall β -cell dysfunction and future type 2 diabetes risk. However, whether these loci exert their action on fasting glucose in the initial stages of diabetes progression (e.g. from normoglycemia to impaired glucose regulation) or later (e.g. from IGT to type 2 diabetes) is not known. In the GLACIER cohort, eleven loci (including the known type 2 diabetes genes *TCF7L2* and *SLC30A8*) were nominally associated with IFG cross-sectionally, and *MTNR1B* and *G6PC2* were also associated with development of IFG in longitudinal analyses [11]. We have recently shown that among type 2 diabetes-associated loci, risk alleles at *MTNR1B*, *GCK* and *SLC30A8* confer a stronger rate of progression from normoglycemia to IFG than from IFG to type 2 diabetes [13]. Here we extend these findings by testing these SNPs from the IGT to type 2 diabetes transition, and by assessing their effects on quantitative

glycemic traits at baseline and one year in a multiethnic cohort of persons with IGT.

We have demonstrated that the three loci with the strongest reported effect on fasting glucose (*MTNR1B*, *GCKR* and *G6PC2*) have consistent effects in the DPP. All three were known to be associated with fasting glucose prior to the MAGIC GWAS meta-analysis [4,5,6,7,26,27,28]. Power may have been limiting to detect the other reported associations [24].

We have also confirmed that the glucose-raising allele at *MTNR1B* is associated with a reduced insulinogenic index, as measured during the initial phase of insulin secretion during an OGTT [9,29]. As shown by Lyssenko and coworkers, the deleterious effects of this allele on β -cell function persist over time; while they noted such worsening over 24 years of follow-up [29], here we see such effects over a much shorter time span (one year). In GLACIER a similar non-significant trend was noted over 10 years of follow-up [11], although a consistent effect was not detected in the Whitehall II study [12]. Because *MTNR1B* does increase risk of type 2 diabetes [8], this pattern of sustained deterioration suggests that identifying these individuals early in their glycemic progression may be beneficial in prevention efforts.

In contrast, the glucose-raising allele at *G6PC2* is associated with superior β -cell function on dynamic testing; this has been shown previously [9,30], and is consistent with the role of this gene product in regulating hepatic glucokinase and its null effect on type 2 diabetes risk [8]. We found no evidence in support of a non-additive interaction between *MTNR1B* and *G6PC2* on fasting glucose at baseline or one year. The strong effect of the *MADD* locus on fasting proinsulin levels is also confirmed [9,31]; because this association is adjusted for concomitant insulin levels, it reflects an increased secretion of insulin precursors out of proportion to the degree of basal insulin resistance. The other nominal associations newly reported here do not withstand correction for the multiple statistical tests performed, and should be considered hypothesis-generating requiring confirmation in independent studies.

In summary, the strongest effects of genetic loci on fasting glucose in non-diabetic individuals of European descent are also evident in a multiethnic cohort with IGT. The deleterious influence of the glucose-raising allele at *MTNR1B* on β -cell function appears to worsen with time, and this effect is evident in as short a time as one year. Genetic testing may identify a subset of patients with IGT more likely to respond to preventive interventions [32].

Supporting Information

Appendix S1 DPP Research Group.
(DOC)

Acknowledgments

The Investigators gratefully acknowledge the commitment and dedication of the participants of the DPP. We thank the MAGIC investigators for access to pre-publication data.

Author Contributions

Conceived and designed the experiments: JCF KAJ PWF ARS WCK. Performed the experiments: JCF JBM. Analyzed the data: KAJ JCF CCM WCK. Contributed reagents/materials/analysis tools: KM EH RG SEK RFA WCK. Wrote the paper: JCF. Reviewed and edited the manuscript: KM EH RG SEK RFA WCK JCF KAJ PWF ARS JBM CCM DD.

References

- Meigs JB, Panhuysen CIM, Myers RH, Wilson PWF, Cupples LA (2002) A genome-wide scan for loci linked to plasma levels of glucose and HbA1c in a community-based sample of Caucasian pedigrees: the Framingham Offspring Study. *Diabetes* 51: 833–840.
- Panhuysen CIM, Cupples LA, Wilson PWF, Herbert AG, Myers RH, et al. (2003) A genome scan for loci linked to quantitative insulin traits in persons without diabetes: the Framingham Offspring Study. *Diabetologia* 46: 579–587.
- Weedon MN, Clark VJ, Qjan Y, Ben-Shlomo Y, Timpson N, et al. (2006) A common haplotype of the glucokinase gene alters fasting glucose and birth weight: association in six studies and population-genetics analyses. *Am J Hum Genet* 79: 991–1001.
- Bouatia-Naji N, Rocheleau G, Van Lommel L, Lemaire K, Schuit F, et al. (2008) A polymorphism within the *G6PC2* gene is associated with fasting plasma glucose levels. *Science* 320: 1085–1088.
- Chen W-M, Erdos MR, Jackson AU, Saxena R, Sanna S, et al. (2008) Association studies in Caucasians identify variants in the *G6PC2/ABCB11* region regulating fasting glucose levels. *J Clin Invest* 118: 2620–2628.
- Prokopenko I, Langenberg C, Florez JC, Saxena R, Soranzo N, et al. (2009) Variants in *MTNR1B* influence fasting glucose levels. *Nat Genet* 41: 77–81.
- Bouatia-Naji N, Bonnafant A, Cavalcanti-Proenca C, Sparso T, Holmkvist J, et al. (2009) A variant near *MTNR1B* is associated with increased fasting plasma glucose levels and type 2 diabetes risk. *Nat Genet* 41: 89–94.
- Dupuis J, Langenberg C, Prokopenko I, Saxena R, Soranzo N, et al. (2010) New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. *Nat Genet* 42: 105–116.
- Ingelsson E, Langenberg C, Hivert MF, Prokopenko I, Lyssenko V, et al. (2010) Detailed physiologic characterization reveals diverse mechanisms for novel genetic Loci regulating glucose and insulin metabolism in humans. *Diabetes* 59: 1266–1275.
- Yang Q, Liu T, Shrader P, Yesupriya A, Chang MH, et al. (2010) Racial/ethnic differences in association of fasting glucose-associated genomic loci with fasting glucose, HOMA-B, and impaired fasting glucose in the U.S. adult population. *Diabetes Care* 33: 2370–2377.
- Renstrom F, Shungin D, Johansson I, Florez JC, Hallmans G, et al. (2011) Genetic predisposition to long-term nondiabetic deteriorations in glucose homeostasis: Ten-year follow-up of the GLACIER study. *Diabetes* 60: 345–354.
- Jensen AC, Barker A, Kumari M, Brunner EJ, Kivimaki M, et al. (2011) Associations of common genetic variants with age-related changes in fasting and postload glucose: Evidence from 18 years of follow-up of the Whitehall II cohort. *Diabetes* 60: 1617–1623.
- Walford GA, Green T, Neale B, Isakova T, Rotter JL, et al. (2011) Common genetic variants differentially influence the transition between clinically-defined states of fasting glucose metabolism. *Diabetologia* (epub ahead of print).
- Saxena R, Hivert MF, Langenberg C, Tanaka T, Pankow JS, et al. (2010) Genetic variation in *GIPR* influences the glucose and insulin responses to an oral glucose challenge. *Nat Genet* 42: 142–148.
- Voight BF, Scott LJ, Steinthorsdottir V, Morris AP, Dina C, et al. (2010) Twelve type 2 diabetes susceptibility loci identified through large-scale association analysis. *Nat Genet* 42: 579–589.
- Jensen AC, Barker A, Kumari M, Brunner EJ, Kivimaki M, et al. (2011) Associations of common genetic variants with age-related changes in fasting and postload glucose: evidence from 18 years of follow-up of the Whitehall II cohort. *Diabetes* 60: 1617–1623.
- The Diabetes Prevention Program Research Group (1999) The Diabetes Prevention Program. Design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care* 22: 623–634.
- The Diabetes Prevention Program Research Group (2000) The Diabetes Prevention Program: baseline characteristics of the randomized cohort. *The Diabetes Prevention Program Research Group. Diabetes Care* 23: 1619–1629.
- The Diabetes Prevention Program Research Group (2002) Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346: 393–403.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, et al. (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28: 412–419.
- Utzschneider KM, Prigeon RL, Faulenbach MV, Tong J, Carr DB, et al. (2009) Oral disposition index predicts the development of future diabetes above and beyond fasting and 2-h glucose levels. *Diabetes Care* 32: 335–341.
- Florez JC, Jablonski KA, Bayley N, Pollin TI, de Bakker PIW, et al. (2006) *TCF7L2* polymorphisms and progression to diabetes in the Diabetes Prevention Program. *N Engl J Med* 355: 241–250.
- Tang K, Fu DJ, Julien D, Braun A, Cantor CR, et al. (1999) Chip-based genotyping by mass spectrometry. *Proc Natl Acad Sci U S A* 96: 10016–10020.
- Moore AF, Jablonski KA, McAteer JB, Saxena R, Pollin TI, et al. (2008) Extension of type 2 diabetes genome-wide association scan results in the Diabetes Prevention Program. *Diabetes* 57: 2503–2510.
- Majithia AR, Jablonski KA, McAteer JB, Mather KJ, Goldberg RB, et al. (2011) Association of the *SLC30A8* missense polymorphism R325W with proinsulin levels at baseline and after lifestyle, metformin or troglitazone intervention in the Diabetes Prevention Program. *Diabetologia* 54: 2570–2574.
- Diabetes Genetics Initiative of Broad Institute of Harvard and MIT, Lund University and Novartis Institutes for BioMedical Research (2007) Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* 316: 1331–1336.
- Sparso T, Andersen G, Nielsen T, Burgdorf KS, Gjesing AP, et al. (2008) The *GCKR* rs780094 polymorphism is associated with elevated fasting serum triacylglycerol, reduced fasting and OGTT-related insulinaemia, and reduced risk of type 2 diabetes. *Diabetologia* 51: 70–75.
- Orho-Melander M, Melander O, Guiducci C, Perez-Martinez P, Corella D, et al. (2008) Common missense variant in the glucokinase regulatory protein gene is associated with increased plasma triglyceride and C-reactive protein but lower fasting glucose concentrations. *Diabetes* 57: 3112–3121.
- Lyssenko V, Nagorny CL, Erdos MR, Wierup N, Jonsson A, et al. (2009) Common variant in *MTNR1B* associated with increased risk of type 2 diabetes and impaired early insulin secretion. *Nat Genet* 41: 82–88.
- Rose CS, Grarup N, Krarup NT, Poulsen P, Wegner L, et al. (2009) A variant in the *G6PC2/ABCB11* locus is associated with increased fasting plasma glucose, increased basal hepatic glucose production and increased insulin release after oral and intravenous glucose loads. *Diabetologia* 52: 2122–2129.
- Strawbridge RJ, Dupuis J, Prokopenko I, Barker A, Ahlqvist E, et al. (2011) Genome-wide association identifies nine common variants associated with fasting proinsulin levels and provides new insights into the pathophysiology of type 2 diabetes. *Diabetes* 60: 2624–2634.
- Hivert MF, Jablonski KA, Perreault L, Saxena R, McAteer JB, et al. (2011) Updated genetic score based on 34 confirmed type 2 diabetes loci is associated with diabetes incidence and regression to normoglycemia in the Diabetes Prevention Program. *Diabetes* 60: 1340–1348.