



Response to Comment on: Sun et al. Protection From Retinopathy and Other Complications in Patients With Type 1 Diabetes of Extreme Duration: The Joslin 50-Year Medalist Study. *Diabetes Care* 2011;34:968–974

Citation

Sun, Jennifer K., Hillary A. Keenan, Vincent M. Monnier, Lloyd Paul Aiello, and George L. King. 2011. Response to comment on: Sun et al. Protection from retinopathy and other complications in patients with type 1 diabetes of extreme duration: the Joslin 50-year medalist study. *Diabetes Care* 2011;34:968–974. *Diabetes Care* 34(9): e149.

Published Version

doi:10.2337/dc11-1188

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10482551>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

 COMMENTS AND
 RESPONSES

**Response to
 Comment on: Sun
 et al. Protection From
 Retinopathy and
 Other Complications
 in Patients With
 Type 1 Diabetes of
 Extreme Duration:
 The Joslin 50-Year
 Medalist Study.
 Diabetes Care
 2011;34:968-974**

We thank Conway et al. (1) for the discussion of their findings regarding skin intrinsic fluorescence in the Pittsburgh Epidemiology of Diabetes Complications (EDC) study cohort. Their results—that skin intrinsic fluorescence values are higher in individuals with no diabetic retinopathy (DR) as compared with more advanced DR—are consistent with our findings in the Medalist cohort in which increased concentrations of certain advanced glycation end

products (carboxymethyl-lysine [CML] and the early glycation product fructose-lysine in the Medalists) were associated with protection from proliferative DR. Our findings of an inverse relationship between CML levels and proliferative retinopathy may suggest low levels of endothelial or inflammatory cell receptor for advanced glycation end product expression or activation (2) in response to its ligand CML in these participants.

In contrast to the Medalist group in which no difference in glycemic control was found between those with and without proliferative DR, the EDC participants without DR had better glycemic control than those with DR. This difference may reflect the fact that the EDC population had a shorter average duration of diabetes than the Medalists (41 vs. 57 years) and thus may have a lower percentage of participants with elevated protective factors against complications. Nonetheless, this report further confirms the existence of a subgroup of individuals who appear to be highly protected against the detrimental effects of chronic hyperglycemia and supports additional characterization of this unique population in hopes of identifying novel protective factors against retinopathy and other microvascular complications of diabetes.

JENNIFER K. SUN, MD, MPH^{1,2,3}
 HILLARY A. KEENAN, PHD^{3,4}

VINCENT M. MONNIER, MD⁵
 LLOYD PAUL AIELLO, MD, PHD^{1,2,3}
 GEORGE L. KING, MD^{3,4}

From the ¹Beetham Eye Institute, Joslin Diabetes Center, Boston, Massachusetts; the ²Department of Ophthalmology, Harvard Medical School, Boston, Massachusetts; the ³Research Division, Joslin Diabetes Center, Boston, Massachusetts; the ⁴Department of Internal Medicine, Harvard Medical School, Boston, Massachusetts; and the ⁵Departments of Pathology and Biochemistry, Case Western Reserve University, Cleveland, Ohio.

Corresponding author: George L. King, george.king@joslin.harvard.edu.

DOI: 10.2337/dc11-1188

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

.....

References

1. Conway BN, Maynard JD, Orchard TJ. Comment on: Sun et al. Protection from retinopathy and other complications in patients with type 1 diabetes of extreme duration: the Joslin 50-Year Medalist Study. *Diabetes Care* 2011; 34:968–974 (Letter). *Diabetes Care* 2011;34:e148. DOI: 10.2337/dc11-0971
2. Pachydaki SI, Tari SR, Lee SE, et al. Upregulation of RAGE and its ligands in proliferative retinal disease. *Exp Eye Res* 2006;82:807–815