



Who is at risk for alcohol-related negative consequences?

Citation

Pedrelli, Paola, Charlotte Brill, and Fava Maurizio. 2012. Who is at risk for alcohol-related negative consequences? *Addiction Science & Clinical Practice* 7(Suppl 1): A42.

Published Version

doi:10.1186/1940-0640-7-S1-A42

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10578929>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

MEETING ABSTRACT

Open Access

Who is at risk for alcohol-related negative consequences?

Paola Pedrelli^{1*}, Charlotte Brill², Fava Maurizio¹

From International Network on Brief Interventions for Alcohol Problems (INEBRIA) Meeting 2011 Boston, MA, USA. 21-23 September 2011

Although a large body of research has documented that severe consequences are associated with alcohol use among college students, this may not be the case for all students who consume alcohol. Identifying those variables associated with risk for negative consequences in this population has important public health implications. Mixed findings are available on the relationship between alcohol-related negative consequences and alcohol consumed, depressive symptoms, and use of illicit substances. We examined the relationship between alcohol-related consequences and amount consumed during a typical and a heavy drinking week, cannabis use in the previous 12 months, and depressive symptoms. Participants consisted of a convenience sample of 85 (53.5% females) undergraduate college students (mean age = 19 years) taking part in an on-campus mental-health screening. Participants reported use of alcohol and of other drugs, mental health symptoms, and negative consequences. Results showed a direct association between alcohol-related negative consequences and total drinks consumed during a typical week ($r = 0.41$; $p < 0.001$) and during a heavy drinking week ($r = 0.58$; $p < 0.001$). Students with any cannabis use in the previous 12 months reported a significant greater number of alcohol-related negative consequences ($t [1, 80] = -2.66$; $p < 0.01$). Alcohol-related negative consequences were not associated with depressive symptoms. In a hierarchical linear regression model including cannabis use, total drinks during a typical week, and total drinks during a heavy drinking week, the latter was the only predictor of alcohol-related negative consequences. The other two variables did not predict any variance over and above the one explained by total drinks during a heavy drinking week. In conclusion, programs aimed at reducing the negative outcomes of alcohol

consumption should aim at screening for heavy rather than typical drinking.

Author details

¹Department of Psychology, Massachusetts General Hospital/Harvard Medical School, Boston, MA, USA. ²Depression Clinical & Research Program, Massachusetts General Hospital, Boston, MA, USA.

Published: 9 October 2012

doi:10.1186/1940-0640-7-S1-A42

Cite this article as: Pedrelli et al.: Who is at risk for alcohol-related negative consequences? *Addiction Science & Clinical Practice* 2012 **7**(Suppl 1):A42.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Psychology, Massachusetts General Hospital/Harvard Medical School, Boston, MA, USA

Full list of author information is available at the end of the article