



The structure of Polycomb-repressed chromatin in the bithorax complex

Citation

Bowman, Sarah K, Aimee M Deaton, Heber Domingues, Welcome Bender, and Robert E Kingston. 2013. The structure of Polycomb-repressed chromatin in the bithorax complex. *Epigenetics & Chromatin* 6(Suppl 1): 022.

Published Version

doi:10.1186/1756-8935-6-S1-022

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10609673>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

ORAL PRESENTATION

Open Access

The structure of Polycomb-repressed chromatin in the bithorax complex

Sarah K Bowman^{1*}, Aimee M Deaton¹, Heber Domingues², Welcome Bender², Robert E Kingston¹

From *Epigenetics and Chromatin: Interactions and processes*
Boston, MA, USA. 11-13 March 2013

The 300 kb of the *Drosophila* bithorax complex (BX-C) is the original model for studying Hox gene repression by Polycomb group proteins. Decades of genetic experiments have led to the hypothesis that Polycomb organizes BX-C chromatin differently in each of the abdominal parasegments, and the structure of the chromatin is critical for maintaining cell identity. Studying the molecular organization of chromatin in individual parasegments has been technically difficult. We solved this problem by developing a sorted nuclei ChIP-seq pipeline. In this system, transgenic embryos produce tagged nuclei in single parasegments. Using FACS, we sorted these tagged nuclei and performed small-scale ChIP-seq. Initial results show that a histone mark catalyzed by Polycomb group proteins, trimethylated histone H3 at lysine 27 (H3K27me3), covers less of the BX-C as we move from PS5 to PS7. The boundaries of the H3K27 methylation correlate precisely with previously identified CTCF binding sites, a protein known to act as a chromatin barrier. Correspondingly, a mark that correlates with gene activation, H3K4me3, appears over the transcription start sites of BX-C genes that are not associated with Polycomb histone methyltransferase activity. Going forward, we will assay the localization of Polycomb group proteins and many other chromatin regulatory proteins to gain a complete picture of the chromatin environment that regulates Hox gene expression in the developing embryo. Still, the current results offer a compelling first look at BX-C chromatin *in vivo*, and provide molecular support to a long-standing genetic hypothesis.

Author details

¹Department of Molecular Biology, Massachusetts General Hospital, and Department of Genetics, Harvard Medical School, Boston, MA, USA.

²Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA, USA.
Full list of author information is available at the end of the article

²Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA, USA.

Published: 18 March 2013

doi:10.1186/1756-8935-6-S1-O22

Cite this article as: Bowman et al.: The structure of Polycomb-repressed chromatin in the bithorax complex. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):O22.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

