



Genome-wide analysis reveals TET-and TDG-mediated 5-methylcytosine oxidation dynamics

Citation

Shen, Li, Hao Wu, Dinh Diep, Ana C D'Alessio, Alan Fung, Kun Zhang, and Yi Zhang. 2013. Genome-wide analysis reveals tet-and tdg-mediated 5-methylcytosine oxidation dynamics. *Epigenetics & Chromatin* 6(Suppl 1): P88.

Published Version

doi:10.1186/1756-8935-6-S1-P88

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10610380>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

POSTER PRESENTATION

Open Access

Genome-wide analysis reveals TET-and TDG-mediated 5-methylcytosine oxidation dynamics

Li Shen^{1,2,3†}, Hao Wu^{1,2,3,5*†}, Dinh Diep⁶, Ana C D'Alessio^{1,2,3}, Alan Fung⁶, Kun Zhang⁶, Yi Zhang^{1,2,3,4}

From *Epigenetics and Chromatin: Interactions and processes*
Boston, MA, USA. 11-13 March 2013

Ten-eleven translocation (Tet) family of DNA dioxygenases converts 5-methylcytosine (5mC) into 5-hydroxymethylcytosine (5hmC), 5-formylcytosine (5fC), and 5-carboxylcytosine (5caC) through iterative oxidation reactions. While 5mC and 5hmC are relatively abundant, 5fC and 5caC are at very low levels in the mammalian genome. Thymine DNA glycosylase (TDG) and base excision repair (BER) pathways can actively remove 5fC/5caC to regenerate unmethylated cytosine, but it is unclear to what extent and at which part of the genome such active demethylation processes take place. Here, we have performed high-throughput sequencing analysis of 5mC/5hmC/5fC/5caC-enriched DNA using modification-specific antibodies and generated genome-wide distribution maps of these cytosine modifications in wild-type and *Tdg*-deficient mouse embryonic stem cells (ESCs). We observe that the steady state 5fC and 5caC are preferentially detected at repetitive sequences in wild-type mouse ESCs. Depletion of TDG causes marked accumulation of 5fC and 5caC at a large number of distal gene regulatory elements and transcriptionally repressed/poised gene promoters, suggesting that Tet/TDG-dependent dynamic cycling of 5mC oxidation states may be involved in regulating the function of these regions. Thus, comprehensive mapping of 5mC oxidation and BER pathway activity in the mammalian genome provides a promising approach for better understanding of biological roles of DNA methylation and demethylation dynamics in development and diseases.

Author details

¹Howard Hughes Medical Institute, Harvard Medical School, WAB- 149G, 200 Longwood Ave., Boston, MA 02115, USA. ²Program in Cellular and Molecular Medicine, Boston Children's Hospital, Harvard Medical School, WAB- 149G, 200 Longwood Ave., Boston, MA 02115, USA. ³Department of Genetics, Harvard Medical School, WAB- 149G, 200 Longwood Ave., Boston, MA 02115, USA. ⁴Harvard Stem Cell Institute, Harvard Medical School, WAB-149G, 200 Longwood Ave., Boston, MA 02115, USA. ⁵Department of Stem Cell and Regenerative Biology, Harvard University, 7 Divinity Street, Cambridge, MA 02138, USA. ⁶Departments of Bioengineering, University of California at San Diego, La Jolla, California, USA.

Published: 18 March 2013

doi:10.1186/1756-8935-6-S1-P88

Cite this article as: Shen et al.: Genome-wide analysis reveals TET-and TDG-mediated 5-methylcytosine oxidation dynamics. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):P88.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



† Contributed equally

¹Howard Hughes Medical Institute, Harvard Medical School, WAB- 149G, 200 Longwood Ave., Boston, MA 02115, USA

Full list of author information is available at the end of the article