



Health Risks from Lead-Based Ammunition in the Environment

The Harvard community has made this
article openly available. [Please share](#) how
this access benefits you. Your story matters

Citation	Bellinger, D. C., J. Burger, T. J. Cade, D. A. Cory-Slechta, M. Finkelstein, H. Hu, M. Kosnett, et al. 2013. "Health Risks from Lead-Based Ammunition in the Environment." <i>Environmental Health Perspectives</i> 121 (6): a178-a179. doi:10.1289/ehp.1306945. http://dx.doi.org/10.1289/ehp.1306945 .
Published Version	doi:10.1289/ehp.1306945
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:11708572
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

Health Risks from Lead-Based Ammunition in the Environment

<http://dx.doi.org/10.1289/ehp.1306945>

Lead is one of the most studied toxicants, and overwhelming scientific evidence demonstrates that lead is toxic to several physiological systems in vertebrates, including the nervous, renal, cardiovascular, reproductive, immune, and hematologic systems (Health Risks from Lead-Based Ammunition in the Environment—A Consensus Statement of Scientists 2013). Furthermore, there is no level of lead exposure in children known to be without adverse effects [Centers for Disease Control and Prevention (CDC) 2012a, 2012b].

In light of this evidence, there is an urgent need to end a major source of lead for animals and humans: spent lead bullets and shotgun pellets. Notably, production of lead-based ammunition in the United States accounted for > 69,000 metric tons consumed in 2012; this is second only to the amount of lead used to manufacture storage batteries (U.S. Geological Survey 2013). However, there are few regulations regarding the release of lead into the environment through discharge of lead-based ammunition. For other major categories of lead consumption, such as lead batteries and sheet lead/lead pipes, environmental discharge and disposal are regulated. Therefore, lead-based ammunition is likely the greatest largely unregulated source of lead that is knowingly discharged into the environment in the United States. In contrast, the release or distribution of other major sources of environmental lead contamination (e.g., leaded gasoline, lead-based paint, lead solder) have been substantially regulated and reduced since the mid-1970s (Health Risks from Lead-Based Ammunition in the Environment—A Consensus Statement of Scientists 2013).

There is a national discussion—polarized at times—of the health risks posed to humans and wildlife from the discharge of lead-based ammunition. To inform this discussion, a group of 30 nationally and internationally recognized scientists with expertise regarding lead and environmental health recently collaborated to create an evidence-based consensus statement (Health Risks from Lead-Based Ammunition in the Environment—A Consensus Statement of Scientists 2013) supporting the reduction and eventual elimination of lead released to the environment through the discharge of lead-based ammunition.

The discharge of lead bullets and shotgun pellets into the environment poses significant health risks to humans and wildlife. The best available scientific evidence demonstrates that the discharge of lead-based ammunition substantially increases environmental lead levels, especially in areas with higher shooting activity (U.S. Environmental Protection Agency 2012) and that the discharge of lead-based ammunition poses risks of elevated lead exposure to gun users (National Research Council 2012). When lead-containing bullets are used to shoot wildlife, they can fragment into hundreds of small pieces, many of which are small enough to be easily ingested by scavenging animals or to be retained in meat prepared for human consumption (Hunt et al. 2009; Knott et al. 2010; Pauli and Burkirk 2007). Consequently, lead-based ammunition may be a significant source of lead exposure in humans that regularly ingest wild game (Hanning et al. 2003; Johansen et al. 2006; Levesque et al. 2003; Tsuji et al. 2008). In addition, lead pellets and fragments have been reported in gastrointestinal tracts of hunters who consume meat from animals shot with lead-based ammunition (Carey 1977; Reddy 1985).

The use of lead pellets in shotgun shells for hunting waterfowl posed a serious threat to wetland birds, and secondarily to bald eagles, in the United States, leading to the U.S. Fish and Wildlife Service's 1991 nationwide regulations requiring use of nontoxic shotgun pellets for hunting waterfowl (Anderson 1992). However, lead poisoning from ingestion of spent lead-based ammunition fragments continues

to pose a particularly serious health threat for scavenging species. These lead-containing fragments remain the principal source of lead exposure to endangered California condors and continue to prevent the successful recovery of these birds in the wild (Church et al. 2006; Finkelstein et al. 2012; Green et al. 2008; Parish et al. 2009; Rideout et al. 2012; Woods et al. 2007). Other wildlife species, such as golden eagles, bald eagles, ravens, turkey vultures, and pumas, are also exposed to the fragments of spent lead ammunition (Burco et al. 2012; Clark and Scheuhammer 2003; Craighead and Bedrosian 2008; Cruz-Martinez et al. 2012; Fisher et al. 2006; Kelly and Johnson 2011; Stauber et al. 2010; Wayland and Bollinger 1999).

No rational deliberation about the use of lead-based ammunition can ignore the overwhelming evidence for the toxic effects of lead, or that the discharge of lead bullets and shot into the environment poses significant risks of lead exposure to humans and wildlife. Given the availability of non-lead ammunition for shooting and hunting (Thomas 2013), the use of lead-based ammunition that introduces lead into the environment can be reduced and eventually eliminated. This seems to be a reasonable and equitable action to protect the health of humans and wildlife.

The authors declare they have no actual or potential competing financial interests.

David C. Bellinger

Harvard Medical School
Harvard School of Public Health
Boston, Massachusetts

Joanna Burger

Environmental and Occupational Health Sciences Institute
Rutgers University
Piscataway, New Jersey

Tom J. Cade

Department of Ecology and Evolutionary Biology (Emeritus)
Cornell University
Ithaca, New York

Deborah A. Cory-Slechta

Department of Environmental Medicine
University of Rochester School of Medicine
Rochester, New York

Myra Finkelstein

Department of Microbiology and Environmental Toxicology
University of California, Santa Cruz
Santa Cruz, California

Howard Hu

Dalla Lana School of Public Health
University of Toronto
Toronto, Ontario, Canada

Michael Kosnett

Division of Clinical Pharmacology & Toxicology
Department of Medicine
University of Colorado School of Medicine
Aurora, Colorado

Philip J. Landrigan

Department of Preventive Medicine
Icahn School of Medicine at Mount Sinai
New York, New York

Bruce Lanphear

Child & Family Research Institute
BC Children's Hospital
Simon Fraser University
Vancouver, British Columbia, Canada

Mark A. Pokras

Wildlife Clinic
Department of Infectious Disease and Global Health
Cummings School of Veterinary Medicine
Tufts University
North Grafton, Massachusetts

Patrick T. Redig

The Raptor Center
Veterinary Clinical Sciences
University of Minnesota
St. Paul, Minnesota

Bruce A. Rideout

Wildlife Disease Laboratories
Institute for Conservation Research
San Diego Zoo Global
San Diego, California

Ellen Silbergeld

Department of Environmental Health Sciences
Johns Hopkins Bloomberg School of Public Health
Baltimore, Maryland

Robert Wright

Department of Preventive Medicine
Icahn School of Medicine at Mount Sinai
New York, New York

Donald R. Smith

Department of Microbiology and Environmental Toxicology
University of California, Santa Cruz
Santa Cruz, California
E-mail: drsmith@ucsc.edu

REFERENCES

- Anderson WL. 1992. Legislation and lawsuits in the United States and their effects on nontoxic shot regulations. In: *Lead Poisoning in Waterfowl* (Pain DJ, ed). IWRB Special Publication 16. Slimbridge, UK:International Waterfowl and Wetlands Research Bureau, 56–60.
- Burco J, Myers AM, Schuler K, Gillin C. 2012. Acute lead toxicosis via ingestion of spent ammunition in a free-ranging cougar (*Puma concolor*). *J Wildl Dis* 48(1):216–219.
- Carey LS. 1977. Lead shot appendicitis in northern native people. *J Can Assoc Radiol* 28:171–174.
- CDC (Centers for Disease Control and Prevention). 2012a. CDC Response to Advisory Committee on Childhood Lead Poisoning Prevention Recommendations in “Low Level Lead Exposure Harms Children: A Renewed Call of Primary Prevention.” Available: http://www.cdc.gov/nceh/lead/ACCLPP/CDC_Response_Lead_Exposure_Recs.pdf [accessed 9 May 2013].
- CDC (Centers for Disease Control and Prevention). 2012b. Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention. Report of the Advisory Committee on Childhood Lead Poisoning Prevention of the Centers for Disease Control and Prevention. Available: http://www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf [accessed 9 May 2013].
- Church ME, Gwiazda R, Risebrough RW, Sorenson K, Chamberlain CP, Farry S, et al. 2006. Ammunition is the principal source of lead accumulated by California condors re-introduced to the wild. *Environ Sci Technol* 40:6143–6150.
- Clark AJ, Scheuhammer AM. 2003. Lead poisoning in upland-foraging birds of prey in Canada. *Ecotoxicology* 12:23–30.
- Craighead D, Bedrosian B. 2008. Blood lead levels of common ravens with access to big-game offal. *J Wildl Manage* 72(1):240–245.
- Cruz-Martinez L, Redig PT, Deen J. 2012. Lead from spent ammunition: a source of exposure and poisoning in bald eagles. *Hum Wildl Interact* 6(1):94–104.
- Finkelstein ME, Doak DF, George D, Burnett J, Brandt J, Church M, et al. 2012. Lead poisoning and the deceptive recovery of the critically endangered California condor. *Proc Natl Acad Sci USA* 109(28):11449–11454.
- Fisher IJ, Pain DJ, Thomas VG. 2006. A review of lead poisoning from ammunition sources in terrestrial birds. *Biol Conser* 131(3):421–432.
- Green RE, Hunt WG, Parish CN, Newton I. 2008. Effectiveness of action to reduce exposure of free-ranging California condors in Arizona and Utah to lead from spent ammunition. *PLoS ONE* 3(12):e4022; doi:10.1371/journal.pone.0004022 [Online 24 December 2008].
- Hanning RM, Sandhu R, MacMillan A, Moss L, Tsuji LJS, Nieboer E. 2003. Impact of blood lead levels of maternal and early infant feeding practices of First Nation Cree in the Mushkegowuk Territory of northern Ontario, Canada. *J Environ Monit* 5:241–245.
- Health Risks from Lead-Based Ammunition in the Environment—A Consensus Statement of Scientists. 2013. Available: <http://escholarship.org/uc/item/6dq3h64x> [accessed 9 May 2013].
- Hunt WG, Watson RT, Oaks JL, Parish CN, Burnham KK, Tucker RL, et al. 2009. Lead bullet fragments in venison from rifle-killed deer: potential for human dietary exposure. *PLoS ONE* 4(4):e5330; doi:10.1371/journal.pone.0005330 [Online 24 April 2009].
- Johansen P, Pedersen HS, Asmund G, Riget F. 2006. Lead shot from hunting as a source of lead in human blood. *Environ Pollut* 142:93–97.
- Kelly TR, Johnson CK. 2011. Lead exposure in free-flying turkey vultures is associated with big game hunting in California. *PLoS ONE* 6(4):e15350; doi:10.1371/journal.pone.0015350 [Online 6 April 2011].
- Knott J, Gilbert J, Hoccom D, Green R. 2010. Implications for wildlife and humans of dietary exposure to lead from fragments of lead rifle bullets in deer shot in the UK. *Sci Total Environ* 409:95–99.
- Levesque B, Duchesne JF, Gariépy C, Rhainds M, Dumas P, Scheuhammer AM, et al. 2003. Monitoring of umbilical cord blood lead levels and sources assessment among the Inuit. *Occup Environ Med* 60:693–695.
- National Research Council. 2012. Potential Health Risks to DOD Firing-Range Personnel from Recurrent Lead Exposure. Washington, DC:National Academies Press.
- Parish CN, Hunt WG, Feltes E, Sieg R, Orr K. 2009. Lead exposure among a reintroduced population of California condors in northern Arizona and southern Utah. In: *Ingestion of Lead from Spent Ammunition: Implications for Wildlife and Humans* (Watson RT, Fuller M, Pokras M, Hunt WG, eds). Boise, ID:The Peregrine Fund, 259–264. Available: <http://www.peregrinefund.org/subsites/conference-lead/PDF/0217%20Parish.pdf> [accessed 9 May 2013].
- Pauli J, Burkirk S. 2007. Recreational shooting of prairie dogs: a portal for lead entering wildlife food chains. *J Wildl Manage* 71(1):103–108.
- Reddy ER. 1985. Retained lead shot in the appendix. *J Can Assoc Radiol* 36:47–48.
- Rideout BA, Stalis I, Papendick R, Pessier A, Puschner B, Finkelstein ME, et al. 2012. Patterns of mortality in free-ranging California condors (*Gymnogyps californianus*). *J Wildl Dis* 48(1):95–112.
- Stauber E, Finch N, Talcott PA, Gay JM. 2010. Lead poisoning of bald (*Haliaeetus leucocephalus*) and golden (*Aquila chrysaetos*) eagles in the U.S. inland Pacific northwest region—an 18-year retrospective study: 1991–2008. *J Avian Med Surg* 24(4):279–287.
- Thomas VG. 2013. Lead-free hunting rifle ammunition: product availability, price, effectiveness, and role in global wildlife conservation. *Ambio* doi:10.1007/s13280-012-0361-7 [Online 4 January 2013].
- Tsuji LJS, Wainman BC, Martin ID, Sutherland C, Weberd JP, Dumas P, et al. 2008. The identification of lead ammunition as a source of lead exposure in First Nations: the use of lead isotope ratios. *Sci Total Environ* 393:291–298.
- U.S. Environmental Protection Agency. 2012. Integrated Science Assessment for Lead (Third External Review Draft). EPA/600/R-10/075C. Washington, DC:U.S. Environmental Protection Agency. Available: <http://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=242655> [accessed 9 May 2013].
- U.S. Geological Survey. 2013. Mineral Industry Surveys: Lead in January 2013. Available: <http://minerals.usgs.gov/minerals/pubs/commodity/lead/mis-201301-lead.pdf> [accessed 9 May 2013].
- Wayland M, Bollinger T. 1999. Lead exposure and poisoning in bald eagles and golden eagles in the Canadian Prairie Provinces. *Environ Poll* 104(3):341–350.
- Woods CP, Heinrich WR, Farry SC, Parish CN, Osborn SAH, Cade TJ. 2007. Survival and reproduction of California condors released in Arizona. In: *California Condors in the 21st Century* (Mee A, Hall LS, eds). Series in Ornithology No. 2. Washington, DC, and Cambridge, MA:American Ornithologists Union and Nuttall Ornithological Club, 57–78. Available: <http://www.peregrinefund.org/docs/pdf/research-library/2007/2007-Woods-condor-survival.pdf> [accessed 9 May 2013].