



PW03-010 - MHC complexity in Behçet's disease

The Harvard community has made this
article openly available. [Please share](#) how
this access benefits you. Your story matters

Citation	Ombrello, MJ, Y Kirino, P de Bakker, F Cosan, DL Kastner, A Gul, and EF Remmers. 2013. "PW03-010 - MHC complexity in Behçet's disease." <i>Pediatric Rheumatology Online Journal</i> 11 (Suppl 1): A236. doi:10.1186/1546-0096-11-S1-A236. http://dx.doi.org/10.1186/1546-0096-11-S1-A236 .
Published Version	doi:10.1186/1546-0096-11-S1-A236
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:12064458
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA



MEETING ABSTRACT

Open Access

PW03-010 - MHC complexity in Behçet's disease

MJ Ombrello^{1*}, Y Kirino², P de Bakker^{3,4}, F Cosan⁵, DL Kastner⁶, A Gul⁵, EF Remmers⁶

From 7th Congress of International Society of Systemic Auto-Inflammatory Diseases (ISSAID)
Lausanne, Switzerland. 22-26 May 2013

Introduction

Family studies support a genetic contribution to Behçet's disease (BD), with a sibling recurrence-risk ratio of 11-52. The class I MHC molecule, *HLA-B*51* (*B*51*), is the strongest known genetic risk factor for BD, however the gene immediately centromeric to *HLA-B*, *MICA*, has also been implicated in BD. Because of strong linkage disequilibrium (LD) between *HLA-B* and *MICA*, their respective contributions to BD susceptibility have been debated. A recent report has proposed that *B*51* is not a BD susceptibility allele, and several studies have identified *B*51*-independent association signals within the MHC.

Objectives

To clarify the relationship between *B*51* and BD, and to test for *B*51*-independent genetic variation within the MHC that influences BD susceptibility.

Methods

Using Illumina Human 370CNV SNP genotypes in a Turkish collection of 1244 BD patients and 1303 geographically-matched healthy subjects, we examined SNP haplotypes and LD patterns across the *HLA-B/MICA* region with Haploview. We performed SNP imputation of the MHC using IMPUTE2 and the 1000 Genomes Phase 1 dataset. We inferred classical HLA types and their amino acids using SNP2HLA. Association testing and regression analyses were performed using SNPTEST and SNP & Variation Suite 7.

Results

We identified a *B*51*(+) *HLA-B/MICA* haplotype that was strongly associated with BD ($p=1.22E-46$, OR 2.8). A *B*51*(-) version of the same haplotype occurred at equal frequencies in cases and controls, demonstrating that *B*51* is essential to the risk haplotype. Further, we found that rs2848713, a variant on the *MICA* end of the

haplotype, conferred additional risk of BD in *B*51*(+) individuals. Through imputation, we generated a set of 32,689 imputed SNPs. The 2 most strongly associated SNPs were 4.8Kb centromeric of *HLA-B* ($p_{\min}=1.4E-50$), but no SNP was more strongly associated with BD than was *B*51* itself ($p=1.3E-55$). Conditioning on *B*51* revealed an association near *HLA-A* ($p_{\min}=5.4E-9$), and upon adding a representative *HLA-A* SNP to the regression model, we detected residual association centromeric of *HLA-B* ($p=1.5E-5$). Analysis of imputed HLA types supported these findings. In addition to the association of BD with *B*51* ($p=2.2E-55$), sequential regression of imputed HLA types identified associations of *HLA-A*03* ($p=1E-8$), *HLA-C*0701* ($p=9.5E-4$), and *HLA-B*15* ($p=1.2E-4$) with BD. Stepwise forward regression of imputed *HLA-B* amino acids identified 6 *HLA-B* residues that together fully accounted for the regional association at *HLA-B*.

Conclusion

This study affirms *B*51* as the strongest risk factor of BD. We have provided strong evidence opposing a *B*51*-independent role for *MICA* variants in BD susceptibility. We have identified significant effects of *HLA-A*03* and *HLA-C*0701*, which protect against BD, and *HLA-B*15*, which confers risk of BD. We have identified a group of *HLA-B* amino acids, most of which reside in the antigen binding groove, that together account for the entire association signal at the *HLA-B* locus.

Disclosure of interest

None declared.

Authors' details

¹Translational Genetics and Genomics Unit, NIAMS, Bethesda, USA.

²Department of Internal Medicine and Clinical Immunology, Yokohama City University, Yokohama City, Japan. ³Division of Genetics, Brigham and

Women's Hospital and Harvard Medical School, Boston, USA. ⁴Departments of Medical Genetics and of Epidemiology, University Medical Center Utrecht,

¹Translational Genetics and Genomics Unit, NIAMS, Bethesda, USA
Full list of author information is available at the end of the article

Utrecht, Netherlands. ⁵Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey. ⁶Inflammatory Disease Section, NHGRI, Bethesda, USA.

Published: 8 November 2013

doi:10.1186/1546-0096-11-S1-A236

Cite this article as: Ombrello *et al.*: PW03-010 - MHC complexity in Behçet's disease. *Pediatric Rheumatology* 2013 **11**(Suppl 1):A236.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

