



# Identification of ALK Gene Alterations in Urothelial Carcinoma

## Citation

Bellmunt, J., S. Selvarajah, S. Rodig, M. Salido, S. de Muga, I. Costa, B. Bellosillo, et al. 2014. "Identification of ALK Gene Alterations in Urothelial Carcinoma." PLoS ONE 9 (8): e103325. doi:10.1371/journal.pone.0103325. <http://dx.doi.org/10.1371/journal.pone.0103325>.

## Published Version

doi:10.1371/journal.pone.0103325

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:12785962>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



# Identification of *ALK* Gene Alterations in Urothelial Carcinoma

Joaquim Bellmunt<sup>1,2\*</sup>, Shamini Selvarajah<sup>1</sup>, Scott Rodig<sup>1</sup>, Marta Salido<sup>3</sup>, Silvia de Muga<sup>3</sup>, Irmgard Costa<sup>4</sup>, Beatriz Bellosillo<sup>3</sup>, Lillian Werner<sup>5</sup>, Stephanie Mullane<sup>1</sup>, André P. Fay<sup>1</sup>, Robert O'Brien<sup>1</sup>, Jordi Barretina<sup>6</sup>, André E. Minoche<sup>7,8,9</sup>, Sabina Signoretti<sup>1</sup>, Clara Montagut<sup>2</sup>, Heinz Himmelbauer<sup>8,9</sup>, David M. Berman<sup>10</sup>, Philip Kantoff<sup>1</sup>, Toni K. Choueiri<sup>1</sup>, Jonathan E. Rosenberg<sup>11</sup>

**1** Bladder Cancer Center, Dana-Farber Cancer Institute/Harvard Medical School, Boston, Massachusetts, United States of America, **2** Hospital del Mar Research Institute-IMIM, Barcelona, Spain, **3** Department of Pathology, Hospital del Mar Research Institute-IMIM, Barcelona, Spain, **4** Hospital Parc Tauli, Sabadell, Spain, **5** Biostatistics and Computational Biology, Harvard Medical School, Dana-Farber Cancer Institute, Boston, Massachusetts, United States of America, **6** Broad Institute, Cambridge, Massachusetts, United States of America, **7** Max Planck Institute for Molecular Genetics, Berlin, Germany, **8** Centre for Genomic Regulation (CRG), Barcelona, Spain, **9** Universitat Pompeu Fabra, Barcelona, Spain, **10** Department of Pathology, Johns Hopkins University, Baltimore, Maryland, United States of America, **11** Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, New York, United States of America

## Abstract

**Background:** Anaplastic lymphoma kinase (*ALK*) genomic alterations have emerged as a potent predictor of benefit from treatment with *ALK* inhibitors in several cancers. Currently, there is no information about *ALK* gene alterations in urothelial carcinoma (UC) and its correlation with clinical or pathologic features and outcome.

**Methods:** Samples from patients with advanced UC and correlative clinical data were collected. Genomic imbalances were investigated by array comparative genomic hybridization (aCGH). *ALK* gene status was evaluated by fluorescence *in situ* hybridization (FISH). *ALK* expression was assessed by immunohistochemistry (IHC) and high-throughput mutation analysis with Oncomap 3 platform. Next generation sequencing was performed using Illumina Genome Analyzer Ix, and Illumina HiSeq 2000 in the FISH positive case.

**Results:** 70 of 96 patients had tissue available for all the tests performed. Arm level copy number gains at chromosome 2 were identified in 17 (24%) patients. Minor copy number alterations (CNAs) in the proximity of *ALK* locus were found in 3 patients by aCGH. By FISH analysis, one of these samples had a deletion of the 5' *ALK*. Whole genome next generation sequencing was inconclusive to confirm the deletion at the level of the *ALK* gene at the coverage level used. We did not observe an association between *ALK* CNA and overall survival, ECOG PS, or development of visceral disease.

**Conclusions:** *ALK* genomic alterations are rare and probably without prognostic implications in UC. The potential for testing *ALK* inhibitors in UC merits further investigation but might be restricted to the identification of an enriched population.

**Citation:** Bellmunt J, Selvarajah S, Rodig S, Salido M, de Muga S, et al. (2014) Identification of *ALK* Gene Alterations in Urothelial Carcinoma. PLoS ONE 9(8): e103325. doi:10.1371/journal.pone.0103325

**Editor:** Renato Franco, Istituto dei tumori Fondazione Pascale, Italy

**Received:** April 15, 2014; **Accepted:** June 26, 2014; **Published:** August 1, 2014

**Copyright:** © 2014 Bellmunt et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper, supporting information files and previous publications cited in this manuscript.

**Funding:** This work has been supported by PI061513 (Spanish Health Ministry Grant "Fondo de Investigacion Sanitaria") and RTICC 06/0020/19 grants. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** Toni K. Choueiri: Consultancy: Pfizer, Novartis; Advisory board: Pfizer, Novartis, Aveo, GlaxoSmithKline, Exelixis; Research: Pfizer; No Speakers bureau. All remaining authors have declared no conflicts of interest for this work. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials.

\* Email: joaquim\_bellmunt@dfci.harvard.edu

## Introduction

Urothelial carcinoma (UC) accounts for 15,210 cancer deaths per year in the United States [1]. Five-year survival for patients with muscle invasive (T2) disease or greater is only 50%.

Advanced UC of the bladder is often associated with mutations and multiple somatic copy number alterations [2]. Comparative genomic hybridization studies of bladder carcinomas and cell lines have revealed a number of recurrent genetic aberrations including amplifications or gains on 8q22-24, 11q13, 17q21, and losses on chromosomes 9, 8p22-23, and 17p6-9 [3,4]. In several clinical

cohorts, some of these genomic alterations have also been associated with pathological stage and outcome [5].

In the recent years, potential new targets for treatment intervention have been described in urothelial tumors. The identification of driving genomic alterations as mutations even if occurring in only a small subset of bladder cancer patients, may lead to the development of patient-specific therapies as has been the case of the recently described mutations in *TSC1* predicting response to mTOR inhibitors like everolimus [6–8]. Another example is the *PIK3CA* gene, mutated in up to 26% of cases in the

series by Ross and colleagues that may predict sensitivity to *PIK3CA*/mTOR inhibitors [9].

The ALK (anaplastic lymphoma Kinase) inhibitor crizotinib, has recently shown high efficacy in the treatment of patients with non-small cell lung cancer (NSCLC) with *ALK* translocation which is present in about 4–7% of the tumors [10–12]. In a phase I study of NSCLC patients with an *ALK* translocation, the response rate was 57% independent of performance status or number of previous treatments with a 70% probability of progression free survival at 6 months [13]. In several other tumor types besides lung cancer, *ALK* genomic alterations have been identified as potential oncogenic drivers, meaning that cancers in different organs can be targeted for treatment with *ALK* inhibitors regardless of their cell of origin.

In UC, *ALK* copy number gain, amplification, translocations, mutations, or expression have not been characterized. We therefore investigated *ALK* protein expression and underlying genetic aberrations in a cohort of patients who received chemotherapy in the setting of metastatic disease, focusing on clinical and prognostic implications.

In the present study we show that *ALK* genomic alterations, such as copy number alterations (CNA) and deletions, occur in UC. Additionally, we attempted to identify the impact of these alterations with clinical and outcome features.

## Material and Methods

### Patients

This project was approved by the local ethics committee (CEIC-IMAS) at Hospital del Mar, and by the Dana-Farber/Harvard Cancer Center (DF/HCC) institutional review board (IRB). Because the majority of patients were died at the time of collecting samples, a waiver of consent was requested and given from IRB of DF/HCC for all participants (requiring complete deidentification of the samples prior the analysis).

A cohort of 96 patients, with metastatic UC treated with platinum-based combination was identified. All patients underwent several treatment regimens, all containing gemcitabine and a platinum compound, with some patients receiving additional paclitaxel as well. Patient clinical data was collected. The final cohort included 70 patients (52 males, 18 females) with available clinical data and sufficient tissue samples to conduct all the genomic studies.

### Tumor Samples

The analysis was performed in formalin-fixed paraffin embedded (FFPE) tissue from UC of the urinary tract. Other molecular studies have been performed and reported in these samples in order to characterize the biology of UC [14]. The specimens were retrospectively retrieved from the pathology archive at Hospital del Mar and Mar Biobank in Barcelona, Spain. Slides were reviewed separately by two genitourinary specialist pathologists (MS, DB). All patients had high grade transitional cell carcinoma and no other histological variant was included in this study. Tumor areas were evaluated by a single pathologist (DB) and tumor bearing 0.6 mm cores were punched for DNA extraction and/or tissue microarray (TMA) construction.

### ALK analysis

*ALK* genomic alterations were evaluated by array comparative genomic hybridization (aCGH), fluorescence *in situ* hybridization (FISH), immunohistochemistry (IHC), mass spectrometry mutation analysis and next-generation sequencing. Description of methods can be found in the appendix (**Methods S1**).

## Statistical analysis

Statistical analysis of clinical data and molecular features was carried out with SAS version 9.2 (SAS Institute Inc, Cary, NC). Patient and clinical characteristics were summarized as number and percentages for categorical variables and median and interquartile ranges for continuous variables. Overall survival (OS) was defined from the date patients received first line chemotherapy for advanced disease until date of death or censored on the last known alive date. *ALK* copy number alteration was defined as having more than a 4 fold change [15]. Fisher exact test was used to assess the associations of *ALK* copy number alteration with ECOG PS and whether patients developed visceral disease. Cox proportional hazard model was used to assess the associations of *ALK* copy number alteration and overall survival in both univariate and multivariate analyses. Kaplan-Meier estimate was used to summarize median overall survival. All the statistical tests were conducted at the two-sided 0.05 level of significance.

## Results

The median OS was 12 months with 45 patients deceased at the time of analysis, with a median follow-up of 23 months. **Table 1** summarizes patient and clinical characteristic for the entire cohort as well as for patients with more than 4 fold copy number gain in the FISH analysis.

### Recurrent chromosomal gains and losses by aCGH

Analysis by aCGH of the 70 patients included in the study identified 95 focal and 21 broad (identified as >50% of the chromosome arm) events. The results of the broad alteration analysis were largely consistent with the current literature [16–18]. We observed frequent losses of chromosomes 5q (43%), 8p (69%), 9 (p: 48%; q: 41%), 10q (41%), 11p (49%), 17p (51%), and 22q (40%) and recurrent gains of chromosomes 3q (46%), 5p (48%), 8q (48%), 19q (34%), and 20 (60%). Three specimens out of 70 harbored minor non-significant alterations (log<sub>2</sub> ratio 0–0.8) in chromosome 2, where *ALK* gene locus is located. This encouraged us to conduct a more in-depth search of *ALK* genomic alterations and to further characterize the 5′*ALK* deletion seen by FISH in one patient.

### FISH analysis of *ALK* gene/copy number gains


To further characterize genomic imbalances on chromosome 2, all samples underwent FISH analysis. One case presented a deletion of the green signal (5′*ALK*), centromeric to the *ALK* gene, and also had gain of the *ALK* gene fusion signals and 3′*ALK* signal (**Figures 1 and 2**). This FISH pattern was interpreted as an *ALK* atypical rearrangement as has been described in *ALK* positive NSCLC because a single orange (3′*ALK*) signal was seen [19]. In these cases it is assumed that the deletion is the result of translocation. Analyses of *EML4* as well as other known fusion partners such as *TGF* and *KIF5* were performed without finding any translocation of these genes. Even so, it is possible that the deletion does not cause the *ALK* translocation and other molecular techniques need to be applied to further characterize the FISH findings.

*ALK* gene copy number gains and amplification were analyzed in all samples. Two cases presented amplification of *ALK*. 90% of samples showed *ALK* copy number gain due to polysomy of chromosome 2. All of them had 3 to 6 copies of CEP2 except one case with high polysomy. Among 70 urothelial tumors, 7 (10%) demonstrated 2F signals (2 intact *ALK* loci), 46 (65.7%) had 3–4F signals present, and 17 (24.3%) had ≥5F signals (range 5F–11F; median 6F) in >10% of nuclei (**Table 2**). The associations of

**Table 1.** Patients and Clinical Characteristics.


	All patients (N = 70)		Patients with copy number alteration (N = 17)	
	N	% or median (q1, q3)	N	% or median (q1, q3)
<b>Age</b>	61	63 (54, 68)	15	66 (58, 68)
<b>Sex</b>				
Male	52	74%	15	88%
Female	18	16%	2	12%
<b>ECOG PS</b>				
0	22	31%	4	24%
1, 2	48	69%	13	76%
<b>Visceral diseases</b>				
No	41	59%	7	41%
Yes	29	41%	10	59%
<b>Pathological stage</b>				
Stage 0 (Ta)	5	7%	2	12%
Stage I (T1)	5	7%	0	0%
Stage II (T2)	36	51%	8	47%
Stage III (T3, T4)	22	31%	7	41%
Stage IV (L, M)	1	1%	0	0%
Missing	1	1%	0	0%

doi:10.1371/journal.pone.0103325.t001




**Figure 1.** 1298case –FISH + for ALK variant (green probe missing).

doi:10.1371/journal.pone.0103325.g001



**Figure 2. 1298case –FISH copy gain (a) & amplified (b).**  
doi:10.1371/journal.pone.0103325.g002



**Figure 3. Comparison of OS between ≥5F patients and 2F+3–4F.**  
doi:10.1371/journal.pone.0103325.g003

ALK copy number alteration with ECOG PS, visceral disease, and OS are summarized in **Tables 3 and 4**. No significant association between ALK copy number alteration and clinical features or overall survival was observed (**Figure 3**).

Comparison of ALK gene copy number gains to clinical and pathological features for the 70 patients are summarized in **Table 1**. There were no differences between ALK gene copy number gains and clinical features in all the subgroups (2F, 3–4F and ≥5F). OS rates for patients with 2F+3–4F and >5F were 12 and 16 months respectively. There was no statistically significant difference between these groups (**Figure 3**).

**ALK protein expression by immunohistochemistry**

To further characterize whether ALK protein expression was affected, immunohistochemistry analysis of all FFPE samples was performed using the Cell Signaling antibody. Immunohistochemistry staining was negative in the tumor with ALK FISH positive test. Similarly, among tumors with ALK gene copy gain or amplification, ALK protein expression was not detected. None of the tumors classified as ALK negative by FISH showed ALK protein expression by immunohistochemistry.

**Table 2. Copy Number Alteration.**

	N	%
2F	7	10
3-4F	46	66
≥5F	17	24

doi:10.1371/journal.pone.0103325.t002

**High-throughput mutational analysis using OncoMap**

To have more accurate information on genetic alteration in these UC samples, mass spectrometry mutation analysis was also performed for all samples. Ninety-six samples were submitted for OncoMap; 87/96 (91%) passed all quality control steps. 79% (69/87) passing samples harbored candidate mutations. In total, 150 candidate mutation calls were made across 47 genes. Overall, 39% (58/150) of candidate mutations in passing samples were conservative and 61% (92/150) were aggressive. No mutations in ALK were found using this platform. ALK P496L candidate mutation was found in one of the sample but was not confirmed with HME.

**Next-generation sequencing of ALK gene**

Since FISH technique gives no information of the specific sequence and the exact size of the deleted fragment in ALK, directed analysis of ALK gene was performed by next generation sequencing (Illumina). Analysis of the region containing P496 only showed base changes at rates below 1%, reflecting the expected sequencing error rate. Thus, only the wild-type sequence for the position P496 was detected and no mutations on ALK were detected by this technique.

We then extended the search space to the centromere with the intention to explore potential deletions according to FISH results. In the new analysis performed on the FISH positive patient, one read of a pair should match within the ALK locus, 29.37 Mb - 32 Mb, and one read should match at some place towards the centromere (>10 kb up to position 93.3 Mb). However, at the coverage level used no deletions could be confirmed with this approach.

**Table 3.** Association of *ALK* copy number alteration with ECOG PS and visceral disease.

	<i>ALK</i> >4 copies		P-value
	No	Yes	
<b>ECOG PS</b>			0.55
0	18	4	
1, 2	35	13	
<b>Visceral disease</b>			0.16
No	34	7	
Yes	19	10	

doi:10.1371/journal.pone.0103325.t003

## Discussion

In the present study we interrogate whether the *ALK* genomic alterations are of potential clinical relevance in patients with UC. Our study shows that *ALK* amplification and copy number gain but not fusions and translocations occurs in UC but is not associated with poor outcome in our patients with already bad prognosis.

*ALK* gene is located in 2p23 and encodes a transmembrane tyrosine kinase receptor involved in the development of nervous system during embryogenesis [20,21]. *ALK* gene was first shown to have a role in cancer as part of the fusion gene nucleophosmin (NPM)-*ALK* in anaplastic large cell lymphomas(9, 10). Preclinical studies show that tumors with aberrant activation of *ALK* tyrosine kinase are oncogene addicted to *ALK* intracellular signaling, and inhibition of the kinase by specific *ALK* targeting drugs results in tumor growth arrest and cell death (25).

The best well studied genomic alteration is the translocation seen in NSCLC patients. The majority of *ALK* rearrangements come from an interstitial deletion and inversion in chromosome 2p resulting in *EML4-ALK* fusion gene product [22–27]. Although translocation is the most commonly identified mechanism for *ALK* activation, amplification and mutation have also been shown to act as oncogenic events [28–30]. The role of amplification and of copy number gain, as well as the role of deletion found in tumors like RMS remains to be determined [27,31–34].

The finding that several tumor types have been identified that have *ALK* as an oncogenic driver regardless of their cell of origin has prompted the creation of the term “*ALKomas*” implying a “beyond organ” concept classification assuming consequently responses to *ALK* inhibitors such as crizotinib [10,35]. Based on that, exploration of this concept is worthwhile in UC even if the frequency happens to be low.

In our cohort, aCGH-A found only some minor focal events in 3/70 specimens harboring non-significant alterations in *ALK* gene locus region. Since copy number gain has been recently associated with poor prognosis in several tumors like RMS, RCC and colorectal cancer (CRC), FISH analysis to assess the impact of copy number variations of *ALK* in our cohort was performed. In our patients, polysomy was frequently found in 90% of the cases [15,34]. The biological relevance of such finding is uncertain but could reflect genomic instability. The OS for patients with (2F+3–4F) vs. >5F was found to be 12 and 16 months respectively, however did not reach statistical significance (Figure 1). Likewise, there were no differences between *ALK* gene copy number gains and clinical features in all the different subgroups (2F, 3–4F and ≥ 5F). A plausible explanation for this lack of a significant difference between these groups is that it could be related to the natural

history and the aggressive phenotype of our analysis cohort (metastatic disease requiring chemotherapy) with other genetic abnormalities beyond *ALK* gene copy number having a greater functional role in oncogenesis. Similarly, arm level *ALK* gene copy number gain as observed in this analysis may be unrelated to the driver oncogenic events.

Generally, patients with *ALK* copy gain have not shown to have detectable *ALK* protein expression as assessed by IHC except for a recent publication by van Gaal and colleagues [27,29,34,36]. In our series, no patient with gene copy gain or amplification tested positive by IHC. This is similar to that observed in CRC where increased *ALK* gene copy number did not translate to increased *ALK* protein expression [37]. However, this is not the case for patients being categorized as FISH positive, where this positivity strongly correlates with IHC. Of note, in lung cancer, a positive *ALK* FISH test and *ALK* IHC have been proposed as screening tools to detect *ALK* alterations being considered sufficiently sensitive to indicate treatment with crizotinib [37]. Moreover, in NSCLC, abnormal FISH signal patterns have varied from a single split signal to more complex signal patterns, such as deletions of the green 5' end of the *ALK* probe, gain of the split or 5'*ALK* signal or both. These variant *ALK* FISH signals usually, but not always, represent an *ALK* translocation and therefore the finding of a loss of the 5'*ALK* signal has been considered to be a presumptive evidence of an *ALK* gene rearrangement [37].

In our series, the patient with a FISH positive result had a variant signal pattern that did not correlate with *ALK* protein expression as assessed by IHC. The case was interpreted as having a deletion in the *ALK* region due to loss of the green 5' end of the *ALK* signal, after excluding the possibility it could be related to alternative translocation partners [Kinesin family 5B (*KIF5B*) and TRK-fused gene (*TFG*)]. In our patient we did not test for the rearrangement of other fusion partners to *ALK* such as C2orf44, *KIF5B*, *NPM1*, *VCL*, *TFG*, *RET*, *ROS*, and *VCL* [38–43]. These genes have all been shown to be partners of *ALK* in lung cancer [44].

Finally, *ALK* Mutations have been described in 10.4% of neuroblastoma samples but not in other pediatric tumors like RMS, Ewing sarcoma, or DSRCT and only occasionally in other solid tumors like CRC [45,46]. In lung cancers, *ALK* mutations appear to develop during clinical treatment with crizotinib and their generation probably renders *EML4-ALK* resistant not only to crizotinib but also to other *ALK* inhibitors [47]. In our series, no *ALK* P496L mutation was observed. In our study the limitations of the platform used limits our conclusions of the mutation analysis. The absence or very low percentage of activating mutation of *ALK* described in the majority of adult solid tumors tested support our analysis that these alterations are not relevant events in UC.

**Table 4.** Comparison of OS between  $\geq 5F$  patients and  $2F+3-4F$ .

	N	Death	Median OS	Hazard ratio	P-value	Adjusted hazard ratio	P-value
<b>AKL &gt;4 copies</b>					0.80		0.38
<b>2F+3-4F</b>	53	34	12	1.1 (0.55, 2.16)		1.36 (0.60, 2.72)	
<b><math>\geq 5F</math></b>	17	11	16	1 (reference)		1 (reference)	

doi:10.1371/journal.pone.0103325.t004

Unfortunately, the suspected deletion in the *ALK* region was not confirmed with the sequencing approach used. Discordantly, mapping read pairs suggesting deletions resolved into correctly mapping read pairs that were in agreement with the insert size of the library when a single mismatch between read and reference genome was tolerated. Thus, these pairs do not support deletions at the *ALK* locus. The average read coverage across the *ALK* region was  $5\times$  and if only a small proportion of cells contained a deletion, we would not have been able to detect it. Because we suspect the deletion was close to the centromere, we might have missed it and might not have been able to confirm it by next generation sequencing.

To summarize, the increasing evidence that *ALK* alterations are seen in tumors from different origins highlights the concept of stratifying tumors according to oncogenic genotypes as opposed to tissue type when considering treatment strategies. The finding of the absence of *ALK* rearrangement together with no activating mutation in *ALK* suggests that these alterations might not be pathogenic events in UC. The utility of testing *ALK* inhibitors in UC is not supported by this data, although in the absence of effective alternative agents testing *ALK* inhibitors may still be warranted.

In conclusion, *ALK* genomic alterations are rare and probably without prognostic implications in UC. The potential for testing *ALK* inhibitors in patients with deletions and copy number

changes UC merits further investigation in a larger expanded cohort of UCs, but might be restricted to the infrequent finding of a FISH positive patient.

**Supporting Information**

**Methods S1 Supplementary Methods.** (DOCX)

**Acknowledgments**

We thank Fundació Cellex (Barcelona) for a generous donation to the Group of Molecular Therapeutics and Biomarkers, Hospital del Mar. We thank the Tumor Bank of the Department of Pathology of Hospital del Mar (RD09/0076/0036), and the Xarxa de Bancs de Tumors sponsored by Pla Director d'Oncologia de Catalunya (XBTC) for providing tissue samples. We also thank Matthew Ducar from Center for Cancer Genome Discovery (CCGD) for helpful comments and suggestions.

**Author Contributions**

Conceived and designed the experiments: JB TC JR. Performed the experiments: SS SR MS SM BB AM SS HH DB. Analyzed the data: JB SS SR MS SM BB LW AF SS CM HH DB PK TC JR. Contributed reagents/materials/analysis tools: JB IC SM RO JB PK TC JR. Contributed to the writing of the manuscript: JB AF SS CM PK TC JR.

**References**

- Siegel R, Naishadham D, Jemal A (2012) Cancer statistics, 2012. *CA Cancer J Clin* 62: 10–29.
- Cancer Genome Atlas Research N (2014) Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature* 507: 315–322.
- Hoglund M (2012) The bladder cancer genome; chromosomal changes as prognostic makers, opportunities, and obstacles. *Urol Oncol* 30: 533–540.
- Lopez V, Gonzalez-Peramato P, Suela J, Serrano A, Algaba F, et al. (2013) Identification of prefolidin amplification (1q23.3–q24.1) in bladder cancer using comparative genomic hybridization (CGH) arrays of urinary DNA. *J Transl Med* 11: 182.
- Blaveri E, Brewer JL, Roydasgupta R, Fridlyand J, DeVries S, et al. (2005) Bladder cancer stage and outcome by array-based comparative genomic hybridization. *Clin Cancer Res* 11: 7012–7022.
- Balbas-Martinez C, Sagraera A, Carrillo-de-Santa-Pau E, Earl J, Marquez M, et al. (2013) Recurrent inactivation of STAG2 in bladder cancer is not associated with aneuploidy. *Nat Genet* 45: 1464–1469.
- Iyer G, Hanrahan AJ, Milowsky MI, Al-Ahmadie H, Scott SN, et al. (2012) Genome sequencing identifies a basis for everolimus sensitivity. *Science* 338: 221.
- Iyer G, Al-Ahmadie H, Schultz N, Hanrahan AJ, Ostrovskaya I, et al. (2013) Prevalence and co-occurrence of actionable genomic alterations in high-grade bladder cancer. *J Clin Oncol* 31: 3133–3140.
- Ross JS, Wang K, Al-Rohil RN, Nazeer T, Sheehan CE, et al. (2014) Advanced urothelial carcinoma: next-generation sequencing reveals diverse genomic alterations and targets of therapy. *Mod Pathol* 27: 271–280.
- Kwak EL, Bang YJ, Camidge DR, Shaw AT, Solomon B, et al. (2010) Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med* 363: 1693–1703.
- Camidge DR, Bang YJ, Kwak EL, Iafrate AJ, Varella-Garcia M, et al. (2012) Activity and safety of crizotinib in patients with ALK-positive non-small-cell lung cancer: updated results from a phase 1 study. *Lancet Oncol* 13: 1011–1019.
- Shaw AT, Yeap BY, Solomon BJ, Riely GJ, Gainor J, et al. (2011) Effect of crizotinib on overall survival in patients with advanced non-small-cell lung cancer harbouring ALK gene rearrangement: a retrospective analysis. *Lancet Oncol* 12: 1004–1012.
- Lee JO, Kim TM, Lee SH, Kim DW, Kim S, et al. (2011) Anaplastic lymphoma kinase translocation: a predictive biomarker of pemetrexed in patients with non-small cell lung cancer. *J Thorac Oncol* 6: 1474–1480.
- Riester M, Werner L, Bellmunt J, Selvarajah S, Guancial EA, et al. (2014) Integrative analysis of 1q23.3 copy-number gain in metastatic urothelial carcinoma. *Clin Cancer Res* 20: 1873–1883.
- Sukov WR, Hodge JC, Lohse CM, Akre MK, Leibovich BC, et al. (2012) ALK alterations in adult renal cell carcinoma: frequency, clinicopathologic features and outcome in a large series of consecutively treated patients. *Mod Pathol* 25: 1516–1525.
- Hurst CD, Platt FM, Taylor CF, Knowles MA (2012) Novel tumor subgroups of urothelial carcinoma of the bladder defined by integrated genomic analysis. *Clin Cancer Res* 18: 5865–5877.
- Panzeri E, Conconi D, Antolini L, Redaelli S, Valsecchi MG, et al. (2011) Chromosomal aberrations in bladder cancer: fresh versus formalin fixed paraffin embedded tissue and targeted FISH versus wide microarray-based CGH analysis. *PLoS One* 6: e24237.
- Tian Z, Kuang R (2010) Integrative classification and analysis of multiple array/CGH datasets with probe alignment. *Bioinformatics* 26: 2313–2320.
- Camidge DR, Kono SA, Flacco A, Tan AC, Doebele RC, et al. (2010) Optimizing the detection of lung cancer patients harboring anaplastic lymphoma kinase (ALK) gene rearrangements potentially suitable for ALK inhibitor treatment. *Clin Cancer Res* 16: 5581–5590.
- Chiarle R, Voena C, Ambrogio C, Piva R, Inghirami G (2008) The anaplastic lymphoma kinase in the pathogenesis of cancer. *Nat Rev Cancer* 8: 11–23.
- Iwahara T, Fujimoto J, Wen D, Cupples R, Bucay N, et al. (1997) Molecular characterization of ALK, a receptor tyrosine kinase expressed specifically in the nervous system. *Oncogene* 14: 439–449.
- Rikova K, Guo A, Zeng Q, Possemato A, Yu J, et al. (2007) Global survey of phosphotyrosine signaling identifies oncogenic kinases in lung cancer. *Cell* 131: 1190–1203.



23. Soda M, Choi YL, Enomoto M, Takada S, Yamashita Y, et al. (2007) Identification of the transforming EML4-ALK fusion gene in non-small-cell lung cancer. *Nature* 448: 561–566.
24. Rodig SJ, Mino-Kenudson M, Dacic S, Yeap BY, Shaw A, et al. (2009) Unique clinicopathologic features characterize ALK-rearranged lung adenocarcinoma in the western population. *Clin Cancer Res* 15: 5216–5223.
25. Shaw AT, Yeap BY, Mino-Kenudson M, Digumarthy SR, Costa DB, et al. (2009) Clinical features and outcome of patients with non-small-cell lung cancer who harbor EML4-ALK. *J Clin Oncol* 27: 4247–4253.
26. Butrynski JE, D'Adamo DR, Hornick JL, Dal Cin P, Antonescu CR, et al. (2010) Crizotinib in ALK-rearranged inflammatory myofibroblastic tumor. *N Engl J Med* 363: 1727–1733.
27. Salido M, Pijuan L, Martínez-Avilés L, Galván AB, Cañadas I, et al. (2011) Increased ALK gene copy number and amplification are frequent in non-small cell lung cancer. *J Thorac Oncol* 6: 21–27.
28. Carén H, Abel F, Kogner P, Martinsson T (2008) High incidence of DNA mutations and gene amplifications of the ALK gene in advanced sporadic neuroblastoma tumours. *Biochem J* 416: 153–159.
29. George RE, Sanda T, Hanna M, Fröhling S, Luther W, et al. (2008) Activating mutations in ALK provide a therapeutic target in neuroblastoma. *Nature* 455: 975–978.
30. Janoueix-Lerosey I, Lequin D, Brugières L, Ribeiro A, de Pontual L, et al. (2008) Somatic and germline activating mutations of the ALK kinase receptor in neuroblastoma. *Nature* 455: 967–970.
31. Montagut C, Galvan AB, Gallen M, Salido M, Sole F, et al. (2010) ALK chromosomal alterations in colon cancer patients. *Journal of Clinical Oncology (Meeting Abstracts)* Vol 28: 10537.
32. Lee JS, Lim SM, Rha SY, Roh JK, Cho YJ, et al. (2014) Prognostic implications of anaplastic lymphoma kinase gene aberrations in rhabdomyosarcoma; an immunohistochemical and fluorescence in situ hybridisation study. *J Clin Pathol* 67: 33–39.
33. Bonvini P, Zin A, Alaggio R, Pawel B, Bisogno G, et al. (2013) High ALK mRNA expression has a negative prognostic significance in rhabdomyosarcoma. *Br J Cancer* 109: 3084–3091.
34. van Gaal JC, Flucke UE, Roelfen MH, de Bont ES, Sleijfer S, et al. (2012) Anaplastic lymphoma kinase aberrations in rhabdomyosarcoma: clinical and prognostic implications. *J Clin Oncol* 30: 308–315.
35. Mano H (2012) ALKoma: a cancer subtype with a shared target. *Cancer Discov* 2: 495–502.
36. Bavi P, Jehan Z, Bu R, Prabhakaran S, Al-Sanea N, et al. (2013) ALK gene amplification is associated with poor prognosis in colorectal carcinoma. *Br J Cancer* 109: 2735–2743.
37. Dai Z, Kelly JC, Meloni-Ehrig A, Slovak ML, Boles D, et al. (2012) Incidence and patterns of ALK FISH abnormalities seen in a large unselected series of lung carcinomas. *Mol Cytogenet* 5: 44.
38. Lipson D, Capelletti M, Yelensky R, Otto G, Parker A, et al. (2012) Identification of new ALK and RET gene fusions from colorectal and lung cancer biopsies. *Nat Med* 18: 382–384.
39. Takeuchi K, Choi YL, Togashi Y, Soda M, Hatano S, et al. (2009) KIF5B-ALK, a novel fusion oncokine identified by an immunohistochemistry-based diagnostic system for ALK-positive lung cancer. *Clin Cancer Res* 15: 3143–3149.
40. Morris SW, Kirstein MN, Valentine MB, Dittmer KG, Shapiro DN, et al. (1994) Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science* 263: 1281–1284.
41. Debelenko LV, Raimondi SC, Daw N, Shivakumar BR, Huang D, et al. (2011) Renal cell carcinoma with novel VCL-ALK fusion: new representative of ALK-associated tumor spectrum. *Mod Pathol* 24: 430–442.
42. Hernandez L, Pinyol M, Hernandez S, Bea S, Pulford K, et al. (1999) TRK-fused gene (TFG) is a new partner of ALK in anaplastic large cell lymphoma producing two structurally different TFG-ALK translocations. *Blood* 94: 3265–3268.
43. Takeuchi K, Soda M, Togashi Y, Suzuki R, Sakata S, et al. (2012) RET, ROS1 and ALK fusions in lung cancer. *Nat Med* 18: 378–381.
44. Barreca A, Lasorsa E, Riera L, Machiorlatti R, Piva R, et al. (2011) Anaplastic lymphoma kinase in human cancer. *J Mol Endocrinol* 47: R11–23.
45. Shukla N, Ameer N, Yilmaz I, Nafa K, Lau CY, et al. (2012) Oncogene mutation profiling of pediatric solid tumors reveals significant subsets of embryonal rhabdomyosarcoma and neuroblastoma with mutated genes in growth signaling pathways. *Clin Cancer Res* 18: 748–757.
46. Bavi P, Jehan Z, Bu R, Prabhakaran S, Al-Sanea N, et al. (2013) ALK gene amplification is associated with poor prognosis in colorectal carcinoma. *Br J Cancer*.
47. Choi YL, Soda M, Yamashita Y, Ueno T, Takashima J, et al. (2010) EML4-ALK mutations in lung cancer that confer resistance to ALK inhibitors. *N Engl J Med* 363: 1734–1739.