



# Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Summons, Roger E., Jan P. Amend, David Bish, Roger Buick, George D. Cody, David J. Des Marais, Gilles Dromart, Jennifer L. Eigenbrode, Andrew H. Knoll, and Dawn Y. Sumner. 2011. "Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group." <i>Astrobiology</i> 11, no. 2: 157–181.
Published Version	<a href="https://doi.org/10.1089/ast.2010.0506">doi:10.1089/ast.2010.0506</a>
Citable link	<a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:13041033">http://nrs.harvard.edu/urn-3:HUL.InstRepos:13041033</a>
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <a href="http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP">http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP</a>

# Astrobiology

Astrobiology Manuscript Central: <http://mc.manuscriptcentral.com/astrobiology>

## Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

Journal:	<i>Astrobiology</i>
Manuscript ID:	AST-2010-0506.R2
Manuscript Type:	News and Views
Date Submitted by the Author:	13-Dec-2010
Complete List of Authors:	Summons, Roger E.; Massachusetts Institute of Technology, Earth, Atmospheric and Planetary Sciences Amend, Jan; Washington University Buick, Roger; University of Washington Bish, David; Indiana University Cody, George; CIW Des Marais, David; NASA Ames Research Center Dromart, Gilles Eigenbrode, Jennifer; NASA Goddard Space Flight Center Knoll, Andrew; Harvard University, OEB
Keyword:	Astrobiology, Atmospheric Gases, Mars, Biosignatures, Life Detection

SCHOLARONE™  
Manuscripts

# Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

Roger E. Summons<sup>1\*</sup>, Jan P. Amend<sup>2</sup>, David Bish<sup>3</sup>, Roger Buick<sup>4</sup>, George D. Cody<sup>5</sup>,  
David J. Des Marais<sup>6</sup>, Gilles Dromart<sup>7</sup>, Jennifer L. Eigenbrode<sup>8</sup>, Andrew H. Knoll<sup>9</sup>,  
Dawn Y. Sumner<sup>10</sup>

1. Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge MA 02139, USA
2. Department of Earth & Planetary Sciences, Washington University, 1 Brookings Dr., Saint Louis MO 63130, USA
3. Department of Geological Sciences, Indiana University, 1001 East 10th St., Bloomington, IN 47405, USA
4. Department of Earth & Space Sciences, University of Washington, 4000 15th Avenue NE, Seattle, WA 98195, USA
5. Geophysical Laboratory, Carnegie Institution of Washington, 5251 Broad Branch Road NW, Washington, DC 20015, USA
6. Exobiology Branch, NASA Ames Research Center, Mail Stop 239-4, Moffett Field, CA 94035, USA
7. Laboratoire de Science de la Terre, UMR CNRS 5570, Université Claude Bernard/Ecole Normale Supérieure de Lyon, 2 rue Raphaël Dubois, 696222 Villeurbanne Cedex, France
8. Solar System Exploration Division, NASA Goddard Space Flight Center, Mail Code 699, 8800 Greenbelt Rd., Greenbelt, MD 20771, USA
9. Departments of Earth and Planetary Sciences and Organismic and Evolutionary Biology, 26 Oxford St., Cambridge MA 02138, USA
10. Geology Department, University of California Davis, 1 Shields Ave, Davis, CA 95616, USA

\* Corresponding Author: rsummons@mit.edu

## Abstract

The Mars Science Laboratory (MSL) has an instrument package capable of making measurements of past and present environmental conditions. The data generated may tell us if Mars is, or ever was, able to support life. However, the knowledge of Mars' past history and the geological processes most likely to preserve a record of that history remain sparse and, in some instances, ambiguous. Physical, chemical and geological processes relevant to biosignature preservation on Earth, especially under conditions early in its history when microbial life predominated, are also imperfectly known. Here we present the report of a working group chartered by the Co-Chairs of NASA's MSL Project Science Group, John P Grotzinger and Michael A Meyer, to review and evaluate potential for biosignature formation and preservation on Mars.

Orbital images confirm that layered rocks achieved kilometer-scale thicknesses in some regions of ancient Mars. Clearly, interplays of sedimentation and erosional processes govern present-day exposures, and our understanding of these processes is incomplete. MSL can document and evaluate patterns of stratigraphic development as well as the sources of layered materials and their subsequent diagenesis. It can also document other potential biosignature repositories such as hydrothermal environments. These capabilities offer an unprecedented opportunity to decipher key aspects of the environmental evolution of Mars' early surface and aspects of the diagenetic processes that have operated since that time.

Considering the MSL instrument payload package, we identified the following classes of biosignatures as within the MSL detection window: organism morphologies (cells, body fossils, casts), biofabrics (including microbial mats), diagnostic organic molecules, isotopic signatures, evidence of biomineralization and bioalteration, spatial patterns in chemistry and biogenic gases. Of these, biogenic organic molecules and biogenic atmospheric gases are considered the most definitive and most readily detectable by MSL.

## Introduction

In the first decade of the twenty-first century, our understanding of Mars and its environmental history has increased dramatically. Orbital measurements provide unprecedented resolution of both physical and chemical features of the Martian surface. The Mars Exploration Rovers Opportunity and Spirit have contributed our first-ever geologist's-eye views of stratigraphic successions on Mars (e.g. Christensen et al., 2004; Squyres et al., 2004, 2006, 2009; Grotzinger et al., 2005; Haskin et al., 2005). Building on this success, the extraordinary instrument package and anticipated roving capability of the Mars Science Laboratory (MSL) position us to use new rover observations to test hypotheses generated on the basis of high-resolution orbital data. For example, orbital images and spectral data show that layered rocks have accumulated to thicknesses greater than a kilometer on some parts of the ancient Martian surface and comprise diverse lithologies (e.g. Bishop et al., 2008; Ehmann et al., 2008a,b). What fundamental processes of particle generation, transport, and cementation made such accumulations possible, and what erosional processes govern present day exposures? MSL has the imaging capability to document and evaluate patterns of stratigraphic development, as well as the geochemical capacity to evaluate the sources of layered materials and their

1  
2  
3 subsequent diagenesis. Bibring et al. (2005) have identified phyllosilicates by remote sensing of  
4 the Martian surface and hypothesized that these record an early, relatively wet epoch in Martian  
5 history, before the time of acid sulfate deposition recorded in the sediments at Meridiani  
6 Planum. MSL has the capacity to confirm phyllosilicate mineralogy and evaluate it in  
7 stratigraphic context, allowing us to understand more fully the significance of these minerals for  
8 reconstructing Martian environmental history. Finally, the hallmark of life is organic matter —do  
9 Martian strata preserve organic molecules that might illuminate the planet's early environmental,  
10 and possibly biological, history? MSL's Sample Analysis at Mars (SAM) instrument suite  
11 provides unprecedented potential to answer this question.  
12  
13  
14

15 Knowledge of our own planet's biological and environmental history has developed through the  
16 integration of many types of observations. Arguably among the most important are: field  
17 mapping, the measurement and correlation of stratigraphic sections, and both paleontological  
18 and geochemical analyses of samples collected from measured sections located within mapped  
19 terrains. Through the strategic choice of a landing site using high-resolution orbital data, MSL  
20 promises a comparable integration for Mars. With this in mind, it is helpful, even necessary, to  
21 consider how geologists evaluate and select field sites on Earth. Surely, site accessibility and  
22 map data influence choice, as do accumulated observations by previous geologists. But there  
23 is one more consideration that governs site selection by geologists, geochemists, and  
24 paleobiologists: what types of deposits are most likely to preserve geological and possible  
25 biological signals of interest? Simply put, NASA will realize the greatest returns on its  
26 investment in the MSL payload if it targets outcrops that maximize the probability of organic  
27 deposition and preservation. It will maximize its return from the payload if it targets stratigraphic  
28 successions that place mineralogical measurements in temporal and paleoenvironmental  
29 context.  
30  
31  
32  
33  
34

### 35 **Working Group Objectives**

36  
37 The Preservation Working Group was assembled to assess whether current understanding of  
38 organic matter preservation on Earth might help guide site selection and both strategic as well  
39 as tactical planning during surface operations for the MSL mission. The Working Group was  
40 asked to provide general guidance on what specific geologic environments would be most  
41 favorable for preservation of potential biosignatures, including special consideration of organic  
42 carbon (Farmer and Des Marais, 1999).  
43  
44

45 Insert Text Box 1 and Box 2 hereabouts  
46

47 It is essential to recognize that, on Earth, and in spite of a vital biosphere, the majority of  
48 biologically derived organic carbon exists as "fossil" organic carbon stored within layered  
49 sedimentary rocks. This sequestered organic matter is by some estimates  $2 \times 10^5$  that the mass  
50 of the carbon stored in the living biosphere. On Mars, it appears reasonable to assume that if  
51 life exists or ever existed, it never evolved to the point of large differentiated, multi-cellular  
52 organisms (e.g., plants) that biosynthesize large quantities of recalcitrant biopolymers (e.g.,  
53 lignin and cellulose) leading to potential accumulations of extensive organic matter rich  
54 sedimentary deposits (e.g., coal). There is general consensus that extant microbial life on Mars  
55  
56  
57  
58  
59  
60

1  
2  
3 would likely exist (if at all) in the subsurface and at low abundance, making it difficult, if not  
4 impossible, to detect using sampling technologies foreseeable over the next decade.  
5

6  
7 Detecting an ancient subsurface biota may be even more difficult unless exhumed records of  
8 ancient subsurface environments can be confidently recognized and are accessible to a rover.  
9 Since all life requires an abundance of electron donors and acceptors to grow and to  
10 accumulate biomass, we would also need analyze the mineralogy and geochemistry of  
11 exhumed rocks to assess their bioenergetic potential. Even though the record of early life on  
12 Earth is reported to contain microstructures (e.g. Rasmussen, 2000; Brasier et al., 2006)  
13 purported to reflect subsurface life, significant doubt remains about their biogenicity. These  
14 remains are rare and notoriously difficult to interpret with confidence given their high degree of  
15 thermal alteration. In contrast, it is possible that the organic remains of extinct microbial life  
16 developed at Mars' surface, or subsurface, may persist and perhaps even be enriched in certain  
17 sedimentary rocks.  
18  
19

20  
21 On Earth, microorganisms commonly exist communally in the water column and in sediment  
22 pore spaces or as attached biofilms. Biomass of these communities may be preserved in the  
23 rock record in concentrations sufficient to detect with the MSL payload elements (see Box 1).  
24 Furthermore, sedimentary processing (e.g., hydrodynamic sorting as occurs in fluvial and deltaic  
25 environments on Earth) may concentrate biologically derived carbonaceous particles into  
26 organic-rich horizons in sedimentary strata. A logical approach to establishing whether life ever  
27 existed on Mars is, therefore, to analyze appropriate sedimentary lithologies, seeking evidence  
28 of "fossilized" organic matter preserved in sedimentary deposits.  
29  
30  
31

32 One issue that may add ambiguity in such a search is the fact that, throughout Martian history,  
33 organic-rich chondritic meteorites have undoubtedly rained upon the Martian surface.  
34 Weathering of chondritic meteoritic debris in an environment with minimal oxygen and no  
35 extensive surface biosphere (i.e., microbes and fungi capable of degrading chondritic organic  
36 matter) may lead to a persistence of extraterrestrial organic particles and, consequently, its  
37 accumulation into certain sedimentary rocks through hydrodynamic sorting. At the same time,  
38 strong oxidants in Martian regolith along with exposure to ionizing radiation might alter or  
39 destroy molecular signatures from meteorites or organisms. Notwithstanding the potential  
40 complexity of interpreting any organic material that may be detected, the simple detection of  
41 organic matter in rock or soil via the MSL mission will constitute a critically important result.  
42  
43  
44

45 The membership and objectives of the Working Group were chartered by the Co-Chairs of  
46 NASA's MSL Project Science Group, John P Grotzinger and Michael A Meyer. The charge to  
47 the Working Group was to outline a search strategy that included consideration of the nature of  
48 the payload elements (PE), a ranked order of biosignatures detectable by those instruments and  
49 an evaluation of the types of environments conducive to the formation and, especially,  
50 preservation of these biosignatures. Four candidate landing sites remain at the time of writing  
51 and this strategy could serve as a guide to members of the Mars exploration community in  
52 deliberating whether potential landing sites have more or less appropriate lithologies for  
53 biosignature preservation. A second objective was to assess, in very general terms, how the  
54 MSL instrumentation may be best exploited for the analysis of any record of organic compounds  
55  
56  
57  
58  
59  
60



(biogenic or abiogenic) that might be preserved in Martian sedimentary rocks. The working group was asked not to address the merits or otherwise of specific Mars localities. Nor were we asked to assess instrument specifications and how those instruments might be operated at the surface of Mars, as these are issues more appropriately addressed by very different groups of specialists. We were guided in our work by two recent publications from the National Academies namely the report on the *Limits of Life in Planetary Systems* (Baross et al., 2007) and an *Astrobiological Strategy for the Exploration of Mars* (Jakosky et al., 2007).

### **Taphonomic Windows**

Taphonomy is the term used by paleontologists to describe the processes by which living organisms become fossils. Taphonomy is commonly portrayed as “post-mortem information loss” because fossilized bones, shells, or microorganisms record only a small subset of the biological information originally present in their makers. With a different perspective, however, we might view taphonomy as “post-mortem information preservation” because, however selective it may be, fossilization provides our only permanent record of past life. In the context of MSL, we can broaden the concept of taphonomy to include the geologic preservation of the full suite of materials we might wish to measure using the rover’s instrument package. What combination of processes, for example, maximizes the probability of preserving organic molecules, body fossils and sedimentary textures?

If we can understand the processes that facilitate preservation, then we can conceive of the likely operation of those processes in time and space. That is, we can define a taphonomic window – the sedimentary and diagenetic circumstances most conducive to preservation. We know that ancient organic molecules will preserve only if buried in sediments and long-term preservation will occur only if buried organics are shielded from oxidizing fluids. On Mars, surface oxidants and UV radiation will have altered or destroyed organic molecules (Benner et al., 2000; Navarro-González et al., 2006; Sumner, 2006). On Mars, surface oxidants and UV radiation will have altered or destroyed organic molecules at or near the surface (Benner et al., 2000; Navarro-González et al., 2006; Sumner, 2006). If we can identify generic features that can be used to characterize potential landing sites with respect to the presence or absence of specific taphonomic windows, then we can help maximize the chances of successful analyses by MSL.

### **Working Group Findings**

A recent review of biosignatures and strategies for their use in life detection can be found in Botta et al., (2008). Our deliberations yielded the ranked order of biosignatures and taphonomic windows that are provided in Tables 1 & 2. However, we also recognized that, even on Earth with its vigorous and multi-billion-year-old biosphere, many of these biosignatures can be ambiguous and are preserved only under rare and exceptional circumstances.

Insert Table 1 hereabouts

Insert Table 2 hereabouts

**Box 1: Biosignatures at a glance – summary of what can be observed with the payload elements (PE) of MSL. Note that environments can be reconstructed from physical and chemical features of ancient sediments that are not considered to be biomarkers. (see tables 1 & 2)**

1	Organism Morphologies (cells, body fossils, casts) PE: MAHLI Minimum size would have to be greater than 100 µm and rock preparation techniques are not available to expose organisms within rock. Martian life is expected to be microbial, so the probability of detection is low Potential as a biosignature: exceptionally high Potential as an environmental indicator: low
2	Biofabrics (including microbial mats) PE: MAHLI, MastCam Accreted structures analogous to those on Earth are detectable; However, few bedding plane surfaces are exposed, so potential surface biosignatures will be difficult to detect Potential as a biosignature: moderate Potential as an environmental indicator: low
3	Diagnostic organic molecules; Organic carbon PE: SAM, ±ChemCam only if very abundant. Detection potential high incl. atmospheric gases Potential as a biosignature: exceptionally high Potential as an environmental indicator: high
4	Isotopic Signatures PE: SAM Contextual knowledge is essential; results can be ambiguous and complex to interpret Potential as a biosignature: moderate Potential as an environmental indicator: low
5	Biom mineralization & bioalteration PE: CheMin, ±MAHLI, ±SAM, ±APXS detection of specific minerals is good morphological pattern may be useful but needs very fine spatial resolution Potential as a biosignature: low Potential as an environmental indicator: low
6	Spatial patterns in chemistry PE: SAM, CheMin, ±ChemCam if very abundant C, N, S elemental distributions; Detection potential on cm scale to facies scale Potential as a biosignature: low on its own Potential as an environmental indicator: low
7	Biogenic Gases (Non-equilibrium) PE: SAM Excellent capacity to detect gases Potential as a biosignature: high (e.g., CH <sub>4</sub> ) Potential as an environmental indicator: low

Site selection decision-making and landed operations of the MSL should support the search for all of the above-mentioned biosignatures. However, it is also clear that accumulations of organic matter above meteoritic background levels would be amongst the most easily detected and least ambiguous. If life ever existed on Mars, it was likely microbial and existed communally in fluids and/or sediments, either free-living or as biofilms. Biomass from such communities, whatever their physiological characteristics, might then be preserved in the Martian sedimentary rock record in sufficient concentration to be detectable with MSL payload elements. Notwithstanding the numerous possibilities for habitable niches on Mars, a first-order approach to evaluating the organic record there would be to seek evidence of sedimentary environments -- the preferred geological setting for study - - that may have elevated concentrations of biologically derived carbonaceous materials (Table 3). These may include standing water environments such as deltaic systems, that promote retention of *in situ* organics and environments that hydrodynamically concentrate organics. An alternative approach would be to seek concentrations of buried crystalline minerals, such as

clays and evaporites which may protect organic matter from the destructive effects of ionizing radiation and strong oxidants. We also identify and order a range of other possibilities for environments conducive to the preservation of the signs of former and extant life (Tables 3 and 4).

### MSL's Unique Combination of Capabilities



The MSL rover houses a remarkable suite of analytical instrumentation within a mobile platform—a distinct advantage for field investigations. Once on the surface of Mars, this rover will support several years of exploration and sample analysis. The analysis of fossil organic matter entrained in sedimentary rocks requires a proven approach, which can be implemented by the SAM instrument. Extensive studies of fossil organic remains in terrestrial rocks demonstrate that in most cases macromolecular constituents make up the overwhelming bulk of

**Box 2: Molecular biosignatures at a glance**

- Enantiomeric excess  
Homochirality characterizes terrestrial biochemicals; strong preference for one enantiomer over the other in a chiral molecule. Interpretation is complicated by the discovery of L- excess in meteoritic amino acids (Engel & Macko 1997, Cronin & Pizzarello 1997, Pizzarello 2006, Glavin & Dworkin 2009)
- Diastereoisomeric preference  
Strong isomer preferences in molecules with more than one center of asymmetry (Summons et al., 2007)
- Structural isomer preference  
Observing a limited subset of the possible stable structural isomers in a complex molecule (Summons et al., 2007)
- Repeating constitutional sub-units or atomic ratios  
Signifies complex molecules constructed from small common building blocks as in terrestrial biochemistry (McKay, 2004; 2007; Summons et al., 2007)
- Systematic isotopic ordering at molecular and intramolecular levels  
As above
- Uneven distribution patterns or clusters (e.g., C-number, concentration,  $\delta^{13}\text{C}$ ) of structurally related compounds.  
As above

geologically preserved organic material. Pyrolytic breakdown of biopolymeric material (or diagenetically modified biopolymeric material) into small, volatile and identifiable fragments in a specially engineered device at the front end of a gas chromatograph-mass spectrometer (GC-MS) is a proven and manageable analytical approach because it can provide a molecular fingerprint of fossil organisms, in many cases yielding critical information regarding the identity of the organism through the presence of characteristic molecular biomarkers. Some generic characteristics of biomarkers are summarized in Box 2. A combination of these features should be present in biomolecules, whatever their

origin, and most of them are detectable with the SAM instrument. It is noteworthy that interstellar organic matter that may also be present in certain sedimentary rocks on Mars, also is amenable to characterization via pyrolysis GC-MS and may be distinguished from biologically derived material through specific kinds of molecular distributions (Sephton & Botta, 2008 and references therein).

1  
2  
3 Aside from MSL's capabilities to provide high-resolution organic molecular information, the  
4 addition of CheMin and ChemCam to the traditional APXS broadens the capability to search for  
5 ancient life on Mars. The ability to determine quantitatively the major mineral and chemical  
6 constituents of prospective rocks will no doubt aid in assessing whether Mars had an active  
7 biosphere at a much earlier time in its history. In terrestrial rocks, the presence of biological  
8 organic matter can effect changes in local mineralogy by a number of different processes.  
9  
10

11 In some cases even if the organic matter has ultimately been destroyed through long-term  
12 exposure to oxidizing fluids, specific mineral associations with the now-absent organics may  
13 remain, providing a more persistent biomarker. Finally, certain minerals that exist on Earth on a  
14 global scale are generally recognized to be the consequence of an active biosphere, e.g.,  
15 biogenic carbonates, hazenite ( $\text{KNaMg}_2(\text{PO}_4)_2 \cdot 14\text{H}_2\text{O}$ ), or brushite ( $\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$ ). Hazen et  
16 al. (2008) list and describe a large variety of mineral species that can be produced by biological  
17 processes. The presence of CheMin in the analytical suite of MSL is, therefore, of profound  
18 importance for the detection of ancient life.  
19  
20  
21

22 The essential factor in addressing whether organic matter may be preserved in the Martian  
23 sedimentary record is the ability to access appropriate lithologies (Tables 3 & 4). In this regard,  
24 mobility is critical. As the MER mission has clearly demonstrated, only an "eyes-on-the-ground"  
25 approach can provide the information required to interpret the lithologies of outcrops.  
26 Integrating this mobile capability with analytical instrumentation provides the means to address  
27 whether organic matter preservation has occurred or not.  
28  
29  
30  
31

32  
33 Insert tables 3 and 4 hereabouts  
34  
35

### 36 **Brief description of the payload elements**

37  
38 Full descriptions of the MSL payload instruments can be found at: [http://msl-  
39 scicorner.jpl.nasa.gov/Instruments/](http://msl-scicorner.jpl.nasa.gov/Instruments/). In addition to the suite of analytical tools there are four  
40 cameras. The Mast Camera (MastCam), the Mars Hand Lens Imager (MAHLI), the Remote  
41 Micro-Imager on ChemCam (RMI) and the Mars Descent Imager (MARDI).  
42  
43

44 MastCam will take color images and color video footage of the Martian terrain. These can be  
45 stitched together to create panoramas of the landscape around the rover. MastCam consists of  
46 two camera systems mounted on a mast extending upward from the rover deck and will be used  
47 to study the Martian landscape, rocks, and soils; to view frost and weather phenomena; and to  
48 support the driving and sampling operations of the rover.  
49  
50

51 MAHLI will provide close-up views of the minerals, textures, and structures in Martian rocks and  
52 the surface layer of rocky debris and dust. The self-focusing, roughly 4-centimeter-wide (1.5-  
53 inch-wide) camera will take color images of features as small as 14 micrometers and will carry  
54 both white light and ultraviolet light sources making the imager functional both day and night.  
55 The ultraviolet light will be used to induce fluorescence to help detect carbonate and evaporite  
56 minerals, both of which would indicate that water helped shape the landscape on Mars.  
57  
58  
59  
60

1  
2  
3 RMI will provide telescopic views of the surfaces to be analyzed by LIBS, and will put LIBS  
4 analyses into their geologic context. However, the RMI can also be used to image textures  
5 whether or not the LIBS is used. The RMI has a field of view of 19 milliradians. Due to  
6 optimization of the telescope for LIBS, the RMI resolution is not pixel-limited, and is  
7 approximately 80 microradians. The RMI can clearly distinguish the submillimeter LIBS spot on  
8 a metal plate at a distance of at least 10 m. Therefore, this camera has the ability to make  
9 discoveries of texturally-based potential biosignatures, such as stromatolites and other textures  
10 suggestive of former microbial processes.  
11  
12

13  
14 MARDI will take color video during the rover's descent toward the surface, providing an  
15 "astronaut's view" of the local environment. As soon as the rover jettisons its heat shield several  
16 kilometers above the surface, the Mars Descent Imager will begin producing a five-frames-per-  
17 second video stream of high-resolution, overhead views of the landing site. It will continue  
18 acquiring images until the rover lands, storing the video data in digital memory. After landing on  
19 Mars, the rover will transfer the data to Earth. MARDI will provide information about the larger  
20 geologic context surrounding the landing site.  
21  
22

23  
24 ChemCam combines laser-induced breakdown spectroscopy with a remote micro-imager that  
25 provides images of the target. It will provide elemental analysis of spatially resolved solid  
26 samples at distances of 1-9 m. ChemCam's primary objective is to determine the chemical  
27 composition of rocks and regolith in order to characterize the materials in the vicinity of the  
28 rover.  
29  
30

31 The Alpha Particle X-Ray Spectrometer (APXS) will measure the abundance of chemical  
32 elements in rocks and soils. It will be placed in contact with rock and soil samples on Mars and  
33 will expose the material to alpha particles and X-rays emitted during the radioactive decay of  
34 curium. The MER rovers have used APXS successfully, but the greatest benefit of APXS will be  
35 when it is integrated with mineralogical analyses made by CheMin.  
36  
37

38 CheMin is a mineralogy instrument that will unequivocally identify and quantify the minerals  
39 present in rocks and soil delivered to it by the Sample Acquisition, Sample Processing and  
40 Handling (SA/SPaH) system. By determining the mineralogy of rocks and soils, CheMin will  
41 enable assessments of the involvement of water in their formation, deposition, or alteration.  
42 CheMin data will also be useful in the search for potential mineral biosignatures, energy sources  
43 for life, or indicators of past habitable environments. CheMin can unequivocally identify and  
44 quantify minerals above its detection limits in complex natural samples such as basalts, multi-  
45 component evaporite systems, and soils.  
46  
47

48  
49 The Sample Analysis at Mars (SAM) Suite Investigation in the MSL Analytical Laboratory is  
50 designed to address the present and past habitability of Mars by exploring molecular and  
51 elemental chemistry relevant to life. SAM evaluates carbon chemistry through a search for  
52 organic compounds, the chemical state of light elements other than carbon, and isotopic tracers  
53 of planetary change. SAM is a suite of three instruments, a Quadrupole Mass Spectrometer  
54 (QMS), a Gas Chromatograph (GC), and a Tunable Laser Spectrometer (TLS). The QMS and  
55 the GC can operate together in a GC-MS mode for chromatographic separation and mass  
56 spectral identification of organic compounds. The TLS obtains precise stable isotope ratios for C  
57  
58  
59  
60

1  
2  
3 and O in carbon dioxide as well as C isotopes and abundance of trace methane.  
4

5 The Radiation Assessment Detector (RAD) will detect and analyze the most hazardous  
6 energetic particle radiation on the surface of Mars. Characterizing and understanding the  
7 radiation environment on the Martian surface is fundamental to quantitatively assessing the  
8 habitability of the planet (both past and present) and is essential for future manned Mars  
9 missions. RAD will address radiation effects on biological potential and past habitability, as well  
10 the contribution to chemical alteration of the regolith due to impinging space radiation.  
11  
12

13 REMS is designed to record six atmospheric parameters: wind speed/direction, pressure,  
14 relative humidity, air temperature, ground temperature, and ultraviolet radiation. All sensors are  
15 located around three elements: two booms attached to the rover Remote Sensing Mast (RSM),  
16 the Ultraviolet Sensor (UVS) assembly located on the rover top deck, and the Instrument  
17 Control Unit (ICU) inside the rover body.  
18  
19

20 The Dynamic Albedo of Neutrons (DAN) instrument is an active/passive neutron spectrometer  
21 that measures the abundance and depth distribution of H- and OH-bearing materials (e.g.,  
22 adsorbed water, hydrated minerals) in a shallow layer (0.5-1 m) of Mars' subsurface along the  
23 path of the MSL rover.  
24  
25

26 The Sample Acquisition/Sample Processing and Handling subsystem (SA/SPaH) for MSL  
27 consists of a powdering drill (Powder Acquisition Drill System or PADS) together with a  
28 scooping, sieving, and portioning device (Collection and Handling for Interior Martian Rock  
29 Analysis or CHIMRA) mounted on a turret at the end of a robotic arm. There is also a dust  
30 removal tool (DRT) for clearing the surface of rocks prior to sampling. The drill enables powder  
31 to be acquired from depths of 20 to 50 mm over a wide range of rock hardness with the top ~15-  
32 20 mm being discarded. The scoop also enables samples of soil to be acquired, sieved and  
33 apportioned. Five bricks of a silicon dioxide ceramic organic check material (OCM) are mounted  
34 in canisters on the front of the rover to help assess end-to-end sample handling and potential  
35 organic contamination at different times during the mission. Each brick, which is sealed under  
36 vacuum in its own canister, can be drilled, sieved and portioned in CHIMRA. The powder is then  
37 delivered to SAM or CheMin following the same pathway as for Martian rock samples. The  
38 bricks are doped with traces of non-natural volatile fluorinated compounds. The sampling tools  
39 and protocols are subject to equivalent and, as far as is known, benign impacts on biosignature  
40 integrity as those used in terrestrial laboratories.  
41  
42  
43  
44  
45

46 These instruments have been developed specifically for the MSL mission. Many aspects of their  
47 specifications and ultimate performance, individually or in concert, are still being evaluated.  
48

#### 49 **Brief synopsis of environmental and physical features detectable with MSL payload**

50 In the MSL payload, SAM can detect traces of organic matter in rocks and sediment. It does so  
51 in a variety of ways—from bulk organic carbon detection (by the difference between combusted  
52 total carbon and inorganic carbon) or non-specific molecular detection by evolved gas analysis  
53 (Bibring et al., 2005) to gas chromatographic (GC) separation and structural identification of  
54 discrete molecules volatilized directly or after reactions with chemical derivatizing reagents that  
55  
56  
57  
58  
59  
60

1  
2  
3 enhance volatility and thermal stability (Tables 2, 5-8). It also detects traces of volatile organics  
4 in the atmosphere, such as C1-C6 hydrocarbons and other gases, and it can precisely measure  
5 carbon isotopic composition of methane and carbon dioxide.  
6

7  
8 The camera systems are designed document color and grain size variations in rocks and  
9 sediments that reflect sedimentary structures, mineral growth processes, weathering and  
10 biofabrics, if present (see Noffke, 2009; Tables 9-12). These features provide essential data for  
11 interpreting the processes forming the rocks as well as their alteration. Images are essential for  
12 sample selection in addition to providing a context for chemical analyses.  
13

14  
15 ChemCam can remotely detect carbon in soils and rocks (>1% carbon subject to final testing  
16 and calibration) but the ChemCam does not directly discriminate between organic and inorganic  
17 carbon (Tables 5 & 6). This is in contrast to the SAM capability of detecting sub ppb levels of  
18 organic compounds that are <535 dalton via Evolved Gas Analysis (EGA) and pyrolysis.  
19

20  
21 CheMin in XRD mode cannot detect organic carbon, but it can detect and identify any crystalline  
22 inorganic hosts of carbon and sulfur, for example (Tables 11 & 12). Used in conjunction with  
23 SAM, it can be particularly useful in providing more accurate discrimination between organic  
24 and inorganic hosts. Thus, the MSL payload is capable of mapping the distribution of organic  
25 carbon and its molecular composition in rocks and sediments in stratigraphic, geomorphic,  
26 and/or chronologic context. This distribution, the types of molecular structures detected, their  
27 redox state, polarity, volatility, and their relationship to other elements and minerals detected by  
28 other MSL payload instruments would provide critical insight to deciphering biosignatures, other  
29 organic matter sources (abiogenic and meteoritic), environmental records, surface processes  
30 and carbon cycling.  
31  
32

33  
34 Insert Tables 4 and 5 hereabouts  
35

### 36 **Carbon isotopes as potential biosignatures on Mars**

37  
38 The patterns of isotopic composition ( $^{13}\text{C}/^{12}\text{C}$  ratios) of Martian crustal and atmospheric  
39 constituents reflect the compositions of their original sources as well as any isotopic  
40 discrimination associated with the network of physical, chemical (and biological?) processes  
41 that created and cycled these constituents (Tables 7 & 8). Fig. 1 depicts the terrestrial C cycle  
42 consisting of C reservoirs (boxes) and processes (arrows) in the atmosphere, crust, and interior.  
43 The caption for Fig. 1 addresses these processes. An illustration of the S cycle would be  
44 analogous to Fig. 1, except sulfide and sulfate reservoirs would be substituted for organic and  
45 inorganic C reservoirs, respectively. The figure is included to illustrate the complexity of known  
46 reservoirs, the timescales over which they interact and range of isotopic values on Earth. This  
47 picture has only emerged gradually over many years through extensive study through direct  
48 observation of the chemical, biological, isotopic and geological aspects of carbon cycling (e.g.  
49 see Des Marais, 2001).  
50  
51  
52  
53  
54  
55

56 Insert Figure 1 Hereabouts  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Insert Table 7 hereabouts

Overall, the terrestrial C-cycle consists of multiple nested cyclic pathways that differ with respect to reservoirs, processes and timescales. Carbon is exchanged between the atmosphere and shallow crust by processes which act on relatively short timescales. Carbon deeper within the crust is cycled more slowly by processes of burial and exhumation under tectonic control. These processes are now probably negligible due to the presumed absence of subduction on Mars. Processes which exchange carbon between the crust and atmosphere of Mars might be somewhat active as indicated by the recent indications of methane in the Martian atmosphere (Mumma et al., 2009). This methane almost certainly has a subsurface source that very likely involves aqueous processes, possibly including life, which we discuss further below.

The  $^{13}\text{C}/^{12}\text{C}$  ratios of the carbon reservoirs reflect isotopic discrimination associated with the above processes. On Earth, isotopic discrimination associated with organic biosynthesis has been principally responsible for determining the  $^{13}\text{C}/^{12}\text{C}$  ratios observed in organic and inorganic crustal reservoirs. Biological processes can change these values by several percent (e.g., Des Marais, 2001). Accordingly  $^{13}\text{C}/^{12}\text{C}$  ratios might serve as biosignatures of any past or present life on Mars if key components of the C-cycling system can be constrained. However Jakosky et al. (1994) calculated that processes that caused loss of C to space were isotopically selective and increased by several percent the  $^{13}\text{C}/^{12}\text{C}$  ratios of the remaining atmospheric and near-surface crustal C reservoirs. These increases might equal or exceed, and thus obscure, any changes in  $^{13}\text{C}/^{12}\text{C}$  ratios due to biological processes. Note, as well, that abiotic processes of organic synthesis involve C-isotopic fractionation, further complicating the isotopic interpretation of any organic compounds that may be detected (Chang et al., 1983). To the extent that carbon situated in deeper interior reservoirs was isolated from these atmospheric escape processes, its  $^{13}\text{C}/^{12}\text{C}$  value would be less affected and therefore lower, and perhaps most closely reflect initial Martian  $^{13}\text{C}/^{12}\text{C}$  ratios. Consequently we cannot be entirely sure whether any measured  $^{13}\text{C}/^{12}\text{C}$  patterns might indicate life or whether they reflect principally the effects of atmospheric escape or other environmentally sensitive equilibrium fractionation processes. Another difficult challenge for interpretation of C-isotopes on Mars is establishing whether organic matter and any inorganic carbon in a given sample have a genetic relationship. If, for example, the OM is meteoritic and carbonate originated hydrothermally, an isotopic separation is likely to be meaningless.

Precise and accurate carbon isotopic compositions of methane and carbon dioxide from the TLS in SAM may provide important benchmarks for understanding planetary-scale carbon cycling on Mars. For atmospheric measurements these experiments are rapid and not resource intensive and, thus, could be included in a regular sampling scheme. Detection and quantification of atmospheric methane by MSL would provide an important verification of the Earth-based and orbital spectroscopic detections of methane at ppb levels (Formisano et al., 2004, Krasnopolsky et al., 2005; Mumma et al., 2009). Temporal variations in methane concentration would be detectable by MSL and could potentially distinguish between episodic



1  
2  
3 release of subsurface methane or methane destruction from photolytic, oxidative or dust  
4 reactions in the modern environment (Atreya et al., 2006). Spatial variations in methane  
5 concentration are not expected to be observable unless MSL is proximal to a subsurface  
6 hydrothermal, volcanic or unstable hydrate point source. Using C-isotopic data to constrain the  
7 origins of atmospheric methane on Mars, however, presents a formidable problem. Abiogenic  
8 methane production associated with serpentinization, that is, aqueous alteration of olivine-  
9 and/or pyroxene-rich rocks, is a process likely to have been prevalent early in Mars history and  
10 which feasibly continues today (Oze and Sharma, 2005). Multiple carbon and hydrogen isotopic  
11 data suggest the methane that is abundant in the fluids emanating from the Lost City  
12 hydrothermal field, a site of contemporary serpentinization on Earth, has an abiogenic origin.  
13 However, it is known that methanogenic Archaea are also active in this system (Bradley et al.,  
14 2009). Accordingly, carbon isotopes are of limited value in discriminating between biological  
15 and abiogenic sources where multiple processes can contribute to a pool of methane and  
16 reservoir affects provide added complications (Bradley and Summons, 2010). This is the  
17 situation on Earth where the biochemistry of methanogenesis is reasonably well understood and  
18 even more uncertainly would accompany methane measurements on Mars. Carbon isotopic  
19 data might discriminate between abiogenic and biogenic methane sources only if the  
20 fractionation between methane and co-existing CO<sub>2</sub> were well in excess of the equilibrium  
21 values modeled for feasible P-T conditions, thereby implying kinetic (i.e., enzymatic) control on  
22 methanogenesis.  
23  
24  
25  
26  
27  
28

29 Variations in C-isotopic compositions of carbon dioxide are less likely as it is the dominant  
30 atmospheric species. SAM design also enables isotopic measurements of trace amounts of bulk  
31 inorganic and organic carbon by comparing CO<sub>2</sub> evolved under inert conditions with CO<sub>2</sub>  
32 generated from combustion with O<sub>2</sub>. Mapping variations in these isotopic compositions could  
33 further our understanding of redox processes, isotopic fractionation pathways (including  
34 biosignatures, abiogenic processes, and possibly meteoritic contributions), and environmental  
35 carbon cycling (Table 8). The TLS can also measure oxygen isotopes of carbon dioxide and  
36 deuterium/hydrogen in water, which may indicate the effects of surface and atmospheric cycling.  
37 Oxygen isotopes of carbon dioxide evolved from carbonates in SAM will help with assessment  
38 of post-depositional alteration by subsurface fluids.  
39  
40  
41

### 42 **Sulfur isotopes as potential biosignatures on Mars**

44 Observed <sup>34</sup>S/<sup>32</sup>S, and <sup>33</sup>S/<sup>32</sup>S values presumably reflect processing by redox reactions in the  
45 atmosphere (e.g. photochemistry, see Farquhar et al., 2000) and crust (e.g. weathering,  
46 hydrothermal and life). On Earth, microbial reduction of sulfate and sulfur disproportionation  
47 reactions occurring at ambient temperatures have created large <sup>34</sup>S/<sup>32</sup>S differences (several  
48 percent) between oxidized and reduced sulfur reservoirs (Canfield, 2001). Because  
49 microorganisms are required to catalyze S-isotopic exchange reactions having significant  
50 fractionations at low ambient temperatures, large <sup>34</sup>S/<sup>32</sup>S differences in sedimentary rocks can  
51 be reliably interpreted as biosignatures in many cases. However, a careful assessment of the  
52 original environment of deposition is always essential. Accordingly, before stable isotopic  
53 patterns can serve as potential biosignatures on Mars, we must characterize the isotopic  
54 composition of major sulfur reservoirs and also understand more fully the consequences of key  
55  
56  
57  
58  
59  
60

1  
2  
3 non-biological processes.  
4

5 Tests on a SAM breadboard QMS instrument have shown that sulfur isotope measurements of  
6 sulfur dioxide evolved from the thermal degradation of sulfate minerals are possible (Franz et  
7 al., 2007). However, these measurements are complicated by overlapping spectra and the  
8 evolution of gas from multiple sulfur species. Unlike the sub-per-mil precision from Earth-based  
9 instrumentation, the SAM QMS may provide percent precision. These data may still be valuable  
10 on Mars, especially if basaltic sulfur, which shows <1% variability on Earth and meteorites,  
11 provides a reference point for comparing sedimentary and hydrothermal sulfur analyses (Table  
12 8). On Earth, sedimentary sulfides show 4% variation, reflecting biological cycling of S, a  
13 sensitive redox element (Canfield, 2001; Farquhar et al., 2003). If a similar record were to exist  
14 on Mars, sulfur isotopic detection by the SAM QMS may detect it. Sulfur isotopic compositions  
15 of sulfate and reduced-sulfur minerals may also provide valuable information of environmental  
16 sulfur cycling by abiogenic hydrothermal (Rye, 2005; Greenwood et al., 2000) and atmospheric  
17 processes (Farquhar et al., 2000).  
18  
19  
20  
21

### 22 **Other isotopic data**

23  
24 Noble gas isotope measurements and elemental ratios are regarded as isotopic tracers of  
25 mantle and atmospheric evolution (Swindle, 2002) and possible comet-borne contributions  
26 (Owen et al., 1992). Although they do not provide direct insight into environmental conditions  
27 preserved in the sedimentary record, they can supplement the larger context of Mars chemical  
28 and environmental evolution. These data may help explain other isotopic records.  
29  
30

### 31 **Environmental conditions**

32  
33 Sedimentary and other near-surface materials can serve as recorders of environmental  
34 conditions that prevailed during previous epochs of Martian history. Insights into ancient  
35 environmental conditions might help to identify potential processes responsible for setting the  
36 observed isotopic patterns. Paleotemperatures often can be inferred by measuring stable  
37 isotopic compositions of pairs of minerals or fluids that equilibrated isotopically when they  
38 formed. Oxygen isotopes have been utilized most frequently to infer paleotemperatures within  
39 the habitable range (<120°C). Elevated temperatures have been inferred using sulfur (>150°C)  
40 and carbon (>300°C) isotopes. Stable isotopic compositions of preserved minerals can also help  
41 to elucidate the nature of fluids associated with their formation. For example, elevated <sup>18</sup>O/<sup>16</sup>O  
42 values might indicate the former presence of brines in evaporitic environments. Many additional  
43 examples can be cited where stable isotopic patterns have helped to constrain  
44 paleoenvironmental conditions.  
45  
46  
47  
48

49 Insert Table 8 hereabouts  
50

### 51 **Preservation potential**

52  
53 To the extent that sediments, cements, and other surface materials have escaped alteration  
54 subsequent to their formation, they can preserve information about earlier environmental  
55 conditions and, potentially, biosignatures (Tables 9 & 10). Preservation can be compromised by  
56 weathering and erosion or by alteration *in situ* by oxidation and migrating fluids, for example.  
57  
58  
59  
60

1  
2  
3 The following minerals can isolate and preserve biosignatures (listed in order of increasing  
4 crustal residence times on Earth): ice, halite, sulfates (e.g. Aubrey et al., 2005; Panieri et al.,  
5 2010), carbonates (e.g. Birgel et al., 2008), phyllosilicates (e.g. Hedges and Keil, 1995;  
6 Butterfield, 1990), silica (e.g. Knoll, 1985), hematite (Fernandez Remolar and Knoll, 2008) and  
7 phosphates (e.g. Xiao & Knoll, 1999; Farmer & Des Marais, 1999). For example, carbonates  
8 deposited as a consequence of microbial metabolisms sometimes hold an excellent record of  
9 those processes as is the case with methane seep limestones (e.g. Birgel et al., 2008) or  
10 hydrothermal systems like Lost City (Bradley et al., 2009). Silica-rich water derived from  
11 hydrothermal systems is another well-established medium that promotes faithful preservation  
12 (e.g. Knoll et al., 1985; Trewin, 1996). In fact, Walter (1996) has identified numerous fossil  
13 hydrothermal systems on Earth, both terrestrial and marine, that are known and potential  
14 repositories of paleobiological information. Preservation is optimized when temperatures remain  
15 low and mineral matrices form during sedimentation/precipitation and reduce the permeability of  
16 the sediments to near zero. Detailed  $^{18}\text{O}/^{16}\text{O}$  values of some of these minerals can help to  
17 assess the extent to which invading fluids caused post-depositional alteration.

### 22 Hydrated mineral phases

23  
24  
25 The presence of hydrated minerals reflects specific chemical conditions, including the activity of  
26 water (Table 12). Specific implications of different hydrous phases vary with the composition  
27 and context of the minerals. Some, such as hydrous Mg-sulfates, require low temperatures  
28 and/or substantial humidity to remain stable. Their hydration states reflect local current  
29 conditions due to their rapid dehydration kinetics, although their formation may reflect older  
30 conditions. Other hydrous minerals, such as the clay minerals, remain metastable for long  
31 periods of time and provide a record of past hydrous activity. Extracting the history of water on  
32 Mars requires careful characterization of hydrous minerals within their environmental context.

### 36 Textural features preserved in sediments and hydrothermal systems

37  
38 Physical, chemical and biological processes all influence the preservation of biosignatures in  
39 hydrothermal and sedimentary systems on Earth, and we can use our understanding of these  
40 processes to predict their impact on possible biosignature preservation on Mars (Tables 9 & 10).

41  
42 Purely physical and chemical processes should be comparable on Earth and Mars. For  
43 subsurface, hydrothermal and sedimentary systems, physical and chemical processes can  
44 provide substantial insights into the history and habitability of the system. For example, in  
45 sedimentary systems, physical processes such as sediment transport produce structures that  
46 are characteristic of specific processes, e.g., types of flows. These can be used to interpret  
47 processes in the depositional environment, providing invaluable constraints on habitability and  
48 guiding the search for biomarkers. There are some non-negligible differences due to the lower  
49 g of Mars, however, these do not affect the overall interpretation of transport-related suites of  
50 sedimentary structures (Grotzinger et al., 2005). Similarly, chemical processes leave distinctive  
51 signatures, whether they are active in the depositional environment or within the rock. They can  
52 be used to evaluate the habitability of the environment and rocks at different points in time,  
53 including the potential for subsurface colonization of rocks of volcanic, hydrothermal or  
54 sedimentary origins.

1  
2  
3 Insert tables 9 & 10 hereabouts  
4  
5  
6  
7

8 On Earth, biological processes are very active in almost all sedimentary and lower-temperature  
9 hydrothermal systems. Biological processes in terrestrial hydrothermal systems provide a  
10 diverse suite of potential biomarkers, including concentrations of organics and minerals that  
11 reflect mats, stromatolites, roll-ups, biofilms, streamers, etc. (Tables 9 & 11). Concentrations of  
12 elements and high concentrations of migrated organics are also characteristics of some  
13 terrestrial hydrothermal systems. Preservation of these indicators of biological activity is  
14 strongly affected by physical and chemical processes. The high temperatures and abundant  
15 water flow tend to degrade biomarkers. Hydrothermal systems commonly experience intense  
16 brecciation and fracturing due to high-pressure fluids. This type of fracturing leaves distinctive  
17 textural features that are best avoided when looking for a good biosignature preservation  
18 window. Chemical processes also affect preservation. Mineral precipitation can entomb  
19 organics and biological processes can influence the distribution of minerals. In hydrothermal  
20 settings, a good preservation window is created by precipitation of non-redox sensitive, low-  
21 porosity minerals. However, minerals can also recrystallize and dissolve, particularly when  
22 exposed to high-temperature fluids, making preservation of biosignatures less likely. When  
23 looking for a good preservation window, one wants to avoid recrystallized areas, those that have  
24 experienced intense oxidation, and those exposed to high temperatures or ionizing radiation  
25 (e.g. Dahl et al., 1988).  
26  
27  
28  
29  
30

31 Sedimentary systems on Earth also have substantial potential for preserving biosignatures  
32 because of the intensity of biological activity, and their natural tendency to concentrate or high-  
33 grade organic matter across hydraulic gradients. Again, these terrestrial processes serve as  
34 models for predicting good preservation windows on Mars (Tables 10 & 12). Biomass  
35 accumulation in terrestrial systems is reflected in mats, stromatolites, roll-ups, wrinkle  
36 structures, etc. For preservation, these accumulations need to incorporate sediment or be  
37 mineralized prior to degradation of the organics. Biological processes can also create fenestrae  
38 (gas-produced pores), affect grain sizes, and influence elemental concentrations. These  
39 features can be well preserved if they are lithified early and do not experience significant  
40 alteration. Physical processes such as dewatering, hydration changes and structural  
41 deformation can destroy these signatures. Similarly, chemical processes such as  
42 recrystallization, redox changes and metamorphism destroy biomarkers. Thus, for biomarker  
43 preservation, one should identify areas with an appropriate sedimentary environment that was  
44 lithified early and experienced minimal post-depositional alteration (Farmer and Des Marais,  
45 1999).  
46  
47  
48  
49

### 50 **Elemental concentrations and mineral distributions in hydrothermal and deep subsurface** 51 **systems** 52

53 Hydrothermal environments can preserve potential biosignatures in the form of elemental  
54 concentrations and mineral distributions (Table 11). Thermally driven aqueous convection can  
55 significantly alter environmental conditions through selective mineral dissolution, alteration, and  
56 precipitation; element leaching and subsequent transport; and oxidation and reduction (redox)  
57  
58  
59  
60

1  
2  
3 chemistry. On Earth, each of these processes can occur with or without biological mediation,  
4 but through careful analysis with multi-faceted approaches, the role of biology can often be  
5 assessed.  
6

7  
8 If life emerged on Mars and prospered in hydrothermal systems, these alteration processes  
9 likely include biotic and abiotic components (Tables 9 & 11). Each yields an array of features  
10 including specific mineralogy that is characteristic of certain hydrothermal conditions and  
11 chemistries, element gradients and zonation within those minerals and spatial distribution of  
12 alteration minerals and precipitates at scales from  $\mu\text{m}$  to km (Table 11). The identification and  
13 chemical analysis of chlorite, amorphous silica or quartz, sulfide minerals, kaolinite and other  
14 clays, hematite and other ferric (hydr)oxides, carbonates, and sulfates can be used to determine  
15 formation temperatures, redox conditions, and pH. For example, specific sulfides (PbS, ZnS,  
16 CuS) and sulfates ( $\text{BaSO}_4$ ,  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) or mineral assemblages (e.g., sulfate and/or sulfide  
17 together with saddle dolomite) can point to specific formation temperatures. The presence of  
18 jarosite, alunite, or kaolinite indicate low-pH environments, and discrimination between  
19 crystalline silica minerals (e.g., quartz or cristobalite) and non-crystalline silica phases, such as  
20 opal-CT, is most effective when chemical data are combined with mineralogical information  
21 (e.g., XRD data). Other minerals bearing S, Fe, Mn, U, As or other redox-sensitive elements  
22 can provide further constraints on Eh and, if present above the minimum detection limits for  
23 CheMin, could be detected by XRD. All of these features would help constrain the possibility,  
24 probability, and physiology of potential life forms on Mars. However, because hydrothermal  
25 systems are dynamic, high-energy environments, the preservation of labile features (e.g.  
26 organic matter, amorphous solids) would undoubtedly require encasement or entombment in  
27 protective minerals or other removal from the destructive forces of oxidation, metamorphism,  
28 and continued hydrothermal activity.  
29  
30  
31  
32  
33  
34  
35  
36

37 Insert Table 11 hereabouts

### 38 39 **Elemental concentrations and mineral distributions in sedimentary systems**

40  
41 Sedimentary environments present both similarities and differences to the hydrothermal case  
42 highlighted above. The preservation potential of environmental signatures is determined  
43 principally by the sedimentary material, the chemical composition and flux of the aqueous  
44 solutions responsible for the sediment transport, the redox state and pH of that solution, and  
45 any post-depositional chemical processes (including putative microbe-mineral interactions) that  
46 may have operated (Table 12). The most informative environmental signatures include  
47 sedimentary structures and redox-sensitive (e.g., S, Fe, and Mn-bearing) and pH-sensitive (e.g.,  
48 jarosite, alunite, kaolinite, carbonates) mineral assemblages and abundances. Many  
49 sedimentary environments on Earth or Mars also host evaporite minerals (including sulfates and  
50 chlorides) and corresponding trace element distributions, as well as oxides, carbonates, sulfides  
51 and, perhaps, phosphate precipitates resulting from microbe-mineral interactions. The  
52 preservation potential of these environmental features is particularly enhanced by early  
53 lithification of the sediments. In addition, the greatest scientific return would likely come from  
54 sedimentary systems, lacustrine or marine, that have not been subjected to significant  
55  
56  
57  
58  
59  
60



1  
2  
3 recrystallization, prolonged strong oxidizing conditions, dissolution and solute removal, major  
4 structural alteration, or significant thermal metamorphism. As closed basins, lakes in particular,  
5 represent terminal receptors for both primary and transported organics together with the clays,  
6 which preserve OM due to strong absorption capacity, low reactivity, and low permeability when  
7 compacted (Farmer and Des Marais, 1999; Meyers and Ishiwatari, 1995).  
8  
9

10  
11  
12 Insert Table 12 hereabouts  
13

### 14 **Critical Mars - Earth contrasts: considerations for MSL decision-making**

15  
16 Any martian biomass must be, and probably would have been, less abundant than biomass on  
17 Earth throughout its history. Land plants dominate modern biomass on Earth, and marine  
18 organic matter is created almost exclusively by photosynthetic biota and it is pervasively  
19 concentrated at the margins of continents. Only a relatively small percentage of all primary  
20 productivity (~0.1%) survives remineralization and becomes preserved in sediments. If life  
21 colonized Mars, its global primary productivity is expected to have been much lower, given the  
22 less clement and more ephemeral surface habitable environments and the presumed absence  
23 of plants. The fraction of any biomass that was preserved in martian sedimentary rocks is, of  
24 course, unknown.  
25  
26  
27

28  
29 Meteorite impacts delivered organic matter to the martian surface. The early solar system had a  
30 greater abundance of debris and thus experienced a higher rate of impacts than later in its  
31 history. The size distribution of impactors may also have been different. Radiometrically dated  
32 samples from the moon make it possible to associate an absolute age with a certain crater  
33 density. The relation is non-linear because the flux of impactors was higher before 3.5 Ga, but  
34 subsequently the flux has apparently remained nearly constant. The following major questions  
35 persist: 1) Did the early impact flux decrease steadily or did an "impact spike" occur at ~4 Ga  
36 (known as the "Late Heavy Bombardment")? 2) There is a large uncertainty (factor of about 2)  
37 for young ages ( $\leq 1$  Ga). Because Mars is closer to the asteroid belt the number of impacts on  
38 Mars is estimated to be  $3 \pm 1.5$  times the number of impacts on the Earth-Moon system.  
39 However the accumulation of meteoritic organics in the martian crust also depends upon the  
40 fraction of material that is actually preserved.  
41  
42  
43

44  
45 The Late Heavy Bombardment is an important consideration in estimating organic matter  
46 preservation on Mars. As far as we know, virtually all of the biogenic organic matter that  
47 persists on Earth today was formed after 3.8 Ga and most of that was formed within the most  
48 recent 500 million years. If a biosphere existed in the first 500 million years of martian history a  
49 substantial fraction of its remains might have been altered or destroyed during the Late Heavy  
50 Bombardment. Smaller, relatively recent impacts might have exhumed any remnants of an early  
51 biosphere.  
52  
53

54  
55 The tectonic regimes of Mars and Earth are distinctly different, as reflected by significantly older  
56 surface ages and relatively minor regional metamorphism on Mars. Organic carbon  
57 accumulation on Earth is, and has been, modulated by a vigorous tectonic cycle, and the  
58  
59  
60



1  
2  
3 residence time of organic carbon in the crust is consistent with the Wilson cycle period (i.e.,  
4 modern plate tectonics). Most importantly, tectonic processes drive biogeochemical cycles by  
5 sustaining nutrient availability and creating the spaces for subaqueous accumulations of  
6 sediment and its entrained organic matter.  
7

8  
9 Mars currently has no global dynamo-driven magnetic field, but strong local crustal fields  
10 indicate that a global field likely existed in the past. However, magnetic field age and strength  
11 are not known to directly influence organic matter formation preservation except that a global  
12 magnetic field would have attenuated the flux of deleterious radiation and reduced losses of  
13 atmospheric species to space. The radiation environment on Earth is conducive to harvesting  
14 solar energy and forming organic matter through photosynthesis. A dense atmosphere, the  
15 pervasiveness of liquid water on Earth's surface and the operation of a magnetic field all can  
16 reduce deleterious radiation. Widespread photosynthesis on Earth has clearly enhanced organic  
17 matter production in environments that also favored its preservation in aqueous sediments.  
18

19  
20  
21 Processes operating in such favorable sedimentary environments can concentrate organic  
22 matter prior to burial. Important concentrating mechanisms include the following: density sorting  
23 during transport, adherence to the fine particles of clay minerals, and ballasting of organics on  
24 biogenic minerals. Differences in the hydrological cycles of Mars and Earth would have affected  
25 any potential concentration or organics during transport.  
26  
27

28  
29 Sedimentary rocks with anomalously high concentrations of organic matter (> few percent) are a  
30 historically pervasive feature of Earth's sedimentary rock record; they occur even in the early  
31 Precambrian record. However organic-rich and biomarker-bearing deposits are typically non-  
32 uniformly distributed and unpredictable. No known example of subaerial fossilized deposits on  
33 Earth have a total organic carbon content exceeding 1% by weight; Mars is expected to be  
34 similar.  
35

36  
37 Aeolian transport processes lead to destruction of organic matter by continually refreshing its  
38 contact with oxidizing agents and UV radiation. The fate of organic matter transported in this  
39 way is expected to be similar on Mars to what it is on Earth or, possibly, worse.  
40

41  
42 Therefore, on modern Earth, biomass has been detected in almost every wet environment,  
43 including very harsh and extreme sites in terms of temperature, pH, water activity, intermittent  
44 desiccation, and pressure. Essentially every wet environment below the upper thermal limit of  
45 life can be considered habitable. However the production, concentration and preservation of  
46 organics have varied substantially and have been controlled by the spatial and temporal  
47 distribution of subaqueous environments and sedimentary processes. The search for martian  
48 biosignatures has become more promising due to the discovery that surface and near-surface  
49 aqueous environments existed on Mars at the same time when biological organic matter was  
50 being preserved in ancient aqueous sediments on Earth.  
51

52  
53 In its quest to find organic-bearing strata, MSL should investigate, ideally, a subaqueously  
54 deposited and rapidly buried suite of strata that represents the longest duration possible, i.e.,  
55 the thickest section in the absence of other age constraints (Farmer and Des Marais, 1999). If  
56 the record at Meridiani Planum is representative of other early martian sedimentary  
57  
58  
59  
60

1  
2  
3 environments, at least some beds might have been deposited in water even if the stratigraphic  
4 succession is predominantly eolian in origin (Grotzinger et al., 2005).  
5  
6

### 7 **Consideration of planetary age**

8  
9 Planetary evolution strongly influences biosignature preservation, particularly on a planet like  
10 Mars that has experienced significant geological and climate variations. Changes in habitability  
11 through time affect the abundance and diversity of potential biosignatures. For example, loss  
12 of the Martian magnetic field strongly affected surface environments through atmospheric loss  
13 and increased radiation; this change significantly degraded surface habitability, reducing the  
14 chances of biosignatures in the rock record. Changes in temperature and moisture also clearly  
15 affect habitability both on the surface and in subsurface aquifers. Hydrothermal activity tied to  
16 volcanism and impacts was more abundant on early Mars and more abundant and continuous  
17 hydrothermal activity is more likely to support a biosphere that could leave signatures. Also, it is  
18 critical to understand long term climate evolution when choosing the best places to look for  
19 biosignatures; rocks deposited during warm and wet intervals, e.g. early Mars, are more  
20 promising for biosignature development. Preservation of any biosignatures also depends on  
21 climate-dependent sedimentary processes. Clay minerals preserve biosignatures in ways that  
22 are fundamentally different to what occurs in carbonate or sulfate minerals (Hedges and Keil,  
23 1995). Therefore, long term changes in the relative abundance of certain minerals affect the  
24 likelihood of specific biosignature preservation in rocks of a particular age. Thus, the specific  
25 evolution of Mars as a planet suggests that the best time interval for the search for  
26 biosignatures would be represented by early to mid-Noachian rocks, when clays were thought to  
27 be forming. This is not to say that clays did not form in younger times, in fact, tests of this  
28 hypothesis could be one outcome of the MSL mission.  
29  
30  
31  
32  
33

### 34 **Potential biological sources of organic carbon on Mars**

35  
36 On Earth today, photoautotrophy is by far the dominant physiology leading to organic matter  
37 synthesis. With a seemingly unlimited solar energy source, photoautotrophs in the ocean and on  
38 land can produce copious amounts of organic matter compared with chemoautotrophs (orders  
39 of magnitude more). In present-day aquatic environments chemoheterotrophs thrive on  
40 chemical energy from the decomposition of organic matter from photosynthetic communities  
41 using electron acceptors that are also regenerated by photoautotrophs.  
42  
43  
44

45 Chemoautotrophs thrive by carbon fixation using electron donors that are also generated or  
46 regenerated via light-harvesting processes. In the absence of photoautotrophy, primary biomass  
47 production would be limited to chemoautotrophs that harvest chemical energy from geological  
48 processes—namely those occurring in hydrothermal vents in the oceans and terrestrial  
49 geothermal springs (see Table 4) and subsurface microenvironments within the fractures and  
50 pores of ultramafic and mafic rocks. The Lost City Hydrothermal Field (Kelley et al., 2005), a  
51 low-temperature marine ultramafic hydrothermal system is one model for an environment that  
52 could support life in the absence of photosynthetic light harvesting (Martin & Russell, 2007;  
53 Martin et al., 2008). Such a system could conceivably occur anywhere that water circulates  
54 through ultramafic rock. Although ultramafic rock is rare on the ocean floor today, occurring  
55 mainly at ultraslow-spreading ridges, it could have been more prevalent in the geological past  
56  
57  
58  
59  
60

1  
2  
3 (Sleep et al., 2004; Sleep and Bird 2007). Molecular and isotopic data indicate that Lost City  
4 fluids sustain a flourishing microbial community of methanogens and sulfate-reducing bacteria  
5 (SRB) (Bradley et al., 2009; Brazelton et al., 2006). Although the SRB would be sulfate-limited  
6 if disconnected from a sulfate supply that is ultimately coupled to oxygenic photosynthesis or  
7 abiotic oxidation reactions occurring at the surface or subsurface, there appear to be no such  
8 constraints on the activity of methanogens that require only hydrogen and CO<sub>2</sub>. In addition,  
9 thermodynamic calculations showed that Lost City-type vent fluids mixed with seawater are  
10 energetically favorable for biomass synthesis (Amend & McCollom, 2009). Thus, the recent  
11 detection of serpentine deposits in Noachian terrains on the surface of Mars (Ehlmann et al.,  
12 2010) identifies the probable past occurrence of hydrogen-producing water-rock reactions and,  
13 therefore, an established set of processes that would both support chemosynthetic life and  
14 preserve a molecular and/or isotopic record of its prior existence (e.g. Kelly et al., 2005; Bradley  
15 et al., 2009)

16  
17  
18  
19  
20 At life's origin, the dominant energy source was unlikely to have been sunlight. Energy flowing  
21 from chemical and thermal disequilibria and particularly from the interaction of hot rocks with  
22 water is more likely. Perhaps the same was the case for early Mars. In aqueous depositional  
23 environments, chemoautotrophs may have been the cornerstone of microbial communities  
24 relying on fermentation and heterotrophy fueled by a weak oxidant flux from chemoautotrophy  
25 or irradiative oxidation. If Mars evolved a biosphere, it may not have progressed to  
26 photoautotrophy or a dependence on photoautotrophy as it did on Earth. Thus, in the  
27 consideration of Martian environments conducive to producing molecular biosignatures,  
28 targeting depositional environments that had a strong chemical energy flux and sustained redox  
29 gradients for long periods by biogeochemical cycling is a most promising strategy.

30  
31  
32  
33 One additional possibility for chemoautotrophy is energy derived from radiolysis of water (Pratt  
34 et al., 2006). The recent discovery of a microbial biome dominated by thermophilic sulfate  
35 reducing bacteria in a c. 3km deep saline aquifer in Archean metabasalt seems to required that  
36 they were sustained by geologically produced sulfate and hydrogen at concentrations sufficient  
37 to maintain biological activity for millions of years (Chivian et al., 2008; Lin et al., 2006).  
38 Radiolysis of water coupled to oxidation of sulfide minerals could have provided the energy  
39 drive for low-biomass, low-diversity subsurface ecosystems.

### 40 41 42 43 **Abiotic sources of organic carbon on Mars**

44  
45 Sources of abiotic organic matter on Mars could have been similar to those hypothesized to  
46 have been present or formed on the early Earth. This would have included organic matter  
47 delivered by meteorites and interplanetary dust (Anders, 1989), organic matter produced as a  
48 result of atmospheric photochemistry (Chang et al., 1983) and organic matter produced during  
49 fluid mixing in hydrothermal systems (Shock & Schulte, 1998). The Lost City hydrothermal  
50 system also serves as an example of another route to abiotic organic compounds. The isotopic  
51 composition and chain length distributions of hydrocarbon gases isolated from Lost City fluids  
52 have been interpreted to signify an abiotic source (Proskurowski et al., 2008; Sherwood Lollar et  
53 al., 2006). Hydrogen produced in high concentration by serpentinization chemistry leads to a  
54 thermodynamic drive for CO<sub>2</sub> reduction (Shock & Schulte, 1998). Methane and higher  
55  
56  
57  
58  
59  
60

1  
2  
3 hydrocarbons may be thus produced abiotically in ultramafic hydrothermal systems by Fischer-  
4 Tropsch type processes that comprise polymerization reactions leading to methane and higher  
5 hydrocarbons (Horita & Berndt, 1999; McCollom & Seewald, 2006; McCollom & Seewald,  
6 2007).

7  
8  
9 Steele and co-workers (2007) have identified macromolecular carbon, in an intimate association  
10 with magnetite, through imaging and Raman spectroscopic studies of carbonate globules in the  
11 Mars meteorite ALH84001. This abiotically-formed organic matter appears to be native to Mars.  
12 It is hypothesized to have formed formation via reactions of the Fe-C-O system for which there  
13 is a terrestrial analogue in the Bockfjorden volcanic complex of Svalbard.  
14  
15  
16  
17

### 18 **Interstellar organic matter**

19  
20 Irrespective of whether extensive abiotic organic synthesis of predominantly hydrocarbons  
21 occurred on the early Earth and, potentially, on Mars, chondritic meteorites and interplanetary  
22 dust particles (IDPs) have delivered abiotic organic matter to the Martian surface.  
23 Carbonaceous chondritic (CC) meteorites consist of up to 2 wt % of organic matter finely  
24 intermixed with matrix silicates (Alexander et al., 2007). It is well known that some CCs contain  
25 relatively high abundances of small, polar organic molecules, e.g. amino acids. The highest  
26 reported concentration of amino acids, however, is not very high, being on the order of 250 ppm  
27 (Martins et al. 2007). The primary form of organic matter in all classes of chondritic meteorites  
28 is Insoluble Organic Matter (IOM), a chemically complex macromolecule (e.g. Cody & Alexander  
29 2005) that is by definition insoluble in any solvent and is in many cases 99 % of the organic  
30 matter in a given chondrite (Botta, 2005; Sephton, 2002; Sephton et al., 2004). Among the  
31 meteorites collected as finds in Antarctica, 0.8 % are CCs, the majority, 77 %, are ordinary  
32 chondrites (OCs) (<http://curator.jsc.nasa.gov/antmet/ppr.cfm>). While ordinary chondrites do  
33 contain IOM, its abundance is considerably lower than that in CCs (Alexander et al. 2007) and,  
34 even in the least metamorphosed OCs, the IOM has been significantly altered through long term  
35 thermal processing yielding highly aromatized macromolecular structures (Cody et al., 2008). If  
36 the distribution of meteorite types on Mars was and is similar to that of Earth, it is reasonable to  
37 assume that the predominant IOM would be exceptionally stable over geological timescales  
38 even under the harsh Martian surface conditions. An instrument analogous to SAM flew on the  
39 Viking mission and did not detect organics via pyrolytic analysis. All evidence suggests that the  
40 Viking technical approach and instrument worked according to its design (Biemann, 1979,  
41 Biemann, 2007) so these Viking results do not exclude the potential presence of abundant IOM,  
42 if the IOM was derived predominantly from OCs. IOM in OCs has already been subjected to  
43 extensive natural pyrolysis in the OC parent body (Cody et al., 2008). Minimal, if any, pyrolysate  
44 remains to be derived, therefore, from thermally metamorphosed IOM. Pyrolysate from  
45 chondritic IOM might be detected only if the abundance of CCs that accumulate on the Martian  
46 surface exceeds what is observed in the Antarctic collections.  
47  
48  
49  
50  
51  
52  
53  
54

55 Finally, there is the issue of Interplanetary Dust Particles (IDPs) that, in case of the Earth,  
56 constitute a much greater influx of extraterrestrial matter. IDPs typically contain considerably  
57 more organic matter than CCs and the organic macromolecule bears significant spectroscopic  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

resemblance to IOM derived from primitive CCs. Lacking the silicate matrix protection afforded to CC IOM, it appears likely that IDPs will be subjected to more degradation than CC IOM on the Martian surface. Under ideal conditions, however, IDPs could survive on Mars, and be detectable via pyrolysis GCMS. The SAM instrument has enhanced sensitivity and experiment flexibility compared to Viking. However, an even more important quality is the mobility and composition of the entire MSL package as this enables optimization of sample selection and handling.

### **Distribution and importance of phyllosilicates for habitability and organic preservation**

It has long been recognized that, on Earth, there is a close association between organic matter and phyllosilicate minerals. This is the case for both modern environments, particularly in large aqueous catchments where fine-grained particulates often rich in phyllosilicate minerals can settle from the water column (lakes and ocean margins), and in ancient sediments (e.g. shale and organic-rich mudstones). Certainly, low porosity and permeability of compacted phyllosilicate sheets aid in entombing and protecting organic matter from oxidizing fluids and biological activity over geological time scales. Abundant experimental data show that many phyllosilicates, particularly smectites, interact strongly with organic molecules and are capable of adsorbing and preserving them. However, the specific mechanisms that give rise to this association are not well understood and are the subject of ongoing investigations. Accordingly, we should be very careful in extending empirical observations made on Earth to the situation on Mars. Rather, the combined ability of the MSL to detect both organic carbon and clay mineral assemblages on Mars offers us an unprecedented opportunity to learn much about this particular issue during the landed operations.

Phyllosilicates such as smectites, chlorites, and kaolin minerals form during the weathering of minerals in soils and in hydrothermal systems. Phyllosilicates all have sheet-like structures and can accommodate a large variety of cations, most commonly including Fe, Mg, and Al in their octahedral sheets and Na, Ca, and K between the layers in the so-called interlayer region. Fine-grained, disordered phyllosilicates are often called clay minerals, and they have high surface areas and the ability to exchange their interlayer cations and adsorb H<sub>2</sub>O molecules, as a result of negatively charged interlayer regions. Many organic compounds can be adsorbed onto surfaces and into the interlayer regions, in some cases forming weak bonds with phyllosilicate surfaces. Adsorption of organic molecules into the interlayer region is particularly important for very low molecular weight compounds, such as amino acids (Hedges & Hare, 1987) and polysaccharides (Dontsova and Bigham, 2005) as well as higher molecular weight material (Mayer, 1994a, b; Kennedy, et al., 2002). Organic molecules compete with other polar species in the environment (e.g., water, cations, etc.) for active sites on phyllosilicate minerals. However, a key factor for enhanced organic matter preservation by phyllosilicates is coincidental timing of organic matter diagenesis and phyllosilicate mineral formation (Hedges & Kiel, 1995). Specifically, it requires synchronous availability of organics and clay mineral formation. To grasp the significance of this timing issue, it is also important to understand how organic matter ends up in sediments and ultimately the rock record.



1  
2  
3 On Earth, most sedimentary and hydrothermal organic matter is macromolecular. In these  
4 terrestrial settings, polymerization to macromolecules may begin at the point of formation (i.e.,  
5 cellular and biopolymer material), during diagenesis in the water column, or in pore waters of  
6 sediments as dissolved organic matter. It is composed of smaller constituents cross-linked  
7 together by covalent bonds into a three-dimensional material. Initially, it is porous, internally and  
8 externally charged, and has varying degrees of hydrated surfaces giving rise to hydrophilic (i.e.,  
9 charged with functional groups having O, N, and S moieties) and hydrophobic regions (i.e.,  
10 dominated by C and H). It also has a high affinity for complexation with trace elements and  
11 sorption to particles (Guo & Santschi, 1997), such as phyllosilicates (Ohashi & Nakazawa,  
12 1996). Once deposited, the chemistry, porosity, and overall activity of macromolecular organic  
13 matter become more stable as it equilibrates with the sedimentary and eventually, lithified,  
14 environment. On Earth, this environment may be oxidizing and lead to overall degradation and  
15 loss of the organic macromolecules, potentially leaving behind the minerals, trace elements, and  
16 morphologies associated with the original material. This is commonly the case for subaerial  
17 environments. However, organic macromolecules stand a better chance of preservation if they  
18 are quickly stabilized in sediments. There are two primary mechanisms for this. One, common in  
19 the anoxic marine realm on Earth, is where hydrophilic functional groups of organic  
20 macromolecules are effectively 'quenched' and replaced by sulfide during diagenesis. The  
21 second mechanism, potentially more important on Mars, is rapid burial and lithification of  
22 organic macromolecules. Organic macromolecular sorption to mineral surfaces significantly aids  
23 in burial of organic matter (e.g., Bock & Mayer, 2000; Kennedy et al., 2006). On Mars, the  
24 environments that may have allowed for this diagenetic complexation between dissolved  
25 organic macromolecules and phyllosilicates would thus provide promising sites for organic  
26 molecular preservation and detection by MSL. These environments must also be proximal to  
27 sources of both phyllosilicates (e.g., weathering regolith or hydrothermal) and organic matter.  
28 Therefore, Martian hydrothermal and stable aquatic sedimentary environments may be very  
29 favorable for both phyllosilicate and organic matter accumulation (Farmer and Des Marais,  
30 1999).

### 31 32 **Assessments of potential carbon accumulation and preservation in sediments and** 33 **hydrothermal systems** 34

35  
36  
37  
38  
39  
40  
41  
42 Terrestrial models of organic carbon accumulation and preservation provide predictive capability  
43 in respect to these processes in the regolith of Mars. However, due to differences in processes  
44 and chemical environments, our estimates of the propensity for early Mars environments to  
45 support organic matter formation and preservation are necessarily crude (Table 3). However,  
46 our understanding of terrestrial environments makes estimating the possibility of particular  
47 observations of potential biomarkers on Mars possible, even if limited. Estimates are thus  
48 scaled as high, medium or low. For example, the potential for biosynthesis of organic matter is  
49 highest in near-surface hydrothermal systems and in deltaic and lacustrine environments. The  
50 potential for hydrodynamic processes to concentrate those organics is ranked as high for the  
51 deltaic and perennial lacustrine environments but lower for the surface and subsurface  
52 hydrothermal systems as well as for evaporitic environments. Three columns toward the right of  
53 the table rank the potential to recognize such environments using remote sensing and criteria  
54 that are geomorphic, mineralogical, or stratigraphic, respectively. Finally, the right-hand column  
55  
56  
57  
58  
59  
60



1  
2  
3 makes an assessment of the MSL instrument package's potential to gather data identifying such  
4 environments.  
5

6  
7 The potential for organic matter formation and accumulation on early Mars is assessed as low in  
8 the volcanic pyroclastic and flow deposits and moderate in hydrothermal environments (Table  
9 4). Any organic matter that accumulated in such places would stand a moderate chance of  
10 being preserved over time. In tables 5 and 6 we evaluate how the MSL payload elements could  
11 be used to confirm environmental features specific to processes needed to form, transport,  
12 concentrate and preserve organic molecules on early Mars. Only environments identified in  
13 Table 1 as having moderate to high potential to support organic carbon formation and  
14 preservation that were at moderate or higher are considered for the sedimentary category (i.e.,  
15 fluvial floodplain, deltaic, lacustrine (perennial), lacustrine (evaporitic). Only hydrothermal  
16 (<100°C) subsurface + surface environments are considered for the hydrothermal category.  
17  
18

### 19 20 **An assessment of processes essential to the preservation of isotopic abundances in** 21 **Mars sediments and hydrothermal systems** 22

23 Interpretation of the stable isotopic compositions of carbon and sulfur in Mars gases, organics  
24 and minerals requires a thorough understanding of the environmental context under which the  
25 fractionation occurred and the degree to which original features may have been preserved.  
26 Such understandings can be gained, to some degree, by using the MSL payload elements as  
27 summarized in Table 7 for hydrothermal environments and in Table 8 for sedimentary  
28 environments. Interpretations of isotopic data are likely to be complex and prone to significant  
29 ambiguity.  
30  
31

### 32 33 **Synthesis** 34

35 The MSL instrument package has the potential to detect biosignatures if they are present at the  
36 landing site on Mars. Our understanding of the formation and preservation of biosignatures on  
37 Earth can guide our expectations of how and where they might have developed and be  
38 preserved on Mars even though the planets have distinct histories. The classes of  
39 biosignatures that could be detected and identified on Mars given appropriate biosignature  
40 formation and preservation include: diagnostic organic molecules, biogenic gases, body fossils,  
41 biofabrics, mineralogy affected by biomineralization and bioalteration, spatial patterns in  
42 chemistry due to metabolic processes, and isotopic signatures reflecting metabolic processes.  
43  
44

45 Diagnostic organic molecules and biogenic gases are the most definitive as biosignatures and  
46 are also readily detectable with the SAM instrument. However, they require sequestration from  
47 oxidative processes for preservation. Clay minerals promote the preservation of diagnostic  
48 organic molecules on Earth, and accumulations of sedimentary clay minerals commonly  
49 preserve organic molecules. Thus, using Earth as a model, a MSL landing site in a sedimentary  
50 basin containing clay minerals is ideal for maximizing the chances of detecting diagnostic  
51 organic molecules. Carbonates and other minerals deposited as a consequence of  
52 microbiological activity (microbialites) can also preserve diagnostic organic molecules.  
53  
54

55  
56  
57 Body fossils and biofabrics can also be definitive biosignatures if they are sufficiently complex  
58  
59  
60

1  
2  
3 and can be observed in context. For detection by MSL, body fossils and biofabrics must be  
4 large enough to be observed by MAHLI. Many terrestrial biofabrics are sufficiently large, but  
5 body fossils of bacteria are not. Biofabrics are easily preserved in hydrothermal and  
6 sedimentary rocks if protected from extensive recrystallization. Thus, detection of biofabrics by  
7 MSL is possible if microbial communities developed on Mars and an appropriate landing site is  
8 chosen. Low temperature hydrothermal and persistently wet sedimentary environments are  
9 most likely to develop and preserve biofabrics.  
10  
11

12  
13 Biomineralization and bioalteration effects on the spatial patterns in chemistry rarely produce  
14 definitive microbial biosignatures due to substantial overlaps in abiotic and biotic processes.  
15 However, preserved disequilibrium mineral distributions or variations in chemistry can indicate  
16 good sites to look for more definitive biosignatures. CheMin, ChemCam and APX can all help  
17 characterize minerals and chemical patterns that have the potential to be biosignatures.  
18  
19

20 Stable isotopic signatures are commonly used in terrestrial materials to characterize the extent  
21 of biological chemical activity. However, such interpretations require a detailed knowledge of  
22 the biogeochemical context, including cycles, of the activity. Without that context, stable  
23 isotopic signatures rarely provide strong evidence for biological activity. SAM can characterize  
24 the isotopic composition of various important materials. If the proper environment is  
25 encountered on Mars, we have the potential to identify stable isotopic biosignatures if present.  
26 However, definitive biosignature detection would require characterization of co-occurring  
27 elemental reservoirs, preferably of carbon and sulfur with different oxidation states. To date, we  
28 have not identified such an environment on Mars.  
29  
30  
31

32 Early Earth is the best analogue we have currently for guiding the search for biosignatures on  
33 Mars. Still, we must be constantly aware of the limits of our understanding of terrestrial  
34 processes of biosignature formation and preservation, especially as they relate to the Earth's  
35 earliest sedimentary record. The MSL is at the heart of the first NASA Astrobiology mission and  
36 provides an extraordinary opportunity to learn more about martian environments and processes,  
37 particularly in localities that might have been inhabited by microorganisms.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

- Alexander, C. M. O'D., Fogel M. L., Yabuta, H., Cody, G. D. (2007) The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochimica et Cosmochimica Acta* 71, 4380-4403.
- Amend, J.P., McCollom, T.M. (2009) Energetics of biomolecule synthesis on early Earth. In *Chemical Evolution II: From the Origins of Life to Modern Society*, Vol. 1025 (eds. L. Zaikowski, J.M. Friedrich, and S.R. Seidel), pp. 63-94. American Chemical Society.
- Anders, E., (1989) Pre-biotic organic matter from comets and asteroids. *Nature* 342, 255-257.
- Anders, E., Hayatsu, R., Studier, M.H. (1973) Organic Compounds in Meteorites: They may have formed in the solar nebula, by catalytic reactions of carbon monoxide, hydrogen, and ammonia. *Science*, 182, 781-790.
- Atreya, S.K. Wong, A.-S., Renno, N.O., Farrell, W. M., Delory, G.T., Sentman, D.D., Cummer, S.A., Marshall, J.R., Rafkin, S.C.R., Catling, D.C. (2006) Oxidant enhancement in Martian dust devils and storms: Implications for life and habitability. *Astrobiology* 6, 439-450.
- Aubrey A., Cleaves H. J., Chalmers J. H., Skelley A. M., Mathies R. A., Grunthaner F. J., Ehrenfreund P., and Bada J. L. (2006) Sulfate minerals and organic compounds on Mars. *Geology* 34, 357-360.
- Baross J. and others, *Limits of Life in Planetary Systems* (2007) Committee on Limits of Life in Planetary Systems, National Academies Press, ISBN-13: 978-0-309-10484-5.
- Benner, S.A., Devine, K.G., Matveeva, L.N., Powell, D.H. (2000) The missing organic molecules on Mars. *Proceedings of the National Academy of Sciences of the United States of America* 97, 2425-2430.
- Bibring, J.-P., Langevin, Y., Gendrin, A., Gondet, B., Poulet, F., Berthe, M., Soufflot, A., Arvidson, R., Mangold, N., Mustard, J., Drossart, P., and the Omega Team (2005) Mars Surface Diversity as Revealed by the OMEGA/Mars Express Observations. *Science* 307,1576-1581.
- Biemann, K. (1979) The implications and limitations of the findings of the Viking organic analysis experiment. *Journal of Molecular Evolution* 14, 65-70.
- Biemann, K. (2007) On the ability of the Viking gas chromatograph-mass spectrometer to detect organic matter. *Proceedings of the National Academy of Sciences of the United States of America* 104, 10310-10313.
- Birgel D., Himmler T., Freiwald A., and Peckmann J. r. (2008) A new constraint on the antiquity of anaerobic oxidation of methane: Late Pennsylvanian seep limestones from southern Namibia. *Geology* 36, 543-546.
- Bock, M.J. and Mayer, L.M. (2000) Mesodensity organo – clay associations in a near-shore

- 1  
2  
3 sediment. *Marine Geology* 163, 65-75.  
4  
5 Botta, O., (2005) Organic Chemistry in Meteorites, Comets, and the Interstellar Medium.  
6 *Proceedings of the International Astronomical Union* 1, 479-488.  
7  
8 Botta, O., Bada, J.L., Gomez-Elvira, J., Javaux, E., Selsis, F. and Summons, R.E. (Eds), (2008)  
9 Strategies of Life Detection. In: *Space Sciences Series of ISSI*, 25. Springer US, Bern.  
10  
11 Bishop, J. L., Dobreá, E. Z. N., McKeown, N. K., Parente, M., Ehlmann, B. L., Michalski, J. R.,  
12 Milliken, R. E., Poulet, F., Swayze, G. A., Mustard, J. F., Murchie, S. L., and Bibring, J.-P.  
13 (2008) Phyllosilicate Diversity and Past Aqueous Activity Revealed at Mawrth Vallis,  
14 Mars. *Science* 321(5890), 830-833.  
15  
16  
17  
18 Bradley, A.S., and Summons, R.E. (2010) Multiple origins of methane at the Lost City  
19 Hydrothermal Field. *Earth and Planetary Science Letters* 297, 34-41.  
20  
21 Bradley, A.S., Hayes, J.M., Summons, R.E. (2009) Extraordinary <sup>13</sup>C enrichment of diether  
22 lipids at the Lost City Hydrothermal Field indicates a carbon-limited ecosystem.  
23 *Geochimica et Cosmochimica Acta* 73, 102-118.  
24  
25  
26 Brasier, M., McLoughlin, N., Green, O., and Wacey, D. (2006) A fresh look at the fossil evidence  
27 for early Archaean cellular life. *Philosophical Transactions of the Royal Society B:*  
28 *Biological Sciences* 361, 887-902.  
29  
30 Brazelton, W.J., Schrenk, M.O., Kelley, D.S., Baross, J.A. (2006) Methane- and sulfur-  
31 metabolizing microbial communities dominate the Lost City Hydrothermal Field  
32 ecosystem. *Applied and Environmental Microbiology* 72, 6257-6270.  
33  
34  
35 Butterfield, N. J. (1990) Organic Preservation of Non-Mineralizing Organisms and the  
36 Taphonomy of the Burgess Shale. *Paleobiology* 16, 272.  
37  
38 Canfield, D. E. (2001) Biogeochemistry of sulfur isotopes. In: J.W. Valley and D.R. Cole, Eds.  
39 Stable Isotope Geochemistry, *Reviews in Mineralogy* 43, 607-636.  
40  
41 Chang, S., Des Marais, D.J., Mack, R., Miller, S.L., Strathearn, G.E. (1983) Prebiotic organic  
42 syntheses and the origin of life. In: J.W. Schopf (Ed.), *Earth's earliest biosphere: Its origin*  
43 *and evolution*, Princeton University Press, Princeton, NJ., pp. 53-92.  
44  
45  
46 Chen J., Walter M.R., Logan G.A., Hinman M.C. and Summons R.E., 2003. The  
47 Paleoproterozoic McArthur River (HYC) Pb/Zn/Ag deposit of northern Australia: organic  
48 geochemistry and ore genesis. *Earth and Planetary Science Letters* 210, 467-479.  
49  
50  
51 Chivian, D., Brodie, E.L., Alm, E.J., Culley, D.E., Dehal, P.S., *et al.*, (2008) Environmental  
52 Genomics Reveals a Single-Species Ecosystem Deep Within Earth. *Science* 322, 275-  
53 278.  
54  
55  
56 Christensen, P. R., Wyatt, M. B., Glotch, T. D., Rogers, A. D., Anwar, S., Arvidson, R. E.,  
57 Bandfield, J. L., Blaney, D. L., Budney, C., Calvin, W. M., Fallacaro, A., Fergason, R. L.,  
58  
59  
60

- 1  
2  
3 Gorelick, N., Graff, T. G., Hamilton, V. E., Hayes, A. G., Johnson, J. R., Knudson, A. T.,  
4 McSween, H. Y., Jr., Mehall, G. L., Mehall, L. K., Moersch, J. E., Morris, R. V., Smith, M.  
5 D., Squyres, S. W., Ruff, S. W., and Wolff, M. J. (2004) Mineralogy at Meridiani Planum  
6 from the Mini-TES Experiment on the Opportunity Rover. *Science* 306, 1733-1739.  
7  
8  
9 Cody G.D., Alexander C. M. O'D. (2005) NMR studies of chemical structural variation of  
10 insoluble organic matter from different carbonaceous chondrite groups. *Geochimica et*  
11 *Cosmochimica Acta* 69, 1085-1097.  
12  
13 Cody G. D., Alexander C. M. O'D., Yabuta H., Kilcoyne A. L. D., Araki T., Ade H., Dera P., Fogel  
14 M. L., Militzer B., Mysen B. (2008) Organic Thermometry for Chondritic Parent Bodies,  
15 *Earth and Planetary Science Letters* 272, 446-455.  
16  
17 Cronin, J.R., Pizzarello, S. (1997) Enantiomeric excesses in meteoritic amino acids. *Science*  
18 275, 951-955.  
19  
20 Dahl, J., Hallberg, R., and Kaplan, I. R. (1988) Effects of irradiation from uranium decay on  
21 extractable organic matter in the Alum Shales of Sweden. *Organic Geochemistry* 12,  
22 559-571.  
23  
24 Des Marais, D.J., (2001) Isotopic evolution of the biogeochemical carbon cycle during the  
25 Precambrian. In: J.W. Valley and D.R. Cole, Eds. *Stable Isotope Geochemistry, Reviews*  
26 *in Mineralogy* 43, 555-578.  
27  
28 Dontsova, K.M., Bigham, J.M. (2005) Anionic Polysaccharide Sorption by Clay Minerals, *Soil*  
29 *Science Society of America Journal* 69, 1026-1035.  
30  
31 Ehlmann, B. L., Mustard, J. F., Fassett, C. I., Schon, S. C., Head, J. W., Des Marais, D. J.,  
32 Grant, J. A., and Murchie, S. L. (2008a) Clay minerals in delta deposits and organic  
33 preservation potential on Mars. *Nature Geoscience* 1(6), 355-358.  
34  
35 Ehlmann, B. L., Mustard, J. F., Murchie, S. L., Poulet, F., Bishop, J. L., Brown, A. J., Calvin, W.  
36 M., Clark, R. N., Des Marais, D.J., Milliken, R. E., Roach, L. H., Roush, T. L., Swayze, G.  
37 A., and Wray, J. J. (2008b) Orbital Identification of Carbonate-Bearing Rocks on Mars.  
38 *Science* 322(5909), 1828-1832.  
39  
40 Ehlmann, B. L., Mustard, J. F., and Murchie, S. L. (2010) Geologic setting of serpentine  
41 deposits on Mars. *Geophysical Research Letters* 37, In Press.  
42  
43 Engel, M.H., Macko, S.A. (1997) Isotopic evidence for extraterrestrial non-racemic amino acids  
44 in the Murchison meteorite. *Nature* 389, 265-268.  
45  
46 Farmer, J. D., Des Marais, D.J. (1999), Exploring for a record of ancient Martian life, *Journal of*  
47 *Geophysical Research* 104, 977-995.  
48  
49 Farquhar, J., Johnston, D.T., Wing, B.A., Habicht, K.S., Canfield, D.E., Airieau, S., Thiemens,  
50 M.H. (2003) Multiple sulphur isotopic interpretations of biosynthetic pathways:  
51 implications for biological signatures in the sulphur isotope record, *Geobiology* 1, 27-36.  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 Farquhar, J., Savarino, J., Jackson, T.L., Thiemens, M.H. (2000) Evidence of atmospheric  
4 sulphur in the Martian regolith from sulphur isotopes in meteorites. *Nature* 404, 50-52.  
5  
6  
7 Fernandez-Remolar, D., Knoll, A.H. (2008) Fossilization potential of iron-bearing minerals in  
8 acidic environments of Rio Tinto, Spain: Implications for Mars exploration. *Icarus* 194,  
9 72-85.  
10  
11 Formisano V., Atreya S., Encrenaz T., Ignatiev N., and Giuranna M. (2004) Detection of  
12 Methane in the Atmosphere of Mars. *Science* 306, 1758-1761.  
13  
14 Franz, H.B., Mahaffy, P.R., Farquhar, J. (2007) Preliminary estimate of sulfur isotope ratio  
15 precision expected with the Sample Analysis at Mars (SAM) Instrument Suite of the 2009  
16 Mars Science Laboratory. *38th Lunar and Planetary Science Conference*, abstract 1874.  
17  
18  
19 Glavin, D.P., Dworkin, J.P. (2009) Enrichment of the amino acid l-isovaline by aqueous  
20 alteration on CI and CM meteorite parent bodies. *Proceedings of the National Academy*  
21 *of Sciences of the United States of America* 106, 5487-5492.  
22  
23  
24 Greenwood, J.P., Mojzsis, S.J., Coath, C.D., (2000) Sulfur isotopic compositions of individual  
25 sulfides in Martian meteorites ALH84001 and Nakhla: implications for crust–regolith  
26 exchange on Mars. *Earth and Planetary Science Letters* 184, 23-35.  
27  
28  
29 Grotzinger J. P., Arvidson R. E., Bell J. F., Calvin W., Clark B. C., Fike D. A., Golombek M.,  
30 Greeley R., Haldemann A., Herkenhoff K. E., Jolliff B. L., Knoll A. H., Malin M., McLennan  
31 S. M., Parker T., Soderblom L., Sohl-Dickstein J. N., Squyres S. W., Tosca N. J., and  
32 Watters W. A. (2005) Stratigraphy and sedimentology of a dry to wet eolian depositional  
33 system, Burns formation, Meridiani Planum, Mars. *Earth and Planetary Science Letters*  
34 240, 11-72.  
35  
36  
37 Guo, L., Santschi, P.H. (1997) Composition and cycling of colloids in marine environments.  
38 *Reviews in Geophysics*, 35, 17-40.  
39  
40 Haskin, L. A., Wang, A., Jolliff, B. L., McSween, H. Y., Clark, B. C., Des Marais, D. J.,  
41 McLennan, S. M., Tosca, N. J., Hurowitz, J. A., Farmer, J. D., Yen, A., Squyres, S. W.,  
42 Arvidson, R. E., Klingelhofer, G., Schroder, C., de Souza, P. A., Ming, D. W., Gellert, R.,  
43 Zipfel, J., Bruckner, J., Bell, J. F., Herkenhoff, K., Christensen, P. R., Ruff, S., Blaney,  
44 D., Gorevan, S., Cabrol, N. A., Crumpler, L., Grant, J., and Soderblom, L. (2005) Water  
45 alteration of rocks and soils on Mars at the Spirit rover site in Gusev crater. *Nature*  
46 436(7047), 66.  
47  
48  
49 Hazen, R.M., Papineau, D., Bleeker, W., Downs, R.T., Ferry, J.M., McCoy, T.J., Sverjensky,  
50 D.A., Yang, H. (2008) Mineral evolution. *American Mineralogist* 93, 1693–1720.  
51  
52  
53 Hedges, J.I., Hare, P.E. (1987) Amino acid adsorption by clay minerals in distilled water.  
54 *Geochimica et Cosmochimica Acta* 51, 255-259.  
55  
56  
57 Hedges, J.I., Keil, R.G. (1995) Sedimentary organic matter preservation: an assessment and  
58 speculative synthesis. *Marine Chemistry* 49, 81-115.  
59  
60

- 1  
2  
3 Horita, J., Berndt, M.E. (1999) Abiogenic Methane Formation and Isotopic Fractionation Under  
4 Hydrothermal Conditions. *Science* 285, 1055-1057.  
5  
6  
7 Jakosky B.M. and others (2007) *An Astrobiological Strategy for the Exploration of Mars*.  
8 Committee on an Astrobiology Strategy for the Exploration of Mars, National Academies  
9 Press, ISBN-13: 978-0-309-10851-5.  
10  
11 Jakosky, B. M., Pepin, R. O. Johnson, R. E., Fox, J. L. (1994) Mars atmospheric loss and  
12 isotopic fractionation by solar-wind-induced sputtering and photochemical escape. *Icarus*  
13 111, 271-288.  
14  
15  
16 Kelley, D.S., Karson, J.A., Fruh-Green, G.L., Yoerger, D.R., Shank, T.M., Butterfield, D.A.,  
17 Hayes, J.M., Schrenk, M.O., Olson, E.J., Proskurowski, G., Jakuba, M., Bradley, A.,  
18 Larson, B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A.S., Brazelton, W.J., Roe,  
19 K., Elend, M.J., Delacour, A., Bernasconi, S.M., Lilley, M.D., Baross, J.A., Summons,  
20 R.E., Sylva, S.P. (2005) A Serpentinite-Hosted Ecosystem: The Lost City Hydrothermal  
21 Field. *Science* 307, 1428-1434.  
22  
23  
24 Kennedy, M.J., Pevear, D.R., and Hill, R.J. (2002) Mineral Surface Control of Organic Carbon in  
25 Black Shale. *Science* 295, 657-660.  
26  
27  
28 Kennedy, M., Droser, M., Mayer, L.M., Pevear, D., Mrofka, D. (2006) Late Precambrian  
29 Oxygenation; Inception of the Clay Mineral Factory. *Science* 11, 1446-1449.  
30  
31 Knoll, A.H. (1985) Exceptional Preservation of Photosynthetic Organisms in Silicified  
32 Carbonates and Silicified Peats. *Philosophical Transactions of the Royal Society of*  
33 *London. Series B, Biological Sciences* 311, 111-122.  
34  
35  
36 Krasnopolsky, V.A., Maillard, J.P., Owen T.C. (2004) Detection of methane in the martian  
37 atmosphere: evidence for life? *Icarus* 172, 537-547.  
38  
39 Lin, L.-H., Wang, P.-L., Rumble, D., Lippmann-Pipke, J., Boice, E., Pratt, L.M., Lollar, B.S.,  
40 Brodie, E.L., Hazen, T.C., Andersen, G.L., DeSantis, T.Z., Moser, D.P., Kershaw, D.,  
41 Onstott, T.C., (2006) Long-Term Sustainability of a High-Energy, Low-Diversity Crustal  
42 Biome. *Science* 314, 479-482.  
43  
44  
45 Martin, W., Baross, J., Kelley, D., Russell, M.J. (2008) Hydrothermal vents and the origin of life.  
46 *Nature Reviews in Microbiology* 6, 805-814.  
47  
48 Martin, W., Russell, M.J. (2007) On the origin of biochemistry at an alkaline hydrothermal vent.  
49 *Philosophical Transactions of the Royal Society B: Biological Sciences* 362, 1887-1926.  
50  
51 Martins, Z., Alexander, C.M.O.D., Orzechowska, G.E., Fogel, M.L., Ehrenfreund, P., (2007)  
52 Indigenous amino acids in primitive CR meteorites. *Meteoritics & Planetary Science* 42,  
53 2125-2182.  
54  
55  
56 Mayer, L.M. (1994a) Relationships between mineral surfaces and organic carbon  
57 concentrations in soils and sediments. *Chemical Geology* 114, 347-363.  
58  
59  
60

- 1  
2  
3 Mayer, L.M. (1994b) Surface area control of organic carbon accumulation in continental shelf  
4 sediments. *Geochimica et Cosmochimica Acta* 58, 1271-1284.  
5  
6  
7 McCollom, T.M., Seewald, J.S. (2006) Carbon isotope composition of organic compounds  
8 produced by abiotic synthesis under hydrothermal conditions. *Earth and Planetary*  
9 *Science Letters* 243, 74-84.  
10  
11 McCollom, T.M., Seewald, J.S. (2007) Abiotic Synthesis of Organic Compounds in Deep-Sea  
12 Hydrothermal Environments. *Chemical Reviews* 107, 382-401.  
13  
14 McKay, C.P. (2004) What Is Life-and How Do We Search for It in Other Worlds? *PLoS Biology*  
15 2, e302.  
16  
17  
18 McKay, C.P. (2007) An Approach to Searching for Life on Mars, Europa, and Enceladus. *Space*  
19 *Science Reviews* 135, 49-54.  
20  
21 Meyers, P.A and Ishiwatari, R. (1995) Organic matter accumulation records in lake sediments.  
22 In: A. Lerman et al., Editors, *Physics and Chemistry of Lakes*, Springer, pp. 279–328.  
23  
24 Mumma, M., Villanueva, G.L., Novak, R.E., Hewagama, T., Bonev, B.P., DiSanti, M. A. Mandell,  
25 A.M., Smith M.D. (2009) Strong Release of Methane on Mars in Northern Summer 2003.  
26 *Science* 323, 1041-1045.  
27  
28  
29 Navarro-Gonzalez, R., Navarro, K.F., Rosa, J.D.L., Iniguez, E., Molina, P., Miranda, L.D.,  
30 Morales, P., Cienfuegos, E., Coll, P., Raulin, F., Amils, R., McKay, C.P. (2006) The  
31 limitations on organic detection in Mars-like soils by thermal volatilization-gas  
32 chromatography-MS and their implications for the Viking results. *Proceedings of the*  
33 *National Academy of Sciences of the United States of America* 103, 16089-16094.  
34  
35  
36 Noffke, N. (2009) The criteria for the biogenicity of microbially induced sedimentary structures  
37 (MISS) in Archean and younger sandy deposits. *Earth Science Reviews* 96, 173-180.  
38  
39  
40 Ohashi, H., Nakazawa, H. (1996) The microstructure of humic acid-montmorillonite composites.  
41 *Clay Minerals* 31, 347-354.  
42  
43 Owen, T., Bar-Nun, A., Kleinfeld, I. (1992) Possible cometary origin of heavy noble gases in the  
44 atmospheres of Venus, Earth and Mars. *Nature*, 358, 43-46.  
45  
46  
47 Oze, C. and Sharma, M. (2005) Have olivine, will Gas: Serpentinization and the abiogenic  
48 production of methane on Mars," *Geophysical Research Letters* 32, L10203-L10206.  
49  
50 Panieri, G., Lugli, S., Manzi, V., Roveri, M., Schrieber, B.C., Palinska, K.A. (2010) *Geobiology* 8,  
51 101-111.  
52  
53 Pizzarello, S. (2006) The chemistry of life's origin: a carbonaceous meteorite perspective.  
54 *Accounts of Chemical Research* 39, 231-7.  
55  
56  
57 Pratt, L.M., Lefticariu, L., Ripley, E.M., Onstott, T.C., (2006) Radiolysis of water as a source of  
58 bioavailable chemical energy. *Geochimica et Cosmochimica Acta* 70 (Issue18,  
59  
60

Supplement 1), A503.

- Proskurowski, G., Lilley, M.D., Seewald, J.S., Frueh-Green, G.L., Olson, E.J., Lupton, J.E., Sylva, S.P., Kelley, D.S. (2008) Abiogenic Hydrocarbon Production at Lost City Hydrothermal Field. *Science* 319, 604-607.
- Rasmussen, B. (2000) Filamentous microfossils in a 3,235-million-year-old volcanogenic massive sulphide deposit. *Nature* 405, 676-679.
- Rye, R.O. (2005) A review of the stable-isotope geochemistry of sulfate minerals in selected igneous environments and related hydrothermal systems. *Chemical Geology* 215, 5-36.
- Sephton, M.A. (2002) Organic compounds in carbonaceous meteorites. *Natural Products Research* 19, 292 - 311.
- Sephton, M.A., Botta O. (2008) Extraterrestrial organic matter and the detection of life. *Space Science Reviews* 135, 25-35.
- Sephton, M.A., Love, G.D., Watson, J.S., Verchovsky, A.B., Wright, I.P., Snape, C.E., Gilmour, I. (2004) Hydropyrolysis of insoluble carbonaceous matter in the Murchison meteorite: new insights into its macromolecular structure. *Geochimica et Cosmochimica Acta* 68, 1385-1393.
- Sherwood Lollar, B., Lacrampe-Couloume, G., Slater, G.F., Ward, J., Moser, D.P., Gihring, T.M., Lin, L.H., Onstott, T.C. (2006) Unraveling abiogenic and biogenic sources of methane in the Earth's deep subsurface. *Chemical Geology* 226, 328-339.
- Shock, E.L., Schulte, M.D. (1998) Organic synthesis during fluid mixing in hydrothermal systems. *Journal of Geophysical Research [Planets]* 103(E12), 28513-28527.
- Sleep, N.H., Bird, D.K. (2007) Niches of the pre-photosynthetic biosphere and geologic preservation of Earth's earliest ecology. *Geobiology* 5, 101-117.
- Sleep, N.H., Meibom, A., Fridriksson, T., Coleman, R.G., Bird, D.K. (2004) H<sub>2</sub>-rich fluids from serpentinization: geochemical and biotic implications. *Proceedings of the National Academy of Sciences of the United States of America* 101, 12818-12823.
- Smith, J.W., Kaplan, I.R. (1970) Endogenous Carbon in Carbonaceous Meteorites. *Science* 167, 1367-1370.
- Squyres, S. W., Arvidson, R. E., Bell, J. F., III, Bruckner, J., Cabrol, N. A., Calvin, W., Carr, M. H., Christensen, P. R., Clark, B. C., Crumpler, L., Des Marais, D. J., d'Uston, C., Economou, T., Farmer, J., Farrand, W., Folkner, W., Golombek, M., Gorevan, S., Grant, J. A., Greeley, R., Grotzinger, J., Haskin, L., Herkenhoff, K. E., Hviid, S., Johnson, J., Klingelhofer, G., Knoll, A., Landis, G., Lemmon, M., Li R., Madsen, M. B., Malin, M. C., McLennan, S. M., McSween, H. Y., Ming, D. W., Moersch, J., Morris, R. V., Parker, T., Rice, J. W., Jr., Richter, L., Rieder, R., Sims, M., Smith, M., Smith, P., Soderblom, L. A., Sullivan, R., Wanke, H., Wdowiak, T., Wolff, M., and Yen, A. (2004) The Spirit Rover's

1  
2  
3 Athena Science Investigation at Gusev Crater, Mars. *Science* 305, 794-799.  
4

5 Squyres, S.W., A.H. Knoll, R.E. Arvidson, B.C. Clark, J.P. Grotzinger, B.L. Jolliff, S.M.  
6 McLennan, N. Tosca, J.F. Bell III, W.M. Calvin, W.H. Farrand, T.D. Glotch, M.P.  
7 Golombek, K.E. Herkenhoff, J.R. Johnson, G. Klingelhofer, H.Y. McSween, and A.S.  
8 Yen (2006) Continuing observations by the Opportunity rover at Meridiani Planum,  
9 Mars. *Science* 313, 1403-1407.  
10  
11

12 Squyres, S.W., et al.(2009) Exploration of Victoria Crater by the rover Opportunity.  
13 *Science* 324, 1058-1061.  
14

15 Steele, A., Fries, M. D., Amundsen, H. E. F., Mysen, B. O., Fogel, M. L., Schweizer, M. (2007)  
16 Comprehensive imaging and Raman spectroscopy of carbonate globules from Martian  
17 meteorite ALH 84001 and a terrestrial analogue from Svalbard. *Meteoritics & Planetary*  
18 *Science* 42, 1549-1566.  
19

20 Summons, R.E., Albrecht, P., McDonald, G., Moldowan, J.M. (2007) Molecular Biosignatures:  
21 Generic Qualities of Organic Compounds that Betray Biological Origins. *Space Science*  
22 *Reviews* 135, 133-157.  
23  
24

25 Swindle, T.D. (2002) Martian Noble Gases. *Reviews in Mineralogy and Geochemistry* 47, 171-  
26 190.  
27  
28



29 Trewin, N. H. (1996) The Rhynie cherts: an early Devonian ecosystem preserved by  
30 hydrothermal activity. *Evolution of Hydrothermal Ecosystems on Earth (and Mars?)*, Ciba  
31 Foundation Symposium #202, pp. 131-45.  
32

33 Walter, M. R. (1996) Ancient hydrothermal ecosystems on earth: a new palaeobiological  
34 frontier. *Evolution of Hydrothermal Ecosystems on Earth (and Mars?)*, Ciba Foundation  
35 Symposium #202, pp. 112-27  
36  
37

38 Xiao S. and Knoll A. H. (1999) Fossil preservation in the Neoproterozoic Doushantuo  
39 phosphorite Lagerstätte, South China. *Lethaia* 32(3), 219-238.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



<b>Biosignatures detectable by MSL</b>	<b>How definitive as a biosignature?</b>	<b>How well it can be measured by MSL?</b>
biogenic organic molecules	highly definitive	readily with SAM
biogenic gases	often definitive	readily with SAM
body fossils	often definitive	with MAHLI if large enough
biofabrics	sometimes definitive	with MAHLI , MastCam
stable isotopic compositions	occasionally; context critical	readily with SAM
biomineralization/ alteration	rarely definitive	detectible with CheMin, ChemCam
spatial chemical patterns	rarely definitive	detectible with CheMin, ChemCam

Table 1

<b>Biosignatures taphonomic window</b>	<b>Confidence in the geological context</b>	<b>How this informs about potential biosignature preservation</b>
atmospheric gases	very high	predictable via chemical modeling
crystalline sedimentary mineral entrapment of organics	very high	can deduce formation mechanism and subsequent history
biofabric lithification	very high	can deduce history from lithology and stratigraphic relationships
body fossil preservation	very high	can deduce history from lithology and stratigraphic relationships
mineral replacement of body fossil	high	can deduce from mineralogy

Table 2

Martian context --> Early Mars Environment	Support biotic OM formation	Support for abiotic OM formation	Support OM conc'n	Support preservation	Potential for recent exhumation*	ID by remote sensing			ID by MSL
						Geo-morphic	Mineral-ogic	Strati-graphic	
Aeolian sediments (sand)	low	low	low	low	low ??	high	n/a	mod	high
altered aeolinites (dust)	very low	low	low	low	low ??	low	n/a	n/a	high
Fluvial channel	low	low	low	low	high	high	n/a	high	high
Fluvial floodplain	low-mod	low	mod	mod	possible ??	high	n/a	high	high
alluvial fan	low	low	low	low	low ??	high	n/a	high	high
Deltaic	high	low	high	high	low ??	high	n/a	high	high
Lacustrine (perennial)	high	low	high	high	high	mod	mod	mod	high
Lacustrine (evaporitic)(Cl)	low	low	high	high-very high	high	mod	high	mod	high
Lacustrine (evaporitic)(SO4)	mod	low	high	high-very high	high	mod	high	mod	high
Regional groundwater pore system	low	low	low	low	high ??	n/a	n/a	n/a	mod
Glacial deposits	low	low	low	low	high	high	n/a	low	high
permafrost	low	low	low	mod	mod	high	n/a	n/a	high
soil (surface fines chemically altered by atmosphere )	low	low	low	low	low	n/a	n/a (albedo and TI)	n/a	high
Regolith/Fractured Bedrock (not soil)	low	low	low	low	low	high	n/a	n/a	high

\* Poorly constrained estimates; require improved knowledge of erosional processes and timescales

Table 3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

Specific cases	Support biotic OM formation	Support for abiotic OM formation	Support OM conc'n	Support preservation	Potential for recent exhumation*	ID by remote sensing			ID by MSL
						Geo-morphic	Mineral-ogic	Strati-graphic	
Pyroclastic Deposits (unaltered)	low	low	low	low	low?	mod	low	high	high
Volcanic flows	very low	low	low	low	low?	high	high	mod	high
aqueous altered at surface	low	low	low	mod-high	low?	low	mod	low	mod
aqueous altered in subsurface	low	low	low	mod-high	low?	low	mod	low	mod
aqueous altered at surface	low	low	low	low-mod	low?	low	mod	low	mod
Volcanics-hydrothermal altered	mod	low	low	mod	mod	mod	mod	low	high
aqueous altered in subsurface	mod	mod	low	low-mod	mod	low	mod	low	mod
Hydrothermal (<100C) subsurface	mod	mod (F/T)	mod-low	mod	mod	mod	mod-high	n/a	high
Hydrothermal (<100C) surface	high	low	mod-high	mod	mod	high	mod-high	low	high
ultramafic subsurface (<100C)	high	low	mod-high	mod	mod	high	mod-high	low	high

\* Poorly constrained estimates;require improved knowledge of erosional processes and timescales

Table 4



	<b>Specific Processes</b>	<b>EGA</b>	<b>GCMS</b>	<b>APXS</b>	<b>CheMin</b>	<b>ChemCam</b>	<b>MAHLI</b>
<b>Abiotic Processes</b>	F/T and catalyzed polymerization	X	X	X	X		
<b>Bioprocess</b>	Redox dependent metabolisms (e.g. H <sub>2</sub> /Fe/S/C - metabolisms)	X	X	X	X	X	
<b>Conc Process</b>	Adsorption to mineral surfaces			X	X	X	
	Immiscibility & migration	X					X
<b>Preservation Process</b>	Adsorption to mineral surfaces			X	X	X	
	mineral encapsulation (trapping)	X				X	X
	thermal polymerization (pyrobitumen formation)	X			X	X	X (UV)

Table 5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

	<b>Specific Processes</b>	<b>EGA</b>	<b>GCMS</b>	<b>APXS</b>	<b>CheMin</b>	<b>ChemCam</b>	<b>MAHLI</b>
	Cosmic (footnote: specific targets for PE are organic molecules and Cr, Ni anomalies)		X	X	X		
	Atmospheric photolytic synthesis		X	X	X		
<b>Bioprocess</b>	chemosynthesis		X	X	X		
	photosynthesis	X (%TOC)	X	X	X		
<b>Conc Process</b>	sorting of transported organics and organics in/on minerals		X <sup>#</sup>	X <sup>#</sup>	X	X	X
<b>Preservation Process</b>	lithification (includes cementation)	X		X	X	X	X
	Adsorption to mineral surfaces			X	X	X	X
	Burial			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X		X	X	X	X

Table 6

	<b>Specific Processes</b>	<b>EGA-TLS</b>	<b>EGA-QMS</b>	<b>GCMS</b>	<b>APXS</b>	<b>CheMin</b>	<b>ChemCam</b>	<b>MAHLI</b>
<b>Fractionation-Related Processes</b>	impacts (excavation for access deep-carbon sources)	X	X					
	atmospheric processes (e.g. photolysis)	X	X?					
	Biology	X	X?			x		
	hydrologic cycle (over martian history indirectly understood via CO2 record)	X			x	x		
	Global C cycling	X			x	x		
	Global S cycling		X		x	x		
	other biogeochemical cycling	X	X		x	x		
	volcanism	X	X		x	x		
<b>Environmental State</b>	atmospheric composition (past & present), temperature, pressure	X	X					
	water inventories, state, and activity	X	X			x		
	Redox state	X	X		X	x		
	pH	X	X		X	x		
	surface materials (in exchange with atmosphere)	X	X			x		
<b>Preservation Process</b>	lithification (includes cementation)	X	X		X	X	X	X
	Adsorption to mineral surfaces	X			X	X	X	X
	Burial	X			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X	X		X	X	X	X

Table 7

	Specific Processes	EGA-TLS	EGA-QMS	GCMS	APXS	CheMin	ChemCam	MAHLI
<b>Processes</b>	impacts (excavation for access deep-carbon sources)	X	X					
	cosmic (influx and loss)	X	X					
	weathering	X	X					
	atmospheric processes (e.g. photolysis)	X	X?					
	Biology	X	X?			X		
	hydraulic cycle (over martian history indirectly understood via CO2 record)	X			X	X		
	C cycling	X			X	X		
	S cycling		X		X	X		
	other biogeochemical cycling	X	X		X	X		
	volcanism	X	X		X	X		
	aeolian (fractionation via oxidation of materials or concentration of materials from particular sources)	X	X			X		
<b>Environmental State</b>	atmospheric composition (past & present), temperature, pressure	X	X					
	water inventories, state, and activity	X	X			X		
	Redox state	X	X		X	X		
	pH	X	X		X	X		
	surface materials (in exchange with atmosphere)	X	X			X		
<b>Preservation Process</b>	lithification (includes cementation)	X	X		X	X	X	X
	Adsorption to mineral surfaces	X			X	X	X	X
	Burial	X			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X	X		X	X	X	X

Table 8

	<b>Processes</b>	<b>Features</b>	<b>Preservation Window</b>
<b>Physical</b>	Brecciation Fracturing	hydraulic breccia/fractures	Avoidance of strong structural deformation and very high thermal metamorphism
<b>Chemical</b>	Mineral Precipitation +- Alteration, Dissolution	cement distribution & composition, alteration crusts, porosity, dissolution surfaces, laminations, terraces, mounds	Avoidance of recrystallization & dissolution, increases in oxidation state (e.g. Fe/Mn, perchlorate, sulfate, clay hydration); avoidance of structural deformation and very high thermal metamorphism
<b>Biological</b>	Biomass Accumulation	"mats, stromatolites, etc; roll-ups, streamers, elemental concentrations, coking	Avoidance of strong structural deformation and very high thermal metamorphism; coke in fractures

Table 9



	<b>Processes</b>	<b>Features</b>	<b>Preservation Window</b>
<b>Physical</b>	Sediment Transport; Soil/permafrost	grain size, sorting, rounding, composition cross strat laminations, cracks/fracturing soft sediment deposition	dewatering, recrystallization of sulfates, hydration changes; structural deformation and very high thermal metamorphism
<b>Chemical</b>	Mineral Precipitation +- Alteration, Dissolution	cement distribution and composition alteration crusts porosity dissolution surfaces concretions displacive recrystallization lamination	Avoidance of recrystallization & dissolution, increases in oxidation state (e.g. Fe/Mn, perchlorate, sulfate, clay hydration); avoidance of structural deformation and very high thermal metamorphism
<b>Biological</b>	Biomass Accumulation	mats, stromatolites, etc roll-ups, wrinkled structures elemental concentrations fenestrae grain size variations	Avoidance of recrystallization, hydration, elemental mobility, organic remineralization; Avoidance of structural deformation and very high thermal metamorphism

Table 10

Table 11: Chemical and Mineralogic Features of Hydrothermal Environments

Processes	Features	Environmental Indicators	Preservation Window	Payload elements	
<b>Environmental</b>	Fluid Convection, selective mineral phase dissolution, alteration, migration, and precipitation; element leaching and transport; Redox chemistry; cooling/heating;	mineral and element zonation/gradients (i.e. characteristic mineralogy associated with variations of hydrothermal environments, such as: chlorite, silica sulfides, kaolinite, hematite, gold, carbonates, sulfates; etc.); spatial distribution of alteration minerals and precipitates; mineral assemblages for redox and temperature*; fracture fills; pH differences; m-km scale mineral zonation		encasement of sensitive phases in less sensitive phases (e.g. silica or organic entombment of minerals); avoidance of oxidation, metamorphism, continued hydrothermal activity	Mastcam, APXS, Chemcam, CheMin, SAM-EGA
<b>Biological</b>	Accelerated redox reactions, organic synthesis, altered mineral assemblages	Altered mineral or elemental abundances, corroded (e.g., bored) minerals, biofabrics, organic matter	Mineral, elemental and textural indicators of aqueous conditions and mineral precipitation	Entombment in minerals formed during cooling or other changes that favor precipitation	MAHLI, APXS, Chemcam, CheMin, SAM-EGA
<b>Physical</b>	Fluid Convection	Hydraulic brecciation; Geomorphic evidence of springs; Overall architecture of system	Heat source	Broad due to large scale	MastCam
	Temperature Changes	Heat flow; Induces variations in chemical properties in time and over spatial scales from m to km	Heat source; Flow patterns	Large scale variations have high preservation potential; small scale changes require avoidance of recrystallization and elemental leaching	Mastcam, MAHLI, ChemCam
<b>Chemical, Potentially Biological</b>	Mineral Precipitation	Evidence of water chemistry & temperature; Springs mounds & terraces; Some mineral precipitation can be localized by organic carbon	Temperature indicators from low T to high T: BaSO <sub>4</sub> , PbS, ZnS, SO <sub>4</sub> /Sulfides together, saddle dolomite, CuS, CaSO <sub>4</sub> ; Low pH indicators: jarosite, abundant kaolinite, alunite vs carbonate	Encasement of sensitive phases in less sensitive phases (e.g. silica or organic entombment of minerals); Avoidance of oxidation, metamorphism, continued hydrothermal activity	MastCam, MAHLI, APXS, ChemCam, CheMin, SAM-EGA
	Mineral Alteration & Dissolution	Evidence of changes in water chemistry			
	Elemental Leaching & Transport	m to km-scale spatial gradients in elemental composition			
	Redox Changes	Important redox indicators: S <sup>2-</sup> , S <sup>0</sup> , S <sup>6+</sup> , Fe <sup>2+</sup> , Fe <sup>3+</sup> , Mn <sup>2+</sup> , Mn <sup>4+</sup> , U <sup>2+</sup> , U <sup>4+</sup> , As <sup>3+</sup> , As <sup>5+</sup>			

**\* Temp mineral assemb.**

mod T SO<sub>4</sub>/sulfide together  
saddle dolomite  
Low T --> high T PbS, ZnS, CuS  
Low T --> high T BaSO<sub>4</sub>--> CaSO<sub>4</sub>

**\* Redox mins**

S<sup>2-</sup> --> S<sup>0</sup>-->S<sup>6+</sup>  
Fe<sup>2+</sup> --> Fe<sup>3+</sup>  
Mn<sup>2+</sup> --> Mn<sup>4+</sup>  
U<sup>2+</sup> --> U<sup>4+</sup>  
As<sup>3+</sup> --> As<sup>5+</sup>

**\* low pH mins**

jarosite, large kaolinite conc., alunite vs carb.

Important factors	Factor related Feature	Preservation Window	Payload elements
Sediment source	detrital mineral assemblages	Early lithification is favorable. Avoid the following: sediments altered by recrystallization and dissolution (especially evaporites), redox changes toward oxidizing conditions, and strong structural alteration and strong thermal metamorphism	Mastcam, MAHLI, APXS, Chemcam, CheMin, SAM-EGA
H <sub>2</sub> O solute chemistry and flux	evaporite assemblages and trace element distribution		
Redox and pH	Redox sensitive (Fe, Mn, S, U, etc.) and pH sensitive (Jarosite, alunite, kaolinite, etc. Vs carbonates) mineral phase and abundance		
bio-mineral interactions	bio-magnetite, bio carbonate, bio-sulfides, oxides, phosphate precipitation; bioleaching and dissolution products		

Table 12