



Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Summons, Roger E., Jan P. Amend, David Bish, Roger Buick, George D. Cody, David J. Des Marais, Gilles Dromart, Jennifer L. Eigenbrode, Andrew H. Knoll, and Dawn Y. Sumner. 2011. "Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group." <i>Astrobiology</i> 11, no. 2: 157–181.
Published Version	doi:10.1089/ast.2010.0506
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:13041033
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP

Astrobiology

Astrobiology Manuscript Central: <http://mc.manuscriptcentral.com/astrobiology>

Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

Journal:	<i>Astrobiology</i>
Manuscript ID:	AST-2010-0506.R2
Manuscript Type:	News and Views
Date Submitted by the Author:	13-Dec-2010
Complete List of Authors:	Summons, Roger E.; Massachusetts Institute of Technology, Earth, Atmospheric and Planetary Sciences Amend, Jan; Washington University Buick, Roger; University of Washington Bish, David; Indiana University Cody, George; CIW Des Marais, David; NASA Ames Research Center Dromart, Gilles Eigenbrode, Jennifer; NASA Goddard Space Flight Center Knoll, Andrew; Harvard University, OEB
Keyword:	Astrobiology, Atmospheric Gases, Mars, Biosignatures, Life Detection

SCHOLARONE™
Manuscripts

Preservation of Martian Organic and Environmental Records: Final Report of the Mars Biosignature Working Group

Roger E. Summons^{1*}, Jan P. Amend², David Bish³, Roger Buick⁴, George D. Cody⁵,
David J. Des Marais⁶, Gilles Dromart⁷, Jennifer L. Eigenbrode⁸, Andrew H. Knoll⁹,
Dawn Y. Sumner¹⁰

1. Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Ave., Cambridge MA 02139, USA
2. Department of Earth & Planetary Sciences, Washington University, 1 Brookings Dr., Saint Louis MO 63130, USA
3. Department of Geological Sciences, Indiana University, 1001 East 10th St., Bloomington, IN 47405, USA
4. Department of Earth & Space Sciences, University of Washington, 4000 15th Avenue NE, Seattle, WA 98195, USA
5. Geophysical Laboratory, Carnegie Institution of Washington, 5251 Broad Branch Road NW, Washington, DC 20015, USA
6. Exobiology Branch, NASA Ames Research Center, Mail Stop 239-4, Moffett Field, CA 94035, USA
7. Laboratoire de Science de la Terre, UMR CNRS 5570, Université Claude Bernard/Ecole Normale Supérieure de Lyon, 2 rue Raphaël Dubois, 696222 Villeurbanne Cedex, France
8. Solar System Exploration Division, NASA Goddard Space Flight Center, Mail Code 699, 8800 Greenbelt Rd., Greenbelt, MD 20771, USA
9. Departments of Earth and Planetary Sciences and Organismic and Evolutionary Biology, 26 Oxford St., Cambridge MA 02138, USA
10. Geology Department, University of California Davis, 1 Shields Ave, Davis, CA 95616, USA

* Corresponding Author: rsummons@mit.edu

Abstract

The Mars Science Laboratory (MSL) has an instrument package capable of making measurements of past and present environmental conditions. The data generated may tell us if Mars is, or ever was, able to support life. However, the knowledge of Mars' past history and the geological processes most likely to preserve a record of that history remain sparse and, in some instances, ambiguous. Physical, chemical and geological processes relevant to biosignature preservation on Earth, especially under conditions early in its history when microbial life predominated, are also imperfectly known. Here we present the report of a working group chartered by the Co-Chairs of NASA's MSL Project Science Group, John P Grotzinger and Michael A Meyer, to review and evaluate potential for biosignature formation and preservation on Mars.

Orbital images confirm that layered rocks achieved kilometer-scale thicknesses in some regions of ancient Mars. Clearly, interplays of sedimentation and erosional processes govern present-day exposures, and our understanding of these processes is incomplete. MSL can document and evaluate patterns of stratigraphic development as well as the sources of layered materials and their subsequent diagenesis. It can also document other potential biosignature repositories such as hydrothermal environments. These capabilities offer an unprecedented opportunity to decipher key aspects of the environmental evolution of Mars' early surface and aspects of the diagenetic processes that have operated since that time.

Considering the MSL instrument payload package, we identified the following classes of biosignatures as within the MSL detection window: organism morphologies (cells, body fossils, casts), biofabrics (including microbial mats), diagnostic organic molecules, isotopic signatures, evidence of biomineralization and bioalteration, spatial patterns in chemistry and biogenic gases. Of these, biogenic organic molecules and biogenic atmospheric gases are considered the most definitive and most readily detectable by MSL.

Introduction

In the first decade of the twenty-first century, our understanding of Mars and its environmental history has increased dramatically. Orbital measurements provide unprecedented resolution of both physical and chemical features of the Martian surface. The Mars Exploration Rovers Opportunity and Spirit have contributed our first-ever geologist's-eye views of stratigraphic successions on Mars (e.g. Christensen et al., 2004; Squyres et al., 2004, 2006, 2009; Grotzinger et al., 2005; Haskin et al., 2005). Building on this success, the extraordinary instrument package and anticipated roving capability of the Mars Science Laboratory (MSL) position us to use new rover observations to test hypotheses generated on the basis of high-resolution orbital data. For example, orbital images and spectral data show that layered rocks have accumulated to thicknesses greater than a kilometer on some parts of the ancient Martian surface and comprise diverse lithologies (e.g. Bishop et al., 2008; Ehmann et al., 2008a,b). What fundamental processes of particle generation, transport, and cementation made such accumulations possible, and what erosional processes govern present day exposures? MSL has the imaging capability to document and evaluate patterns of stratigraphic development, as well as the geochemical capacity to evaluate the sources of layered materials and their

1
2
3 subsequent diagenesis. Bibring et al. (2005) have identified phyllosilicates by remote sensing of
4 the Martian surface and hypothesized that these record an early, relatively wet epoch in Martian
5 history, before the time of acid sulfate deposition recorded in the sediments at Meridiani
6 Planum. MSL has the capacity to confirm phyllosilicate mineralogy and evaluate it in
7 stratigraphic context, allowing us to understand more fully the significance of these minerals for
8 reconstructing Martian environmental history. Finally, the hallmark of life is organic matter —do
9 Martian strata preserve organic molecules that might illuminate the planet's early environmental,
10 and possibly biological, history? MSL's Sample Analysis at Mars (SAM) instrument suite
11 provides unprecedented potential to answer this question.
12
13
14

15 Knowledge of our own planet's biological and environmental history has developed through the
16 integration of many types of observations. Arguably among the most important are: field
17 mapping, the measurement and correlation of stratigraphic sections, and both paleontological
18 and geochemical analyses of samples collected from measured sections located within mapped
19 terrains. Through the strategic choice of a landing site using high-resolution orbital data, MSL
20 promises a comparable integration for Mars. With this in mind, it is helpful, even necessary, to
21 consider how geologists evaluate and select field sites on Earth. Surely, site accessibility and
22 map data influence choice, as do accumulated observations by previous geologists. But there
23 is one more consideration that governs site selection by geologists, geochemists, and
24 paleobiologists: what types of deposits are most likely to preserve geological and possible
25 biological signals of interest? Simply put, NASA will realize the greatest returns on its
26 investment in the MSL payload if it targets outcrops that maximize the probability of organic
27 deposition and preservation. It will maximize its return from the payload if it targets stratigraphic
28 successions that place mineralogical measurements in temporal and paleoenvironmental
29 context.
30
31
32
33
34

35 **Working Group Objectives**

36
37 The Preservation Working Group was assembled to assess whether current understanding of
38 organic matter preservation on Earth might help guide site selection and both strategic as well
39 as tactical planning during surface operations for the MSL mission. The Working Group was
40 asked to provide general guidance on what specific geologic environments would be most
41 favorable for preservation of potential biosignatures, including special consideration of organic
42 carbon (Farmer and Des Marais, 1999).
43
44

45 Insert Text Box 1 and Box 2 hereabouts
46

47 It is essential to recognize that, on Earth, and in spite of a vital biosphere, the majority of
48 biologically derived organic carbon exists as "fossil" organic carbon stored within layered
49 sedimentary rocks. This sequestered organic matter is by some estimates 2×10^5 that the mass
50 of the carbon stored in the living biosphere. On Mars, it appears reasonable to assume that if
51 life exists or ever existed, it never evolved to the point of large differentiated, multi-cellular
52 organisms (e.g., plants) that biosynthesize large quantities of recalcitrant biopolymers (e.g.,
53 lignin and cellulose) leading to potential accumulations of extensive organic matter rich
54 sedimentary deposits (e.g., coal). There is general consensus that extant microbial life on Mars
55
56
57
58
59
60

1
2
3 would likely exist (if at all) in the subsurface and at low abundance, making it difficult, if not
4 impossible, to detect using sampling technologies foreseeable over the next decade.
5

6
7 Detecting an ancient subsurface biota may be even more difficult unless exhumed records of
8 ancient subsurface environments can be confidently recognized and are accessible to a rover.
9 Since all life requires an abundance of electron donors and acceptors to grow and to
10 accumulate biomass, we would also need analyze the mineralogy and geochemistry of
11 exhumed rocks to assess their bioenergetic potential. Even though the record of early life on
12 Earth is reported to contain microstructures (e.g. Rasmussen, 2000; Brasier et al., 2006)
13 purported to reflect subsurface life, significant doubt remains about their biogenicity. These
14 remains are rare and notoriously difficult to interpret with confidence given their high degree of
15 thermal alteration. In contrast, it is possible that the organic remains of extinct microbial life
16 developed at Mars' surface, or subsurface, may persist and perhaps even be enriched in certain
17 sedimentary rocks.
18
19

20
21 On Earth, microorganisms commonly exist communally in the water column and in sediment
22 pore spaces or as attached biofilms. Biomass of these communities may be preserved in the
23 rock record in concentrations sufficient to detect with the MSL payload elements (see Box 1).
24 Furthermore, sedimentary processing (e.g., hydrodynamic sorting as occurs in fluvial and deltaic
25 environments on Earth) may concentrate biologically derived carbonaceous particles into
26 organic-rich horizons in sedimentary strata. A logical approach to establishing whether life ever
27 existed on Mars is, therefore, to analyze appropriate sedimentary lithologies, seeking evidence
28 of "fossilized" organic matter preserved in sedimentary deposits.
29
30
31

32 One issue that may add ambiguity in such a search is the fact that, throughout Martian history,
33 organic-rich chondritic meteorites have undoubtedly rained upon the Martian surface.
34 Weathering of chondritic meteoritic debris in an environment with minimal oxygen and no
35 extensive surface biosphere (i.e., microbes and fungi capable of degrading chondritic organic
36 matter) may lead to a persistence of extraterrestrial organic particles and, consequently, its
37 accumulation into certain sedimentary rocks through hydrodynamic sorting. At the same time,
38 strong oxidants in Martian regolith along with exposure to ionizing radiation might alter or
39 destroy molecular signatures from meteorites or organisms. Notwithstanding the potential
40 complexity of interpreting any organic material that may be detected, the simple detection of
41 organic matter in rock or soil via the MSL mission will constitute a critically important result.
42
43
44

45 The membership and objectives of the Working Group were chartered by the Co-Chairs of
46 NASA's MSL Project Science Group, John P Grotzinger and Michael A Meyer. The charge to
47 the Working Group was to outline a search strategy that included consideration of the nature of
48 the payload elements (PE), a ranked order of biosignatures detectable by those instruments and
49 an evaluation of the types of environments conducive to the formation and, especially,
50 preservation of these biosignatures. Four candidate landing sites remain at the time of writing
51 and this strategy could serve as a guide to members of the Mars exploration community in
52 deliberating whether potential landing sites have more or less appropriate lithologies for
53 biosignature preservation. A second objective was to assess, in very general terms, how the
54 MSL instrumentation may be best exploited for the analysis of any record of organic compounds
55
56
57
58
59
60

(biogenic or abiogenic) that might be preserved in Martian sedimentary rocks. The working group was asked not to address the merits or otherwise of specific Mars localities. Nor were we asked to assess instrument specifications and how those instruments might be operated at the surface of Mars, as these are issues more appropriately addressed by very different groups of specialists. We were guided in our work by two recent publications from the National Academies namely the report on the *Limits of Life in Planetary Systems* (Baross et al., 2007) and an *Astrobiological Strategy for the Exploration of Mars* (Jakosky et al., 2007).

Taphonomic Windows

Taphonomy is the term used by paleontologists to describe the processes by which living organisms become fossils. Taphonomy is commonly portrayed as “post-mortem information loss” because fossilized bones, shells, or microorganisms record only a small subset of the biological information originally present in their makers. With a different perspective, however, we might view taphonomy as “post-mortem information preservation” because, however selective it may be, fossilization provides our only permanent record of past life. In the context of MSL, we can broaden the concept of taphonomy to include the geologic preservation of the full suite of materials we might wish to measure using the rover’s instrument package. What combination of processes, for example, maximizes the probability of preserving organic molecules, body fossils and sedimentary textures?

If we can understand the processes that facilitate preservation, then we can conceive of the likely operation of those processes in time and space. That is, we can define a taphonomic window – the sedimentary and diagenetic circumstances most conducive to preservation. We know that ancient organic molecules will preserve only if buried in sediments and long-term preservation will occur only if buried organics are shielded from oxidizing fluids. On Mars, surface oxidants and UV radiation will have altered or destroyed organic molecules (Benner et al., 2000; Navarro-González et al., 2006; Sumner, 2006). On Mars, surface oxidants and UV radiation will have altered or destroyed organic molecules at or near the surface (Benner et al., 2000; Navarro-González et al., 2006; Sumner, 2006). If we can identify generic features that can be used to characterize potential landing sites with respect to the presence or absence of specific taphonomic windows, then we can help maximize the chances of successful analyses by MSL.

Working Group Findings

A recent review of biosignatures and strategies for their use in life detection can be found in Botta et al., (2008). Our deliberations yielded the ranked order of biosignatures and taphonomic windows that are provided in Tables 1 & 2. However, we also recognized that, even on Earth with its vigorous and multi-billion-year-old biosphere, many of these biosignatures can be ambiguous and are preserved only under rare and exceptional circumstances.

Insert Table 1 hereabouts

Insert Table 2 hereabouts

Box 1: Biosignatures at a glance – summary of what can be observed with the payload elements (PE) of MSL. Note that environments can be reconstructed from physical and chemical features of ancient sediments that are not considered to be biomarkers. (see tables 1 & 2)

1	Organism Morphologies (cells, body fossils, casts) PE: MAHLI Minimum size would have to be greater than 100 μm and rock preparation techniques are not available to expose organisms within rock. Martian life is expected to be microbial, so the probability of detection is low Potential as a biosignature: exceptionally high Potential as an environmental indicator: low
2	Biofabrics (including microbial mats) PE: MAHLI, MastCam Accreted structures analogous to those on Earth are detectable; However, few bedding plane surfaces are exposed, so potential surface biosignatures will be difficult to detect Potential as a biosignature: moderate Potential as an environmental indicator: low
3	Diagnostic organic molecules; Organic carbon PE: SAM, \pm ChemCam only if very abundant. Detection potential high incl. atmospheric gases Potential as a biosignature: exceptionally high Potential as an environmental indicator: high
4	Isotopic Signatures PE: SAM Contextual knowledge is essential; results can be ambiguous and complex to interpret Potential as a biosignature: moderate Potential as an environmental indicator: low
5	Biom mineralization & bioalteration PE: CheMin, \pm MAHLI, \pm SAM, \pm APXS detection of specific minerals is good morphological pattern may be useful but needs very fine spatial resolution Potential as a biosignature: low Potential as an environmental indicator: low
6	Spatial patterns in chemistry PE: SAM, CheMin, \pm ChemCam if very abundant C, N, S elemental distributions; Detection potential on cm scale to facies scale Potential as a biosignature: low on its own Potential as an environmental indicator: low
7	Biogenic Gases (Non-equilibrium) PE: SAM Excellent capacity to detect gases Potential as a biosignature: high (e.g., CH_4) Potential as an environmental indicator: low

Site selection decision-making and landed operations of the MSL should support the search for all of the above-mentioned biosignatures. However, it is also clear that accumulations of organic matter above meteoritic background levels would be amongst the most easily detected and least ambiguous. If life ever existed on Mars, it was likely microbial and existed communally in fluids and/or sediments, either free-living or as biofilms. Biomass from such communities, whatever their physiological characteristics, might then be preserved in the Martian sedimentary rock record in sufficient concentration to be detectable with MSL payload elements. Notwithstanding the numerous possibilities for habitable niches on Mars, a first-order approach to evaluating the organic record there would be to seek evidence of sedimentary environments -- the preferred geological setting for study - - that may have elevated concentrations of biologically derived carbonaceous materials (Table 3). These may include standing water environments such as deltaic systems, that promote retention of *in situ* organics and environments that hydrodynamically concentrate organics. An alternative approach would be to seek concentrations of buried crystalline minerals, such as

clays and evaporites which may protect organic matter from the destructive effects of ionizing radiation and strong oxidants. We also identify and order a range of other possibilities for environments conducive to the preservation of the signs of former and extant life (Tables 3 and 4).

MSL's Unique Combination of Capabilities

The MSL rover houses a remarkable suite of analytical instrumentation within a mobile platform—a distinct advantage for field investigations. Once on the surface of Mars, this rover will support several years of exploration and sample analysis. The analysis of fossil organic matter entrained in sedimentary rocks requires a proven approach, which can be implemented by the SAM instrument. Extensive studies of fossil organic remains in terrestrial rocks demonstrate that in most cases macromolecular constituents make up the overwhelming bulk of

Box 2: Molecular biosignatures at a glance

- Enantiomeric excess
Homochirality characterizes terrestrial biochemicals; strong preference for one enantiomer over the other in a chiral molecule. Interpretation is complicated by the discovery of L- excess in meteoritic amino acids (Engel & Macko 1997, Cronin & Pizzarello 1997, Pizzarello 2006, Glavin & Dworkin 2009)
- Diastereoisomeric preference
Strong isomer preferences in molecules with more than one center of asymmetry (Summons et al., 2007)
- Structural isomer preference
Observing a limited subset of the possible stable structural isomers in a complex molecule (Summons et al., 2007)
- Repeating constitutional sub-units or atomic ratios
Signifies complex molecules constructed from small common building blocks as in terrestrial biochemistry (McKay, 2004; 2007; Summons et al., 2007)
- Systematic isotopic ordering at molecular and intramolecular levels
As above
- Uneven distribution patterns or clusters (e.g., C-number, concentration, $\delta^{13}\text{C}$) of structurally related compounds.
As above

geologically preserved organic material. Pyrolytic breakdown of biopolymeric material (or diagenetically modified biopolymeric material) into small, volatile and identifiable fragments in a specially engineered device at the front end of a gas chromatograph-mass spectrometer (GC-MS) is a proven and manageable analytical approach because it can provide a molecular fingerprint of fossil organisms, in many cases yielding critical information regarding the identity of the organism through the presence of characteristic molecular biomarkers. Some generic characteristics of biomarkers are summarized in Box 2. A combination of these features should be present in biomolecules, whatever their

origin, and most of them are detectable with the SAM instrument. It is noteworthy that interstellar organic matter that may also be present in certain sedimentary rocks on Mars, also is amenable to characterization via pyrolysis GC-MS and may be distinguished from biologically derived material through specific kinds of molecular distributions (Sephton & Botta, 2008 and references therein).

1
2
3 Aside from MSL's capabilities to provide high-resolution organic molecular information, the
4 addition of CheMin and ChemCam to the traditional APXS broadens the capability to search for
5 ancient life on Mars. The ability to determine quantitatively the major mineral and chemical
6 constituents of prospective rocks will no doubt aid in assessing whether Mars had an active
7 biosphere at a much earlier time in its history. In terrestrial rocks, the presence of biological
8 organic matter can effect changes in local mineralogy by a number of different processes.
9
10

11 In some cases even if the organic matter has ultimately been destroyed through long-term
12 exposure to oxidizing fluids, specific mineral associations with the now-absent organics may
13 remain, providing a more persistent biomarker. Finally, certain minerals that exist on Earth on a
14 global scale are generally recognized to be the consequence of an active biosphere, e.g.,
15 biogenic carbonates, hazenite ($\text{KNaMg}_2(\text{PO}_4)_2 \cdot 14\text{H}_2\text{O}$), or brushite ($\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$). Hazen et
16 al. (2008) list and describe a large variety of mineral species that can be produced by biological
17 processes. The presence of CheMin in the analytical suite of MSL is, therefore, of profound
18 importance for the detection of ancient life.
19
20
21

22 The essential factor in addressing whether organic matter may be preserved in the Martian
23 sedimentary record is the ability to access appropriate lithologies (Tables 3 & 4). In this regard,
24 mobility is critical. As the MER mission has clearly demonstrated, only an "eyes-on-the-ground"
25 approach can provide the information required to interpret the lithologies of outcrops.
26 Integrating this mobile capability with analytical instrumentation provides the means to address
27 whether organic matter preservation has occurred or not.
28
29
30
31
32

33 Insert tables 3 and 4 hereabouts
34
35

36 **Brief description of the payload elements**

37 Full descriptions of the MSL payload instruments can be found at: [http://msl-
38 scicorner.jpl.nasa.gov/Instruments/](http://msl-scicorner.jpl.nasa.gov/Instruments/). In addition to the suite of analytical tools there are four
39 cameras. The Mast Camera (MastCam), the Mars Hand Lens Imager (MAHLI), the Remote
40 Micro-Imager on ChemCam (RMI) and the Mars Descent Imager (MARDI).
41
42
43

44 MastCam will take color images and color video footage of the Martian terrain. These can be
45 stitched together to create panoramas of the landscape around the rover. MastCam consists of
46 two camera systems mounted on a mast extending upward from the rover deck and will be used
47 to study the Martian landscape, rocks, and soils; to view frost and weather phenomena; and to
48 support the driving and sampling operations of the rover.
49
50

51 MAHLI will provide close-up views of the minerals, textures, and structures in Martian rocks and
52 the surface layer of rocky debris and dust. The self-focusing, roughly 4-centimeter-wide (1.5-
53 inch-wide) camera will take color images of features as small as 14 micrometers and will carry
54 both white light and ultraviolet light sources making the imager functional both day and night.
55 The ultraviolet light will be used to induce fluorescence to help detect carbonate and evaporite
56 minerals, both of which would indicate that water helped shape the landscape on Mars.
57
58
59
60

1
2
3 RMI will provide telescopic views of the surfaces to be analyzed by LIBS, and will put LIBS
4 analyses into their geologic context. However, the RMI can also be used to image textures
5 whether or not the LIBS is used. The RMI has a field of view of 19 milliradians. Due to
6 optimization of the telescope for LIBS, the RMI resolution is not pixel-limited, and is
7 approximately 80 microradians. The RMI can clearly distinguish the submillimeter LIBS spot on
8 a metal plate at a distance of at least 10 m. Therefore, this camera has the ability to make
9 discoveries of texturally-based potential biosignatures, such as stromatolites and other textures
10 suggestive of former microbial processes.
11
12

13
14 MARDI will take color video during the rover's descent toward the surface, providing an
15 "astronaut's view" of the local environment. As soon as the rover jettisons its heat shield several
16 kilometers above the surface, the Mars Descent Imager will begin producing a five-frames-per-
17 second video stream of high-resolution, overhead views of the landing site. It will continue
18 acquiring images until the rover lands, storing the video data in digital memory. After landing on
19 Mars, the rover will transfer the data to Earth. MARDI will provide information about the larger
20 geologic context surrounding the landing site.
21
22

23
24 ChemCam combines laser-induced breakdown spectroscopy with a remote micro-imager that
25 provides images of the target. It will provide elemental analysis of spatially resolved solid
26 samples at distances of 1-9 m. ChemCam's primary objective is to determine the chemical
27 composition of rocks and regolith in order to characterize the materials in the vicinity of the
28 rover.
29
30

31 The Alpha Particle X-Ray Spectrometer (APXS) will measure the abundance of chemical
32 elements in rocks and soils. It will be placed in contact with rock and soil samples on Mars and
33 will expose the material to alpha particles and X-rays emitted during the radioactive decay of
34 curium. The MER rovers have used APXS successfully, but the greatest benefit of APXS will be
35 when it is integrated with mineralogical analyses made by CheMin.
36
37

38 CheMin is a mineralogy instrument that will unequivocally identify and quantify the minerals
39 present in rocks and soil delivered to it by the Sample Acquisition, Sample Processing and
40 Handling (SA/SPaH) system. By determining the mineralogy of rocks and soils, CheMin will
41 enable assessments of the involvement of water in their formation, deposition, or alteration.
42 CheMin data will also be useful in the search for potential mineral biosignatures, energy sources
43 for life, or indicators of past habitable environments. CheMin can unequivocally identify and
44 quantify minerals above its detection limits in complex natural samples such as basalts, multi-
45 component evaporite systems, and soils.
46
47

48
49 The Sample Analysis at Mars (SAM) Suite Investigation in the MSL Analytical Laboratory is
50 designed to address the present and past habitability of Mars by exploring molecular and
51 elemental chemistry relevant to life. SAM evaluates carbon chemistry through a search for
52 organic compounds, the chemical state of light elements other than carbon, and isotopic tracers
53 of planetary change. SAM is a suite of three instruments, a Quadrupole Mass Spectrometer
54 (QMS), a Gas Chromatograph (GC), and a Tunable Laser Spectrometer (TLS). The QMS and
55 the GC can operate together in a GC-MS mode for chromatographic separation and mass
56 spectral identification of organic compounds. The TLS obtains precise stable isotope ratios for C
57
58
59
60

1
2
3 and O in carbon dioxide as well as C isotopes and abundance of trace methane.
4

5 The Radiation Assessment Detector (RAD) will detect and analyze the most hazardous
6 energetic particle radiation on the surface of Mars. Characterizing and understanding the
7 radiation environment on the Martian surface is fundamental to quantitatively assessing the
8 habitability of the planet (both past and present) and is essential for future manned Mars
9 missions. RAD will address radiation effects on biological potential and past habitability, as well
10 the contribution to chemical alteration of the regolith due to impinging space radiation.
11
12

13 REMS is designed to record six atmospheric parameters: wind speed/direction, pressure,
14 relative humidity, air temperature, ground temperature, and ultraviolet radiation. All sensors are
15 located around three elements: two booms attached to the rover Remote Sensing Mast (RSM),
16 the Ultraviolet Sensor (UVS) assembly located on the rover top deck, and the Instrument
17 Control Unit (ICU) inside the rover body.
18
19

20 The Dynamic Albedo of Neutrons (DAN) instrument is an active/passive neutron spectrometer
21 that measures the abundance and depth distribution of H- and OH-bearing materials (e.g.,
22 adsorbed water, hydrated minerals) in a shallow layer (0.5-1 m) of Mars' subsurface along the
23 path of the MSL rover.
24
25

26 The Sample Acquisition/Sample Processing and Handling subsystem (SA/SPaH) for MSL
27 consists of a powdering drill (Powder Acquisition Drill System or PADS) together with a
28 scooping, sieving, and portioning device (Collection and Handling for Interior Martian Rock
29 Analysis or CHIMRA) mounted on a turret at the end of a robotic arm. There is also a dust
30 removal tool (DRT) for clearing the surface of rocks prior to sampling. The drill enables powder
31 to be acquired from depths of 20 to 50 mm over a wide range of rock hardness with the top ~15-
32 20 mm being discarded. The scoop also enables samples of soil to be acquired, sieved and
33 apportioned. Five bricks of a silicon dioxide ceramic organic check material (OCM) are mounted
34 in canisters on the front of the rover to help assess end-to-end sample handling and potential
35 organic contamination at different times during the mission. Each brick, which is sealed under
36 vacuum in its own canister, can be drilled, sieved and portioned in CHIMRA. The powder is then
37 delivered to SAM or CheMin following the same pathway as for Martian rock samples. The
38 bricks are doped with traces of non-natural volatile fluorinated compounds. The sampling tools
39 and protocols are subject to equivalent and, as far as is known, benign impacts on biosignature
40 integrity as those used in terrestrial laboratories.
41
42
43
44
45

46 These instruments have been developed specifically for the MSL mission. Many aspects of their
47 specifications and ultimate performance, individually or in concert, are still being evaluated.
48

49 **Brief synopsis of environmental and physical features detectable with MSL payload**

50 In the MSL payload, SAM can detect traces of organic matter in rocks and sediment. It does so
51 in a variety of ways—from bulk organic carbon detection (by the difference between combusted
52 total carbon and inorganic carbon) or non-specific molecular detection by evolved gas analysis
53 (Bibring et al., 2005) to gas chromatographic (GC) separation and structural identification of
54 discrete molecules volatilized directly or after reactions with chemical derivatizing reagents that
55
56
57
58
59
60

1
2
3 enhance volatility and thermal stability (Tables 2, 5-8). It also detects traces of volatile organics
4 in the atmosphere, such as C1-C6 hydrocarbons and other gases, and it can precisely measure
5 carbon isotopic composition of methane and carbon dioxide.
6

7
8 The camera systems are designed document color and grain size variations in rocks and
9 sediments that reflect sedimentary structures, mineral growth processes, weathering and
10 biofabrics, if present (see Noffke, 2009; Tables 9-12). These features provide essential data for
11 interpreting the processes forming the rocks as well as their alteration. Images are essential for
12 sample selection in addition to providing a context for chemical analyses.
13

14
15 ChemCam can remotely detect carbon in soils and rocks (>1% carbon subject to final testing
16 and calibration) but the ChemCam does not directly discriminate between organic and inorganic
17 carbon (Tables 5 & 6). This is in contrast to the SAM capability of detecting sub ppb levels of
18 organic compounds that are <535 dalton via Evolved Gas Analysis (EGA) and pyrolysis.
19

20
21 CheMin in XRD mode cannot detect organic carbon, but it can detect and identify any crystalline
22 inorganic hosts of carbon and sulfur, for example (Tables 11 & 12). Used in conjunction with
23 SAM, it can be particularly useful in providing more accurate discrimination between organic
24 and inorganic hosts. Thus, the MSL payload is capable of mapping the distribution of organic
25 carbon and its molecular composition in rocks and sediments in stratigraphic, geomorphic,
26 and/or chronologic context. This distribution, the types of molecular structures detected, their
27 redox state, polarity, volatility, and their relationship to other elements and minerals detected by
28 other MSL payload instruments would provide critical insight to deciphering biosignatures, other
29 organic matter sources (abiogenic and meteoritic), environmental records, surface processes
30 and carbon cycling.
31
32

33
34 Insert Tables 4 and 5 hereabouts
35

36 **Carbon isotopes as potential biosignatures on Mars**

37
38 The patterns of isotopic composition ($^{13}\text{C}/^{12}\text{C}$ ratios) of Martian crustal and atmospheric
39 constituents reflect the compositions of their original sources as well as any isotopic
40 discrimination associated with the network of physical, chemical (and biological?) processes
41 that created and cycled these constituents (Tables 7 & 8). Fig. 1 depicts the terrestrial C cycle
42 consisting of C reservoirs (boxes) and processes (arrows) in the atmosphere, crust, and interior.
43 The caption for Fig. 1 addresses these processes. An illustration of the S cycle would be
44 analogous to Fig. 1, except sulfide and sulfate reservoirs would be substituted for organic and
45 inorganic C reservoirs, respectively. The figure is included to illustrate the complexity of known
46 reservoirs, the timescales over which they interact and range of isotopic values on Earth. This
47 picture has only emerged gradually over many years through extensive study through direct
48 observation of the chemical, biological, isotopic and geological aspects of carbon cycling (e.g.
49 see Des Marais, 2001).
50
51
52
53
54

55
56 Insert Figure 1 Hereabouts
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Insert Table 7 hereabouts

Overall, the terrestrial C-cycle consists of multiple nested cyclic pathways that differ with respect to reservoirs, processes and timescales. Carbon is exchanged between the atmosphere and shallow crust by processes which act on relatively short timescales. Carbon deeper within the crust is cycled more slowly by processes of burial and exhumation under tectonic control. These processes are now probably negligible due to the presumed absence of subduction on Mars. Processes which exchange carbon between the crust and atmosphere of Mars might be somewhat active as indicated by the recent indications of methane in the Martian atmosphere (Mumma et al., 2009). This methane almost certainly has a subsurface source that very likely involves aqueous processes, possibly including life, which we discuss further below.

The $^{13}\text{C}/^{12}\text{C}$ ratios of the carbon reservoirs reflect isotopic discrimination associated with the above processes. On Earth, isotopic discrimination associated with organic biosynthesis has been principally responsible for determining the $^{13}\text{C}/^{12}\text{C}$ ratios observed in organic and inorganic crustal reservoirs. Biological processes can change these values by several percent (e.g., Des Marais, 2001). Accordingly $^{13}\text{C}/^{12}\text{C}$ ratios might serve as biosignatures of any past or present life on Mars if key components of the C-cycling system can be constrained. However Jakosky et al. (1994) calculated that processes that caused loss of C to space were isotopically selective and increased by several percent the $^{13}\text{C}/^{12}\text{C}$ ratios of the remaining atmospheric and near-surface crustal C reservoirs. These increases might equal or exceed, and thus obscure, any changes in $^{13}\text{C}/^{12}\text{C}$ ratios due to biological processes. Note, as well, that abiotic processes of organic synthesis involve C-isotopic fractionation, further complicating the isotopic interpretation of any organic compounds that may be detected (Chang et al., 1983). To the extent that carbon situated in deeper interior reservoirs was isolated from these atmospheric escape processes, its $^{13}\text{C}/^{12}\text{C}$ value would be less affected and therefore lower, and perhaps most closely reflect initial Martian $^{13}\text{C}/^{12}\text{C}$ ratios. Consequently we cannot be entirely sure whether any measured $^{13}\text{C}/^{12}\text{C}$ patterns might indicate life or whether they reflect principally the effects of atmospheric escape or other environmentally sensitive equilibrium fractionation processes. Another difficult challenge for interpretation of C-isotopes on Mars is establishing whether organic matter and any inorganic carbon in a given sample have a genetic relationship. If, for example, the OM is meteoritic and carbonate originated hydrothermally, an isotopic separation is likely to be meaningless.

Precise and accurate carbon isotopic compositions of methane and carbon dioxide from the TLS in SAM may provide important benchmarks for understanding planetary-scale carbon cycling on Mars. For atmospheric measurements these experiments are rapid and not resource intensive and, thus, could be included in a regular sampling scheme. Detection and quantification of atmospheric methane by MSL would provide an important verification of the Earth-based and orbital spectroscopic detections of methane at ppb levels (Formisano et al., 2004, Krasnopolsky et al., 2005; Mumma et al., 2009). Temporal variations in methane concentration would be detectable by MSL and could potentially distinguish between episodic

1
2
3 release of subsurface methane or methane destruction from photolytic, oxidative or dust
4 reactions in the modern environment (Atreya et al., 2006). Spatial variations in methane
5 concentration are not expected to be observable unless MSL is proximal to a subsurface
6 hydrothermal, volcanic or unstable hydrate point source. Using C-isotopic data to constrain the
7 origins of atmospheric methane on Mars, however, presents a formidable problem. Abiogenic
8 methane production associated with serpentinization, that is, aqueous alteration of olivine-
9 and/or pyroxene-rich rocks, is a process likely to have been prevalent early in Mars history and
10 which feasibly continues today (Oze and Sharma, 2005). Multiple carbon and hydrogen isotopic
11 data suggest the methane that is abundant in the fluids emanating from the Lost City
12 hydrothermal field, a site of contemporary serpentinization on Earth, has an abiogenic origin.
13 However, it is known that methanogenic Archaea are also active in this system (Bradley et al.,
14 2009). Accordingly, carbon isotopes are of limited value in discriminating between biological
15 and abiogenic sources where multiple processes can contribute to a pool of methane and
16 reservoir affects provide added complications (Bradley and Summons, 2010). This is the
17 situation on Earth where the biochemistry of methanogenesis is reasonably well understood and
18 even more uncertainly would accompany methane measurements on Mars. Carbon isotopic
19 data might discriminate between abiogenic and biogenic methane sources only if the
20 fractionation between methane and co-existing CO₂ were well in excess of the equilibrium
21 values modeled for feasible P-T conditions, thereby implying kinetic (i.e., enzymatic) control on
22 methanogenesis.
23
24
25
26
27
28

29 Variations in C-isotopic compositions of carbon dioxide are less likely as it is the dominant
30 atmospheric species. SAM design also enables isotopic measurements of trace amounts of bulk
31 inorganic and organic carbon by comparing CO₂ evolved under inert conditions with CO₂
32 generated from combustion with O₂. Mapping variations in these isotopic compositions could
33 further our understanding of redox processes, isotopic fractionation pathways (including
34 biosignatures, abiogenic processes, and possibly meteoritic contributions), and environmental
35 carbon cycling (Table 8). The TLS can also measure oxygen isotopes of carbon dioxide and
36 deuterium/hydrogen in water, which may indicate the effects of surface and atmospheric cycling.
37 Oxygen isotopes of carbon dioxide evolved from carbonates in SAM will help with assessment
38 of post-depositional alteration by subsurface fluids.
39
40
41

42 **Sulfur isotopes as potential biosignatures on Mars**

44 Observed ³⁴S/³²S, and ³³S/³²S values presumably reflect processing by redox reactions in the
45 atmosphere (e.g. photochemistry, see Farquhar et al., 2000) and crust (e.g. weathering,
46 hydrothermal and life). On Earth, microbial reduction of sulfate and sulfur disproportionation
47 reactions occurring at ambient temperatures have created large ³⁴S/³²S differences (several
48 percent) between oxidized and reduced sulfur reservoirs (Canfield, 2001). Because
49 microorganisms are required to catalyze S-isotopic exchange reactions having significant
50 fractionations at low ambient temperatures, large ³⁴S/³²S differences in sedimentary rocks can
51 be reliably interpreted as biosignatures in many cases. However, a careful assessment of the
52 original environment of deposition is always essential. Accordingly, before stable isotopic
53 patterns can serve as potential biosignatures on Mars, we must characterize the isotopic
54 composition of major sulfur reservoirs and also understand more fully the consequences of key
55
56
57
58
59
60

1
2
3 non-biological processes.
4

5 Tests on a SAM breadboard QMS instrument have shown that sulfur isotope measurements of
6 sulfur dioxide evolved from the thermal degradation of sulfate minerals are possible (Franz et
7 al., 2007). However, these measurements are complicated by overlapping spectra and the
8 evolution of gas from multiple sulfur species. Unlike the sub-per-mil precision from Earth-based
9 instrumentation, the SAM QMS may provide percent precision. These data may still be valuable
10 on Mars, especially if basaltic sulfur, which shows <1% variability on Earth and meteorites,
11 provides a reference point for comparing sedimentary and hydrothermal sulfur analyses (Table
12 8). On Earth, sedimentary sulfides show 4% variation, reflecting biological cycling of S, a
13 sensitive redox element (Canfield, 2001; Farquhar et al., 2003). If a similar record were to exist
14 on Mars, sulfur isotopic detection by the SAM QMS may detect it. Sulfur isotopic compositions
15 of sulfate and reduced-sulfur minerals may also provide valuable information of environmental
16 sulfur cycling by abiogenic hydrothermal (Rye, 2005; Greenwood et al., 2000) and atmospheric
17 processes (Farquhar et al., 2000).
18
19
20
21

22 **Other isotopic data**

23
24 Noble gas isotope measurements and elemental ratios are regarded as isotopic tracers of
25 mantle and atmospheric evolution (Swindle, 2002) and possible comet-borne contributions
26 (Owen et al., 1992). Although they do not provide direct insight into environmental conditions
27 preserved in the sedimentary record, they can supplement the larger context of Mars chemical
28 and environmental evolution. These data may help explain other isotopic records.
29
30

31 **Environmental conditions**

32
33 Sedimentary and other near-surface materials can serve as recorders of environmental
34 conditions that prevailed during previous epochs of Martian history. Insights into ancient
35 environmental conditions might help to identify potential processes responsible for setting the
36 observed isotopic patterns. Paleotemperatures often can be inferred by measuring stable
37 isotopic compositions of pairs of minerals or fluids that equilibrated isotopically when they
38 formed. Oxygen isotopes have been utilized most frequently to infer paleotemperatures within
39 the habitable range (<120°C). Elevated temperatures have been inferred using sulfur (>150°C)
40 and carbon (>300°C) isotopes. Stable isotopic compositions of preserved minerals can also help
41 to elucidate the nature of fluids associated with their formation. For example, elevated ¹⁸O/¹⁶O
42 values might indicate the former presence of brines in evaporitic environments. Many additional
43 examples can be cited where stable isotopic patterns have helped to constrain
44 paleoenvironmental conditions.
45
46
47
48

49 Insert Table 8 hereabouts
50

51 **Preservation potential**

52
53 To the extent that sediments, cements, and other surface materials have escaped alteration
54 subsequent to their formation, they can preserve information about earlier environmental
55 conditions and, potentially, biosignatures (Tables 9 & 10). Preservation can be compromised by
56 weathering and erosion or by alteration *in situ* by oxidation and migrating fluids, for example.
57
58
59
60

1
2
3 The following minerals can isolate and preserve biosignatures (listed in order of increasing
4 crustal residence times on Earth): ice, halite, sulfates (e.g. Aubrey et al., 2005; Panieri et al.,
5 2010), carbonates (e.g. Birgel et al., 2008), phyllosilicates (e.g. Hedges and Keil, 1995;
6 Butterfield, 1990), silica (e.g. Knoll, 1985), hematite (Fernandez Remolar and Knoll, 2008) and
7 phosphates (e.g. Xiao & Knoll, 1999; Farmer & Des Marais, 1999). For example, carbonates
8 deposited as a consequence of microbial metabolisms sometimes hold an excellent record of
9 those processes as is the case with methane seep limestones (e.g. Birgel et al., 2008) or
10 hydrothermal systems like Lost City (Bradley et al., 2009). Silica-rich water derived from
11 hydrothermal systems is another well-established medium that promotes faithful preservation
12 (e.g. Knoll et al., 1985; Trewin, 1996). In fact, Walter (1996) has identified numerous fossil
13 hydrothermal systems on Earth, both terrestrial and marine, that are known and potential
14 repositories of paleobiological information. Preservation is optimized when temperatures remain
15 low and mineral matrices form during sedimentation/precipitation and reduce the permeability of
16 the sediments to near zero. Detailed $^{18}\text{O}/^{16}\text{O}$ values of some of these minerals can help to
17 assess the extent to which invading fluids caused post-depositional alteration.

22 Hydrated mineral phases

23
24 The presence of hydrated minerals reflects specific chemical conditions, including the activity of
25 water (Table 12). Specific implications of different hydrous phases vary with the composition
26 and context of the minerals. Some, such as hydrous Mg-sulfates, require low temperatures
27 and/or substantial humidity to remain stable. Their hydration states reflect local current
28 conditions due to their rapid dehydration kinetics, although their formation may reflect older
29 conditions. Other hydrous minerals, such as the clay minerals, remain metastable for long
30 periods of time and provide a record of past hydrous activity. Extracting the history of water on
31 Mars requires careful characterization of hydrous minerals within their environmental context.

36 Textural features preserved in sediments and hydrothermal systems

37
38 Physical, chemical and biological processes all influence the preservation of biosignatures in
39 hydrothermal and sedimentary systems on Earth, and we can use our understanding of these
40 processes to predict their impact on possible biosignature preservation on Mars (Tables 9 & 10).

41
42 Purely physical and chemical processes should be comparable on Earth and Mars. For
43 subsurface, hydrothermal and sedimentary systems, physical and chemical processes can
44 provide substantial insights into the history and habitability of the system. For example, in
45 sedimentary systems, physical processes such as sediment transport produce structures that
46 are characteristic of specific processes, e.g., types of flows. These can be used to interpret
47 processes in the depositional environment, providing invaluable constraints on habitability and
48 guiding the search for biomarkers. There are some non-negligible differences due to the lower
49 g of Mars, however, these do not affect the overall interpretation of transport-related suites of
50 sedimentary structures (Grotzinger et al., 2005). Similarly, chemical processes leave distinctive
51 signatures, whether they are active in the depositional environment or within the rock. They can
52 be used to evaluate the habitability of the environment and rocks at different points in time,
53 including the potential for subsurface colonization of rocks of volcanic, hydrothermal or
54 sedimentary origins.

1
2
3 Insert tables 9 & 10 hereabouts
4
5
6
7

8 On Earth, biological processes are very active in almost all sedimentary and lower-temperature
9 hydrothermal systems. Biological processes in terrestrial hydrothermal systems provide a
10 diverse suite of potential biomarkers, including concentrations of organics and minerals that
11 reflect mats, stromatolites, roll-ups, biofilms, streamers, etc. (Tables 9 & 11). Concentrations of
12 elements and high concentrations of migrated organics are also characteristics of some
13 terrestrial hydrothermal systems. Preservation of these indicators of biological activity is
14 strongly affected by physical and chemical processes. The high temperatures and abundant
15 water flow tend to degrade biomarkers. Hydrothermal systems commonly experience intense
16 brecciation and fracturing due to high-pressure fluids. This type of fracturing leaves distinctive
17 textural features that are best avoided when looking for a good biosignature preservation
18 window. Chemical processes also affect preservation. Mineral precipitation can entomb
19 organics and biological processes can influence the distribution of minerals. In hydrothermal
20 settings, a good preservation window is created by precipitation of non-redox sensitive, low-
21 porosity minerals. However, minerals can also recrystallize and dissolve, particularly when
22 exposed to high-temperature fluids, making preservation of biosignatures less likely. When
23 looking for a good preservation window, one wants to avoid recrystallized areas, those that have
24 experienced intense oxidation, and those exposed to high temperatures or ionizing radiation
25 (e.g. Dahl et al., 1988).
26
27
28
29
30

31 Sedimentary systems on Earth also have substantial potential for preserving biosignatures
32 because of the intensity of biological activity, and their natural tendency to concentrate or high-
33 grade organic matter across hydraulic gradients. Again, these terrestrial processes serve as
34 models for predicting good preservation windows on Mars (Tables 10 & 12). Biomass
35 accumulation in terrestrial systems is reflected in mats, stromatolites, roll-ups, wrinkle
36 structures, etc. For preservation, these accumulations need to incorporate sediment or be
37 mineralized prior to degradation of the organics. Biological processes can also create fenestrae
38 (gas-produced pores), affect grain sizes, and influence elemental concentrations. These
39 features can be well preserved if they are lithified early and do not experience significant
40 alteration. Physical processes such as dewatering, hydration changes and structural
41 deformation can destroy these signatures. Similarly, chemical processes such as
42 recrystallization, redox changes and metamorphism destroy biomarkers. Thus, for biomarker
43 preservation, one should identify areas with an appropriate sedimentary environment that was
44 lithified early and experienced minimal post-depositional alteration (Farmer and Des Marais,
45 1999).
46
47
48
49

50 **Elemental concentrations and mineral distributions in hydrothermal and deep subsurface** 51 **systems** 52

53 Hydrothermal environments can preserve potential biosignatures in the form of elemental
54 concentrations and mineral distributions (Table 11). Thermally driven aqueous convection can
55 significantly alter environmental conditions through selective mineral dissolution, alteration, and
56 precipitation; element leaching and subsequent transport; and oxidation and reduction (redox)
57
58
59
60

1
2
3 chemistry. On Earth, each of these processes can occur with or without biological mediation,
4 but through careful analysis with multi-faceted approaches, the role of biology can often be
5 assessed.
6

7
8 If life emerged on Mars and prospered in hydrothermal systems, these alteration processes
9 likely include biotic and abiotic components (Tables 9 & 11). Each yields an array of features
10 including specific mineralogy that is characteristic of certain hydrothermal conditions and
11 chemistries, element gradients and zonation within those minerals and spatial distribution of
12 alteration minerals and precipitates at scales from μm to km (Table 11). The identification and
13 chemical analysis of chlorite, amorphous silica or quartz, sulfide minerals, kaolinite and other
14 clays, hematite and other ferric (hydr)oxides, carbonates, and sulfates can be used to determine
15 formation temperatures, redox conditions, and pH. For example, specific sulfides (PbS, ZnS,
16 CuS) and sulfates (BaSO_4 , $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) or mineral assemblages (e.g., sulfate and/or sulfide
17 together with saddle dolomite) can point to specific formation temperatures. The presence of
18 jarosite, alunite, or kaolinite indicate low-pH environments, and discrimination between
19 crystalline silica minerals (e.g., quartz or cristobalite) and non-crystalline silica phases, such as
20 opal-CT, is most effective when chemical data are combined with mineralogical information
21 (e.g., XRD data). Other minerals bearing S, Fe, Mn, U, As or other redox-sensitive elements
22 can provide further constraints on Eh and, if present above the minimum detection limits for
23 CheMin, could be detected by XRD. All of these features would help constrain the possibility,
24 probability, and physiology of potential life forms on Mars. However, because hydrothermal
25 systems are dynamic, high-energy environments, the preservation of labile features (e.g.
26 organic matter, amorphous solids) would undoubtedly require encasement or entombment in
27 protective minerals or other removal from the destructive forces of oxidation, metamorphism,
28 and continued hydrothermal activity.
29
30
31
32
33
34
35
36

37 Insert Table 11 hereabouts

38 39 **Elemental concentrations and mineral distributions in sedimentary systems**

40
41 Sedimentary environments present both similarities and differences to the hydrothermal case
42 highlighted above. The preservation potential of environmental signatures is determined
43 principally by the sedimentary material, the chemical composition and flux of the aqueous
44 solutions responsible for the sediment transport, the redox state and pH of that solution, and
45 any post-depositional chemical processes (including putative microbe-mineral interactions) that
46 may have operated (Table 12). The most informative environmental signatures include
47 sedimentary structures and redox-sensitive (e.g., S, Fe, and Mn-bearing) and pH-sensitive (e.g.,
48 jarosite, alunite, kaolinite, carbonates) mineral assemblages and abundances. Many
49 sedimentary environments on Earth or Mars also host evaporite minerals (including sulfates and
50 chlorides) and corresponding trace element distributions, as well as oxides, carbonates, sulfides
51 and, perhaps, phosphate precipitates resulting from microbe-mineral interactions. The
52 preservation potential of these environmental features is particularly enhanced by early
53 lithification of the sediments. In addition, the greatest scientific return would likely come from
54 sedimentary systems, lacustrine or marine, that have not been subjected to significant
55
56
57
58
59
60

1
2
3 recrystallization, prolonged strong oxidizing conditions, dissolution and solute removal, major
4 structural alteration, or significant thermal metamorphism. As closed basins, lakes in particular,
5 represent terminal receptors for both primary and transported organics together with the clays,
6 which preserve OM due to strong absorption capacity, low reactivity, and low permeability when
7 compacted (Farmer and Des Marais, 1999; Meyers and Ishiwatari, 1995).
8
9

10
11
12 Insert Table 12 hereabouts
13

14 **Critical Mars - Earth contrasts: considerations for MSL decision-making**

15

16
17 Any martian biomass must be, and probably would have been, less abundant than biomass on
18 Earth throughout its history. Land plants dominate modern biomass on Earth, and marine
19 organic matter is created almost exclusively by photosynthetic biota and it is pervasively
20 concentrated at the margins of continents. Only a relatively small percentage of all primary
21 productivity (~0.1%) survives remineralization and becomes preserved in sediments. If life
22 colonized Mars, its global primary productivity is expected to have been much lower, given the
23 less clement and more ephemeral surface habitable environments and the presumed absence
24 of plants. The fraction of any biomass that was preserved in martian sedimentary rocks is, of
25 course, unknown.
26
27

28
29 Meteorite impacts delivered organic matter to the martian surface. The early solar system had a
30 greater abundance of debris and thus experienced a higher rate of impacts than later in its
31 history. The size distribution of impactors may also have been different. Radiometrically dated
32 samples from the moon make it possible to associate an absolute age with a certain crater
33 density. The relation is non-linear because the flux of impactors was higher before 3.5 Ga, but
34 subsequently the flux has apparently remained nearly constant. The following major questions
35 persist: 1) Did the early impact flux decrease steadily or did an “impact spike” occur at ~4 Ga
36 (known as the “Late Heavy Bombardment”)? 2) There is a large uncertainty (factor of about 2)
37 for young ages (≤ 1 Ga). Because Mars is closer to the asteroid belt the number of impacts on
38 Mars is estimated to be 3 ± 1.5 times the number of impacts on the Earth-Moon system.
39 However the accumulation of meteoritic organics in the martian crust also depends upon the
40 fraction of material that is actually preserved.
41
42

43
44 The Late Heavy Bombardment is an important consideration in estimating organic matter
45 preservation on Mars. As far as we know, virtually all of the biogenic organic matter that
46 persists on Earth today was formed after 3.8 Ga and most of that was formed within the most
47 recent 500 million years. If a biosphere existed in the first 500 million years of martian history a
48 substantial fraction of its remains might have been altered or destroyed during the Late Heavy
49 Bombardment. Smaller, relatively recent impacts might have exhumed any remnants of an early
50 biosphere.
51
52

53
54 The tectonic regimes of Mars and Earth are distinctly different, as reflected by significantly older
55 surface ages and relatively minor regional metamorphism on Mars. Organic carbon
56 accumulation on Earth is, and has been, modulated by a vigorous tectonic cycle, and the
57
58
59

1
2
3 residence time of organic carbon in the crust is consistent with the Wilson cycle period (i.e.,
4 modern plate tectonics). Most importantly, tectonic processes drive biogeochemical cycles by
5 sustaining nutrient availability and creating the spaces for subaqueous accumulations of
6 sediment and its entrained organic matter.
7

8
9 Mars currently has no global dynamo-driven magnetic field, but strong local crustal fields
10 indicate that a global field likely existed in the past. However, magnetic field age and strength
11 are not known to directly influence organic matter formation preservation except that a global
12 magnetic field would have attenuated the flux of deleterious radiation and reduced losses of
13 atmospheric species to space. The radiation environment on Earth is conducive to harvesting
14 solar energy and forming organic matter through photosynthesis. A dense atmosphere, the
15 pervasiveness of liquid water on Earth's surface and the operation of a magnetic field all can
16 reduce deleterious radiation. Widespread photosynthesis on Earth has clearly enhanced organic
17 matter production in environments that also favored its preservation in aqueous sediments.
18

19
20
21 Processes operating in such favorable sedimentary environments can concentrate organic
22 matter prior to burial. Important concentrating mechanisms include the following: density sorting
23 during transport, adherence to the fine particles of clay minerals, and ballasting of organics on
24 biogenic minerals. Differences in the hydrological cycles of Mars and Earth would have affected
25 any potential concentration or organics during transport.
26

27
28 Sedimentary rocks with anomalously high concentrations of organic matter (> few percent) are a
29 historically pervasive feature of Earth's sedimentary rock record; they occur even in the early
30 Precambrian record. However organic-rich and biomarker-bearing deposits are typically non-
31 uniformly distributed and unpredictable. No known example of subaerial fossilized deposits on
32 Earth have a total organic carbon content exceeding 1% by weight; Mars is expected to be
33 similar.
34

35
36
37 Aeolian transport processes lead to destruction of organic matter by continually refreshing its
38 contact with oxidizing agents and UV radiation. The fate of organic matter transported in this
39 way is expected to be similar on Mars to what it is on Earth or, possibly, worse.
40

41
42 Therefore, on modern Earth, biomass has been detected in almost every wet environment,
43 including very harsh and extreme sites in terms of temperature, pH, water activity, intermittent
44 desiccation, and pressure. Essentially every wet environment below the upper thermal limit of
45 life can be considered habitable. However the production, concentration and preservation of
46 organics have varied substantially and have been controlled by the spatial and temporal
47 distribution of subaqueous environments and sedimentary processes. The search for martian
48 biosignatures has become more promising due to the discovery that surface and near-surface
49 aqueous environments existed on Mars at the same time when biological organic matter was
50 being preserved in ancient aqueous sediments on Earth.
51

52
53 In its quest to find organic-bearing strata, MSL should investigate, ideally, a subaqueously
54 deposited and rapidly buried suite of strata that represents the longest duration possible, i.e.,
55 the thickest section in the absence of other age constraints (Farmer and Des Marais, 1999). If
56 the record at Meridiani Planum is representative of other early martian sedimentary
57
58
59
60

1
2
3 environments, at least some beds might have been deposited in water even if the stratigraphic
4 succession is predominantly eolian in origin (Grotzinger et al., 2005).
5
6

7 **Consideration of planetary age**

8
9 Planetary evolution strongly influences biosignature preservation, particularly on a planet like
10 Mars that has experienced significant geological and climate variations. Changes in habitability
11 through time affect the abundance and diversity of potential biosignatures. For example, loss
12 of the Martian magnetic field strongly affected surface environments through atmospheric loss
13 and increased radiation; this change significantly degraded surface habitability, reducing the
14 chances of biosignatures in the rock record. Changes in temperature and moisture also clearly
15 affect habitability both on the surface and in subsurface aquifers. Hydrothermal activity tied to
16 volcanism and impacts was more abundant on early Mars and more abundant and continuous
17 hydrothermal activity is more likely to support a biosphere that could leave signatures. Also, it is
18 critical to understand long term climate evolution when choosing the best places to look for
19 biosignatures; rocks deposited during warm and wet intervals, e.g. early Mars, are more
20 promising for biosignature development. Preservation of any biosignatures also depends on
21 climate-dependent sedimentary processes. Clay minerals preserve biosignatures in ways that
22 are fundamentally different to what occurs in carbonate or sulfate minerals (Hedges and Keil,
23 1995). Therefore, long term changes in the relative abundance of certain minerals affect the
24 likelihood of specific biosignature preservation in rocks of a particular age. Thus, the specific
25 evolution of Mars as a planet suggests that the best time interval for the search for
26 biosignatures would be represented by early to mid-Noachian rocks, when clays were thought to
27 be forming. This is not to say that clays did not form in younger times, in fact, tests of this
28 hypothesis could be one outcome of the MSL mission.
29
30
31
32
33

34 **Potential biological sources of organic carbon on Mars**

35
36 On Earth today, photoautotrophy is by far the dominant physiology leading to organic matter
37 synthesis. With a seemingly unlimited solar energy source, photoautotrophs in the ocean and on
38 land can produce copious amounts of organic matter compared with chemoautotrophs (orders
39 of magnitude more). In present-day aquatic environments chemoheterotrophs thrive on
40 chemical energy from the decomposition of organic matter from photosynthetic communities
41 using electron acceptors that are also regenerated by photoautotrophs.
42
43
44

45 Chemoautotrophs thrive by carbon fixation using electron donors that are also generated or
46 regenerated via light-harvesting processes. In the absence of photoautotrophy, primary biomass
47 production would be limited to chemoautotrophs that harvest chemical energy from geological
48 processes—namely those occurring in hydrothermal vents in the oceans and terrestrial
49 geothermal springs (see Table 4) and subsurface microenvironments within the fractures and
50 pores of ultramafic and mafic rocks. The Lost City Hydrothermal Field (Kelley et al., 2005), a
51 low-temperature marine ultramafic hydrothermal system is one model for an environment that
52 could support life in the absence of photosynthetic light harvesting (Martin & Russell, 2007;
53 Martin et al., 2008). Such a system could conceivably occur anywhere that water circulates
54 through ultramafic rock. Although ultramafic rock is rare on the ocean floor today, occurring
55 mainly at ultraslow-spreading ridges, it could have been more prevalent in the geological past
56
57
58
59
60

1
2
3 (Sleep et al., 2004; Sleep and Bird 2007). Molecular and isotopic data indicate that Lost City
4 fluids sustain a flourishing microbial community of methanogens and sulfate-reducing bacteria
5 (SRB) (Bradley et al., 2009; Brazelton et al., 2006). Although the SRB would be sulfate-limited
6 if disconnected from a sulfate supply that is ultimately coupled to oxygenic photosynthesis or
7 abiotic oxidation reactions occurring at the surface or subsurface, there appear to be no such
8 constraints on the activity of methanogens that require only hydrogen and CO₂. In addition,
9 thermodynamic calculations showed that Lost City-type vent fluids mixed with seawater are
10 energetically favorable for biomass synthesis (Amend & McCollom, 2009). Thus, the recent
11 detection of serpentine deposits in Noachian terrains on the surface of Mars (Ehlmann et al.,
12 2010) identifies the probable past occurrence of hydrogen-producing water-rock reactions and,
13 therefore, an established set of processes that would both support chemosynthetic life and
14 preserve a molecular and/or isotopic record of its prior existence (e.g. Kelly et al., 2005; Bradley
15 et al., 2009)

16
17
18
19
20 At life's origin, the dominant energy source was unlikely to have been sunlight. Energy flowing
21 from chemical and thermal disequilibria and particularly from the interaction of hot rocks with
22 water is more likely. Perhaps the same was the case for early Mars. In aqueous depositional
23 environments, chemoautotrophs may have been the cornerstone of microbial communities
24 relying on fermentation and heterotrophy fueled by a weak oxidant flux from chemoautotrophy
25 or irradiative oxidation. If Mars evolved a biosphere, it may not have progressed to
26 photoautotrophy or a dependence on photoautotrophy as it did on Earth. Thus, in the
27 consideration of Martian environments conducive to producing molecular biosignatures,
28 targeting depositional environments that had a strong chemical energy flux and sustained redox
29 gradients for long periods by biogeochemical cycling is a most promising strategy.

30
31
32
33
34 One additional possibility for chemoautotrophy is energy derived from radiolysis of water (Pratt
35 et al., 2006). The recent discovery of a microbial biome dominated by thermophilic sulfate
36 reducing bacteria in a c. 3km deep saline aquifer in Archean metabasalt seems to required that
37 they were sustained by geologically produced sulfate and hydrogen at concentrations sufficient
38 to maintain biological activity for millions of years (Chivian et al., 2008; Lin et al., 2006).
39 Radiolysis of water coupled to oxidation of sulfide minerals could have provided the energy
40 drive for low-biomass, low-diversity subsurface ecosystems.

41 42 43 **Abiotic sources of organic carbon on Mars**

44
45 Sources of abiotic organic matter on Mars could have been similar to those hypothesized to
46 have been present or formed on the early Earth. This would have included organic matter
47 delivered by meteorites and interplanetary dust (Anders, 1989), organic matter produced as a
48 result of atmospheric photochemistry (Chang et al., 1983) and organic matter produced during
49 fluid mixing in hydrothermal systems (Shock & Schulte, 1998). The Lost City hydrothermal
50 system also serves as an example of another route to abiotic organic compounds. The isotopic
51 composition and chain length distributions of hydrocarbon gases isolated from Lost City fluids
52 have been interpreted to signify an abiotic source (Proskurowski et al., 2008; Sherwood Lollar et
53 al., 2006). Hydrogen produced in high concentration by serpentinization chemistry leads to a
54 thermodynamic drive for CO₂ reduction (Shock & Schulte, 1998). Methane and higher
55
56
57
58
59
60

1
2
3 hydrocarbons may be thus produced abiotically in ultramafic hydrothermal systems by Fischer-
4 Tropsch type processes that comprise polymerization reactions leading to methane and higher
5 hydrocarbons (Horita & Berndt, 1999; McCollom & Seewald, 2006; McCollom & Seewald,
6 2007).

7
8
9 Steele and co-workers (2007) have identified macromolecular carbon, in an intimate association
10 with magnetite, through imaging and Raman spectroscopic studies of carbonate globules in the
11 Mars meteorite ALH84001. This abiotically-formed organic matter appears to be native to Mars.
12 It is hypothesized to have formed formation via reactions of the Fe-C-O system for which there
13 is a terrestrial analogue in the Bockfjorden volcanic complex of Svalbard.
14
15
16
17

18 **Interstellar organic matter**

19
20 Irrespective of whether extensive abiotic organic synthesis of predominantly hydrocarbons
21 occurred on the early Earth and, potentially, on Mars, chondritic meteorites and interplanetary
22 dust particles (IDPs) have delivered abiotic organic matter to the Martian surface.
23 Carbonaceous chondritic (CC) meteorites consist of up to 2 wt % of organic matter finely
24 intermixed with matrix silicates (Alexander et al., 2007). It is well known that some CCs contain
25 relatively high abundances of small, polar organic molecules, e.g. amino acids. The highest
26 reported concentration of amino acids, however, is not very high, being on the order of 250 ppm
27 (Martins et al. 2007). The primary form of organic matter in all classes of chondritic meteorites
28 is Insoluble Organic Matter (IOM), a chemically complex macromolecule (e.g. Cody & Alexander
29 2005) that is by definition insoluble in any solvent and is in many cases 99 % of the organic
30 matter in a given chondrite (Botta, 2005; Sephton, 2002; Sephton et al., 2004). Among the
31 meteorites collected as finds in Antarctica, 0.8 % are CCs, the majority, 77 %, are ordinary
32 chondrites (OCs) (<http://curator.jsc.nasa.gov/antmet/ppr.cfm>). While ordinary chondrites do
33 contain IOM, its abundance is considerably lower than that in CCs (Alexander et al. 2007) and,
34 even in the least metamorphosed OCs, the IOM has been significantly altered through long term
35 thermal processing yielding highly aromatized macromolecular structures (Cody et al., 2008). If
36 the distribution of meteorite types on Mars was and is similar to that of Earth, it is reasonable to
37 assume that the predominant IOM would be exceptionally stable over geological timescales
38 even under the harsh Martian surface conditions. An instrument analogous to SAM flew on the
39 Viking mission and did not detect organics via pyrolytic analysis. All evidence suggests that the
40 Viking technical approach and instrument worked according to its design (Biemann, 1979,
41 Biemann, 2007) so these Viking results do not exclude the potential presence of abundant IOM,
42 if the IOM was derived predominantly from OCs. IOM in OCs has already been subjected to
43 extensive natural pyrolysis in the OC parent body (Cody et al., 2008). Minimal, if any, pyrolysate
44 remains to be derived, therefore, from thermally metamorphosed IOM. Pyrolysate from
45 chondritic IOM might be detected only if the abundance of CCs that accumulate on the Martian
46 surface exceeds what is observed in the Antarctic collections.
47
48
49
50
51
52
53
54

55 Finally, there is the issue of Interplanetary Dust Particles (IDPs) that, in case of the Earth,
56 constitute a much greater influx of extraterrestrial matter. IDPs typically contain considerably
57 more organic matter than CCs and the organic macromolecule bears significant spectroscopic
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

resemblance to IOM derived from primitive CCs. Lacking the silicate matrix protection afforded to CC IOM, it appears likely that IDPs will be subjected to more degradation than CC IOM on the Martian surface. Under ideal conditions, however, IDPs could survive on Mars, and be detectable via pyrolysis GCMS. The SAM instrument has enhanced sensitivity and experiment flexibility compared to Viking. However, an even more important quality is the mobility and composition of the entire MSL package as this enables optimization of sample selection and handling.

Distribution and importance of phyllosilicates for habitability and organic preservation

It has long been recognized that, on Earth, there is a close association between organic matter and phyllosilicate minerals. This is the case for both modern environments, particularly in large aqueous catchments where fine-grained particulates often rich in phyllosilicate minerals can settle from the water column (lakes and ocean margins), and in ancient sediments (e.g. shale and organic-rich mudstones). Certainly, low porosity and permeability of compacted phyllosilicate sheets aid in entombing and protecting organic matter from oxidizing fluids and biological activity over geological time scales. Abundant experimental data show that many phyllosilicates, particularly smectites, interact strongly with organic molecules and are capable of adsorbing and preserving them. However, the specific mechanisms that give rise to this association are not well understood and are the subject of ongoing investigations. Accordingly, we should be very careful in extending empirical observations made on Earth to the situation on Mars. Rather, the combined ability of the MSL to detect both organic carbon and clay mineral assemblages on Mars offers us an unprecedented opportunity to learn much about this particular issue during the landed operations.

Phyllosilicates such as smectites, chlorites, and kaolin minerals form during the weathering of minerals in soils and in hydrothermal systems. Phyllosilicates all have sheet-like structures and can accommodate a large variety of cations, most commonly including Fe, Mg, and Al in their octahedral sheets and Na, Ca, and K between the layers in the so-called interlayer region. Fine-grained, disordered phyllosilicates are often called clay minerals, and they have high surface areas and the ability to exchange their interlayer cations and adsorb H₂O molecules, as a result of negatively charged interlayer regions. Many organic compounds can be adsorbed onto surfaces and into the interlayer regions, in some cases forming weak bonds with phyllosilicate surfaces. Adsorption of organic molecules into the interlayer region is particularly important for very low molecular weight compounds, such as amino acids (Hedges & Hare, 1987) and polysaccharides (Dontsova and Bigham, 2005) as well as higher molecular weight material (Mayer, 1994a, b; Kennedy, et al., 2002). Organic molecules compete with other polar species in the environment (e.g., water, cations, etc.) for active sites on phyllosilicate minerals. However, a key factor for enhanced organic matter preservation by phyllosilicates is coincidental timing of organic matter diagenesis and phyllosilicate mineral formation (Hedges & Kiel, 1995). Specifically, it requires synchronous availability of organics and clay mineral formation. To grasp the significance of this timing issue, it is also important to understand how organic matter ends up in sediments and ultimately the rock record.

1
2
3 On Earth, most sedimentary and hydrothermal organic matter is macromolecular. In these
4 terrestrial settings, polymerization to macromolecules may begin at the point of formation (i.e.,
5 cellular and biopolymer material), during diagenesis in the water column, or in pore waters of
6 sediments as dissolved organic matter. It is composed of smaller constituents cross-linked
7 together by covalent bonds into a three-dimensional material. Initially, it is porous, internally and
8 externally charged, and has varying degrees of hydrated surfaces giving rise to hydrophilic (i.e.,
9 charged with functional groups having O, N, and S moieties) and hydrophobic regions (i.e.,
10 dominated by C and H). It also has a high affinity for complexation with trace elements and
11 sorption to particles (Guo & Santschi, 1997), such as phyllosilicates (Ohashi & Nakazawa,
12 1996). Once deposited, the chemistry, porosity, and overall activity of macromolecular organic
13 matter become more stable as it equilibrates with the sedimentary and eventually, lithified,
14 environment. On Earth, this environment may be oxidizing and lead to overall degradation and
15 loss of the organic macromolecules, potentially leaving behind the minerals, trace elements, and
16 morphologies associated with the original material. This is commonly the case for subaerial
17 environments. However, organic macromolecules stand a better chance of preservation if they
18 are quickly stabilized in sediments. There are two primary mechanisms for this. One, common in
19 the anoxic marine realm on Earth, is where hydrophilic functional groups of organic
20 macromolecules are effectively 'quenched' and replaced by sulfide during diagenesis. The
21 second mechanism, potentially more important on Mars, is rapid burial and lithification of
22 organic macromolecules. Organic macromolecular sorption to mineral surfaces significantly aids
23 in burial of organic matter (e.g., Bock & Mayer, 2000; Kennedy et al., 2006). On Mars, the
24 environments that may have allowed for this diagenetic complexation between dissolved
25 organic macromolecules and phyllosilicates would thus provide promising sites for organic
26 molecular preservation and detection by MSL. These environments must also be proximal to
27 sources of both phyllosilicates (e.g., weathering regolith or hydrothermal) and organic matter.
28 Therefore, Martian hydrothermal and stable aquatic sedimentary environments may be very
29 favorable for both phyllosilicate and organic matter accumulation (Farmer and Des Marais,
30 1999).

31 32 33 34 35 36 37 38 39 **Assessments of potential carbon accumulation and preservation in sediments and** 40 **hydrothermal systems**

41
42 Terrestrial models of organic carbon accumulation and preservation provide predictive capability
43 in respect to these processes in the regolith of Mars. However, due to differences in processes
44 and chemical environments, our estimates of the propensity for early Mars environments to
45 support organic matter formation and preservation are necessarily crude (Table 3). However,
46 our understanding of terrestrial environments makes estimating the possibility of particular
47 observations of potential biomarkers on Mars possible, even if limited. Estimates are thus
48 scaled as high, medium or low. For example, the potential for biosynthesis of organic matter is
49 highest in near-surface hydrothermal systems and in deltaic and lacustrine environments. The
50 potential for hydrodynamic processes to concentrate those organics is ranked as high for the
51 deltaic and perennial lacustrine environments but lower for the surface and subsurface
52 hydrothermal systems as well as for evaporitic environments. Three columns toward the right of
53 the table rank the potential to recognize such environments using remote sensing and criteria
54 that are geomorphic, mineralogical, or stratigraphic, respectively. Finally, the right-hand column
55
56
57
58
59
60

1
2
3 makes an assessment of the MSL instrument package's potential to gather data identifying such
4 environments.
5

6
7 The potential for organic matter formation and accumulation on early Mars is assessed as low in
8 the volcanic pyroclastic and flow deposits and moderate in hydrothermal environments (Table
9 4). Any organic matter that accumulated in such places would stand a moderate chance of
10 being preserved over time. In tables 5 and 6 we evaluate how the MSL payload elements could
11 be used to confirm environmental features specific to processes needed to form, transport,
12 concentrate and preserve organic molecules on early Mars. Only environments identified in
13 Table 1 as having moderate to high potential to support organic carbon formation and
14 preservation that were at moderate or higher are considered for the sedimentary category (i.e.,
15 fluvial floodplain, deltaic, lacustrine (perennial), lacustrine (evaporitic). Only hydrothermal
16 (<100°C) subsurface + surface environments are considered for the hydrothermal category.
17
18

19 20 **An assessment of processes essential to the preservation of isotopic abundances in** 21 **Mars sediments and hydrothermal systems** 22

23 Interpretation of the stable isotopic compositions of carbon and sulfur in Mars gases, organics
24 and minerals requires a thorough understanding of the environmental context under which the
25 fractionation occurred and the degree to which original features may have been preserved.
26 Such understandings can be gained, to some degree, by using the MSL payload elements as
27 summarized in Table 7 for hydrothermal environments and in Table 8 for sedimentary
28 environments. Interpretations of isotopic data are likely to be complex and prone to significant
29 ambiguity.
30
31

32 33 **Synthesis** 34

35 The MSL instrument package has the potential to detect biosignatures if they are present at the
36 landing site on Mars. Our understanding of the formation and preservation of biosignatures on
37 Earth can guide our expectations of how and where they might have developed and be
38 preserved on Mars even though the planets have distinct histories. The classes of
39 biosignatures that could be detected and identified on Mars given appropriate biosignature
40 formation and preservation include: diagnostic organic molecules, biogenic gases, body fossils,
41 biofabrics, mineralogy affected by biomineralization and bioalteration, spatial patterns in
42 chemistry due to metabolic processes, and isotopic signatures reflecting metabolic processes.
43
44

45 Diagnostic organic molecules and biogenic gases are the most definitive as biosignatures and
46 are also readily detectable with the SAM instrument. However, they require sequestration from
47 oxidative processes for preservation. Clay minerals promote the preservation of diagnostic
48 organic molecules on Earth, and accumulations of sedimentary clay minerals commonly
49 preserve organic molecules. Thus, using Earth as a model, a MSL landing site in a sedimentary
50 basin containing clay minerals is ideal for maximizing the chances of detecting diagnostic
51 organic molecules. Carbonates and other minerals deposited as a consequence of
52 microbiological activity (microbialites) can also preserve diagnostic organic molecules.
53
54

55
56
57 Body fossils and biofabrics can also be definitive biosignatures if they are sufficiently complex
58
59
60

1
2
3 and can be observed in context. For detection by MSL, body fossils and biofabrics must be
4 large enough to be observed by MAHLI. Many terrestrial biofabrics are sufficiently large, but
5 body fossils of bacteria are not. Biofabrics are easily preserved in hydrothermal and
6 sedimentary rocks if protected from extensive recrystallization. Thus, detection of biofabrics by
7 MSL is possible if microbial communities developed on Mars and an appropriate landing site is
8 chosen. Low temperature hydrothermal and persistently wet sedimentary environments are
9 most likely to develop and preserve biofabrics.
10
11

12
13 Biomineralization and bioalteration effects on the spatial patterns in chemistry rarely produce
14 definitive microbial biosignatures due to substantial overlaps in abiotic and biotic processes.
15 However, preserved disequilibrium mineral distributions or variations in chemistry can indicate
16 good sites to look for more definitive biosignatures. CheMin, ChemCam and APX can all help
17 characterize minerals and chemical patterns that have the potential to be biosignatures.
18
19

20 Stable isotopic signatures are commonly used in terrestrial materials to characterize the extent
21 of biological chemical activity. However, such interpretations require a detailed knowledge of
22 the biogeochemical context, including cycles, of the activity. Without that context, stable
23 isotopic signatures rarely provide strong evidence for biological activity. SAM can characterize
24 the isotopic composition of various important materials. If the proper environment is
25 encountered on Mars, we have the potential to identify stable isotopic biosignatures if present.
26 However, definitive biosignature detection would require characterization of co-occurring
27 elemental reservoirs, preferably of carbon and sulfur with different oxidation states. To date, we
28 have not identified such an environment on Mars.
29
30
31

32 Early Earth is the best analogue we have currently for guiding the search for biosignatures on
33 Mars. Still, we must be constantly aware of the limits of our understanding of terrestrial
34 processes of biosignature formation and preservation, especially as they relate to the Earth's
35 earliest sedimentary record. The MSL is at the heart of the first NASA Astrobiology mission and
36 provides an extraordinary opportunity to learn more about martian environments and processes,
37 particularly in localities that might have been inhabited by microorganisms.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Alexander, C. M. O'D., Fogel M. L., Yabuta, H., Cody, G. D. (2007) The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochimica et Cosmochimica Acta* 71, 4380-4403.
- Amend, J.P., McCollom, T.M. (2009) Energetics of biomolecule synthesis on early Earth. In *Chemical Evolution II: From the Origins of Life to Modern Society*, Vol. 1025 (eds. L. Zaikowski, J.M. Friedrich, and S.R. Seidel), pp. 63-94. American Chemical Society.
- Anders, E., (1989) Pre-biotic organic matter from comets and asteroids. *Nature* 342, 255-257.
- Anders, E., Hayatsu, R., Studier, M.H. (1973) Organic Compounds in Meteorites: They may have formed in the solar nebula, by catalytic reactions of carbon monoxide, hydrogen, and ammonia. *Science*, 182, 781-790.
- Atreya, S.K. Wong, A.-S., Renno, N.O., Farrell, W. M., Delory, G.T., Sentman, D.D., Cummer, S.A., Marshall, J.R., Rafkin, S.C.R., Catling, D.C. (2006) Oxidant enhancement in Martian dust devils and storms: Implications for life and habitability. *Astrobiology* 6, 439-450.
- Aubrey A., Cleaves H. J., Chalmers J. H., Skelley A. M., Mathies R. A., Grunthaner F. J., Ehrenfreund P., and Bada J. L. (2006) Sulfate minerals and organic compounds on Mars. *Geology* 34, 357-360.
- Baross J. and others, *Limits of Life in Planetary Systems* (2007) Committee on Limits of Life in Planetary Systems, National Academies Press, ISBN-13: 978-0-309-10484-5.
- Benner, S.A., Devine, K.G., Matveeva, L.N., Powell, D.H. (2000) The missing organic molecules on Mars. *Proceedings of the National Academy of Sciences of the United States of America* 97, 2425-2430.
- Bibring, J.-P., Langevin, Y., Gendrin, A., Gondet, B., Poulet, F., Berthe, M., Soufflot, A., Arvidson, R., Mangold, N., Mustard, J., Drossart, P., and the Omega Team (2005) Mars Surface Diversity as Revealed by the OMEGA/Mars Express Observations. *Science* 307,1576-1581.
- Biemann, K. (1979) The implications and limitations of the findings of the Viking organic analysis experiment. *Journal of Molecular Evolution* 14, 65-70.
- Biemann, K. (2007) On the ability of the Viking gas chromatograph-mass spectrometer to detect organic matter. *Proceedings of the National Academy of Sciences of the United States of America* 104, 10310-10313.
- Birgel D., Himmler T., Freiwald A., and Peckmann J. r. (2008) A new constraint on the antiquity of anaerobic oxidation of methane: Late Pennsylvanian seep limestones from southern Namibia. *Geology* 36, 543-546.
- Bock, M.J. and Mayer, L.M. (2000) Mesodensity organo – clay associations in a near-shore

- 1
2
3 sediment. *Marine Geology* 163, 65-75.
4
5 Botta, O., (2005) Organic Chemistry in Meteorites, Comets, and the Interstellar Medium.
6 *Proceedings of the International Astronomical Union* 1, 479-488.
7
8 Botta, O., Bada, J.L., Gomez-Elvira, J., Javaux, E., Selsis, F. and Summons, R.E. (Eds), (2008)
9 Strategies of Life Detection. In: *Space Sciences Series of ISSI*, 25. Springer US, Bern.
10
11 Bishop, J. L., Dobreá, E. Z. N., McKeown, N. K., Parente, M., Ehlmann, B. L., Michalski, J. R.,
12 Milliken, R. E., Poulet, F., Swayze, G. A., Mustard, J. F., Murchie, S. L., and Bibring, J.-P.
13 (2008) Phyllosilicate Diversity and Past Aqueous Activity Revealed at Mawrth Vallis,
14 Mars. *Science* 321(5890), 830-833.
15
16
17
18 Bradley, A.S., and Summons, R.E. (2010) Multiple origins of methane at the Lost City
19 Hydrothermal Field. *Earth and Planetary Science Letters* 297, 34-41.
20
21 Bradley, A.S., Hayes, J.M., Summons, R.E. (2009) Extraordinary ¹³C enrichment of diether
22 lipids at the Lost City Hydrothermal Field indicates a carbon-limited ecosystem.
23 *Geochimica et Cosmochimica Acta* 73, 102-118.
24
25
26 Brasier, M., McLoughlin, N., Green, O., and Wacey, D. (2006) A fresh look at the fossil evidence
27 for early Archaean cellular life. *Philosophical Transactions of the Royal Society B:*
28 *Biological Sciences* 361, 887-902.
29
30
31 Brazelton, W.J., Schrenk, M.O., Kelley, D.S., Baross, J.A. (2006) Methane- and sulfur-
32 metabolizing microbial communities dominate the Lost City Hydrothermal Field
33 ecosystem. *Applied and Environmental Microbiology* 72, 6257-6270.
34
35
36 Butterfield, N. J. (1990) Organic Preservation of Non-Mineralizing Organisms and the
37 Taphonomy of the Burgess Shale. *Paleobiology* 16, 272.
38
39 Canfield, D. E. (2001) Biogeochemistry of sulfur isotopes. In: J.W. Valley and D.R. Cole, Eds.
40 Stable Isotope Geochemistry, *Reviews in Mineralogy* 43, 607-636.
41
42 Chang, S., Des Marais, D.J., Mack, R., Miller, S.L., Strathearn, G.E. (1983) Prebiotic organic
43 syntheses and the origin of life. In: J.W. Schopf (Ed.), *Earth's earliest biosphere: Its origin*
44 *and evolution*, Princeton University Press, Princeton, NJ., pp. 53-92.
45
46
47 Chen J., Walter M.R., Logan G.A., Hinman M.C. and Summons R.E., 2003. The
48 Paleoproterozoic McArthur River (HYC) Pb/Zn/Ag deposit of northern Australia: organic
49 geochemistry and ore genesis. *Earth and Planetary Science Letters* 210, 467-479.
50
51
52 Chivian, D., Brodie, E.L., Alm, E.J., Culley, D.E., Dehal, P.S., *et al.*, (2008) Environmental
53 Genomics Reveals a Single-Species Ecosystem Deep Within Earth. *Science* 322, 275-
54 278.
55
56
57 Christensen, P. R., Wyatt, M. B., Glotch, T. D., Rogers, A. D., Anwar, S., Arvidson, R. E.,
58 Bandfield, J. L., Blaney, D. L., Budney, C., Calvin, W. M., Fallacaro, A., Fergason, R. L.,
59
60

1
2
3 Gorelick, N., Graff, T. G., Hamilton, V. E., Hayes, A. G., Johnson, J. R., Knudson, A. T.,
4 McSween, H. Y., Jr., Mehall, G. L., Mehall, L. K., Moersch, J. E., Morris, R. V., Smith, M.
5 D., Squyres, S. W., Ruff, S. W., and Wolff, M. J. (2004) Mineralogy at Meridiani Planum
6 from the Mini-TES Experiment on the Opportunity Rover. *Science* 306, 1733-1739.
7
8

9 Cody G.D., Alexander C. M. O'D. (2005) NMR studies of chemical structural variation of
10 insoluble organic matter from different carbonaceous chondrite groups. *Geochimica et*
11 *Cosmochimica Acta* 69, 1085-1097.
12

13
14 Cody G. D., Alexander C. M. O'D., Yabuta H., Kilcoyne A. L. D., Araki T., Ade H., Dera P., Fogel
15 M. L., Militzer B., Mysen B. (2008) Organic Thermometry for Chondritic Parent Bodies,
16 *Earth and Planetary Science Letters* 272, 446-455.
17

18
19 Cronin, J.R., Pizzarello, S. (1997) Enantiomeric excesses in meteoritic amino acids. *Science*
20 275, 951-955.
21

22
23 Dahl, J., Hallberg, R., and Kaplan, I. R. (1988) Effects of irradiation from uranium decay on
24 extractable organic matter in the Alum Shales of Sweden. *Organic Geochemistry* 12,
25 559-571.
26

27
28 Des Marais, D.J., (2001) Isotopic evolution of the biogeochemical carbon cycle during the
29 Precambrian. In: J.W. Valley and D.R. Cole, Eds. *Stable Isotope Geochemistry, Reviews*
30 *in Mineralogy* 43, 555-578.
31

32
33 Dontsova, K.M., Bigham, J.M. (2005) Anionic Polysaccharide Sorption by Clay Minerals, *Soil*
34 *Science Society of America Journal* 69, 1026-1035.
35

36
37 Ehlmann, B. L., Mustard, J. F., Fassett, C. I., Schon, S. C., Head, J. W., Des Marais, D. J.,
38 Grant, J. A., and Murchie, S. L. (2008a) Clay minerals in delta deposits and organic
39 preservation potential on Mars. *Nature Geoscience* 1(6), 355-358.
40

41
42 Ehlmann, B. L., Mustard, J. F., Murchie, S. L., Poulet, F., Bishop, J. L., Brown, A. J., Calvin, W.
43 M., Clark, R. N., Des Marais, D.J., Milliken, R. E., Roach, L. H., Roush, T. L., Swayze, G.
44 A., and Wray, J. J. (2008b) Orbital Identification of Carbonate-Bearing Rocks on Mars.
45 *Science* 322(5909), 1828-1832.
46

47
48 Ehlmann, B. L., Mustard, J. F., and Murchie, S. L. (2010) Geologic setting of serpentine
49 deposits on Mars. *Geophysical Research Letters* 37, In Press.
50

51
52 Engel, M.H., Macko, S.A. (1997) Isotopic evidence for extraterrestrial non-racemic amino acids
53 in the Murchison meteorite. *Nature* 389, 265-268.
54

55
56 Farmer, J. D., Des Marais, D.J. (1999), Exploring for a record of ancient Martian life, *Journal of*
57 *Geophysical Research* 104, 977-995.
58

59
60 Farquhar, J., Johnston, D.T., Wing, B.A., Habicht, K.S., Canfield, D.E., Airieau, S., Thiemens,
M.H. (2003) Multiple sulphur isotopic interpretations of biosynthetic pathways:
implications for biological signatures in the sulphur isotope record, *Geobiology* 1, 27-36.

- 1
2
3 Farquhar, J., Savarino, J., Jackson, T.L., Thiemens, M.H. (2000) Evidence of atmospheric
4 sulphur in the Martian regolith from sulphur isotopes in meteorites. *Nature* 404, 50-52.
5
6
7 Fernandez-Remolar, D., Knoll, A.H. (2008) Fossilization potential of iron-bearing minerals in
8 acidic environments of Rio Tinto, Spain: Implications for Mars exploration. *Icarus* 194,
9 72-85.
10
11 Formisano V., Atreya S., Encrenaz T., Ignatiev N., and Giuranna M. (2004) Detection of
12 Methane in the Atmosphere of Mars. *Science* 306, 1758-1761.
13
14 Franz, H.B., Mahaffy, P.R., Farquhar, J. (2007) Preliminary estimate of sulfur isotope ratio
15 precision expected with the Sample Analysis at Mars (SAM) Instrument Suite of the 2009
16 Mars Science Laboratory. *38th Lunar and Planetary Science Conference*, abstract 1874.
17
18
19 Glavin, D.P., Dworkin, J.P. (2009) Enrichment of the amino acid l-isovaline by aqueous
20 alteration on CI and CM meteorite parent bodies. *Proceedings of the National Academy*
21 *of Sciences of the United States of America* 106, 5487-5492.
22
23
24 Greenwood, J.P., Mojzsis, S.J., Coath, C.D., (2000) Sulfur isotopic compositions of individual
25 sulfides in Martian meteorites ALH84001 and Nakhla: implications for crust–regolith
26 exchange on Mars. *Earth and Planetary Science Letters* 184, 23-35.
27
28
29 Grotzinger J. P., Arvidson R. E., Bell J. F., Calvin W., Clark B. C., Fike D. A., Golombek M.,
30 Greeley R., Haldemann A., Herkenhoff K. E., Jolliff B. L., Knoll A. H., Malin M., McLennan
31 S. M., Parker T., Soderblom L., Sohl-Dickstein J. N., Squyres S. W., Tosca N. J., and
32 Watters W. A. (2005) Stratigraphy and sedimentology of a dry to wet eolian depositional
33 system, Burns formation, Meridiani Planum, Mars. *Earth and Planetary Science Letters*
34 240, 11-72.
35
36
37 Guo, L., Santschi, P.H. (1997) Composition and cycling of colloids in marine environments.
38 *Reviews in Geophysics*, 35, 17-40.
39
40 Haskin, L. A., Wang, A., Jolliff, B. L., McSween, H. Y., Clark, B. C., Des Marais, D. J.,
41 McLennan, S. M., Tosca, N. J., Hurowitz, J. A., Farmer, J. D., Yen, A., Squyres, S. W.,
42 Arvidson, R. E., Klingelhofer, G., Schroder, C., de Souza, P. A., Ming, D. W., Gellert, R.,
43 Zipfel, J., Bruckner, J., Bell, J. F., Herkenhoff, K., Christensen, P. R., Ruff, S., Blaney,
44 D., Gorevan, S., Cabrol, N. A., Crumpler, L., Grant, J., and Soderblom, L. (2005) Water
45 alteration of rocks and soils on Mars at the Spirit rover site in Gusev crater. *Nature*
46 436(7047), 66.
47
48
49 Hazen, R.M., Papineau, D., Bleeker, W., Downs, R.T., Ferry, J.M., McCoy, T.J., Sverjensky,
50 D.A., Yang, H. (2008) Mineral evolution. *American Mineralogist* 93, 1693–1720.
51
52
53 Hedges, J.I., Hare, P.E. (1987) Amino acid adsorption by clay minerals in distilled water.
54 *Geochimica et Cosmochimica Acta* 51, 255-259.
55
56
57 Hedges, J.I., Keil, R.G. (1995) Sedimentary organic matter preservation: an assessment and
58 speculative synthesis. *Marine Chemistry* 49, 81-115.
59
60

- 1
2
3 Horita, J., Berndt, M.E. (1999) Abiogenic Methane Formation and Isotopic Fractionation Under
4 Hydrothermal Conditions. *Science* 285, 1055-1057.
5
6
7 Jakosky B.M. and others (2007) *An Astrobiological Strategy for the Exploration of Mars*.
8 Committee on an Astrobiology Strategy for the Exploration of Mars, National Academies
9 Press, ISBN-13: 978-0-309-10851-5.
10
11 Jakosky, B. M., Pepin, R. O. Johnson, R. E., Fox, J. L. (1994) Mars atmospheric loss and
12 isotopic fractionation by solar-wind-induced sputtering and photochemical escape. *Icarus*
13 111, 271-288.
14
15
16 Kelley, D.S., Karson, J.A., Fruh-Green, G.L., Yoerger, D.R., Shank, T.M., Butterfield, D.A.,
17 Hayes, J.M., Schrenk, M.O., Olson, E.J., Proskurowski, G., Jakuba, M., Bradley, A.,
18 Larson, B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A.S., Brazelton, W.J., Roe,
19 K., Elend, M.J., Delacour, A., Bernasconi, S.M., Lilley, M.D., Baross, J.A., Summons,
20 R.E., Sylva, S.P. (2005) A Serpentinite-Hosted Ecosystem: The Lost City Hydrothermal
21 Field. *Science* 307, 1428-1434.
22
23
24 Kennedy, M.J., Pevear, D.R., and Hill, R.J. (2002) Mineral Surface Control of Organic Carbon in
25 Black Shale. *Science* 295, 657-660.
26
27
28 Kennedy, M., Droser, M., Mayer, L.M., Pevear, D., Mrofka, D. (2006) Late Precambrian
29 Oxygenation; Inception of the Clay Mineral Factory. *Science* 11, 1446-1449.
30
31 Knoll, A.H. (1985) Exceptional Preservation of Photosynthetic Organisms in Silicified
32 Carbonates and Silicified Peats. *Philosophical Transactions of the Royal Society of*
33 *London. Series B, Biological Sciences* 311, 111-122.
34
35
36 Krasnopolsky, V.A., Maillard, J.P., Owen T.C. (2004) Detection of methane in the martian
37 atmosphere: evidence for life? *Icarus* 172, 537-547.
38
39 Lin, L.-H., Wang, P.-L., Rumble, D., Lippmann-Pipke, J., Boice, E., Pratt, L.M., Lollar, B.S.,
40 Brodie, E.L., Hazen, T.C., Andersen, G.L., DeSantis, T.Z., Moser, D.P., Kershaw, D.,
41 Onstott, T.C., (2006) Long-Term Sustainability of a High-Energy, Low-Diversity Crustal
42 Biome. *Science* 314, 479-482.
43
44
45 Martin, W., Baross, J., Kelley, D., Russell, M.J. (2008) Hydrothermal vents and the origin of life.
46 *Nature Reviews in Microbiology* 6, 805-814.
47
48 Martin, W., Russell, M.J. (2007) On the origin of biochemistry at an alkaline hydrothermal vent.
49 *Philosophical Transactions of the Royal Society B: Biological Sciences* 362, 1887-1926.
50
51 Martins, Z., Alexander, C.M.O.D., Orzechowska, G.E., Fogel, M.L., Ehrenfreund, P., (2007)
52 Indigenous amino acids in primitive CR meteorites. *Meteoritics & Planetary Science* 42,
53 2125-2182.
54
55
56 Mayer, L.M. (1994a) Relationships between mineral surfaces and organic carbon
57 concentrations in soils and sediments. *Chemical Geology* 114, 347-363.
58
59
60

- 1
2
3 Mayer, L.M. (1994b) Surface area control of organic carbon accumulation in continental shelf
4 sediments. *Geochimica et Cosmochimica Acta* 58, 1271-1284.
5
6
7 McCollom, T.M., Seewald, J.S. (2006) Carbon isotope composition of organic compounds
8 produced by abiotic synthesis under hydrothermal conditions. *Earth and Planetary*
9 *Science Letters* 243, 74-84.
10
11 McCollom, T.M., Seewald, J.S. (2007) Abiotic Synthesis of Organic Compounds in Deep-Sea
12 Hydrothermal Environments. *Chemical Reviews* 107, 382-401.
13
14 McKay, C.P. (2004) What Is Life-and How Do We Search for It in Other Worlds? *PLoS Biology*
15 2, e302.
16
17
18 McKay, C.P. (2007) An Approach to Searching for Life on Mars, Europa, and Enceladus. *Space*
19 *Science Reviews* 135, 49-54.
20
21 Meyers, P.A and Ishiwatari, R. (1995) Organic matter accumulation records in lake sediments.
22 In: A. Lerman et al., Editors, *Physics and Chemistry of Lakes*, Springer, pp. 279–328.
23
24 Mumma, M., Villanueva, G.L., Novak, R.E., Hewagama, T., Bonev, B.P., DiSanti, M. A. Mandell,
25 A.M., Smith M.D. (2009) Strong Release of Methane on Mars in Northern Summer 2003.
26 *Science* 323, 1041-1045.
27
28
29 Navarro-Gonzalez, R., Navarro, K.F., Rosa, J.D.L., Iniguez, E., Molina, P., Miranda, L.D.,
30 Morales, P., Cienfuegos, E., Coll, P., Raulin, F., Amils, R., McKay, C.P. (2006) The
31 limitations on organic detection in Mars-like soils by thermal volatilization-gas
32 chromatography-MS and their implications for the Viking results. *Proceedings of the*
33 *National Academy of Sciences of the United States of America* 103, 16089-16094.
34
35
36 Noffke, N. (2009) The criteria for the biogenicity of microbially induced sedimentary structures
37 (MISS) in Archean and younger sandy deposits. *Earth Science Reviews* 96, 173-180.
38
39
40 Ohashi, H., Nakazawa, H. (1996) The microstructure of humic acid-montmorillonite composites.
41 *Clay Minerals* 31, 347-354.
42
43 Owen, T., Bar-Nun, A., Kleinfield, I. (1992) Possible cometary origin of heavy noble gases in the
44 atmospheres of Venus, Earth and Mars. *Nature*, 358, 43-46.
45
46
47 Oze, C.and Sharma, M. (2005) Have olivine, will Gas: Serpentinization and the abiogenic
48 production of methane on Mars," *Geophysical Research Letters* 32, L10203-L10206.
49
50 Panieri, G., Lugli, S., Manzi, V., Roveri, M., Schrieber, B.C., Palinska, K.A. (2010) *Geobiology* 8,
51 101-111.
52
53 Pizzarello, S. (2006) The chemistry of life's origin: a carbonaceous meteorite perspective.
54 *Accounts of Chemical Research* 39, 231-7.
55
56
57 Pratt, L.M., Lefticariu, L., Ripley, E.M., Onstott, T.C., (2006) Radiolysis of water as a source of
58 bioavailable chemical energy. *Geochimica et Cosmochimica Acta* 70 (Issue18,
59
60

- 1
2
3 Supplement 1), A503.
4
5 Proskurowski, G., Lilley, M.D., Seewald, J.S., Frueh-Green, G.L., Olson, E.J., Lupton, J.E.,
6 Sylva, S.P., Kelley, D.S. (2008) Abiogenic Hydrocarbon Production at Lost City
7 Hydrothermal Field. *Science* 319, 604-607.
8
9
10 Rasmussen, B. (2000) Filamentous microfossils in a 3,235-million-year-old volcanogenic
11 massive sulphide deposit. *Nature* 405, 676-679.
12
13 Rye, R.O. (2005) A review of the stable-isotope geochemistry of sulfate minerals in selected
14 igneous environments and related hydrothermal systems. *Chemical Geology* 215, 5-36.
15
16 Sephton, M.A. (2002) Organic compounds in carbonaceous meteorites. *Natural Products*
17 *Research* 19, 292 - 311.
18
19
20 Sephton, M.A., Botta O. (2008) Extraterrestrial organic matter and the detection of life. *Space*
21 *Science Reviews* 135, 25-35.
22
23 Sephton, M.A., Love, G.D., Watson, J.S., Verchovsky, A.B., Wright, I.P., Snape, C.E., Gilmour,
24 I. (2004) Hydroxylation of insoluble carbonaceous matter in the Murchison meteorite:
25 new insights into its macromolecular structure. *Geochimica et Cosmochimica Acta* 68,
26 1385-1393.
27
28
29 Sherwood Lollar, B., Lacrampe-Couloume, G., Slater, G.F., Ward, J., Moser, D.P., Gihring,
30 T.M., Lin, L.H., Onstott, T.C. (2006) Unraveling abiogenic and biogenic sources of
31 methane in the Earth's deep subsurface. *Chemical Geology* 226, 328-339.
32
33
34 Shock, E.L., Schulte, M.D. (1998) Organic synthesis during fluid mixing in hydrothermal
35 systems. *Journal of Geophysical Research [Planets]* 103(E12), 28513-28527.
36
37
38 Sleep, N.H., Bird, D.K. (2007) Niches of the pre-photosynthetic biosphere and geologic
39 preservation of Earth's earliest ecology. *Geobiology* 5, 101-117.
40
41
42 Sleep, N.H., Meibom, A., Fridriksson, T., Coleman, R.G., Bird, D.K. (2004) H₂-rich fluids from
43 serpentinization: geochemical and biotic implications. *Proceedings of the National*
44 *Academy of Sciences of the United States of America* 101, 12818-12823.
45
46
47 Smith, J.W., Kaplan, I.R. (1970) Endogenous Carbon in Carbonaceous Meteorites. *Science*
48 167, 1367-1370.
49
50 Squyres, S. W., Arvidson, R. E., Bell, J. F., III, Bruckner, J., Cabrol, N. A., Calvin, W., Carr, M.
51 H., Christensen, P. R., Clark, B. C., Crumpler, L., Des Marais, D. J., d'Uston, C.,
52 Economou, T., Farmer, J., Farrand, W., Folkner, W., Golombek, M., Gorevan, S., Grant,
53 J. A., Greeley, R., Grotzinger, J., Haskin, L., Herkenhoff, K. E., Hviid, S., Johnson, J.,
54 Klingelhofer, G., Knoll, A., Landis, G., Lemmon, M., Li R., Madsen, M. B., Malin, M. C.,
55 McLennan, S. M., McSween, H. Y., Ming, D. W., Moersch, J., Morris, R. V., Parker, T.,
56 Rice, J. W., Jr., Richter, L., Rieder, R., Sims, M., Smith, M., Smith, P., Soderblom, L. A.,
57 Sullivan, R., Wanke, H., Wdowiak, T., Wolff, M., and Yen, A. (2004) The Spirit Rover's
58
59
60

1
2
3 Athena Science Investigation at Gusev Crater, Mars. *Science* 305, 794-799.
4

5 Squyres, S.W., A.H. Knoll, R.E. Arvidson, B.C. Clark, J.P. Grotzinger, B.L. Jolliff, S.M.
6 McLennan, N. Tosca, J.F. Bell III, W.M. Calvin, W.H. Farrand, T.D. Glotch, M.P.
7 Golombek, K.E. Herkenhoff, J.R. Johnson, G. Klingelhofer, H.Y. McSween, and A.S.
8 Yen (2006) Continuing observations by the Opportunity rover at Meridiani Planum,
9 Mars. *Science* 313, 1403-1407.
10
11

12 Squyres, S.W., et al.(2009) Exploration of Victoria Crater by the rover Opportunity.
13 *Science* 324, 1058-1061.
14

15
16 Steele, A., Fries, M. D., Amundsen, H. E. F., Mysen, B. O., Fogel, M. L., Schweizer, M. (2007)
17 Comprehensive imaging and Raman spectroscopy of carbonate globules from Martian
18 meteorite ALH 84001 and a terrestrial analogue from Svalbard. *Meteoritics & Planetary*
19 *Science* 42, 1549-1566.
20

21 Summons, R.E., Albrecht, P., McDonald, G., Moldowan, J.M. (2007) Molecular Biosignatures:
22 Generic Qualities of Organic Compounds that Betray Biological Origins. *Space Science*
23 *Reviews* 135, 133-157.
24



25
26 Swindle, T.D. (2002) Martian Noble Gases. *Reviews in Mineralogy and Geochemistry* 47, 171-
27 190.
28

29 Trewin, N. H. (1996) The Rhynie cherts: an early Devonian ecosystem preserved by
30 hydrothermal activity. *Evolution of Hydrothermal Ecosystems on Earth (and Mars?)*, Ciba
31 Foundation Symposium #202, pp. 131-45.
32

33
34 Walter, M. R. (1996) Ancient hydrothermal ecosystems on earth: a new palaeobiological
35 frontier. *Evolution of Hydrothermal Ecosystems on Earth (and Mars?)*, Ciba Foundation
36 Symposium #202, pp. 112-27
37

38 Xiao S. and Knoll A. H. (1999) Fossil preservation in the Neoproterozoic Doushantuo
39 phosphorite Lagerstätte, South China. *Lethaia* 32(3), 219-238.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Biosignatures detectable by MSL	How definitive as a biosignature?	How well it can be measured by MSL?
biogenic organic molecules	highly definitive	readily with SAM
biogenic gases	often definitive	readily with SAM
body fossils	often definitive	with MAHLI if large enough
biofabrics	sometimes definitive	with MAHLI , MastCam
stable isotopic compositions	occasionally; context critical	readily with SAM
biomineralization/ alteration	rarely definitive	detectible with CheMin, ChemCam
spatial chemical patterns	rarely definitive	detectible with CheMin, ChemCam

Table 1

Biosignatures taphonomic window	Confidence in the geological context	How this informs about potential biosignature preservation
atmospheric gases	very high	predictable via chemical modeling
crystalline sedimentary mineral entrapment of organics	very high	can deduce formation mechanism and subsequent history
biofabric lithification	very high	can deduce history from lithology and stratigraphic relationships
body fossil preservation	very high	can deduce history from lithology and stratigraphic relationships
mineral replacement of body fossil	high	can deduce from mineralogy

Table 2

Martian context --> Early Mars Environment	Support biotic OM formation	Support for abiotic OM formation	Support OM conc'n	Support preservation	Potential for recent exhumation*	ID by remote sensing			ID by MSL
						Geo-morphic	Mineral-ogic	Strati-graphic	
Aeolian sediments (sand)	low	low	low	low	low ??	high	n/a	mod	high
altered aeolinites (dust)	very low	low	low	low	low ??	low	n/a	n/a	high
Fluvial channel	low	low	low	low	high	high	n/a	high	high
Fluvial floodplain	low-mod	low	mod	mod	possible ??	high	n/a	high	high
alluvial fan	low	low	low	low	low ??	high	n/a	high	high
Deltaic	high	low	high	high	low ??	high	n/a	high	high
Lacustrine (perennial)	high	low	high	high	high	mod	mod	mod	high
Lacustrine (evaporitic)(Cl)	low	low	high	high-very high	high	mod	high	mod	high
Lacustrine (evaporitic)(SO4)	mod	low	high	high-very high	high	mod	high	mod	high
Regional groundwater pore system	low	low	low	low	high ??	n/a	n/a	n/a	mod
Glacial deposits	low	low	low	low	high	high	n/a	low	high
permafrost	low	low	low	mod	mod	high	n/a	n/a	high
soil (surface fines chemically altered by atmosphere)	low	low	low	low	low	n/a	n/a (albedo and TI)	n/a	high
Regolith/Fractured Bedrock (not soil)	low	low	low	low	low	high	n/a	n/a	high

* Poorly constrained estimates; require improved knowledge of erosional processes and timescales

Table 3

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Specific cases	Support biotic OM formation	Support for abiotic OM formation	Support OM conc'n	Support preservation	Potential for recent exhumation*	ID by remote sensing			ID by MSL
						Geo-morphic	Mineral-ogic	Strati-graphic	
Pyroclastic Deposits (unaltered)	low	low	low	low	low?	mod	low	high	high
Volcanic flows	very low	low	low	low	low?	high	high	mod	high
aqueous altered at surface	low	low	low	mod-high	low?	low	mod	low	mod
aqueous altered in subsurface	low	low	low	mod-high	low?	low	mod	low	mod
aqueous altered at surface	low	low	low	low-mod	low?	low	mod	low	mod
Volcanics-hydrothermal altered	mod	low	low	mod	mod	mod	mod	low	high
aqueous altered in subsurface	mod	mod	low	low-mod	mod	low	mod	low	mod
Hydrothermal (<100C) subsurface	mod	mod (F/T)	mod-low	mod	mod	mod	mod-high	n/a	high
Hydrothermal (<100C) surface	high	low	mod-high	mod	mod	high	mod-high	low	high
ultramafic subsurface (<100C)	high	low	mod-high	mod	mod	high	mod-high	low	high

* Poorly constrained estimates;require improved knowledge of erosional processes and timescales

Table 4

	Specific Processes	EGA	GCMS	APXS	CheMin	ChemCam	MAHLI
Abiotic Processes	F/T and catalyzed polymerization	X	X	X	X		
Bioprocess	Redox dependent metabolisms (e.g. H ₂ /Fe/S/C - metabolisms)	X	X	X	X	X	
Conc Process	Adsorption to mineral surfaces			X	X	X	
	Immiscibility & migration	X					X
Preservation Process	Adsorption to mineral surfaces			X	X	X	
	mineral encapsulation (trapping)	X				X	X
	thermal polymerization (pyrobitumen formation)	X			X	X	X (UV)

Table 5

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

	Specific Processes	EGA	GCMS	APXS	CheMin	ChemCam	MAHLI
	Cosmic (footnote:specific targets for PE are organic molecules and Cr, Ni anomalies)		X	X	X		
	Atmospheric photolytic synthesis		X	X	X		
Bioprocess	chemosynthesis		X	X	X		
	photosynthesis	X (%TOC)	X	X	X		
Conc Process	sorting of transported organics and organics in/on minerals		X [#]	X [#]	X	X	X
Preservation Process	lithification (includes cementation)	X		X	X	X	X
	Adsorption to mineral surfaces			X	X	X	X
	Burial			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X		X	X	X	X

Table 6

	Specific Processes	EGA-TLS	EGA-QMS	GCMS	APXS	CheMin	ChemCam	MAHLI
Fractionation-Related Processes	impacts (excavation for access deep-carbon sources)	X	X					
	atmospheric processes (e.g. photolysis)	X	X?					
	Biology	X	X?			x		
	hydrologic cycle (over martian history indirectly understood via CO2 record)	X			x	x		
	Global C cycling	X			x	x		
	Global S cycling		X		x	x		
	other biogeochemical cycling	X	X		x	x		
	volcanism	X	X		x	x		
Environmental State	atmospheric composition (past & present), temperature, pressure	X	X					
	water inventories, state, and activity	X	X			x		
	Redox state	X	X		X	x		
	pH	X	X		X	x		
	surface materials (in exchange with atmosphere)	X	X			x		
Preservation Process	lithification (includes cementation)	X	X		X	X	X	X
	Adsorption to mineral surfaces	X			X	X	X	X
	Burial	X			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X	X		X	X	X	X

Table 7

	Specific Processes	EGA-TLS	EGA-QMS	GCMS	APXS	CheMin	ChemCam	MAHLI
Processes	impacts (excavation for access deep-carbon sources)	X	X					
	cosmic (influx and loss)	X	X					
	weathering	X	X					
	atmospheric processes (e.g. photolysis)	X	X?					
	Biology	X	X?			X		
	hydraulic cycle (over martian history indirectly understood via CO2 record)	X			X	X		
	C cycling	X			X	X		
	S cycling		X		X	X		
	other biogeochemical cycling	X	X		X	X		
	volcanism	X	X		X	X		
	aeolian (fractionation via oxidation of materials or concentration of materials from particular sources)	X	X			X		
Environmental State	atmospheric composition (past & present), temperature, pressure	X	X					
	water inventories, state, and activity	X	X			X		
	Redox state	X	X		X	X		
	pH	X	X		X	X		
	surface materials (in exchange with atmosphere)	X	X			X		
Preservation Process	lithification (includes cementation)	X	X		X	X	X	X
	Adsorption to mineral surfaces	X			X	X	X	X
	Burial	X			X	X	X	X
	Co-precipitation (sorption) of mineral/organics	X	X		X	X	X	X

Table 8

	Processes	Features	Preservation Window
Physical	Brecciation Fracturing	hydraulic breccia/fractures	Avoidance of strong structural deformation and very high thermal metamorphism
Chemical	Mineral Precipitation +- Alteration, Dissolution	cement distribution & composition, alteration crusts, porosity, dissolution surfaces, laminations, terraces, mounds	Avoidance of recrystallization & dissolution, increases in oxidation state (e.g. Fe/Mn, perchlorate, sulfate, clay hydration); avoidance of structural deformation and very high thermal metamorphism
Biological	Biomass Accumulation	"mats, stromatolites, etc; roll-ups, streamers, elemental concentrations, coking	Avoidance of strong structural deformation and very high thermal metamorphism; coke in fractures

Table 9

	Processes	Features	Preservation Window
Physical	Sediment Transport; Soil/permafrost	grain size, sorting, rounding, composition cross strat laminations, cracks/fracturing soft sediment deposition	dewatering, recrystallization of sulfates, hydration changes; structural deformation and very high thermal metamorphism
Chemical	Mineral Precipitation +- Alteration, Dissolution	cement distribution and composition alteration crusts porosity dissolution surfaces concretions displacive recrystallization lamination	Avoidance of recrystallization & dissolution, increases in oxidation state (e.g. Fe/Mn, perchlorate, sulfate, clay hydration); avoidance of structural deformation and very high thermal metamorphism
Biological	Biomass Accumulation	mats, stromatolites, etc roll-ups, wrinkled structures elemental concentrations fenestrae grain size variations	Avoidance of recrystallization, hydration, elemental mobility, organic remineralization; Avoidance of structural deformation and very high thermal metamorphism

Table 10

Table 11: Chemical and Mineralogic Features of Hydrothermal Environments

Processes	Features	Environmental Indicators	Preservation Window	Payload elements	
Environmental	Fluid Convection, selective mineral phase dissolution, alteration, migration, and precipitation; element leaching and transport; Redox chemistry; cooling/heating;	mineral and element zonation/gradients (i.e. characteristic mineralogy associated with variations of hydrothermal environments, such as: chlorite, silica sulfides, kaolinite, hematite, gold, carbonates, sulfates; etc.); spatial distribution of alteration minerals and precipitates; mineral assemblages for redox and temperature*; fracture fills; pH differences; m-km scale mineral zonation		encasement of sensitive phases in less sensitive phases (e.g. silica or organic entombment of minerals); avoidance of oxidation, metamorphism, continued hydrothermal activity	Mastcam, APXS, Chemcam, CheMin, SAM-EGA
Biological	Accelerated redox reactions, organic synthesis, altered mineral assemblages	Altered mineral or elemental abundances, corroded (e.g., bored) minerals, biofabrics, organic matter	Mineral, elemental and textural indicators of aqueous conditions and mineral precipitation	Entombment in minerals formed during cooling or other changes that favor precipitation	MAHLI, APXS, Chemcam, CheMin, SAM-EGA
Physical	Fluid Convection	Hydraulic brecciation; Geomorphic evidence of springs; Overall architecture of system	Heat source	Broad due to large scale	MastCam
	Temperature Changes	Heat flow; Induces variations in chemical properties in time and over spatial scales from m to km	Heat source; Flow patterns	Large scale variations have high preservation potential; small scale changes require avoidance of recrystallization and elemental leaching	Mastcam, MAHLI, ChemCam
Chemical, Potentially Biological	Mineral Precipitation	Evidence of water chemistry & temperature; Springs mounds & terraces; Some mineral precipitation can be localized by organic carbon	Temperature indicators from low T to high T: BaSO ₄ , PbS, ZnS, SO ₄ /Sulfides together, saddle dolomite, CuS, CaSO ₄ ; Low pH indicators: jarosite, abundant kaolinite, alunite vs carbonate	Encasement of sensitive phases in less sensitive phases (e.g. silica or organic entombment of minerals); Avoidance of oxidation, metamorphism, continued hydrothermal activity	MastCam, MAHLI, APXS, ChemCam, CheMin, SAM-EGA
	Mineral Alteration & Dissolution	Evidence of changes in water chemistry			
	Elemental Leaching & Transport	m to km-scale spatial gradients in elemental composition			
	Redox Changes	Important redox indicators: S ²⁻ , S ⁰ , S ⁶⁺ , Fe ²⁺ , Fe ³⁺ , Mn ²⁺ , Mn ⁴⁺ , U ²⁺ , U ⁴⁺ , As ³⁺ , As ⁵⁺			

*** Temp mineral assemb.**

mod T SO₄/sulfide together
saddle dolomite
Low T --> high T PbS, ZnS, CuS
Low T --> high T BaSO₄--> CaSO₄

*** Redox mins**

S²⁻ --> S⁰-->S⁶⁺
Fe²⁺ --> Fe³⁺
Mn²⁺ --> Mn⁴⁺
U²⁺ --> U⁴⁺
As³⁺ --> As⁵⁺

*** low pH mins**

jarosite, large kaolinite conc., alunite vs carb.

Important factors	Factor related Feature	Preservation Window	Payload elements
Sediment source	detrital mineral assemblages	Early lithification is favorable. Avoid the following: sediments altered by recrystallization and dissolution (especially evaporites), redox changes toward oxidizing conditions, and strong structural alteration and strong thermal metamorphism	Mastcam, MAHLI, APXS, Chemcam, CheMin, SAM-EGA
H ₂ O solute chemistry and flux	evaporite assemblages and trace element distribution		
Redox and pH	Redox sensitive (Fe, Mn, S, U, etc.) and pH sensitive (Jarosite, alunite, kaolinite, etc. Vs carbonates) mineral phase and abundance		
bio-mineral interactions	bio-magnetite, bio carbonate, bio-sulfides, oxides, phosphate precipitation; bioleaching and dissolution products		

Table 12