



# Temporal and Spatial Expression Patterns of Bone Morphogenetic Protein 3 in Developing Zebrafish

## Citation

Ito-Amano, Midori, Yukio Nakamura, Mika Morisaki, Xinjun He, Masanori Hayashi, Ramida Watanapokasin, and Hiroyuki Kato. 2014. "Temporal and Spatial Expression Patterns of Bone Morphogenetic Protein 3 in Developing Zebrafish." *The Open Rheumatology Journal* 8 (1): 69-72. doi:10.2174/1874312901408010069. <http://dx.doi.org/10.2174/1874312901408010069>.

## Published Version

doi:10.2174/1874312901408010069

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:13347483>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

# Temporal and Spatial Expression Patterns of Bone Morphogenetic Protein 3 in Developing Zebrafish

Midori Ito-Amano<sup>1</sup>, Yukio Nakamura<sup>\*1</sup>, Mika Morisaki<sup>1</sup>, Xinjun He<sup>2</sup>, Masanori Hayashi<sup>1</sup>, Ramida Watanapokasin<sup>3</sup> and Hiroyuki Kato<sup>1</sup>

<sup>1</sup>Department of Orthopaedic Surgery, Shinshu University School of Medicine, Asahi 3-1-1, Matsumoto 390-8621, Japan

<sup>2</sup>Department of Stem Cell and Regenerative Biology, Harvard University, Cambridge, Massachusetts 02138, USA

<sup>3</sup>Department of Biochemistry, Faculty of Medicine, Srinakharinwirot University, Bangkok 10110, Thailand

**Abstract:** Bone morphogenetic proteins (BMPs) are important elements in bone biology. We herein report the expression profiles of zebrafish *bmp3* (*zbmp3*) as demonstrated by real-time PCR and *in situ* hybridization. The expression of *zbmp3* was highly detectable by real-time PCR from 1 day post-fertilization (1 dpf) to 2 weeks post-fertilization (2 wpf) and peaked at 1 wpf. For *in situ* hybridization experiments, *zbmp3* was expressed in the otic vesicle at 1 dpf, 2 dpf, 3 dpf, and 5 dpf. It was also expressed in the pharyngeal arches, including the opercle, branchiostegal ray, and pectoral fins, at 2 dpf. Our results suggest that *zbmp3* may play an important role in the skeletal biology of developing zebrafish.

**Keywords:** Bmp3, Expression patterns, Zebrafish.

## INTRODUCTION

Bone morphogenetic proteins (BMPs) are members of the transforming growth factor-beta (TGF-beta) superfamily. Although most BMPs are positive regulators of bone biology, BMP3 is an antagonist of osteogenic BMPs [1]. The *BMP3* gene is highly conserved across species, suggesting that it is evolutionarily important for development in mammals [2]. A previous study uncovered that murine *Bmp3* was expressed in the cartilaginous cells of otic capsule sections using *in situ* hybridization [3]. However, as there have been few reports on the temporal and spatial expression profiles of *Bmp3* in whole animals, we examined the expression patterns of *bmp3* in developing vertebrates using a zebrafish model system.

## MATERIALS AND METHODS

### Extraction of Total RNA from Zebrafish Embryos and Cloning of Zebrafish *bmp3* (*zbmp3*)

Total RNA from zebrafish embryos and larvae was extracted at selected time points using an Isogen extraction kit (*Wako Co.*, Japan). Full-length *zbmp3* (GenBank accession no. NM\_001077765) was cloned into a pCR-Blunt2-TOPO vector (*Invitrogen*, Japan) between EcoRI sites. The primers used to amplify the *zbmp3* full-length cDNA were 5'-ATGGATCGCTGTCAGCGCCTGTTTGTCTCC-3' (forward) and 5'-TTACCGACAGGCGCAGGAGTCCACTGTCATG-3' (reverse). The PCR product size was 1359 bp.

### Real-Time Polymerase Chain Reaction (PCR)

The primers used to amplify the *zbmp3* cDNA for real-time PCR experiments were 5'-AAGGGCCATTTGGGAACCAT-3' (forward) and 5'-TGTGGCTGCTGTTGTGAAGA-3' (reverse). The PCR product size was 130 bp. Real-time PCR using SSo Advanced SYBR Green supermix (*BioRad*, Japan) with 1-cell stage embryos (1 cell) and at 3 hours post-fertilization (hpf), 8-9 hpf, 1 day post-fertilization (dpf), 2 dpf, 3 dpf, 1 week post-fertilization (wpf), and 2 wpf were performed as described previously [4].


### *In Situ* Hybridization

*In situ* hybridization was performed as earlier described [5] on 1, 2, 3, and 5 dpf embryos and larvae using full-length *zbmp3*. Digoxigenin-labeled RNA probes were generated using an Sp6 labeling kit (*Roche*, USA) according to the manufacturer's instructions. Signals were detected with an ALP-conjugated anti-digoxigenin antibody (*Roche*) and visualized using 4-nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl-phosphate. The reaction was stopped by replacing the substrate with several rinses of PBT (PBS plus 0.1% Tween 20). Embryos were stored in the dark at 4°C in PBS containing 0.02% sodium azide until they were photographed.

## RESULTS

As shown in Fig. (1), the real-time PCR expression level of *zbmp3* was very low at 1 cell, 3 hpf, and 8-9 hpf time points. However, *zbmp3* expression was dramatically increased from 1 dpf to 2 wpf and peaked at 1wpf.

\*Address correspondence to this author at the Department of Orthopaedic Surgery, Shinshu University School of Medicine, Asahi 3-1-1, Matsumoto 390-8621, Japan; Tel: +81-263-37-2659; Fax: +81-263-35-8844; E-mail: [yxn14@aol.jp](mailto:yxn14@aol.jp)




**Fig. (1).** Expression study of *zbmp3* by real-time PCR using samples from 1 cell, 3 hpf, 8-9 hpf, 1 dpf, 2 dpf, 3 dpf, 1 wpf, and 2 wpf time points.

For *in situ* hybridization experiments, *zbmp3* was not detectable before 1 dpf, but was specifically expressed in the otic vesicle at 1 dpf (Fig. 2a; black arrows), 2 dpf (Fig. 2b, c; black arrows), 3 dpf (Fig. 2d, e; black arrows), and 5 dpf (Fig. 2f, g; black arrows). In addition, as Cheah *et al.* have reported previously, *zbmp3* was also expressed in the opercle (op) and branchiostegal ray (bsr) (Fig. 2c; black arrowhead) and pectoral fins (Fig. 2c; white arrowhead) at 2 dpf. Each experiment was repeated at least twice. A sense (forward) probe of *zbmp3* was used as a control, for which no specific signal was detected.


## DISCUSSION

We observed increased and localized expression of *zbmp3* at specific time points in the present zebrafish development model study. Our results suggest that *bmp3* may be a key player in skeletogenesis in vertebrates.

Schoenebeck *et al.* have reported that *zbmp3* was expressed in the pharyngeal arches and pectoral fins of zebrafish at 48-72 hpf. However, *zbmp3* expression in the otic vesicle was not remarkable in their study as it was in ours (Fig. 2a-g) [6]. The reason for this disparity is not clear.



**Fig. (2).** *In situ* hybridization of *zbmp3* in developing zebrafish. (a) Lateral view of a 1 dpf zebrafish. Arrow indicates otic vesicle. (b) Dorsal view of a 2 dpf zebrafish. Arrows indicate otic vesicles and arrowheads show pectoral fins. (c) Lateral view of a 2 dpf zebrafish. Arrow indicates otic vesicle, black arrowhead shows opercle and branchiostegal ray, and white arrowhead shows pectoral fin. (d) Lateral view of a 3 dpf zebrafish. Arrow indicates otic vesicle. (e) Dorsal view of a 3 dpf zebrafish. Arrows indicate otic vesicles. (f) Lateral view of a 5 dpf zebrafish. Arrow indicates otic vesicle. (g) Dorsal view of a 5 dpf zebrafish. Arrows indicate otic vesicles.



**Fig. (3).** Synteny analysis for *zbmp3* with human *BMP3*. Zebrafish and human *bmp3* genomic loci were analyzed and their neighbor genes were listed in order of occurrence in zebrafish and human genomes. Shared neighbor genes were linked between zebrafish and humans to reveal evolutionary conservation of this specific genomic locus. The *bmp3* gene is indicated by a black background on zebrafish chromosome 5 and human chromosome 4.

Whereas Schoenebeck used an Oregon AB line [6], we adopted Cyprinidae, Rasborinae (*Danio Rerio*). Therefore, it is possible that a difference in genetic background may influence *zbmp3* expression patterns in zebrafish lines.

The vertebrate inner ear has two major functions: 1) maintenance of body balance and 2) auditory perception. The otic vesicle plays a role in the development of ear interaction with the hindbrain [7]. Since *zbmp3* was strongly expressed in the otic vesicle in zebrafish, it may influence the development of the zebrafish ear.

Runx2 is a transcriptional factor required for osteoblastic differentiation and the development of hypertrophic chondrocytes. In zebrafish, there are 2 orthologs of *Runx2*, *runx2a* and *runx2b* [8]. These genes were found to be expressed in the developing zebrafish skeleton, including the pharyngeal arch, op, and pectoral fins, which coincided with *zbmp3* expression patterns in the present study. These results suggest a functional overlap among *runx2a*, *runx2b*, and *bmp3* in zebrafish.

To our knowledge, there have been no reports on the relationship between *BMP3* and rheumatological diseases. However, other *BMP* family members, such as *BMP4*, *BMP6*, and *Runx2*, are possible biomarkers of bone metabolism in several forms of arthritis since their expression patterns are different in *OA* and *RA* groups from those in controls [9]. Based on our findings, further studies are warranted on the role of *BMP3* in patients with rheumatic disease.

Gene duplication has been observed in the zebrafish genome due to an additional whole genome duplication event that is specific to the teleost fish lineage [10]. To study the conservation of *Bmp3* function in vertebrates, we lastly performed a synteny analysis for *zbmp3* with human *BMP3* as shown in Fig. (3). We found only one *zbmp3* gene, on chromosome 5, in the zebrafish genome, although a chromosomal segment duplicate on chromosome 10 that

harbored a deleted *zbmp3* duplicate was detected as well. Thus, one *zbmp3* duplicate appears to have been lost in chromosome 10 during zebrafish genome evolution.

## CONCLUSION

Our results suggest that *zbmp3* plays a role in the arch skeletal system in developing zebrafish. Functional analyses using genetically modified zebrafish are needed to better understand the *in vivo* biological functions of *zbmp3*. The relationship of *BMP3* with *OA* and *RA* may also require further analysis.

## AUTHOR CONTRIBUTIONS

Y.N. directed this study; M.I.A. and M.M. performed experiments and drafted the article; X.H., M.H., M.M., R.W. and H.K. contributed to conception and design, interpreted the data, and revised the article. All authors approved the final version.

## CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

## ACKNOWLEDGEMENTS

Declared none.

## REFERENCES

- [1] Daluiski A, Engstrand T, Bahamonde ME, *et al.* Bone morphogenetic protein-3 is a negative regulator of bone density. *Nat Genet* 2001; 27: 84-8.
- [2] Lowery JW, Lavigne AW, Kokabu S, *et al.* Comparative genomics identifies the mouse *Bmp3* promoter and an upstream evolutionary conserved region (ECR) in mammals. *PLoS One* 2013; 8: Epub 2013.

- [3] Stankovic KM, Adachi O, Tsuji K, *et al.* Differences in gene expression between the otic capsule and other bones. *Hear Res* 2010; 265: 83-9.
- [4] Chandrasekar G, Vesterlund L, Hultenby K, *et al.* The zebrafish orthologue of the dyslexia candidate gene DYX1C1 is essential for cilia growth and function. *PLoS One* 2013; 8: Epub 2013.
- [5] Nakamura Y, Weidinger G, Liang JO, *et al.* The CCN family member Wisp3, mutant in progressive pseudorheumatoid dysplasia, modulates BMP and Wnt signaling. *J Clin Invest* 2007; 117: 3075-86.
- [6] Schoenebeck JJ, Hutchinson SA, Byers A, *et al.* Variation of BMP3 contributes to dog breed skull diversity. *PLoS Genet* 2012; 8: e1002849.
- [7] Malicki J, Schier AF, Solnica-Krezel L, *et al.* Mutations affecting development of the zebrafish ear. *Development*. 1996; 123: 275-83.
- [8] Flores MV, Tsang VW, Hu W, *et al.* Duplicate zebrafish runx2 orthologues are expressed in developing skeletal elements. *Gene Expr Patterns* 2004; 4: 573-81.
- [9] Grcevic D, Jajic Z, Kovacic N, *et al.* Peripheral blood expression profiles of bone morphogenetic proteins, tumor necrosis factor-superfamily molecules, and transcription factor Runx2 could be used as markers of the form of arthritis, disease activity, and therapeutic responsiveness. *J Rheumatol* 2010; 37: 246-56.
- [10] Catchen JM, Braasch I, Postlethwait JH. Conserved synteny and the zebrafish genome. *Methods Cell Biol* 2011; 104: 259-85.

---

Received: February 4, 2014

Revised: July 18, 2014

Accepted: July 19, 2014

© Ito-Amano *et al.*; Licensee *Bentham Open*.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.