



The BrAIST study and the implications for scoliosis screening: our duty for raising awareness and advocacy

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Grivas, Theodoros B, Tomasz Kotwicki, Toru Maruyama, Joseph OBrien, Hubert Labelle, and Timothy Hresko. 2014. "The BrAIST study and the implications for scoliosis screening: our duty for raising awareness and advocacy." <i>Scoliosis</i> 9 (Suppl 1): 042. doi:10.1186/1748-7161-9-S1-042. http://dx.doi.org/10.1186/1748-7161-9-S1-042 .
Published Version	doi:10.1186/1748-7161-9-S1-042
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:13890696
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

ORAL PRESENTATION

Open Access

The BrAIST study and the implications for scoliosis screening: our duty for raising awareness and advocacy

Theodoros B Grivas^{1*}, Tomasz Kotwicki², Toru Maruyama³, Joseph OBrien⁴, Hubert Labelle⁵, Timothy Hresko⁶

From 11th International Conference on Conservative Management of Spinal Deformities - SOSORT 2014 Annual Meeting
Wiesbaden, Germany. 8-10 May 2014

Background

A recently published NIH-funded study named “Bracing in Adolescent Idiopathic Scoliosis Trial” (BrAIST) affirmed the efficacy of bracing for moderate adolescent idiopathic scoliosis (AIS) and reduced surgical recommendations in braced patients. The implications of this study are numerous. BrAIST affirms the value of bracing, but also confirms the importance of wear time, and points to the need to assess compliance with compulsory utilization of brace monitors. Additionally, one of the key implications of the study relates to early detection, possibly heralding a new era for scoliosis screening policy adoption.

Aim

To engage and enlist the feedback and potential advocacy, on behalf AIS patients, of all persons involved in relevant state agencies, scientific organizations and health care professions.

Methods

After a short presentation of BrAIST results, its implications will be highlighted on a) brace treatment, b) the importance of compliance, c) the value of screening programs. Subsequently the current concepts and recommendations on screening will be analyzed.

Design

Other (Advocacy/Awareness raising report).

Results

This established bracing effectiveness, reducing the number of patients who progress to surgery, resulting in cost savings and great benefit for scoliotics, also calls for raising awareness and advocacy of all the involved related professionals. Awareness will be increased by familiarizing people with the a) history and geography of screening policies, b) current evidence about screening, c) impact of screening on frequency of surgical treatment and its negative discontinuation effects on patients, d) the analysis of evolving aim of screening, e) the encouragement for policy statement publications from professional organizations and governmental agencies regarding scoliosis screening, f) by legislating national program and g) by presentations in pertinent scientific meetings. The advocacy will be benefited by a) providing guidelines on setting up these programs, b) recommendations for improvement of a screening program and c) popularizing its additional benefits on its contribution to clinical research on IS aetiology.

Conclusions

Due to the fact that the implementation of screening programs is inextricably bonded to non-operative IS treatment, it is believed that this reported “BrAIST” trial will have further impact on IS management swinging the pendulum.

Authors' details

¹Department of Orthopedics and Traumatology Tzani General Hospital Tzani and Afendouli 1, Piraeus, Greece. ²Department of Paediatric Orthopaedics University of Medical Sciences, Poznan, Poland. ³Department of Orthopedic Surgery Saitama Medical Center at Saitama Medical University, Tokyo, Japan. ⁴National Scoliosis Foundation, Boston, USA. ⁵University of Montreal Ste-Justine Mother and Child University Hospital, Montreal, Canada. ⁶Harvard Medical School, Boston Children Hospital, USA.

¹Department of Orthopedics and Traumatology Tzani General Hospital Tzani and Afendouli 1, Piraeus, Greece
Full list of author information is available at the end of the article

Published: 4 December 2014

doi:10.1186/1748-7161-9-S1-O42

Cite this article as: Grivas *et al.*: The BraIST study and the implications for scoliosis screening: our duty for raising awareness and advocacy. *Scoliosis* 2014 **9**(Suppl 1):O42.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

