



Methionine restriction extends lifespan of *Drosophila melanogaster* under conditions of low amino acid status

Citation

Lee, Byung Cheon, Alaattin Kaya, Siming Ma, Gwansu Kim, Maxim V. Gerashchenko, Sun Hee Yim, Zhen Hu, Lawrence G. Harshman, and Vadim N. Gladyshev. 2014. "Methionine restriction extends lifespan of *Drosophila melanogaster* under conditions of low amino acid status." *Nature communications* 5 (1): 3592. doi:10.1038/ncomms4592. <http://dx.doi.org/10.1038/ncomms4592>.

Published Version

doi:10.1038/ncomms4592

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:14351048>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



Published in final edited form as:

Nat Commun. ; 5: 3592. doi:10.1038/ncomms4592.

Methionine restriction extends lifespan of *Drosophila melanogaster* under conditions of low amino acid status

Byung Cheon Lee¹, Alaattin Kaya¹, Siming Ma¹, Gwansu Kim², Maxim V. Gerashchenko¹, Sun Hee Yim¹, Zhen Hu³, Lawrence G. Harshman³, and Vadim N. Gladyshev^{1,*}

¹Division of Genetics, Department of Medicine, Brigham & Women's Hospital and Harvard Medical School, Boston, MA 02115 USA

²Department of Statistics, Korea University, Seoul, South Korea

³School of Biological Sciences, University of Nebraska-Lincoln, Lincoln, NE, USA

Abstract

Reduced methionine (Met) intake can extend lifespan of rodents, but whether this regimen represents a general strategy for regulating aging has been controversial. Here we report that Met restriction extends lifespan in both fruit flies and yeast, and that this effect requires low amino acid status. Met restriction in *Drosophila* mimicks the effect of dietary restriction and is associated with decreased reproduction. However, under conditions of high amino acid status, Met restriction is ineffective and the trade-off between longevity and reproduction is not observed.

Overexpression of InRDN or Tsc2 inhibits lifespan extension by Met restriction, suggesting the role of TOR signaling in the Met control of longevity. Overall, this study defines the specific roles of Met and amino acid imbalance in aging and suggests that Met restriction is a general strategy for lifespan extension.

INTRODUCTION

Calorie restriction (CR) as a means of extending lifespan has been used extensively in a variety of model organisms, such as yeast, nematodes, fruit flies, rats, and mice [1–5]. Though the underlying mechanisms are not well understood, previous studies revealed the role of nutrient-sensing acting through mTOR, which regulates metabolic processes and affects the rate of aging [6–9]. In spite of numerous studies involving CR, the dietary components responsible for this effect remain poorly characterized in most model systems. In fruit flies, restriction in dietary yeast extended lifespan, whereas a decrease in dietary sugar was ineffective, suggesting that specific component(s) of yeast include factor(s) that

* Address correspondence to Vadim N. Gladyshev, Division of Genetics, Department of Medicine, Brigham & Women's Hospital and Harvard Medical School, New Research Building, Room 435, 77 Avenue Louis Pasteur, Boston, MA 02115 USA. Phone: (617) 525-5122. vgladyshev@rics.bwh.harvard.edu.

Author Contributions

The author(s) have made the following declarations about their contributions: Conceived and designed experiments: BCL, VNG. Performed experiments: BCL, AK, ZH, SHY. Performed statistical analyses: BCL, GK, SM. Analyzed data: BCL, LGH, VNG. Wrote the paper: BCL, VNG.

COMPETING FINANCIAL INTERESTS

The authors declare that no competing financial interests exist.

mediate the effect of dietary restriction [10–12]. It was also reported that methionine (Met) restriction mimics lifespan extension by CR in mice, providing one of the first examples of how a specific dietary factor could function to extend lifespan [13–17].

In previous studies involving CR and protein restriction in fruit flies, either dietary sugar and yeast dilution, or only dilution of dietary yeast, were used. Although these regimens were useful in addressing several critical questions, the identification of nutritional factors involved was not possible because one cannot remove a component from dietary yeast without changing other components [10]. To identify the specific nutrient factors that modulate longevity, several studies used semi-defined or fully defined diets. For example, the use of a semi-defined diet containing sugar and casein substituting for dietary protein revealed that the restriction in casein mimicked amino acid restriction, thereby maximizing lifespan of Canton-S flies [18]. A recent study developed a fully defined diet that could be used to address the role of Met and other compounds in lifespan in fruit flies [19]. However, the initial studies that used this diet found no changes in mean and maximum lifespan of non-virgin females when Met was restricted.

Nutrient imbalance, particularly amino acid imbalance, also received attention with regard to its influence on physiological function and modulation of aging. It was reported that disproportionate levels of amino acids may be associated with the incidence of adverse effects in a rat model. In particular, Met was defined as the most toxic amino acid when used in excess [20]. Restricted use of amino acids was suggested to be responsible for the lifespan extension caused by protein restriction [18]; on the other hand, Met restriction was reported to be ineffective in extending lifespan of *Drosophila* [19,21]. Nevertheless, the roles of Met and its imbalance in aging remain largely uncharacterized, although potentially promising as lifespan extension by Met restriction was demonstrated in both rats and mice. If this effect is conserved across metazoa and beyond, other organisms including a commonly used model organism of aging, fruit flies, should also be subject to lifespan extension by Met restriction. The use of defined diets varying in amino acids may provide an experimental system to examine the consequences of amino acid imbalance and its effect on aging and other physiological functions.

The role of Met in aging was also examined in studies that linked reproduction and longevity. It was reported that amino acid imbalance caused by changes in Met levels differentially affected reproduction and longevity in *Drosophila*. This finding argued against the trade-off theory, wherein reproduction is inversely proportional to longevity [21]. It was unclear, however, how general this finding is, as only some conditions were examined.

In the present study, we used the fully defined diet to characterize the parameters of aging and reproduction in fruit flies and yeast using Met diets varying in the levels of other amino acids. We describe the relationship between dietary Met, egg production, and lifespan, and explain how Met status impacts on lifespan depending on the status of other amino acids and acting through TOR signaling, establishing critical and specific roles of Met in aging.

RESULTS

Sex-specific regulation of lifespan by high Met diet

High Met intake is known to be toxic, presenting adverse effects on feeding, growth, pregnancy rate, and sulfur metabolism in animals [22–26]. Thus, it is important to define the toxic range of dietary Met prior to its use in lifespan studies. In our study, we adopted a fully defined diet [19], so that the levels of Met and its interaction with other amino acids and dietary components could be thoroughly examined. The 1.0X diet corresponding to 15% dietary yeast was supplemented with various amounts of Met and used to analyze lifespan and egg production in Canton-S flies (Fig. 1 and Supplementary Tables 1 and 2). Reduction of Met from 100 mM to 10 mM increased both mean lifespan (21.65%, $p < 0.0001$, Log-Rank test) and reproduction (25.00%, $p = 0.039$, Student's t-test), suggesting that lifespan was increased by lowering Met toxicity. On the other hand, the reduction of Met from 10 mM to 1.5 mM did not affect lifespan ($p = 0.9647$, Log-Rank test) and decreased reproduction (−53.90%, $p < 0.01$, Student's t-test), suggesting that Met toxicity was not involved at lower levels of this amino acid.

To further define the Met toxicity range, we modified the original diet to reduce the levels of all amino acids ten-fold (the 0.1X diet), varied Met (0, 1, and 10 mM) in this diet, and applied these dietary conditions to examine lifespan of Canton-S flies in conjunction with egg production (Fig. 1 and Supplementary Tables 1 and 2). Mean lifespan was increased when Met was decreased from 10 mM to 1 mM (8.94%, $p = 0.0026$, Log-Rank test), and this effect was accompanied by an increase in egg production (20.17%, $p = 0.048$, Student's t-test), suggesting that a decrease in Met toxicity contributes to an increase in lifespan here as well. To further examine the toxic effect of Met, the 0.1X diet containing 0.15, 1, or 10 mM Met was applied to examine lifespan of Canton-S, Oregon-R, and w^{1118} flies under mixed-sex conditions (Fig. 2 and 3 and Supplementary Table 3). Mean lifespan of female flies increased when Met was reduced from 10 to 1 mM (Canton-S: 10.48%, $p = 0.0308$; Oregon-R: 10.98%, $p = 0.0119$; w^{1118} : 23.60%, $p < 0.0001$, Log-Rank test), whereas no difference was observed in males (Canton-S: $p = 0.1239$; Oregon-R: $p = 0.5754$; w^{1118} : $p = 0.8623$, Log-Rank test). Furthermore, no effect on lifespan was seen in both males and females when Met was reduced from 1 mM to 0.15 mM. Thus, the 10 mM Met-0.1X diet was toxic to females only (it decreased both lifespan and reproduction). Our data suggested that the 1 mM Met-0.1X diet (a ten-fold dilution from the original 10 mM-1X diet) was balanced. Interestingly, dietary Met toxicity was observed only in 100 mM Met-1X and 10 mM Met-0.1X diets that clearly represent Met imbalance (high Met levels when the levels of other amino acids are low). On the other hand, lifespan was not changed between the 10 mM Met-1X and 1.5 mM Met-1X diets. These findings show that the imbalance of Met relative to other amino acids (rather than the presence of high Met) is the reason for the reduction in both longevity and reproduction. These findings established the toxic range of Met and showed that its toxicity is limited to females and depends on its interaction with other amino acids.

Lifespan extension by Met and amino acid restriction

It was recently found that neither Met restriction nor Met excess affected the lifespan of mated female flies in the context of normal or CR diets [19,21]. In contrast, metformin-

induced Met restriction extended the lifespan of nematodes [27], and the lifespan could also be extended by Met restriction in mice and rats [14–16]. This discrepancy in the effect of Met restriction in model organisms raised questions regarding evolutionary conservation and general applicability of the Met restriction strategy. To examine this issue, we focused on the interaction of Met with other amino acids. This interaction brings complexity to lifespan studies, which has not been previously characterized, e.g., Met restriction may reduce translation and reproduction thereby increasing lifespan, whereas it may also lead to an accumulation of unused amino acids thereby decreasing lifespan. To define the role of Met in aging as a function of amino acid status, we employed diets with different levels of amino acids (0.4X corresponding to 6% dietary yeast, 0.7X corresponding to 10.5% dietary yeast, and 1.0X corresponding to 15% dietary yeast) and different Met concentrations (0, 0.15, and 1 mM). We subjected mated Canton-S females to these diets and determined their lifespan and egg production (Fig. 4 and Supplementary Tables 4–6).

With regard to the role of amino acid status, we found that lifespan could be extended by amino acid restriction at any Met concentrations (Fig. 4a–f and Supplementary Tables 4–6). For example, when comparing 0.4X and 1.0X diets, mean lifespan increased 22.09% at 0 mM Met ($p < 0.0001$, Log-Rank test), 64.1% at 0.15 mM Met ($p < 0.0001$, Log-Rank test), and 20.17% at 1 mM Met ($p < 0.0001$, Log-Rank test). It was previously reported that a lower intake of casein that substitutes for dietary yeast protein increases fruit fly lifespan, suggesting that amino acid restriction can extend lifespan and that this effect may possibly explain lifespan extension by dietary restriction [18,21]. Our findings also support the idea that amino acid restriction could extend lifespan of mated female flies maintained on the fully defined diet. However, an amino acid restriction that does not affect reproduction, body mass, and total protein (e.g., see the data for 0 mM Met in Fig. 4 and 5 and Supplementary Tables 4–6) could still lead to lifespan extension. Although food consumption increased at low amino acid levels, it could not contribute to the lifespan extension by amino acid restriction due to its negative regulation of aging (Fig. 5). Overall, amino acid restriction extended lifespan at any Met level (including Met deficiency, when it did not affect reproduction).

With regard to the effect of Met status, Met restriction extended lifespan depending on the status of other amino acids (Fig. 4a–c, 4g–i and Supplementary Tables 4–6). For example, when comparing 1.0X and 0.7X diets, Met restriction decreased (or did not change) mean lifespan, whereas on the 0.15 mM Met–0.4X diet the lifespan was 12.53% higher than on the 1 mM Met–0.4X diet ($p < 0.0001$, Log-Rank test). Moreover, Met restriction was accompanied by decreased reproduction, suggesting that the observed lifespan extension was mechanistically different from the effect of decreased Met toxicity. We further analyzed food consumption rate, determining whether Met status influenced eating behavior that in turn affected lifespan (Fig. 5). Interestingly, food consumption was not affected by dietary Met. Thus, lifespan can be extended in fruit flies by Met restriction, as it can in rodents, but this effect requires low amino acid status.

Interplay between lifespan and reproduction

The cost-of-reproduction is an important factor that may affect lifespan as a part of evolutionary trade-offs involving maintenance and reproduction. However, it was reported that under CR conditions the addition of Met did not affect lifespan, yet increased reproduction, suggesting exception from the trade-off theory [21]. To understand this phenomenon, we quantified egg production by Canton-S flies and compared it with mean lifespan of mated female flies in two ways: using three different levels of amino acids at a constant Met concentration (Fig. 4d–f and Supplementary Tables 4–7); and using three Met concentrations at a constant amino acid level (Fig. 4g–i and Supplementary Tables 4–7). As expected, Met status could limit reproduction. In particular, there was no difference in reproduction between 0 mM Met-0.4X and 0 mM Met-1.0X diets ($p=0.828$, Student's t-test) (Fig. 4d and Supplementary Tables 4–6). Moreover, an increase in Met levels correlated with an increase in lifetime egg production on each amino acid diet (Fig. 4g–i and Supplementary Tables 4–6).

We also examined how Met status affected gene expression by generating microarray data for the examined conditions. Gene clustering following stringent filtering with permutation revealed that most genes downregulated by Met restriction in either 0.4X or 1.0X diets were linked to egg production (Fig. 6 and Supplementary Data 1), whereas this was not the case under conditions of amino acid restriction, even when less stringent filtering was applied (Fig. 6b and Supplementary Data 1). These data suggest that Met plays a key regulatory role in reproduction, whereas the gene expression analyses suggested that the status of other amino acids was more relevant to regulation of functions, such as heme binding, iron ion binding, microsome, monooxygenase activity, and oxidoreductase activity (Fig. 6). On the other hand, mean lifespan decreased with the increased supply of amino acids at any Met concentration, whereas reproduction was not changed with the increased supply of amino acids during Met deficiency (Fig. 4d–f and Supplementary Tables 4–6). Thus, the data suggest that the change in reproduction may not underlie lifespan extension by amino acid restriction, resulting in the appearance of exception from the trade-off under conditions of limited Met supply.

When comparing changes in lifespan and reproduction between 0.15 mM and 1 mM Met diets (and the same amino acid status), mean lifespan increased 12.53% on the 0.15 mM Met-0.4X diet ($p<0.0001$, Log-Rank test), decreased 19.44% on the 0.15 mM Met-1X diet ($p<0.0001$, Log-Rank test), and showed no difference on the 0.15 mM Met-0.7X diet ($p=0.9842$, Log-Rank test), whereas reproduction decreased 103.8% on the 0.15 mM Met-0.4X diet, 135.45% on the 0.15 mM Met-0.7X diet, and 135.38% on the 0.15 mM Met-1.0X diet ($p<0.01$, Student's t-test) (Fig. 4g–i and Supplementary Tables 4–7). Accordingly, a decrease in reproduction due to Met deficiency did not coincide with the increase in lifespan under conditions of high amino acid status (e.g., 0.7X and 1.0X diets), whereas a decrease in reproduction due to Met deficiency inversely correlated with the lifespan increase on the low amino acid status (e.g., 0.4X diet; Pearson coefficient: $-0.963 \sim -0.864$, $p<0.0021 \sim 0.0265$) (Supplementary Table 7). Thus, lifespan extension by Met restriction under conditions of low amino acid status was accompanied by decreased

reproduction, suggesting that the trade-off may be observed, at least under certain conditions.

Lifespan extension by Met restriction in virgin flies

We tested if the observed lifespan extension of mated female flies by Met deficiency also applies to flies under other conditions. To examine these possibilities, virgin Canton-S flies were subjected to the same diets with varying Met and amino acid levels (Fig. 7a–d and Supplementary Tables 8 and 9). Indeed, the effects observed with mated female flies were reproduced with the virgin flies. Mean lifespan of virgin flies reared on the 0.15 mM Met-0.4X diet was ~19% higher than on the 1 mM Met-0.4X diet, and lifespan on the 0.15 mM Met-0.7X diet was ~10% higher than on the 1 mM Met-0.7X diet (0.15 mM Met- and 1 mM Met-0.4X; $p<0.0001$, 0.15 mM Met- and 1 mM Met-1.0X; $p=0.0001$, Log-Rank test). On the other hand, Met reduction from 1 mM to 0.15 mM on the 1.0X diets did not increase lifespan (-8.31%, $p=0.0085$, Log-Rank test). This data supports the finding that the lifespan extension by Met restriction is maximized at low levels of other amino acids.

Extension of yeast lifespan by Met restriction

We also examined another common model organism of aging, the budding yeast. Yeast replicative lifespan was assayed in cells grown on the medium containing various Met levels (0.1, 0.5, 0.05, 0.005, or 1 mg/ml) (Fig. 7e–h and Supplementary Tables 10 and 11). Two lowest Met levels (0.05 and 0.005 mg/ml Met) were most effective in extending yeast lifespan, suggesting that the findings in fruit flies may apply to yeast. Based on these results, we further changed the amino acid status for 0.5 and 0.05 mg/ml Met conditions and observed a significant increase (4.4 generations; $p=0.004$, Rank-sum test) in mean lifespan upon reducing Met. Met decrease was also effective in extending lifespan in other media (0.2X, 1X, and 5X), but the low amino acid status was the most effective. Thus, low amino acid status maximizes the lifespan extension by Met restriction in yeast, suggesting both a general applicability of Met restriction in extending lifespan and the dependence of this effect on low amino acid status.

Lifespan control by Met restriction requires the TOR pathway

It was reported that the effect of dietary restriction on lifespan was due to inhibition of TOR signaling [7–9]. We reasoned that the lifespan extension by Met restriction may also involve TOR inhibition. To test this possibility, we used two transgenic lines of fruit flies containing an upstream activating sequence linked to a dominant negative form of *Drosophila* insulin receptor (*UAS-dInRDN*) or to *Drosophila* Tsc2 (*UAS-dTsc2*). Mated females that overexpress either InRDN or Tsc2 were subjected to four diets (1 mM Met-1X, 0.2 mM Met-1X, 1mM Met-0.4X, 0.2 mM Met-0.4X) and their lifespan was analyzed (Fig. 8 and Supplementary Tables 12 and 13). We found that mean lifespan of control flies (+/*UAS-dInRDN* and +/*UAS-dTsc2*) was differentially regulated by Met restriction upon changes in amino acid status (Fig. 8a–h and Supplementary Tables 12 and 13). For example, Met decrease from 1 mM to 0.2 mM in the 1.0X diet decreased lifespan in both control lines (+/*UAS-dInRDN*, -12.82%, $p<0.0001$; +/*UAS-dTsc2*, -17.85%, $p<0.0001$, Log-Rank test), whereas it extended lifespan when flies were on the 0.4X diet (+/*UAS-dInRDN*, 15.86%,

$p < 0.0001$; $+/UAS-dTsc2$, 4.03%, $p = 0.01$, Log-Rank test). Interestingly, Met decrease from 1 mM to 0.2 mM on the 0.4X diet did not extend lifespan of *da-Gal4>UAS-dInRDN* (-2.88%, $p = 0.10$, Log-Rank test) and *da-Gal4>UAS-dTsc2* (-21.88%, $p < 0.0001$, Log-Rank test) lines (Fig. 8e–h and Supplementary Tables 12 and 13). These data suggest that overexpression of InRDN or Tsc2, which inactivate TOR signaling, inhibited lifespan extension by Met restriction. Thus, similar to the CR case the lifespan extension by Met restriction under conditions of low amino acids utilizes the inhibition of TOR signaling.

As shown above, an imbalance in amino acids suppressed the lifespan extension effect of Met restriction (Fig. 4 and Supplementary Tables 4–6). If Met restriction inhibits TOR signaling, the high amino acid status may affect TOR signaling, blocking the lifespan extension by Met restriction. Surprisingly, overexpression of InRDN or Tsc2 did not extend lifespan on 0.2 mM- and 1 mM-1.0X diets (*da-Gal4>UAS-dInRDN* on the 0.2 mM Met-1.0X, -7.25%, $p = 0.07$; *da-Gal4>UAS-dInRDN* on the 1 mM Met-1.0X, -3.11%, $p = 0.98$; *da-Gal4>UAS-dTsc2* on the 0.2 mM Met-1.0X, -7.65%, $p = 0.02$; *da-Gal4>UAS-dTsc2* on the 1 mM Met-1.0X, -6.98%, $p = 0.15$, Log-Rank test) (Fig. 8e–h and Supplementary Tables 12 and 13). On the other hand, overexpression of InRDN or Tsc2 led to lifespan extension on the 1 mM Met-0.4X diet (*da-Gal4>UAS-InRDN* on the 1 mM Met-0.4X, 15.14%, $p < 0.0001$; *da-Gal4>UAS-Tsc2* on the 1 mM Met-0.4X, 18.65%, $p < 0.0001$, Log-Rank test), whereas no lifespan extension was observed on the 0.2 mM Met-0.4X (*da-Gal4>UAS-dInRDN* on the 0.2 mM Met-0.4X, -3.49%, $p = 0.07$; *da-Gal4>UAS-dTsc2* on the 0.2 mM Met-0.4X, -10.90%, $p < 0.0001$, Log-Rank test) (Fig. 8e–h and Supplementary Tables 12 and 13). This data indicates that delaying the aging process by inhibiting TOR signaling depends on the interaction of Met with other amino acids. Thus, the reason Met restriction was not effective in increasing lifespan under conditions of high amino acids is due interference of the effects of Met restriction, TOR inhibition and amino acid imbalance.

DISCUSSION

It is thought that nutritional deficiency decreases fecundity and increases lifespan through redistribution of nutrient resources, and that food supplies restore egg production leading to a reduction in lifespan [28]. However, a recent study reported that the addition of Met under conditions of dietary dilution increased reproduction without affecting lifespan [21]. Our results involving fully defined diets explain this unexpected finding and relate it to both the trade-off idea and the still controversial role of Met restriction in regulating lifespan. Indeed, Met deficiency on most diets with defined and variable amino acid status (0.7X and 1.0X) reduced reproduction, but it was not associated with lifespan extension. On the other hand, we found that Met deficiency can lead to a decrease, increase, or no change in lifespan depending on the status of other dietary amino acids. Excess amino acids may lead to ineffectiveness of nutritional resource redistribution under conditions of Met restriction, whereby a decrease in reproduction may not necessarily lead to lifespan extension. Defining the mechanisms linking cost-of-reproduction to lifespan has been a long-held question in the aging field. As such, the observed diversity of responses that characterize the trade-off and involve the interplay between Met and other amino acids offers important mechanistic insights with regard to regulation of the trade-off and the aging process.

By varying both total amino acid levels and the levels of Met, we demonstrated that the lifespan extension by Met restriction requires low amino acid status, and this finding was also held in yeast. Together with the previous data in rodents, our data suggest that Met restriction may be viewed as a general strategy to extend lifespan. The data further suggest that the lifespan regulation by Met restriction involves TOR signaling and that high levels of amino acids suppress the lifespan extension conferred by Met restriction or overexpression of InRDN or Tsc2. TORC1 (TOR complex 1) is a sensor of amino acids [29–32], so it is not surprising that accumulation of amino acids activates TOR. However, a question remains regarding the mechanism of the differential inhibition of the Met effect upon changes in amino acid status. We suggest that free amino acids may activate TOR, fully or partially, independent of Tsc1-Tsc2, while Met acts on TOR through the Tsc1-Tsc2 complex (Fig. 8i). Recently, the RagA/B-RagC/D GTPase complex was found to tether Rags to the lysosome in response to amino acid sensing by mTORC1 and couple this signaling to proteins downstream of mTORC1. This signal transduction for activation of TORC1, mediated by sensing amino acids, is independent of the Tsc1-Tsc2 complex [30,31]. Hence, it is possible that saturated activation of TORC1 by amino acid accumulation may occur independently of Tsc1-Tsc2 and not be reversed by InRDN/Tsc2 overexpression or Met restriction. However, because Met is also an amino acid, it, when at low levels, may inhibit TORC1 independent of Tsc1-Tsc2. We hypothesize that the lifespan extension by Met restriction may involve other mechanisms for the following reasons: First, reduced reproduction and protein synthesis in response to Met restriction may lead to an accumulation of other amino acids, and thus free amino acids may be increased by Met restriction. This suggests that Met restriction may not use the same mechanism of amino acid restriction for lifespan extension. Second, Met is a key regulator of reproduction and protein synthesis. Thus, Met may be involved in other regulatory functions, such as control of energy and stress. Third, TORC1 can transmit energy and stress signals sensed by the Tsc1-Tsc2 complex. Fourth, microarray data show that Met restriction-induced genes are involved in the response to toxins, lipid metabolism, and steroid metabolism, all of which are relevant to regulation of energy and stress (Fig. 6c). Although this hypothesis needs further testing, there is increasing evidence that the mechanism of Met restriction may be independent of general amino acid sensing by the TORC1 complex (Fig. 8i).

METHODS

Fly stocks and husbandry

D. melanogaster (Canton-S, Oregon-R, and w^{1118} strain) lines were from the laboratory of Dr. Lawrence G. Harshman (University of Nebraska, Lincoln, NE). *UAS-dTsc2*, *UAS-dInRDN*, w^{1118} , and *da-GAL4* strains were from the laboratory of Dr. Pankaj Kapahi (Buck Institute for Research on Aging, Novato, CA) [9]. Flies were maintained on corn meal food, commonly used in our laboratory, and kept in a temperature-controlled chamber at 25 °C with 12 h light/dark cycle and approximately 60% humidity for all experiments. Newly emerged flies were transferred to fresh corn meal food and allowed to mate for 1–2 days. Virgin flies were collected at 18 °C within 16 h after eclosion and then kept on corn meal food until lifespan analyses. Three day old mated flies were collected using etherization, sorted by sex and then some of them transferred to cages on the designed diet for lifespan

study. Three day old virgin flies were also transferred to cages on the designed diet and their lifespan was analyzed. Unless otherwise stated, experimental flies were held on the designed diet and transferred to fresh vials without anesthesia every three days.

Fly culture media

In order to maintain strains and collect flies for lifespan studies, we used corn meal food (85.7 g corn meal “Aunt Jemina” (The Quaker Oats Company, Chicago, IL), 50 ml golden A unsulfured molasses (Groeb Farms Inc, Onsted, MI), 71.4 g Torula yeast (MP Biomedicals, Solon, OH), 2.86 g p-hydroxybenzoic acid methyl ester (Sigma), 6.4 g agar (MoorAgar Inc, Loomis, CA) and 5.7 ml propionic acid (Sigma) per liter water). In order to test lifespan of flies, fully defined medium was used based on the fruit fly basal mix developed in a recent study [19] with modification in amino acid content. This diet contained 62.08 g Diet TD. 04310 (corresponds to the 0.1X diet) and/or 101.07g Diet TD.10417 (corresponds to the 1.0X diet) (Harlan Teklad), 100 mg lecitin from soybean (Sigma), 500 mg ribonucleic acid from Torula yeast (Sigma), 100 g of dextrose, 20 g agar, 2.85 ml propionic acid, 0.255 ml of phosphoric acid (Sigma), and indicated amounts of Met (Sigma) per liter of water. To prepare 0.4X and 0.7X diets, the two basal mixes were mixed to adjust amino acid concentrations.

Lifespan and egg production analyses

Newly hatched flies were collected within 18 h at 18 °C and kept on corn meal food at density of 35 animals per vial. To prepare mated female flies, flies were incubated for 24 h under the mixed sex conditions. In a typical lifespan trial, three day old flies were placed in cages. The cages were made of plastic 1 L cups with grommets for changing food vials and slits for removal of dead flies by aspiration. 35–70 males and/or 35–70 females were kept in each cage for testing lifespan on the fully defined medium. 3–6 replica cages were used for each lifespan. Fresh food was supplied into cages and dead flies were removed by aspiration and counted every 3 days. The control and experimental group trials were always performed concurrently. Flies were maintained in a temperature-controlled incubator at 25 °C with 12 h light/dark cycle and 50–60% humidity.

Flies were also prepared for the reproduction assay following the outline of the aging experiment. Then, egg production by animals maintained on diets containing four different levels of amino acids (0.1X, 0.4X, 0.7X, 1.0X) with varying Met (0, 0.15, 1, 1.5, 10, 100 mM) was counted, for eggs produced within 24 h, every three days until animals stopped laying eggs. Each vial contained 5 females and 5 males, and 4–5 replicate vials on each diet were used to count egg production. Finally, the data were represented as cumulative egg production for 30 days as well as lifetime egg production.

Yeast aging replicative lifespan assays

Wild-type strain BY4741 (*MATa his3 leu2 met15 ura3*) was used in replicative aging assays. Nitrogen based Synthetic Defined Yeast Media (SD media) including an amino acid mix lacking Met was used to assay lifespan under conditions of Met restriction. Met varied from 0.005 mg/ml to 1 mg/ml, whereas all other components were kept constant. To analyze how amino acid status affects lifespan, nitrogen base dropout media (YNB media) and an amino

acid supplement mix without Met (CSM-Met) (MP Biomedicals) were mixed prior to plate preparation. YNB concentration was 6.7 g/l and the concentration of amino acid mix without Met varied from 0.2X to 5X. Met was added as indicated concentrations. Since our wild-type strain was an auxotroph for uracil, histidine and leucine, we also added these amino acids separately and kept them constant. For replicative aging assays, cells were grown on fresh medium overnight. For each strain, 30 daughter cells (starter mothers) were collected and assayed on agar plates. New buds (daughters) from these starter mothers were removed and discarded as they formed. This process continued until cells stopped dividing. Lifespan was determined as the total number of daughter cells that each mother cell generated.

Food consumption

Mated female flies were prepared following the outline of the aging experiment and reared on diets with different amino acid levels (0.1X, 0.4X, 0.7X, 1.0X) and varying in Met (0, 0.15, 1, 10 mM). For the feeding assay, 0.1% (w/v) erioglaucine (FD&C Blue #1, Sigma) was added to each defined diet, and 6-day old and 21-day old flies were fed these diets for 6 h or 30 min (12 noon – 6 pm or 12 noon – 12:30 pm) [33–35]. Five female flies per group (20 groups were used for 30 min incubation) were homogenized in 200 µl of PBS, and the homogenate was centrifuged at 13,000 rpm for 15 min. Absorbance of the supernatant was measured at 625 nm, with 675 nm used as a reference, and then compared with the absorbance of the control food, prepared in the same way, to calculate consumed food mass.

Analysis of total protein and Met concentration

21-day old mated female flies were prepared following the outline of the aging experiment on diets with three different levels of amino acids (0.4X, 0.7X, 1.0X) and varying Met (0, 0.15, 1, 10 mM). 25 flies per group were used to determine weight. These flies were then homogenized in PBS in order to measure total protein by using the BCA protein assay reagent (Pierce).

To measure endogenous Met concentration in mated female flies, we followed a previously established procedure [36,37]. Briefly, homogenized samples were precipitated by mixing with 5% trichloroacetic acid. After incubation at 4 °C for 10 min and centrifugation at 13,000 rpm for 15 min, the supernatant (50 µl) was diluted 10 fold with distilled water and prepared for OPA derivatization. The derivatization reagent was freshly prepared. Sample solutions (2–5 µl) were mixed with the OPA derivatization reagent to 100 µl final volume. Following a 2 min reaction at room temperature, 50 µl of the mixture were injected onto a Zorbax Eclipse XDB-C18 column (4.6×150 mm) and analyzed by HPLC. Detection was by fluorescence of Met derivatives using a Waters 474 scanning fluorescence detector with excitation at 330 nm and emission at 445 nm.

Fly samples for microarray analysis

Newly hatched flies were collected within 18 h at 18 °C and kept on corn meal food at density of 35 animals per vial. To prepare mated female flies, collected flies were incubated for 24 h under the mixed sex conditions. Three day old flies were placed in vials. 20 females were kept in each vial on the fully defined medium for 18 days. 10 replica vials were used for sample collection. Fresh food was supplied every 3 days and dead flies were removed by

aspiration. Flies were maintained in a temperature-controlled incubator at 25 °C with 12 h light/dark cycle and 50–60% humidity.

Mated female Canton-S flies raised on four different defined media were subjected to RNA extraction. Total RNA was extracted from 20 flies by Trizol (Invitrogen), according to manufacturer's suggestions. Extracted RNA was additionally purified by either MEGAclear of RNAqueous kit (Ambion) to ensure highest purity for microarrays. Microarray analysis was performed by using *Drosophila* Genome 2.0 at the microarray core of *Dana-Farber Cancer Institute*. Fluorescence intensity of each transcript was normalized. Expression data from Affymetrix microarrays were analyzed with dCHIP software with default settings [38,39]. Changes in transcript levels were compared among four sample sets of RNA. To examine changes in expression, the data were filtered with 50 permutation testing to assess False Discovery Rate (FDR) as follows: 1) $E/B > 1.2$ or $B/E > 1.2$ and 2) $E-B > 100$ or $B-E > 100$ (E: Experiment, B: Baseline) or for more stringent comparison: 1) $E/B > 1.8$ (or 1.5) or $B/E > 1.8$ (or 1.5) and 2) $E-B > 100$ or $B-E > 100$ (E: Experiment, B: Baseline).

Statistics

Statistical analyses were performed using JMP (version 10) software (SAS Institute Inc.). Log-rank tests were used for *Drosophila* survival analysis. Wilcoxon rank-sum test were used for the analyses of yeast replicative lifespan. Comparisons of means were made using Student's *t*-test.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by the National Institutes of Health grants AG021518 and AG038004 to VNG. We thank Dr. Pankaj Kapahi (Buck Institute for Research on Aging, Novato, CA) for providing transgenic flies.

References

1. Fernandes G, Yunis EJ, Good RA. Influence of diet on survival of mice. *Proc Natl Acad Sci U S A*. 1976; 73:1279–1283. [PubMed: 1063408]
2. Bosshardt DK, Paul WJ. Caloric restriction and protein metabolism in the growing mouse. *J Nutr*. 1948; 36:773–783. [PubMed: 18100940]
3. Koubova J, Guarente L. How does calorie restriction work? *Genes Dev*. 2003; 17:313–321. [PubMed: 12569120]
4. Lakowski B, Hekimi S. The genetics of caloric restriction in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*. 1998; 95:13091–13096. [PubMed: 9789046]
5. Lin SJ, et al. Calorie restriction extends *Saccharomyces cerevisiae* lifespan by increasing respiration. *Nature*. 2002; 418:344–348. [PubMed: 12124627]
6. Honjoh S, Yamamoto T, Uno M, Nishida E. Signalling through RHEB-1 mediates intermittent fasting-induced longevity in *C. elegans*. *Nature*. 2009; 457:726–730. [PubMed: 19079239]
7. Kaeberlein M, Kapahi P. Cell signaling. Aging is RSKy business. *Science*. 2009; 326:55–56. [PubMed: 19797648]
8. Kaeberlein M, et al. Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science*. 2005; 310:1193–1196. [PubMed: 16293764]

9. Kapahi P, et al. Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr Biol*. 2004; 14:885–890. [PubMed: 15186745]
10. Piper MD, Partridge L. Dietary restriction in *Drosophila*: delayed aging or experimental artefact? *PLoS Genet*. 2007; 3:e57. [PubMed: 17465680]
11. Mair W, Piper MD, Partridge L. Calories do not explain extension of life span by dietary restriction in *Drosophila*. *PLoS Biol*. 2005; 3:e223. [PubMed: 16000018]
12. Lee KP, et al. Lifespan and reproduction in *Drosophila*: New insights from nutritional geometry. *Proc Natl Acad Sci U S A*. 2008; 105:2498–2503. [PubMed: 18268352]
13. Lopez-Torres M, Barja G. Lowered methionine ingestion as responsible for the decrease in rodent mitochondrial oxidative stress in protein and dietary restriction possible implications for humans. *Biochim Biophys Acta*. 2008; 1780:1337–1347. [PubMed: 18252204]
14. Orentreich N, Matias JR, DeFelice A, Zimmerman JA. Low methionine ingestion by rats extends life span. *J Nutr*. 1993; 123:269–274. [PubMed: 8429371]
15. Richie JP Jr, et al. Methionine restriction increases blood glutathione and longevity in F344 rats. *FASEB J*. 1994; 8:1302–1307. [PubMed: 8001743]
16. Sun L, Sadighi Akha AA, Miller RA, Harper JM. Life-span extension in mice by preweaning food restriction and by methionine restriction in middle age. *J Gerontol A Biol Sci Med Sci*. 2009; 64:711–722. [PubMed: 19414512]
17. Zimmerman JA, Malloy V, Krajcik R, Orentreich N. Nutritional control of aging. *Exp Gerontol*. 2003; 38:47–52. [PubMed: 12543260]
18. Min KJ, Tatar M. Restriction of amino acids extends lifespan in *Drosophila melanogaster*. *Mech Ageing Dev*. 2006; 127:643–646. [PubMed: 16616772]
19. Troen AM, et al. Lifespan modification by glucose and methionine in *Drosophila melanogaster* fed a chemically defined diet. *Age (Dordr)*. 2007; 29:29–39. [PubMed: 19424828]
20. Harper AE, Benevenga NJ, Wohlueter RM. Effects of ingestion of disproportionate amounts of amino acids. *Physiol Rev*. 1970; 50:428–558. [PubMed: 4912906]
21. Grandison RC, Piper MD, Partridge L. Amino-acid imbalance explains extension of lifespan by dietary restriction in *Drosophila*. *Nature*. 2009; 462:1061–1064. [PubMed: 19956092]
22. Benevenga NJ. Toxicities of methionine and other amino acids. *J Agric Food Chem*. 1974; 22:2–9. [PubMed: 4590578]
23. Abe M, et al. Methionine imbalance and toxicity in calves. *J Anim Sci*. 2000; 78:2722–2730. [PubMed: 11048939]
24. Muramatsu K, Odagiri H, Morishita S, Takeuchi H. Effect of excess levels of individual amino acids on growth of rats fed casein diets. *J Nutr*. 1971; 101:1117–1125. [PubMed: 5096135]
25. Dioguardi FS. Clinical use of amino acids as dietary supplement: pros and cons. *J Cachex Sarcopenia Muscle*. 2011; 2:75–80.
26. Bass TM, et al. Optimization of dietary restriction protocols in *Drosophila*. *J Gerontol A Biol Sci Med Sci*. 2007; 62:1071–1081. [PubMed: 17921418]
27. Cabreiro F, et al. Metformin retards aging in *C. elegans* by altering microbial folate and methionine metabolism. *Cell*. 2013; 153:228–239. [PubMed: 23540700]
28. Partridge L, Gems D, Withers DJ. Sex and death: what is the connection? *Cell*. 2005; 120:461–472. [PubMed: 15734679]
29. Zoncu R, et al. mTORC1 senses lysosomal amino acids through an inside-out mechanism that requires the vacuolar H(+)-ATPase. *Science*. 2011; 334:678–683. [PubMed: 22053050]
30. Sengupta S, Peterson TR, Sabatini DM. Regulation of the mTOR complex 1 pathway by nutrients, growth factors, and stress. *Mol Cell*. 2010; 40:310–322. [PubMed: 20965424]
31. Kim J, Guan KL. Amino acid signaling in TOR activation. *Annu Rev Biochem*. 2011; 80:1001–1032. [PubMed: 21548787]
32. Kim SG, Buel GR, Blenis J. Nutrient regulation of the mTOR Complex 1 signaling pathway. *Mol Cells*. 2013; 35:463–473. [PubMed: 23694989]
33. Skorupa DA, Dervisevendic A, Zwiener J, Pletcher SD. Dietary composition specifies consumption, obesity, and lifespan in *Drosophila melanogaster*. *Aging Cell*. 2008; 7:478–490. [PubMed: 18485125]

34. Libert S, et al. Regulation of *Drosophila* life span by olfaction and food-derived odors. *Science*. 2007; 315:1133–1137. [PubMed: 17272684]
35. Wong R, Piper MD, Wertheim B, Partridge L. Quantification of food intake in *Drosophila*. *PLoS One*. 2009; 4:e6063. [PubMed: 19557170]
36. Lee BC, Le DT, Gladyshev VN. Mammals reduce methionine-S-sulfoxide with MsrA and are unable to reduce methionine-R-sulfoxide, and this function can be restored with a yeast reductase. *J Biol Chem*. 2008; 283:28361–28369. [PubMed: 18697736]
37. Shchedrina VA, et al. Overexpression of methionine-R-sulfoxide reductases has no influence on fruit fly aging. *Mech Ageing Dev*. 2009; 130:429–443. [PubMed: 19409408]
38. Armstrong SA, et al. MLL translocations specify a distinct gene expression profile that distinguishes a unique leukemia. *Nat Genet*. 2002; 30:41–47. [PubMed: 11731795]
39. Li C. Automating dChip: toward reproducible sharing of microarray data analysis. *BMC Bioinformatics*. 2008; 9:231. [PubMed: 18466620]




Figure 1.

Imbalance of amino acids and high levels of Met decrease lifespan and reproduction. a) Survivorship curve, mean lifespan, and lifetime eggs production of mated female Canton-S flies reared on the 0.1X or 1.0X amino acids diets containing 0, 1, 1.5, 10, or 100 mM Met were analyzed. Bar graphs show lifetime egg production (left axis) (mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test) and line graphs mean lifespan (right axis) (mean \pm s.e.m., $n = 144 \sim 205$, ***: $p < 0.0001$, Log-rank test). See Supplementary Tables 1 and 2 for n number and statistics of individual dietary groups. b) Cumulative eggs production of mated female Canton-S flies reared on the 0.1X diet containing 0, 0.15, 1, or 10 mM Met for 30 days (mean \pm s.d., $n = 40$). c) Cumulative eggs production of mated female Canton-S flies reared on the 1.0X diet containing 0, 1.5, 10, or 100 mM Met for 30 days (mean \pm s.d., $n = 30$).




Figure 2.

Survivorship curves of fruit flies reared on the 0.1X diet varying in Met levels. a) Female Canton-S flies, b) male Canton-S flies, c) female Oregon-R flies, d) male Oregon-R flies, e) female w^{1118} flies and f) male w^{1118} flies were reared on the 0.1X diet containing 0 mM, 0.15 mM, 1 mM, or 10 mM Met are shown in the survivorship curves.






Figure 3.

Sex-specific decrease of lifespan by imbalance of amino acids and high levels of Met. a–c) Female and male Canton-S, Oregon-R, and *w¹¹¹⁸* flies reared on the 0.1X diet containing 1 or 10 mM Met. a) Canton-S (mean±s.e.m., n = 184 ~ 196), b) Oregon-R (mean±s.e.m., n = 198 ~ 206), and c) *w¹¹¹⁸* (mean±s.e.m., n = 81 ~ 94) flies were analyzed for mean lifespan and examined for differences among dietary groups (***: $p < 0.0001$, Log-rank test). d–f) Female and male Canton-S, Oregon-R, and *w¹¹¹⁸* flies reared on the 0.1X diet containing 0.15 or 1 mM Met. d) Canton-S (mean±s.e.m., n = 187 ~ 196), e) Oregon-R (mean±s.e.m., n = 198 ~ 203), and f) *w¹¹¹⁸* (mean±s.e.m., n = 91 ~ 99) flies were analyzed for mean lifespan and analyzed for differences among dietary groups (***: $p < 0.0001$, Log-rank test). See Supplementary Table 3 for n number and statistics of individual strains and dietary group.


**Figure 4.**

Lifespan extension of mated female flies by Met restriction and low amino acids. a–c) Survivorship curves of mated female Canton-S flies reared on a) 1.0X diet containing 0, 0.15, or 1 mM Met, b) 0.7X diet containing 0, 0.15, or 1 mM Met, or c) 0.4X diet containing 0, 0.15, or 1 mM Met, are shown. Mean lifespan and lifetime eggs production of mated female Canton-S flies on d) 0.4X, 0.7X, or 1.0X diets containing 0 mM Met (mean lifespan: mean \pm s.e.m., $n = 134 \sim 151$, ***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), e) 0.4X, 0.7X, or 1.0X diets containing 0.15 mM Met (mean lifespan: mean \pm s.e.m., $n = 123 \sim 150$, ***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), f) 0.4X, 0.7X, or 1.0X diets containing 1 mM Met (mean lifespan: mean \pm s.e.m., $n = 144 \sim 150$, ***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), g) 0.4X diet containing 0, 0.15, or 1 mM Met (mean lifespan: mean \pm s.e.m., $n = 123 \sim 150$, ***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), h) 0.7X diet containing 0, 0.15, or 1 mM Met (mean lifespan: mean \pm s.e.m., $n = 150 \sim 151$, ***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), i) 1.0X diet containing 0, 0.15, or 1 mM Met (mean lifespan: mean \pm s.e.m., $n = 134 \sim 144$,


***: $p < 0.0001$, Log-rank test) (Lifetime eggs production: mean \pm s.d., $n = 40$, *: $p < 0.05$, **: $p < 0.01$, Student's t -test), are shown. Bar graphs show lifetime egg production (right axis) and line graphs mean lifespan (left axis). See Supplementary Tables 4–6 for n number and statistics of individual dietary groups.

**Figure 5.**


Analysis of eating behavior and endogenous Met concentration of mated female flies. a–e) Flies were reared on 0.4X, 0.7X, or 1.0X diets containing 0, 0.15, or 1 mM Met, and then collected on 6th or 21st days. a) Food consumption for 30 min on the 6th day (mean±s.d., n = 100) and b) food consumption for 6 hours on the 6th and 21st days (mean±s.d., n = 20) were analyzed via a feeding assay using erioglaucone. c) Body mass (per fly) (mean±s.d., n = 75), d) Total protein (per fly) (mean±s.d., n = 75), e) Amounts of endogenous Met normalized to body mass (mean±s.d., n = 75) were analyzed on the 21st day. All experiments were repeated three to ten times independently and statistics was assessed by using Student's t-test (*: $p < 0.05$ and **: $p < 0.01$).

**Figure 6.**

Gene clusters whose expression is regulated by Met restriction and amino acid levels. a–c) Mated female Canton-S flies ($n = 30$) were raised on the 0.4X amino acid diet containing 0.15 mM or 1 mM Met, or the 1X amino acid diet containing 0.15 mM or 1 mM Met for 18 days, and gene expression was examined by microarray analyses. a) Gene clusters, whose expression was reduced by Met restriction in both amino acid diets, and b) decreased by low levels of amino acids in both Met diets are shown. c) Gene clusters that were upregulated by Met restriction in both amino acid diets are shown.

**Figure 7.**

Lifespan extension of virgin female flies and yeast by Met restriction. a–c) Survivorship curves of virgin Canton-S flies reared on a) 0.4X diet containing 0, 0.15, or 1 mM Met, b) 0.7X diet containing 0, 0.15, or 1 mM Met, or c) 1.0X diet containing 0, 0.15, or 1 mM Met, are shown. d) Mean lifespan of virgin flies on these diets was compared (mean±s.e.m., $n = 92 \sim 111$, ***: $p < 0.0001$, Log-rank test). See Supplementary Tables 8 and 9 for n number and statistics of individual dietary groups. e) Survivorship curves and f) mean lifespan of yeast grown on defined medium containing 1, 0.1, 0.5, 0.05, or 0.005 mg/ml Met (mean ±s.e.m., $n = 30$, **: $p < 0.01$, Rank-sum test). g) Survivorship curve and h) mean lifespan of yeast grown on defined media containing 0.5 or 0.05 mg/ml Met under conditions of 0.2X or 1.0X amino acid status (mean±s.e.m., $n = 30$, **: $p < 0.01$, Rank-sum test). See Supplementary Tables 10 and 11 for statistics of individual dietary groups.

**Figure 8.**

Role of TOR signaling in lifespan extension by Met restriction. a–d) Survivorship curves of *da-GAL4>dInRDN* and *+>dInRDN* reared on a) 1.0X diet containing 0.2 or 1 mM Met, b) 0.4X diet containing 0.2 or 1 mM Met. Survivorship curves of *da-GAL4>dTsc2* and *+>dTsc2* reared on c) 1.0X diet containing 0.2 or 1 mM Met, d) 0.4X diet containing 0.2 or 1 mM Met. 0.2 and 1 correspond to 0.2 mM Met and 1 mM Met, respectively. e–f) Mean lifespan of *da-GAL4>dInRDN* and *+>dInRDN* reared on e) 1.0X diet containing 0.2 or 1 mM Met (mean±s.e.m., n = 81 ~ 106, ***: $p < 0.0001$, Log-rank test), f) 0.4X diet containing 0.2 or 1 mM Met (mean±s.e.m., n = 88 ~ 106, ***: $p < 0.0001$, Log-rank test). g–h) Mean lifespan of *da-GAL4>dTsc2* and *+>dTsc2* reared on g) 1.0X diet containing 0.2 or 1 mM Met (mean±s.e.m., n = 104 ~ 149, ***: $p < 0.0001$, Log-rank test), h) 0.4X diet containing 0.2 or 1 mM Met (mean±s.e.m., n = 111 ~ 152, ***: $p < 0.0001$, Log-rank test). See Supplementary Tables 12 and 13 for n number and statistics of individual strains and dietary groups. i) Model of TORC1 activation by Met and amino acids.