



Clinical presentation and markers of cutaneous versus systemic mastocytosis: the mastocytosis center at brigham and women's hospital in Boston

Citation

Kanamori, Kelly Yoshimi, Pedro Giavina-Bianchi, Nathalia Pessoa Simis, Cem Akin, and Mariana Castells. 2015. "Clinical presentation and markers of cutaneous versus systemic mastocytosis: the mastocytosis center at brigham and women's hospital in Boston." *The World Allergy Organization Journal* 8 (Suppl 1): A141. doi:10.1186/1939-4551-8-S1-A141. <http://dx.doi.org/10.1186/1939-4551-8-S1-A141>.

Published Version

[doi:10.1186/1939-4551-8-S1-A141](http://dx.doi.org/10.1186/1939-4551-8-S1-A141)

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:15034967>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

MEETING ABSTRACT

Open Access

Clinical presentation and markers of cutaneous versus systemic mastocytosis: the mastocytosis center at brigham and women's hospital in Boston

Kelly Yoshimi Kanamori^{1*}, Pedro Giavina-Bianchi¹, Nathalia Pessoa Simis¹, Cem Akin^{2,3}, Mariana Castells^{2,3}

From 3rd WAO International Scientific Conference (WISC) 2014
Rio de Janeiro, Brazil. 6-9 December 2014

Background

Mastocytosis is a group of heterogeneous diseases characterized by an abnormal expansion and accumulation of mast cells (MCs) in different tissues. While systemic mastocytosis (SM) is defined by the accumulation of MCs in bone marrow (BM) and different tissues, cutaneous mastocytosis (CM) is characterized as an accumulation of MC in the skin with no other organ involvement. How gender differences, tryptase levels and episodes of anaphylaxis affect cutaneous versus systemic mastocytosis is not well understood.

Methods

We report a case series of 109 patients with SM and 59 with CM from the Mastocytosis Center at Brigham and Women's Hospital in Boston. The diagnosis of SM and CM were done according to the International Classification of Mastocytosis from WHO. We reviewed gender, tryptase levels and episodes of anaphylaxis in both populations.

Results

There was a female predominance which was more pronounced in SM (68,8%) than in CM (52,7%). The mean serum tryptase level was 11.5 ng/ml in CM, and 14.9% of CM patients had serum tryptase levels above 20 ng/ml. In SM patients the mean serum tryptase level was 97.2 ng/ml and 84% of the patients presented levels above 20 ng/ml. The symptoms of CM included pruritus, flushing, urticaria, and dermatographism and 10.8% of

the patients (number) had episodes of anaphylaxis. The rate of anaphylaxis in SM patients was found to be 29.4% similar to the previously reported.

Conclusions

A female predominance was seen in both cutaneous and systemic mastocytosis in contrast to other studies indicating that there is no gender predominance. Elevations of serum tryptase above 20 ng/ml and anaphylaxis were seen in a small proportion of patients with cutaneous mastocytosis, raising the question of the potential progression to systemic mastocytosis and the need to closely monitor this subset of patients.

Authors' details

¹University of São Paulo, Brazil. ²Harvard Medical School, USA. ³Brigham & Womens Hospital, USA.

Published: 8 April 2015

doi:10.1186/1939-4551-8-S1-A141

Cite this article as: Kanamori et al.: Clinical presentation and markers of cutaneous versus systemic mastocytosis: the mastocytosis center at brigham and women's hospital in Boston. *World Allergy Organization Journal* 2015 8(Suppl 1):A141.

¹University of São Paulo, Brazil

Full list of author information is available at the end of the article