



# Asthma features in elderly in a tertiary outpatients facility

## Citation

De Castro, Nathalia Siqueira Robert, Carolina Tavares De Alcântara, Kelly Yoshimi Kanamori, Jorge Kalil, Bruna Gama Saliba, Rosana Câmara Agondi, Pedro Giavina-Bianchi, Marcelo Vivolo Aun, and Carla Bisaccioni. 2015. "Asthma features in elderly in a tertiary outpatients facility." *The World Allergy Organization Journal* 8 (Suppl 1): A187. doi:10.1186/1939-4551-8-S1-A187. <http://dx.doi.org/10.1186/1939-4551-8-S1-A187>.

## Published Version

doi:10.1186/1939-4551-8-S1-A187

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:15035019>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

MEETING ABSTRACT

Open Access

# Asthma features in elderly in a tertiary outpatients facility

Nathalia Siqueira Robert De Castro<sup>1\*</sup>, Carolina Tavares De Alcântara<sup>2</sup>, Kelly Yoshimi Kanamori<sup>2</sup>, Jorge Kalil<sup>2</sup>, Bruna Gama Saliba<sup>2</sup>, Rosana Câmara Agondi<sup>1</sup>, Pedro Giavina-Bianchi<sup>3</sup>, Marcelo Vivolo Aun<sup>2</sup>, Carla Bisaccioni<sup>1</sup>

From 3rd WAO International Scientific Conference (WISC) 2014  
Rio de Janeiro, Brazil. 6-9 December 2014

## Background

The elderly comprise approximately 10% of the population. Asthma can affect about 10% of these patients; however, the diagnosis is underestimated. The sensitization to at least one aeroallergen in elderly with asthma ranges from 28% to 74%. The aim of this study was to evaluate the characteristics of asthma in patients aged over 60 years, including the frequency of allergic asthma.

## Methods

Cross-sectional study that evaluated 193 elderly patients with asthma followed at an outpatient tertiary service. Patients were considered when asthma met the criteria of GINA 2012 - anamnesis and pulmonary function demonstrating reversibility. These patients were evaluated for gender, current age, age at onset of asthma, lung function and asthma classification. The assessment of atopy was performed by searching for total IgE and specific IgE (*in vivo* or *in vitro*) for house dust mites, molds, cat and dog dander, cockroach and grass.

## Results

Of these 193 patients, 155 (80.3%) were female, the mean age was 69.3 years and the mean age of onset of asthma was 31.8 years. Forty-one patients (21%) had onset of symptoms before 12 years. Spirometry presented mean FEV1 of 64.5%. The classification of asthma showed 75% of patients with severe persistent asthma, 17% with moderate persistent asthma and 8%, mild persistent asthma. In relation to atopy, mean total IgE was 314.3 IU/mL. Positive specific IgE was demonstrated in 70% of patients, although, the group which the age of onset was between 41 and 59 years it was

42%. The majority was sensitized to house dust mites (91% of positive tests).

## Conclusions

In this study, patients whose asthma had an early-onset ( $\leq 12$  years) had more severe outcome. We noted that patients which asthma onset was between 41 to 59 years (25.9% of them) showed a less sensitization to aeroallergens (42% positive). We found also that 31% of elderly patients showed sensitization to molds, compared to literature where the mean was around 20%.

## Authors' details

<sup>1</sup>Hospital Das Clínicas - University of São Paulo, Brazil. <sup>2</sup>University of São Paulo, Brazil. <sup>3</sup>Brigham and Women's Hospital, Harvard Medical School, USA.

Published: 8 April 2015

doi:10.1186/1939-4551-8-S1-A187

Cite this article as: De Castro et al: Asthma features in elderly in a tertiary outpatients facility. *World Allergy Organization Journal* 2015 **8**(Suppl 1):A187.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Hospital Das Clínicas - University of São Paulo, Brazil  
Full list of author information is available at the end of the article