



Teenage smoking behaviour following a high-school smoking ban in Chile: interrupted time-series analysis

Citation

Feigl, Andrea B, Joshua A Salomon, Goodarz Danaei, Eric L Ding, and Esteban Calvo. 2015. "Teenage smoking behaviour following a high-school smoking ban in Chile: interrupted time-series analysis." *Bulletin of the World Health Organization* 93 (7): 468-475. doi:10.2471/BLT.14.146092. <http://dx.doi.org/10.2471/BLT.14.146092>.

Published Version

[doi:10.2471/BLT.14.146092](https://doi.org/10.2471/BLT.14.146092)

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:17820709>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Teenage smoking behaviour following a high-school smoking ban in Chile: interrupted time-series analysis

Andrea B Feigl,^a Joshua A Salomon,^a Goodarz Danaei,^a Eric L Ding^a & Esteban Calvo^b

Objective To evaluate the effect of a smoking ban in high schools on smoking behaviour among Chilean students.

Methods We conducted an interrupted time-series analysis, using repeated cross-sectional data from Chile's school population survey (2000–2011) for high-school students aged 12–18 years and a control group of persons aged 19–24 years. Poisson regression models were used to assess trends in smoking behaviour before and after the policy changes. The outcome measures were self-reported smoking prevalence (any smoking in the past month) and high frequency of smoking (smoking 15 days or more per month).

Findings From 2005 to 2011, the prevalence of smoking declined among high-school students by 6.8% per year compared with 3.6% decline per year in the control group. The decline in the target group was 2.9% (95% confidence interval, CI: 0.18 to 5.00) greater. We estimated that 5–6 years after enforcing the law, smoking prevalence among high-school students was 13.7% lower as a result of the ban. The impact of the smoking ban was primarily driven by declines in smoking prevalence among students in grades 8 to 10. The smoking ban did not significantly alter the frequency of smoking.

Conclusion The 2005 school smoking ban reduced smoking prevalence among younger high-school students in Chile. Further interventions targeting older individuals and frequent smokers may be needed.

Abstracts in **عربي**, **中文**, **Français**, **Русский** and **Español** at the end of each article.

Introduction

The detrimental health effects of tobacco smoking and use are widely documented.^{1–7} Every year, first-hand smoking causes five million deaths globally⁸ and 80% of smoking-related deaths occur in low- and middle-income countries.^{8,9} To reduce such mortality and morbidity, policy-makers can enforce policies such as public smoking bans,¹⁰ subsidized cessation programmes,^{11–13} tobacco warning labels^{14–16} and tax increases.^{17–19} However, the effect of these legislative changes differ based on context – compliance with smoking bans might be higher in countries with lower smoking prevalence.²⁰

There is a high prevalence of tobacco smoking in Chile, with 44% of adult males and 38% of adult females smoking in 2011, which was the highest prevalence in Latin America at that time.²¹ In 2009, Chile also had the second highest teenage smoking prevalence based on the Global Youth Tobacco Survey, with an annual smoking prevalence of 34.2%.⁹ The Chilean government ratified the WHO Framework Convention on Tobacco Control in 2005²² and passed tobacco legislation that took effect on 1 January 2006.²³

One provision in the legislation is a smoking ban, including a cigarette sales ban, within 300 metres of all high schools, which was the only provision that was enforced with 100% reported compliance (further information available from author). Preliminary evidence suggests that between 2005 and 2011, there was a decline in smoking prevalence among high-school students, but not in the general adult population.²⁴ Here we assess changes in smoking prevalence among high-school students before and after the smoking ban.

Methods

Study design

We used interrupted time-series to evaluate trends in smoking prevalence before and after the smoking ban in the target

population (high-school students aged 12–18 years) and compared it to the general population aged 19–24 years, which we assumed to be unaffected by the ban. We assumed a similar institutional environment in both groups, since in Chile, over 40% of individuals aged 19–24 years attended a university or a secondary institution in 2008.²⁵ To ensure that the control group did not include anyone subjected to the ban, the control population in 2000/2001 to 2006/2007 included individuals aged 19–24 years; in 2008/2009 individuals aged 20–24 years; and in 2010/2011 individuals aged 22–24 years.

Data

For the target population, we used data from the school population survey – a biennial and regionally representative survey on substance abuse and addictive behaviour in the Chilean school population, including the last grade of primary school (8th grade, mean age 13.5 years) and the high-school population (9th to 12th grade, ages 14–18 years). The school surveys were conducted in odd years; data from 2001–2011 were used in the analyses.²⁶ The school population survey employed a stratified, probabilistic two-stage sampling procedure.

For the control population, we used data from the general population survey,²⁶ conducted in even years, for 2000–2010. We combined data from the school and the general population surveys for before (2000/2001, 2002/2003 and 2004/2005) and after (2006/2007, 2008/2009 and 2010/2011) the implementation of the law. For each biennium, we analysed an average sample size of 50 000 individuals in the target group and 2300 individuals in the control group (further details available from the author).

Statistical analyses

To compare trends in smoking prevalence, we used self-reported, past 30-day smoking prevalence (yes/no) as the primary

^a Department of Global Health and Population, Harvard TH Chan School of Public Health, 677 Huntington Ave, 02115 Boston, MA, United States of America (USA).

^b School of Business and Economics, Universidad Diego Portales, Santiago, Chile.

Correspondence to Andrea B Feigl (email: abfeigl@mail.harvard.edu).

(Submitted: 22 August 2014 – Revised version received: 23 December 2014 – Accepted: 13 January 2015 – Published online: 29 April 2015)

outcome. As a secondary outcome, we created a frequent smoking variable indicating smoking 15 or more days per month during the last month (yes/no).

To model the outcomes, we used Poisson regression models to estimate prevalence ratios.^{9,27-29} The models were run separately for control and target groups due to different variances arising from different samples and sample frames. In multivariable regression models, we controlled for age, sex and region and included an interaction term for age and sex. For the target group we also controlled for school type, grade-level and grade point average, and in the control group for socioeconomic status and method of survey administration. Absence of data in covariates was generally lower than 2% and the complete case method was chosen for the main analysis. We then conducted a two-sample *t*-test with unequal variances conducted on the estimated coefficient for post-intervention change in the target group compared to the coefficient in the control group. All analyses were performed using Stata version 12 (StataCorp LP, College Station, United States of America).

We used a 2-stage model with time-dependent spline term for trend before and after 2005 (Table 1). At stage 1, we ran separate robust Poisson regression models for each intervention and control group, specified as follows:

$$Y_{ij} = \alpha + \beta_1 * t_{ij} + \beta_2 t_{2j} + \beta_{3...n}(\text{covariates})_{ij} + \varepsilon_{ij} \tag{1}$$

where *Y* represents the binary smoking variable of interest, for individual *i* at time *j*; ε represents standard errors clustered at the municipal level (high schools) and regional level (universities) and *t* represents the time period. In the first stage, the models (model a and model b) were run separately for each group.

At stage 2, we were interested in the β_2 coefficients of each model, reflecting the annual rate of change in smoking prevalence during the post-intervention period, relative to the annual rate of change in the pre-intervention period.

Table 1. Values for time period variable in first stage Poisson regressions to estimate trends in tobacco smoking, Chile, 2000–2011

| Time period | Survey period | | | | | |
|-------------|---------------|-----------|-----------|-----------|-----------|-----------|
| | 2000/2001 | 2002/2003 | 2004/2005 | 2006/2007 | 2008/2009 | 2010/2011 |
| 1 | 1 | 3 | 5 | 7 | 9 | 11 |
| 2 | 0 | 0 | 0 | 2 | 4 | 6 |

The first null hypothesis (H_01) that was tested in the analyses was:

$$\beta_2 a = \beta_2 b \tag{2}$$

the second null hypothesis (H_02) was:

$$\beta_2 a = 0 \tag{3}$$

the third null hypothesis (H_03) was:

$$\beta_2 b = 0 \tag{4}$$

H_01 was tested via a two-sample *t*-test with unequal variances, and H_02 and H_03 were tested via the significance test of the coefficient in each model.

We conducted several sensitivity analyses: (i) including all aged 19–24 years in the control population; (ii) running analogous analyses using past 30-day cannabis use as the main outcome (to test for the specificity of the school smoking ban on smoking behaviour versus addictive behaviour in general); (iii) including only those municipalities measured in every survey; (iv) excluding individuals with less than high-school education in the control group; (v) including additional control variables (alcohol consumption, and religion); (vi) including additional control variables (alcohol consumption, religion, and paternal education); (vii) adjusting for survey weights; (viii) adjusting for complex survey design; (ix) using the general population survey for individuals aged 19–64 years as the control population; and (x) using the missing indicator method to account for missing data.³⁰ For smoking frequency, we conducted the two sensitivity analyses – adjusting for complex survey design and stratifying the analysis by grade level in the target group – to investigate the robustness of the result.

Ethics approval

The study was exempt from ethics approval as it only used secondary data from surveys that had followed conventional ethics guidelines and recorded no personal identifiers.

Results

We obtained data on 319 798 individuals surveyed between 2000 and 2011. Characteristics for the target and control groups are presented in Table 2. Unadjusted for covariates, 30-day smoking prevalence among high-school students was highest in 2000/2001, with a prevalence of 41.9% (23 822/56 817) and slightly declining to 40.1% (23 678/59 101) in 2004/2005. After the implementation of the law, the prevalence declined to 25.7% (8596/33 509) in 2010/2011. Smoking prevalence in the control group was higher overall, with the highest in 2002/2003 (58.2%; 1132/1945). Before the law came into effect the prevalence was 57.3% (1104/1927) which then steadily declined to 44.9% (405/902) by 2010/2011. Immediately after 2005, there was a greater decline in smoking prevalence among high-school students than among the control population (40.1% to 34.9% and 57.3% to 54.8%, respectively).

Smoking frequency was higher in the control group than in the target group. In 2002/2003, high-school individuals smoked an average of 16.5 days per month, which declined to 15.0 days per month in 2006/2007 and further dropped to 13.0 days per month in 2010/2011. The number of days smoked per month in the control group was relatively stable between 2000 and 2011, with a peak of 21.4 days during 2004/2005 and by 2010/2011 this had declined to 19.9 days. In all groups and grades, average smoking frequency was highest in 2002/2003 and lowest in 2010/2011. Smoking prevalence and fre-

Table 2. Characteristics of study populations, Chile, 2000–2011

| Characteristic | High-school population aged 12–18 years | | | | | | Control population aged 19–24 years, ^a | | | | | |
|--|---|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|---|------------------------|------------------------|------------------------|------------------------|----------------------|
| | 2000/2001 n = 56 817 | 2002/2003 n = 57 032 | 2004/2005 n = 59 101 | 2006/2007 n = 51 432 | 2008/2009 n = 48 213 | 2010/2011 n = 33 509 | 2000/2001 n = 5 466 | 2002/2003 n = 1 945 | 2004/2005 n = 1 927 | 2006/2007 n = 1 937 | 2008/2009 n = 1 517 | 2010/2011 n = 902 |
| Age, years (SD) | 15.5 (1.5) | 15.6 (1.5) | 15.2 (1.5) | 15.4 (1.5) | 15.4 (1.5) | 15.5 (1.5) | 21.5 (1.7) | 21.6 (1.7) | 21.6 (1.7) | 21.5 (1.7) | 22.0 (1.43) | 23.0 (0.80) |
| Number of men, (%) | 27 953 (49.2) | 28 117 (49.3) | 29 019 (49.1) | 25 458 (49.5) | 23 673 (49.1) | 16 520 (49.3) | 2 603 (47.6) | 1 093 (56.2) | 1 045 (54.2) | 1 038 (53.6) | 712 (46.9) | 409 (45.3) |
| Numbers of smokers in the past 30 days, (%) | 23 822 (41.9) | 22 194 (38.9) | 23 678 (40.1) | 17 950 (34.9) | 15 910 (33.0) | 8 596 (25.7) | 3 012 (55.1) | 1 132 (58.2) | 1 104 (57.3) | 1 061 (54.8) | 769 (50.7) | 405 (44.9) |
| Number of days smoked per month | NA | 16.5 | 14.1 | 15.0 | 14.5 | 13.0 | NA | 21.3 | 21.4 | 19.8 | 20.9 | 19.9 |
| Number of high frequency smokers, ^b (%) | NA | 11 076 (19.4) | 9 675 (16.4) | 7 951 (15.5) | 6 928 (14.4) | 3 364 (10.0) | NA | 832 (42.8) | 809 (42.0) | 724 (37.4) | 529 (34.9) | 280 (31.0) |
| Number of cannabis users in past 30 days, (%) | 4 432 (7.8) | 3 992 (7.0) | 3 605 (6.1) | 4 269 (8.3) | 3 857 (8.0) | 3 250 (9.7) | 364 (6.7) | 134 (6.9) | 137 (7.1) | 140 (7.2) | 119 (7.8) | 56 (6.2) |

NA: not applicable; SD: standard deviation.

^a The control population in 2000/2001 to 2006/2007 includes individuals aged 19–24 years, in 2008/2009 individuals aged 20–24 years, and in 2010/2011 individuals aged 22–24 years.

^b Smoking more than 15 days per month.

quency increased with increasing school grade (data available from author).

Smoking prevalence

Adjusting for covariates in the multivariable regression model, smoking prevalence significantly increased among high-school students over the period 2000–2005, from an adjusted smoking prevalence of 37.4% in 2000/2001, to an adjusted smoking prevalence of 38.6% in 2004/05 ($P = 0.014$); there was no significant trend observed in the control group. After 2005, there was a significant decline in smoking prevalence in the target group ($P < 0.001$), resulting in an adjusted smoking prevalence of 24.9% in 2010/11 in the high-school population. The prevalence ratio of the smoking prevalence change before versus after the law in the control versus the target group was 0.971 annually (95% confidence interval, CI: 0.950 to 0.992), or 0.863 (95% CI: 0.774 to 0.961) over the course of the intervention (Table 3). Thus, smoking prevalence after 2005 declined significantly faster in the target group than in the control group.

Since the descriptive analysis revealed a difference in smoking prevalence among different high-school grades (data available from author), a grade-based stratified analysis was done to determine the grades in which the law had the greatest effect. The smoking ban was most effective among students in the lowest grade, leading to a 7.2% (95% CI: 4.3 to 10.1) annual improvement in smoking trend in this population versus the control group. The ban was least effective among students in the two highest grades, where no significant effect on smoking prevalence was detected (Table 4).

Smoking frequency

Before the ban, frequent smoking in the target group declined by 2.5% annually from 2000 to 2005 (95% CI: -4.1 to -0.8); while in the control group no significant trend was observed. After 2005, the difference in smoking frequency between the target and control group was not significant ($P = 0.58$; Table 5).

Sensitivity analyses

When examining the changes in cannabis use before and after the implementation of the law, after 2005 cannabis use increased significantly in the target group, but declined in the control population. Hence, after 2005, past 30-day

Table 3. Changes in smoking prevalence before and after the implementation of a high-school smoking ban in Chile, 2000–2011

| Policy period | Annual prevalence change, PR ^a (95% CI) | | | |
|---|--|----------------------------|----------------------------------|------------------------|
| | Target group ^b | Control group ^c | Intervention effect ^d | |
| | | | Per year | Five years |
| Pre-policy (2000–2005) | 1.006 (1.001 to 1.012) | 1.010 (0.999 to 1.02) | 0.971 (0.950 to 0.992) | 0.863 (0.774 to 0.961) |
| Post-policy (2006–2011) | 0.932 (0.927 to 0.937) | 0.964 (0.953 to 0.974) | | |
| Difference between pre-policy and post-policy | 0.926 (0.917 to 0.934) | 0.953 (0.935 to 0.973) | | |

CI: confidence interval; PR: prevalence rate ratio.

^a PRs represent the annual decrease in smoking prevalence.

^b High-school population aged 12–18 years. Adjusted for age, sex, region, school-type, course, and sex * age.

^c Aged 19–24 years. Adjusted for age, sex, region, socioeconomic status, survey method, and sex * age.

^d Difference in changes of smoking prevalence between target and control group.

cannabis use increased by a 14% (95% CI: 11 to 18) greater rate in the target population compared to the control population (data available from author).

Restricting the analysis to municipalities that were included in all survey years yielded identical effect size estimates as the main analysis (data available from author). When including only those with a high-school diploma in the control population, the effect size for the smoking ban increased to 4.5% (95% CI: 1.3 to 7.5). Adjusting for alcohol consumption, religion and paternal education in the target group also increased the effect size estimate to 4.8% (95% CI: 2.4 to 6.4). The results for smoking frequency were robust to inclusion of complex survey design and educational grade (data available from author).

Discussion

Here we demonstrate that smoke-free zones and tobacco sales restrictions in high schools can be effective in the Chilean context. We observed a greater decline in smoking prevalence in the target group than in the control group after the implementation of the law. The average intervention effect was driven by the decline in smoking prevalence among younger high-school students. In contrast, the smoking ban proved ineffective in lowering prevalence among older students, as well as in reducing the frequency of smoking. These results suggest that the smoking ban prevented smoking initiation and selectively targeted low-frequency smokers. These findings are consistent with the theory that non-smokers and less frequent smokers are most receptive to tobacco-control policies.³¹

The similar decline in frequent smoking in both groups suggests that the ban did not affect how often people smoke and therefore better-targeted programmes and policies are needed for frequent smokers. Examples of effective programmes include smoking-cessation counselling in high schools and free prescription of nicotine patches.¹¹ Legislation featuring such programmes has been proposed, but not yet passed or funded in Chile.

Several sensitivity analyses supported the robustness of our results. When we included all individuals aged 19–24 years in the control group, we found that the effect estimate was lower since the control group in 2008 and 2010 also included people previously targeted by the law. In contrast, when including only those with a high-school diploma in the control group, the effect size increased. A possible explanation is that those with at least a high-school diploma continued, rather than altered, their smoking behaviour between 19 and 24 years of age. Applying cannabis use as the main outcome suggested that relative to the control group, cannabis use in high-school students increased after the implementation of the law. This is further evidence that the decline in smoking prevalence among high-school students is attributable to the smoking ban. Our findings confirm results from previous studies that increased difficulty of obtaining cigarettes was associated with less positive views about smoking among adolescents³² and lower smoking prevalence among Chilean teenagers.³³

In addition to the immediate effect of the law, the long-term effect on disease burden should also be considered. Emerging evidence suggests that adolescence is the most critical period in life for development of long-term ad-

dictive behaviours.^{34–36} In the USA, it has been shown that laws on youth access to tobacco might have long-term effects on smoking prevalence, however, these effects might be limited to women only.³⁷ To estimate whether the smoking ban would have a long-term effect on morbidity, one can use a life table approach such as the 1960–1972 American cohort. In this cohort, the mortality rate in men aged 35–39 years, who have never smoked regularly, was 1.34 per 1000 person-years, versus 2.55 in those who smoked one or more packs of cigarettes per day.³⁸ In the age group 40–44 years, mortality in men who never smoked was 1.93 per 1000 person-years, versus 4.59 in frequent smokers (smoking more than 20 cigarettes per day).³⁸ Overall mortality rates in women were lower in both smokers and non-smokers, but still substantially higher in frequent smokers than non-smokers. Assuming that the mortality rates from this American cohort apply to the current Chilean high-school population and that the effect of the ban on prevalence persists long-term, the Chilean high-school smoking ban could potentially reduce mortality by more than twofold 20 years from now. This reduction would result in better health outcomes for a large number of people as well as reducing health expenditure on smoking-related diseases like lung cancer or chronic obstructive pulmonary disease.^{39–43} However, there is uncertainty as to how long this effect on smoking behaviour will last, and long-term assessments are needed.

This study has limitations. First, individuals were not randomly assigned to the target group and control group. However, all provisions of the 2005 law uniformly affected Chile's general population (data available from author),

Table 4. Changes in smoking prevalence before and after the implementation of a high-school smoking ban in Chile, 2000–2011 by school grade

| Policy period | Annual prevalence change, PR ^a (95% CI) | | | | | Control group ^c |
|---|--|------------------------|------------------------|------------------------|------------------------|----------------------------|
| | 8th | 9th | 10th | 11th | 12th | |
| Pre-policy (2000–2005) | 1.02 (1.001 to 1.03) | 1.01 (0.999 to 1.02) | 1.01 (0.999 to 1.02) | 0.998 (0.990 to 1.01) | 0.995 (0.941 to 1.00) | 1.010 (0.999 to 1.02) |
| Post-policy (2006–2011) | 0.898 (0.886 to 0.910) | 0.930 (0.923 to 0.938) | 0.930 (0.923 to 0.937) | 0.940 (0.934 to 0.947) | 0.948 (0.941 to 0.956) | 0.964 (0.953 to 0.974) |
| Difference between pre-policy and post-policy | 0.885 (0.864 to 0.906) | 0.922 (0.908 to 0.937) | 0.923 (0.911 to 0.936) | 0.942 (0.930 to 0.955) | 0.942 (0.930 to 0.955) | 0.953 (0.935 to 0.973) |
| Intervention effect ^d | 0.928 (0.899 to 0.957) | 0.967 (0.943 to 0.992) | 0.968 (0.945 to 0.992) | 0.988 (0.965 to 1.01) | 1.00 (0.975 to 1.02) | NA |

CI: confidence interval; NA: not applicable; PR: prevalence rate ratio.
^a PRs represent the annual decrease in smoking prevalence.

^b High-school population aged 12–18 years. Adjusted for age, sex, region, school-type, course, and sex * age.
^c Aged 19–24 years. Adjusted for age, sex, region, socioeconomic status, survey method, and sex * age.

Table 5. Change in prevalence of high frequency smoking^b before and after the implementation of a high-school smoking ban in Chile, 2000–2011

| Policy period | Annual prevalence change, PR ^b (95% CI) | | Intervention effect ^e |
|---|--|----------------------------|----------------------------------|
| | Target group ^c | Control group ^d | |
| Pre-policy (2000–2005) | 0.975 (0.959 to 0.992) | 0.992 (0.959 to 1.03) | 0.986 (0.938 to 1.036) |
| Post-policy (2006–2011) | 0.924 (0.916 to 0.932) | 0.953 (0.938 to 0.969) | |
| Difference between pre-policy and post-policy | 0.947 (0.926 to 0.969) | 0.961 (0.919 to 1.00) | |

CI: confidence interval; PR: prevalence rate ratio.
^a Smoking more than 15 days per month.
^b PRs represent the annual decrease in smoking prevalence.

^c High-school population aged 12–18 years. Adjusted for age, sex, region, school-type, course, and sex * age.
^d Aged 19–24 years. Adjusted for age, sex, region, socioeconomic status, survey method, and sex * age.
^e Difference in changes of smoking prevalence between target and control group.

except for the high-school smoking ban. Thus, even if the law affected the control group, the same section of the law also affected the target group. Second, the average age of smoking initiation was 13.2 years in Chile and older individuals are less likely to start or quit smoking. Hence, individuals in the control group (19–24 years) – both smokers and non-smokers – may have a lower propensity to change their behaviour in response to a similar ban. This could have led to an upward bias in the effect size estimates. However, in a sensitivity analysis using the general population aged 25–64 years as a control group, the effect sizes were not significantly higher. Third, the cross sectional design of our study means that we cannot identify specific outcomes of the ban at an individual level. We cannot distinguish, for example, whether the ban prevented people from starting smoking or encouraged current smokers to stop. Neither can we estimate the effect of the ban on overall levels of tobacco consumption per capita. Fourth, there were no data on how schools enforced the ban. Additional data on compliance with the smoking ban at Chilean high schools – similar to a study conducted in India⁴⁴ – could have further strengthened our conclusions.

Smoking prevalence is often highly correlated with socioeconomic status.⁴⁵ In our study, socioeconomic status variables in both target and control group were highly collinear with neighbourhood. For this reason, socioeconomic status variables were omitted from the main analyses. Future studies could explore equity concerns relating to the high-school smoking ban.

In conclusion, smoking bans in schools can be effective in reducing the number of teenagers smoking but not in decreasing the frequency of smoking. These are relevant lessons for both national and global policy-makers. Chile's 2013 tobacco control legislation extended the smoking bans to include bars, restaurants and universities, which may help to further decrease smoking prevalence. However, both the 2005 and 2013 tobacco legislation offered little help to those in need of cessation counselling. Future policies should include accessible cessation programmes, funds for enforcing smoke-free public spaces and other provisions to help people stop smoking. ■

Acknowledgements

The authors thank Thomas Bossert (HSPH), Álvaro Castillo (SENDA), Jessica Cohen (HSPH), Matías Irrarrazaval (SENDA), Esteban Pizarro (SENDA),

Maria Valenzuela-Schmidt (Universidad de Chile) and Macarena Vivent (UDP).

Latin American Studies and the Harvard Michael von Clemm travel grant.

Funding: ABF received travel grants from the Harvard David Rockefeller Center for

Competing interests: None declared.

ملخص

سلوك تدخين المراهقين في أعقاب حظر التدخين في المدارس الثانوية في شيلي: تحليل سلاسل زمنية متقطعة
الغرض تقييم تأثير حظر التدخين في المدارس الثانوية في سلوك التدخين بين الطلاب في شيلي. الطريقة لقد أجرينا تحليل سلاسل زمنية متقطعة، وذلك باستخدام بيانات مقطعية عرضية متكررة من مسح جمهور الطلاب في المدرسة في شيلي (في الفترة ما بين عام 2000 إلى عام 2011) لطلاب المدارس الثانوية الذين تتراوح أعمارهم ما بين 12 إلى 18 عامًا ومجموعة الشاهد من الأشخاص الذين تتراوح أعمارهم ما بين 19 إلى 24 عامًا. واستخدمت نماذج التحوف لبواسون لتقييم النزعات في سلوك التدخين قبل وبعد تغيرات السياسة. كانت قياسات الحصائل عبارة عن معدل انتشار التدخين المبلغ عنه ذاتيًا (أي حالة تدخين في الشهر الماضي) وارتفاع وتيرة التدخين (التدخين لمدة 15 يومًا أو أكثر في الشهر). النتائج في الفترة من عام 2005 إلى عام 2011، انخفض معدل انتشار التدخين بين طلاب المدارس الثانوية بنسبة 6.8٪ سنويًا مقارنة مع انخفاض بنسبة 3.6٪ سنويًا في مجموعة الشاهد. كان الانخفاض في المجموعة المستهدفة بنسبة 2.9٪ (بنسبة أرجحية مقدارها 95٪: 0.18 إلى 5.00) أكبر. ولقد قدرنا أنه بعد تطبيق القانون لمدة من 5 إلى 6 أعوام، كان معدل انتشار التدخين بين طلاب المدارس الثانوية أقل بنسبة 13.7٪ كنتيجة لهذا الحظر. ونتج تأثير حظر التدخين بشكل رئيسي عن الانخفاض في معدلات انتشار التدخين بين الطلاب في الصفوف من 8 إلى 10. ولم يغير حظر التدخين كثيرًا من وتيرة التدخين. الاستنتاج أدى حظر التدخين في المدارس في عام 2005 إلى انخفاض معدل انتشار التدخين بين طلاب المدارس الثانوية الأصغر في شيلي. قد تكون هناك حاجة إلى إجراء المزيد من التدخلات التي تستهدف الأفراد الأكبر سنًا والمدخنين بشكل متكرر.

摘要**智利颁布了高中禁烟令后青少年的吸烟行为：断续时间序列分析**

目的 旨在评估高中禁烟令对智利学生吸烟行为的影响。

方法 我们开展了一项断续时间序列分析，所采用的重复性横断面数据来源于智利学校人口调查（2000–2011年），其中的调查对象为年龄介于12至18岁的高中学生，同时以年龄介于19至24岁的人群作为对照组。泊松回归模型用于评估吸烟行为在政策改变之前和之后的趋势。效果衡量指标为自报吸烟流行率（在过去一个月中的任何吸烟行为）以及频繁程度高的吸烟行为（每个月吸烟15天或更多）。

结果 从2005到2011年，高中学生的吸烟流行率每年下降6.8%，而对照组每年下降3.6%。目标群体的下降率为2.9%（95%置信区间，CI:0.18至5.00）以上。我们估计在强制执行法律后的5至6年内，高中学生的吸烟流行率会因该禁令而降低13.7%。禁烟令的影响主要是由于吸烟流行率在8至10年级的学生中下降而引起的。禁烟令并没有显著改变吸烟的频繁程度。

结论 2005年颁布的学校禁烟令使智利境内较为年轻的高中学生降低了吸烟流行率。我们可能需要针对较为年长的个人和吸烟频繁者实施进一步的干预措施。

Résumé**Comportement tabagique des adolescents à la suite d'une interdiction de fumer dans les établissements d'enseignement secondaire du Chili: analyse de séries chronologiques interrompues**

Objectif Évaluer les conséquences d'une interdiction de fumer dans les établissements d'enseignement secondaire sur le comportement tabagique des élèves chiliens.

Méthodes Nous avons effectué une analyse de séries chronologiques interrompues à l'aide de données transversales répétées provenant d'une enquête sur la population scolaire du Chili (2000-2011) menée auprès d'élèves du secondaire âgés de 12 à 18 ans et d'un groupe de contrôle constitué d'individus âgés de 19 à 24 ans. Des modèles de régression Poisson ont été utilisés pour évaluer l'évolution du comportement tabagique avant et après les changements d'orientation. Les mesures de résultat reposaient sur les auto-déclarations de prévalence du tabagisme (toute consommation de tabac au cours du mois passé) et de fréquence élevée de

consommation de tabac (consommation de tabac 15 jours ou plus par mois).

Résultats De 2005 à 2011, la prévalence du tabagisme a diminué de 6,8 % par an chez les élèves du secondaire; en comparaison, elle a diminué de 3,6 % par an dans le groupe de contrôle. La diminution observée dans le groupe cible était supérieure de 2,9 % (intervalle de confiance, IC, à 95 %: 0,18 à 5,00). Nous avons estimé que 5-6 ans après l'entrée en vigueur de la loi, la prévalence du tabagisme chez les élèves du secondaire avait diminué de 13,7 % grâce à l'interdiction. L'impact de l'interdiction de fumer s'est principalement traduit par une diminution de la prévalence du tabagisme chez les élèves des 8e, 9e et 10e années. L'interdiction de fumer n'a pas considérablement changé la fréquence de consommation de tabac.

Conclusion L'interdiction de fumer dans les écoles, qui date de 2005, a diminué la prévalence du tabagisme chez les plus jeunes élèves des établissements d'enseignement secondaire du Chili. D'autres

interventions ciblant des individus plus âgés et des fumeurs fréquents pourraient être nécessaires.

Резюме

Курение среди подростков после запрета в старших классах школ в Чили: анализ методом прерванного временного ряда

Цель Оценить, каким образом запрет на курение в старших классах чилийских школ отразился на курении учеников.

Методы Был проведен анализ статистических данных, полученных в результате многократных опросов старшеклассников (12–18 лет) и контрольной группы взрослых (19–24 года) в Чили в 2000–2011 гг., методом прерванного временного ряда. Для оценки поведенческих тенденций в отношении курения до изменения государственной политики и после этого применялись регрессионные модели Пуассона. В качестве критериев оценки использовались личные показания опрошенных о самом факте курения (курили ли опрошенные в течение прошлого месяца) и о высокой частоте курения (курение в течение как минимум 15 дней в прошлом месяце).

Результаты В период с 2005 по 2011 год распространенность курения среди школьников старших классов каждый год

снижалась на 6,8%. В контрольной группе снижение составило 3,6%. В целевой группе снижение было выше на 2,9% (95% доверительный интервал, ДИ: от 0,18 до 5,00). По нашим оценкам, спустя 5–6 лет после введения закона в действие распространенность курения среди старшеклассников в результате запрета снизилась на 13,7%. Действие запрета на курение в основном проявилось в том, что стали меньше курить школьники 8–10 классов. Запрет на курение не оказал существенного влияния на частоту курения.

Вывод Запрет на курение в школах в 2005 году снизил распространенность курения среди чилийских старшеклассников младшего возраста. Для более взрослых и часто курящих лиц могут потребоваться дополнительные интервенционные меры.

Resumen

El consumo de tabaco entre adolescentes tras la prohibición de fumar en los institutos de Chile: análisis de series temporales interrumpidas

Objetivo Evaluar el efecto de la prohibición de fumar en los institutos sobre el consumo de tabaco entre los alumnos chilenos.

Métodos Se llevó a cabo un análisis de series temporales interrumpidas utilizando datos transversales repetidos obtenidos de la encuesta de la población escolar de Chile (2000–2011) en alumnos de instituto de entre 12 y 18 años y con un grupo de control de personas de entre 19 y 24 años. Se utilizaron modelos de regresión de Poisson para evaluar las tendencias en el consumo de tabaco antes y después de los cambios en la política. Los indicadores de resultados eran la prevalencia autodeclarada del consumo de tabaco (cualquier consumo de tabaco en el último mes) y la alta frecuencia del consumo de tabaco (fumar 15 o más días al mes).

Resultados De 2005 a 2011, la prevalencia del consumo de tabaco se redujo en un 6,8% anual entre los estudiantes de instituto, en

comparación con la reducción del 3,6% anual en el grupo de control. El descenso en el grupo objetivo fue de un 2,9% (intervalo de confianza, IC, del 95%: de 0,18 a 5,00) superior. Se estimó que 5–6 años después de la aplicación de la ley, la prevalencia del consumo de tabaco entre los alumnos de instituto era un 13,7% menor como consecuencia de la prohibición. El impacto de la prohibición de fumar se notó principalmente en la disminución del consumo de tabaco entre los alumnos de instituto de octavo a décimo grado. La prohibición de fumar no alteró significativamente la frecuencia del consumo de tabaco.

Conclusión La prohibición de fumar en las escuelas de 2005 redujo la prevalencia del consumo de tabaco entre los alumnos de instituto más jóvenes en Chile. Es posible que sean necesarias intervenciones adicionales dirigidas a individuos de más edad y a fumadores frecuentes.

References

- Eriksen M, Mackay JM, Schluger N, Islami F, Drope J. The tobacco atlas. 5th ed. Atlanta: The American Cancer Society; 2015. Available from: www.tobaccoatlas.org [cited 2015 Apr 20].
- Doll R, Hill AB. Smoking and carcinoma of the lung; preliminary report. *BMJ*. 1950 Sep 30;2(4682):739–48. doi: http://dx.doi.org/10.1136/bmj.2.4682.739 PMID: 14772469
- Lin HH, Ezzati M, Murray M. Tobacco smoke, indoor air pollution and tuberculosis: a systematic review and meta-analysis. *PLoS Med*. 2007 Jan;4(1):e20. doi: http://dx.doi.org/10.1371/journal.pmed.0040020 PMID: 17227135
- Mackay JM, Bettcher DW, Minhas R, Schotte K. Successes and new emerging challenges in tobacco control: addressing the vector. *Tob Control*. 2012 Mar;21(2):77–9. doi: http://dx.doi.org/10.1136/tobaccocontrol-2012-050433 PMID: 22345225
- Magrath I, Litvak J. Cancer in developing countries: opportunity and challenge. *J Natl Cancer Inst*. 1993 Jun 2;85(11):862–74. doi: http://dx.doi.org/10.1093/jnci/85.11.862 PMID: 8492315
- Murray CJ, Lopez AD. Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study. *Lancet*. 1997 May 17;349(9063):1436–42. doi: http://dx.doi.org/10.1016/S0140-6736(96)07495-8 PMID: 9164317
- Proctor RN. The history of the discovery of the cigarette-lung cancer link: evidentiary traditions, corporate denial, global toll. *Tob Control*. 2012 Mar;21(2):87–91. doi: http://dx.doi.org/10.1136/tobaccocontrol-2011-050338 PMID: 22345227
- Mortality attributable to tobacco. Geneva: World Health Organization; 2012.
- World tobacco atlas. Atlanta: The American Cancer Society; 2012.
- Callinan JE, Clarke A, Doherty K, Kelleher C. Legislative smoking bans for reducing secondhand smoke exposure, smoking prevalence and tobacco consumption. *Cochrane Database Syst Rev*. 2010; (4):CD005992. PMID: 20393945
- Suls JM, Luger TM, Curry SJ, Mermelstein RJ, Sporer AK, An LC. Efficacy of smoking-cessation interventions for young adults: a meta-analysis. *Am J Prev Med*. 2012 Jun;42(6):655–62. doi: http://dx.doi.org/10.1016/j.amepre.2012.02.013 PMID: 22608385

12. Myung SK, McDonnell DD, Kazinets G, Seo HG, Moskowitz JM. Effects of Web- and computer-based smoking cessation programs: meta-analysis of randomized controlled trials. *Arch Intern Med*. 2009 May 25;169(10):929–37. doi: <http://dx.doi.org/10.1001/archinternmed.2009.109> PMID: 19468084
13. Manske S, Miller S, Moyer C, Phaneuf MR, Cameron R. Best practice in group-based smoking cessation: results of a literature review applying effectiveness, plausibility, and practicality criteria. *Am J Health Promot*. 2004 Jul-Aug;18(6):409–23. doi: <http://dx.doi.org/10.4278/0890-1171-18.6.409> PMID: 15293927
14. Hammond D, Fong GT, McDonald PW, Cameron R, Brown KS. Impact of the graphic Canadian warning labels on adult smoking behaviour. *Tob Control*. 2003 Dec;12(4):391–5. doi: <http://dx.doi.org/10.1136/tc.12.4.391> PMID: 14660774
15. Hammond D, Fong GT, McNeill A, Borland R, Cummings KM. Effectiveness of cigarette warning labels in informing smokers about the risks of smoking: findings from the International Tobacco Control (ITC) Four Country Survey. *Tob Control*. 2006 Jun;15 Suppl 3:iii19–25. doi: <http://dx.doi.org/10.1136/tc.2005.012294> PMID: 16754942
16. Thrasher JF, Allen B, Reynales-Shigematsu LM, Anaya R, Lazcano-Ponce E, Hernández-Avila M. Análisis del impacto en fumadores mexicanos de los avisos gráficos en las cajetillas de cigarros. *Salud Publica Mex*. 2006;48 Suppl 1:565–74. Spanish. PMID: 17684691
17. Chaloupka F, Grossman M. Price, tobacco control policies, and youth smoking. Cambridge: National Bureau of Economic Research; 1996. doi: <http://dx.doi.org/http://dx.doi.org/10.3386/w5740>doi: <http://dx.doi.org/10.3386/w5740>
18. Chaloupka FJ, Straif K, Leon ME; Working Group, International Agency for Research on Cancer. Effectiveness of tax and price policies in tobacco control. *Tob Control*. 2011 May;20(3):235–8. doi: <http://dx.doi.org/10.1136/tc.2010.039982> PMID: 21115556
19. Chaloupka FJ, Wechsler H. Price, tobacco control policies and smoking among young adults. *J Health Econ*. 1997 Jun;16(3):359–73. doi: [http://dx.doi.org/10.1016/S0167-6296\(96\)00530-9](http://dx.doi.org/10.1016/S0167-6296(96)00530-9) PMID: 10169306
20. Levy DT, Chaloupka F, Gitchell J. The effects of tobacco control policies on smoking rates: a tobacco control scorecard. *J Public Health Manag Pract*. 2004 Jul-Aug;10(4):338–53. doi: <http://dx.doi.org/10.1097/00124784-200407000-00011> PMID: 15235381
21. World Development Indicators. Washington: World Bank; 2015.
22. Parties and Signatories to the WHO Framework Convention on Tobacco Control [Internet]. Geneva: World Health Organization; 2012. Available from: http://www.who.int/fctc/signatories_parties/en/ [cited 2012 February 15].
23. Historia de la ley N° 20.105: modifica la ley 19.419, en materias relativas a la publicidad y el consumo del tabaco. Santiago: Biblioteca del Congreso Nacional de Chile; 2006. Spanish.
24. Noveno estudio nacional de drogas en población general de Chile: Observatorio Chileno de drogas. Santiago: Ministerio del Interior y Seguridad Publica, Gobierno de Chile; 2012. Available from: <http://www.senda.gob.cl/observatorio/estudios/poblacion-general/> [cited 2015 Apr 23]. Spanish.
25. Educacion: encuesta casen. Santiago: Ministerio de Planificacion, Gobierno de Chile; 2009. Spanish.
26. Estudios: Observatorio Chileno de Drogas [Internet]. Santiago: Ministerio del Interior y Seguridad Publica, Gobierno de Chile; 2012. Available from: <http://www.senda.gob.cl/observatorio/estudios/> [cited 2015 Apr 23]. Spanish.
27. Spiegelman D, Hertzmark E. Easy SAS calculations for risk or prevalence ratios and differences. *Am J Epidemiol*. 2005 Aug 1;162(3):199–200. doi: <http://dx.doi.org/10.1093/aje/kwi188> PMID: 15987728
28. Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol*. 2004 Apr 1;159(7):702–6. doi: <http://dx.doi.org/10.1093/aje/kwh090> PMID: 15033648
29. McNutt LA, Wu C, Xue X, Hafner JP. Estimating the relative risk in cohort studies and clinical trials of common outcomes. *Am J Epidemiol*. 2003 May 15;157(10):940–3. doi: <http://dx.doi.org/10.1093/aje/kwg074> PMID: 12746247
30. Seaman SR, White IR. Review of inverse probability weighting for dealing with missing data. *Stat Methods Med Res*. 2013 Jun;22(3):278–95. doi: <http://dx.doi.org/10.1177/0962280210395740> PMID: 21220355
31. Becker GS, Murphy KM. A theory of rational addiction. *J Polit Econ*. 1988;96(4):675. doi: <http://dx.doi.org/10.1086/261558>
32. Lorenzo-Blanco EI, Bares C, Delva J. Correlates of Chilean adolescents' negative attitudes toward cigarettes: the role of gender, peer, parental, and environmental factors. *Nicotine Tob Res*. 2012 Feb;14(2):142–52. doi: <http://dx.doi.org/10.1093/ntr/nt152> PMID: 22157230
33. Horner P, Grogan-Kaylor A, Delva J, Bares CB, Andrade F, Castillo M. The association of family and peer factors with tobacco, alcohol, and marijuana use among Chilean adolescents in neighborhood context. *Subst Abuse Rehabil*. 2011 Sep 1;2(1):163–72. doi: <http://dx.doi.org/10.2147/SAR.S20507> PMID: 22224067
34. Chambers RA, Taylor JR, Potenza MN. Developmental neurocircuitry of motivation in adolescence: a critical period of addiction vulnerability. *Am J Psychiatry*. 2003 Jun;160(6):1041–52. doi: <http://dx.doi.org/10.1176/appi.ajp.160.6.1041> PMID: 12777258
35. Crews F, He J, Hodge C. Adolescent cortical development: a critical period of vulnerability for addiction. *Pharmacol Biochem Behav*. 2007 Feb;86(2):189–99. doi: <http://dx.doi.org/10.1016/j.pbb.2006.12.001> PMID: 17222895
36. Slotkin TA. Nicotine and the adolescent brain: insights from an animal model. *Neurotoxicol Teratol*. 2002 May-Jun;24(3):369–84. doi: [http://dx.doi.org/10.1016/S0892-0362\(02\)00199-X](http://dx.doi.org/10.1016/S0892-0362(02)00199-X) PMID: 12009492
37. Gruzca RA, Plunk AD, Hipp PR, Cavazos-Rehg P, Krauss MJ, Brownson RC, et al. Long-term effects of laws governing youth access to tobacco. *Am J Public Health*. 2013 Aug;103(8):1493–9. doi: <http://dx.doi.org/10.2105/AJPH.2012.301123> PMID: 23763414
38. Lew EA, Garfinkel L. Differences in mortality and longevity by sex, smoking habits and health status. *Trans Soc Actuar*. 1987;39:107–30.
39. Bloom DE, Chisholm D, Jane-Llopis E, Prettner K, Stein A, Feigl A. From burden to best buys: reducing the economic impact of non-communicable diseases in low- and middle-income countries. Geneva: World Economic Forum; 2011.
40. Lopez Varela MV, Montes de Oca M, Halbert RJ, Muiño A, Perez-Padilla R, Tálamo C, et al.; PLATINO Team. Sex-related differences in COPD in five Latin American cities: the PLATINO study. *Eur Respir J*. 2010 Nov;36(5):1034–41. doi: <http://dx.doi.org/10.1183/09031936.00165409> PMID: 20378599
41. Rutten-van Mölken M. Raising the awareness: projecting the future burden of COPD with the BOLD model. *Eur Respir J*. 2009 Oct;34(4):787–9. doi: <http://dx.doi.org/10.1183/09031936.00108409> PMID: 19797666
42. Economic impact of COPD and cost effective solutions. Milton: Access Economics and The Australian Lung Foundation; 2008.
43. Menezes AM, Perez-Padilla R, Hallal PC, Jardim JR, Muiño A, Lopez MV, et al.; PLATINO Team. Worldwide burden of COPD in high- and low-income countries. Part II. Burden of chronic obstructive lung disease in Latin America: the PLATINO study. *Int J Tuberc Lung Dis*. 2008 Jul;12(7):709–12. PMID: 18544192
44. Goel S, Ravindra K, Singh RJ, Sharma D. Effective smoke-free policies in achieving a high level of compliance with smoke-free law: experiences from a district of North India. *Tob Control*. 2014 Jul;23(4):291–4. doi: <http://dx.doi.org/10.1136/tobaccocontrol-2012-050673> PMID: 23322311
45. Hosseinpoor AR, Parker LA, Tursan d'Espaignet E, Chatterji S. Socioeconomic inequality in smoking in low-income and middle-income countries: results from the World Health Survey. *PLoS ONE*. 2012;7(8):e42843. doi: <http://dx.doi.org/10.1371/journal.pone.0042843> PMID: 22952617