



Androgen-sensitive hypertension associated with soluble guanylate cyclase alpha1 deficiency is mediated by 20-HETE

Citation

Vandenwijnngaert, S., A. C. Dordea, V. Garcia, R. E. Tainsh, D. I. Nathan, M. J. Raheer, K. Allen, et al. 2015. "Androgen-sensitive hypertension associated with soluble guanylate cyclase alpha1 deficiency is mediated by 20-HETE." *BMC Pharmacology & Toxicology* 16 (Suppl 1): A97. doi:10.1186/2050-6511-16-S1-A97. <http://dx.doi.org/10.1186/2050-6511-16-S1-A97>.

Published Version

doi:10.1186/2050-6511-16-S1-A97

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:22857027>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

MEETING ABSTRACT

Open Access

Androgen-sensitive hypertension associated with soluble guanylate cyclase alpha1 deficiency is mediated by 20-HETE

Sara Vandenwijngaert¹, Ana C Dordea¹, Victor Garcia², Robert E Tainsh¹, Daniel I Nathan¹, Michael J Raheer¹, Kaitlin Allen¹, Fan Zhang², Wolfgang S Lieb¹, Sarah Mikelman¹, Andrew Kirby³, Christine Stevens³, Robrecht Thoonen¹, Allyson Hindle¹, Patrick Y Sips⁴, Rajeev Malhotra¹, Mark J Daly³, Peter Brouckaert⁵, Kenneth D Bloch¹, Michal Schwartzman², Emmanuel S Buys^{1*}

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications Trier, Germany. 19-21 June 2015

Background

Dysregulated nitric oxide (NO) signaling contributes to the pathogenesis of hypertension. Previously, we reported gender- and strain-specific hypertension in mice deficient in the $\alpha 1$ -subunit of the NO receptor soluble guanylate cyclase ($sGC\alpha 1^{-/-}$): male mice on an Sv129/J (S6) but not a C57BL6/J (B6) background are hypertensive.

Methods and results

Via linkage analysis, we identified a quantitative trait locus (QTL) associated with elevated blood pressure in male $sGC\alpha 1^{-/-}$ S6 mice. This QTL encompasses *CYP4a12a*, encoding the predominant murine synthase of the vasoconstrictor 20-hydroxyeicosatetraenoic acid (20-HETE). Renal expression of *CYP4a12a* was strain-, gender-, and testosterone-dependent: *CYP4a12a* gene expression was higher in male WT and $sGC\alpha 1^{-/-}$ S6 mice than in female S6 mice, or than in male and female, WT and $sGC\alpha 1^{-/-}$ B6 mice, higher in testosterone-treated S6 mice than in vehicle-treated S6 mice, and higher in sham-operated S6 mice than orchietomized S6 mice. Also, 20-HETE levels were higher in renal preglomerular microvessels of male $sGC\alpha 1^{-/-}$ S6 than of $sGC\alpha 1^{-/-}$ B6 mice. Furthermore, the 20-HETE antagonist 20-6,15-HEDGE lowered blood pressure in male $sGC\alpha 1^{-/-}$ S6 but not WT mice. Finally, the more significant impairment of acetylcholine-induced relaxation of renal interlobar arteries in male $sGC\alpha 1^{-/-}$ S6

than $sGC\alpha 1^{-/-}$ B6 mice, in male $sGC\alpha 1^{-/-}$ S6 than WT S6 mice, and in male sham-operated $sGC\alpha 1^{-/-}$ S6 mice than orchietomized $sGC\alpha 1^{-/-}$ S6 mice was rescued by 20-6,15-HEDGE.

Conclusion

Gender- and strain-specific hypertension and vascular dysfunction in $sGC\alpha 1^{-/-}$ S6 mice is associated with elevated *CYP4a12a* expression and 20-HETE levels, and is abrogated by antagonizing 20-HETE. These results corroborate our hypothesis that testosterone-induced *CYP4a12a* expression and a concomitant increase in 20-HETE production contribute to the hypertension associated with impaired NO-cGMP signaling and that *CYP4a12a* represents a candidate blood pressure modifying gene in the context of deficient NO-sGC signaling.

Authors' details

¹Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA. ²Department of Pharmacology, New York Medical College, Valhalla, NY 10595, USA. ³Center for Human Genetic Research, Department of Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA. ⁴Division of Cardiovascular Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA 02115, USA. ⁵Department for Biomedical Molecular Biology, Ghent University, Ghent, 9000, Belgium.

Published: 2 September 2015

doi:10.1186/2050-6511-16-S1-A97

Cite this article as: Vandenwijngaert et al.: Androgen-sensitive hypertension associated with soluble guanylate cyclase alpha1 deficiency is mediated by 20-HETE. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A97.

* Correspondence: ebuys@mgh.harvard.edu

¹Department of Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA
Full list of author information is available at the end of the article