



Citrus Fruit Intake Substantially Reduces the Risk of Esophageal Cancer: A Meta-Analysis of Epidemiologic Studies

Citation

Wang, Anqiang, Chengpei Zhu, Lilan Fu, Xueshuai Wan, Xiaobo Yang, Haohai Zhang, Ruoyu Miao, Lian He, Xinting Sang, and Haitao Zhao. 2015. "Citrus Fruit Intake Substantially Reduces the Risk of Esophageal Cancer: A Meta-Analysis of Epidemiologic Studies." *Medicine* 94 (39): e1390. doi:10.1097/MD.0000000000001390. <http://dx.doi.org/10.1097/MD.0000000000001390>.

Published Version

doi:10.1097/MD.0000000000001390

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:23473975>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Citrus Fruit Intake Substantially Reduces the Risk of Esophageal Cancer

A Meta-Analysis of Epidemiologic Studies

Anqiang Wang, MD, Chengpei Zhu, MD, Lilan Fu, MD, Xueshuai Wan, MD, Xiaobo Yang, MD, Haohai Zhang, MD, Ruoyu Miao, MD, Lian He, MD, Xinting Sang, MD, and Haitao Zhao, MD

Abstract: Many epidemiologic studies indicate a potential association between fruit and vegetable intake and various cancers. The purpose of this meta-analysis is to investigate the association between citrus fruit intake and esophageal cancer risk. The authors conducted a comprehensive search on PubMed, EMBASE, and the Cochrane Library from inception until July 2014. Studies presenting information about citrus intake and esophageal cancer were analyzed. The authors extracted the categories of citrus intake, study-specific odds ratio or relative risk, and the *P* value and associated 95% confidence intervals for the highest versus lowest dietary intake of citrus fruit level. The association was quantified using meta-analysis of standard errors with a random-effects model. Thirteen case-control studies and 6 cohort studies were eligible for inclusion. Citrus intake may significantly reduce risk of esophageal cancer (summary odds ratio = 0.63; 95% confidence interval = 0.52–0.75; *P* = 0), without notable publication bias (intercept = –0.79, *P* = 0.288) and with significant heterogeneity across studies (*I*² = 52%). The results from epidemiologic studies suggest an inverse association between citrus fruit intake and esophageal cancer risk. The significant effect is consistent between case-control and cohort studies. Larger prospective studies with rigorous methodology should be considered to validate the association between citrus fruits and esophageal cancer.

(*Medicine* 94(39):e1390)

Abbreviations: CIs = confidence intervals, EAC = esophageal adenocarcinoma, OR = odds ratio, RR = relative risk, SCC = squamous cell carcinoma, SEs = standard errors.

INTRODUCTION

Esophageal cancer, including squamous cell carcinoma (SCC) and esophageal adenocarcinoma (EAC), is a serious

Editor: Yung-Hsiang Chen.

Received: March 22, 2015; revised and accepted: July 24, 2015.

From the Department of Liver Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China (AW, CZ, LF, XW, XY, HZ, LH, XS, HZ); and Liver Center and The Transplant Institute, Department of Medicine, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA (RM).

Correspondence: Haitao Zhao, Xinting Sang, MD, Chinese Academy of Medical Sciences and Peking Union Medical College (CAMS & PUMC), Beijing, China (e-mail: zhaoh@pumch.cn).

Drs Anqiang Wang, Chengpei Zhu, and Lilan Fu contributed equally to the writing of this article.

The authors declare that they have no conflicts of interest.

Copyright © 2015 Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution License 4.0, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. ISSN: 0025-7974

DOI: 10.1097/MD.0000000000001390

malignancy with a poor prognosis in the majority of cases.^{1,2} SCC is the predominant form of esophageal carcinoma worldwide, but a shift in epidemiology has been seen in some countries and regions like Australia, UK, USA, and western Europe, where the incidence of EAC has exceeded that of SCC.³ Every year, >450,000 people worldwide are diagnosed with esophageal cancer and the incidence is rapidly increasing.^{3,4} It is the eighth most common cancer, and the sixth most common cause of cancer-related deaths worldwide with developing nations making up >80% of total cases and deaths.^{5,6} The mortality from these cancers is high and the response to treatments during advanced stages is poor, so effectively reducing the chances of exposure to relative risk factors will have an important impact on the incidence of esophageal cancer.

Cigarettes, red meat, alcohol, hot tea, pickled vegetables, low intake of fresh fruits and vegetables, and low socioeconomic status are associated with a higher risk of SCC.^{7–10} Barrett esophagus is clearly recognized as a risk factor for EAC, with other factors including gastroesophageal reflux disease, acid-suppressive medication use, obesity, tobacco use, and processed meat.^{11–14} Some foods can reduce the incidence of esophageal cancer.^{9,15–18} Many researchers conducted meta-analyses on diet and esophageal cancer. The study by Coleman et al¹⁹ suggested that dietary fiber may protect against esophageal carcinogenesis, especially esophageal adenocarcinoma. Zhu et al²⁰ found that meat consumption is associated with the risk of esophageal cancer. The intake of red meat is likely to increase the esophageal SCC risk and the processed meat may increase esophageal adenocarcinoma risk; however, the consumption of fish may not be associated with esophageal cancer incidence. This phenomenon may be explained by the effects of various micronutrients such as folate, B vitamins, antioxidants, lutein, and carotenoids.^{21–24}

Citrus fruits include oranges, tangerines, grapefruits, lemons, and limes. They include several components, including flavonoids, folate, carotenoids, and vitamin C,^{25–26} which have protective effects against cancer. Previous studies have suggested that citrus intake may improve the incidence of various cancers including pancreatic, breast, and prostate cancers.^{27–29} Consequently, we hypothesize that citrus intake is associated with a reduced risk of esophageal cancer. Epidemiologic evidence from cohort and case-control studies on this association has not yet been summarized. Therefore, we conducted a meta-analysis to explore this hypothesis.

STUDY CHARACTERISTICS

Search Strategy

A computerized search of the English language literature on citrus fruits and esophageal cancer yielded no relevant

publications from inception to July 2014. We, therefore, decided to use the key words “fruit” and “citrus.” The search terms were ([esophagus] OR [esophageal]) AND ([cancer] OR [tumor] OR [carcinoma]) AND (‘citrus’itrus OR ‘fruit’ruits). We limited the search to human adults without language restrictions. We searched the 3 major electronic databases: PubMed, EMBASE, and The Cochrane Library. Additionally, we reviewed the references from retrieved articles for additional studies. Furthermore, ethical approval was not necessary because our article is a review.

Study Selection

The included studies²⁹ had to be epidemiologic studies such as case–control and cohort studies. The studies concerning human that addressed the association between citrus intake and incidence of esophageal cancer were collected; however, if the study provides no original data or insufficient information on the odds ratio (OR) or relative risk (RR), and their corresponding 95% confidence intervals (CIs), we excluded it. The studies not measuring the intake of citrus fruits or citrus juice at the individual level are not eligible. The instrument of assessment of citrus intake is questionnaire. Two independent reviewers read the abstracts or full-text articles to assess the eligibility of studies in a standardized manner. We resolved the disagreement by consensus.

Data Abstraction

We extracted important information from all eligible studies. They included study design, country of origin, years of publication, origin of control, number of cases and control, sex distribution, types of citrus fruits, types of cancer, comparison of exposure level, and potential confounding variables adjusted. The estimates of OR/RR, their associated 95% CIs, and *P* values were also extracted by us. If separate researches based on the same population were published, we selected the article containing more complete information for inclusion.

Statistical Analyses

We extracted the study specific OR/RR and 95% CIs for highest versus lowest intake of citrus fruits from every study. And we calculated the standard error (SE) of the log OR/RR by using the following equation: $SE = (\ln[OR/RR_{upper}] - \ln[OR/RR_{lower}]) \div 3.92$. Then, we summarized the overall OR and CI by using general variance-based methods³⁰ of RevMan 5.0. For studies that provided OR/RR by cancer subtypes,^{15,31} we used a random-effects model to obtain a pooled estimate from the individual study (Table 1). We adopted the Newcastle-Ottawa Scale to evaluate research quality and defined them as high, middle, and low quality by score 7 to 9, 4 to 6, 1 to 3, respectively. The Grades of Recommendation, Assessment, Development, and Evaluation working group system of rating quality of evidence also were used to evaluate the research quality.

The value of I^2 was used to evaluate the extent of heterogeneity derived from study differences rather than chance.⁴⁴ The smaller value I^2 suggested less obvious heterogeneity. We used the random-effects model to calculate the summary OR and its 95% CI⁴⁵ with suspecting heterogeneity. We evaluated the impact of the changes on pooled ORs by study design, cancer subtypes, geographical location, source of controls, research quality, and some adjusted confounders such as alcohol and body mass index as prior hypotheses to explain heterogeneity through subgroup analyses and meta-regression analyses. Sensitivity analyses were conducted by removing 1 study from

all studies to evaluate the impact on the pooled ORs and heterogeneity. We can, therefore, evaluate whether the results are stable. In an attempt to detect publication bias, we visually examined asymmetry in a funnel plot. We conducted Begg and Egger test to assess whether there is an obvious publication. We considered the funnel plot to be asymmetrical if the intercept of the regression line deviated from zero with $P < 0.10$. If the test suggests an obvious publication bias, we would conduct the trim and fill analysis to further verify.

We used the Cochrane Collaboration software (Oxford, UK) to analyze the extracted data with fixed or random-effects model analysis.⁴⁶ STATA (StataCorp, College Station, TX) was used to conduct the Egger and Begg regression asymmetry test by using the metabis command.⁴⁷ We conducted the trim and fill analysis to observe whether the results are stable and evaluate the publication bias.

RESULTS

Search Results

The computerized search yielded 433 references, of which 112 were included after abstract review. Citation search identified another 715 articles. Of the 827 articles that were obtained for full-text review, we excluded 808 articles based on the exclusion criteria. In particular, the result of Tuyns et al⁴⁸ published in 1987 was replaced by Tuyns et al³² published in 1983, as it shared the same database. The result of De Stefani et al⁴⁹ published in 2003 was replaced by De Stefani et al⁴⁰ published in 2005, as the latter expanded the sample size based on the former population.

A total of 19 articles were included in the meta-analysis, including 6 cohort studies^{15,16,61,41-43} and 13 case–control studies^{9,10,17,18,32-39} (Figure 1).

TABLE 1. Logarithmic OR or RR (Log[OR/RR]) and Its SE for the Meta-Analysis

Author	Categories	Log (OR/RR)	SE
Tuyns 1983 ³²	Total	−0.4780358	0.1101486
Brown 1988 ³³	Men	−0.6931472	0.2802582
Cheng et al 1992 ¹⁰	Total	−2.419119	0.6944997
Castelletto, 1994 ³⁴	Total	0.4700036	0.3455474
Brown et al 1995 ³⁵	Men	−0.3566749	1.2199227
Zhang 1997 ³⁶	Total	−0.1053605	0.1972423
Brown et al 1998 ¹⁸	Men	−0.2231435	1.97403993
Launoy 1998 ³⁷	Men	−0.6161861	0.2530941
Levi 2000 ¹⁷	Total	−1.514128	0.4570815
Bosetti 2000 ³⁸	Total	−0.8675005	0.2662765
Chen 2002 ³⁹	Total	−0.7339692	0.4224383
De Stefani 2005 ⁴⁰	Total	−1.272966	0.3314497
Sapkota 2008 ⁹	Total	−0.2744368	0.2610686
Boeing 2006 ⁴¹	Total	−0.2744368	0.2029495
Gonzalez 2006 ⁴²	Total	−0.3147107	0.3205151
Freedman 2007 ³¹	Total	−0.2484614	0.2482107
Yamaji 2008 ¹⁶	Men	−0.2484614	0.2441614
Li et al 2010 ⁴³	Total	−0.3424903	0.2531607
Steevens 2011 ¹⁵	Men	−0.2231435	0.2729166

OR = odds ratio, RR = relative risk; SE = standard error.

*The estimate was obtained by fixed-effect model using the hazard ratio given by sex control.




FIGURE 1. Flowchart of the searching and review of literatures.

Study Characteristics

Some details of the selected studies are shown in Tables 2 and 3. All articles were published in English. Six studies were conducted among residents of the United States,^{18,31,33,35,36,39} 1 in Italy,³⁸ 2 in Japan,^{16,43} 2 in France,^{32,37} 3 in Europe,^{9,41,42} and the remaining 5 in China,¹⁰ Argentina,³⁴ Switzerland,¹⁷ Uruguay,³² and the Netherlands.¹⁵ Two of the studies recruited participants in the 1980s, 5 in the 1990s, and 12 between 2000 and 2011.

The factor of age was adjusted in all of the studies except Brown et al⁵⁰ The confounding variables that were adjusted in different studies were presented in detail in Tables 2 and 3. For all of the studies, the relationship between intake of citrus fruits and esophageal cancer was not primary hypothesis and the citrus fruits were often included in a broader dietary evaluation. The ranges of adjusted ORs/RRs were from 0.089 to 1.6 and only 5 studies^{31,32,33,37,39} reached the usual threshold of $P=0.05$ in the association between citrus fruits and esophageal cancer.

Heterogeneity and Pooled Results

There was no significant heterogeneity among the study results ($I^2=52\%$; $P=0.005$). Overall summary OR using the random-effects model showed a 37%, statistically significant reduction in risk of esophageal cancer associated with citrus fruits intake (summary OR = 0.63; 95% CI = 0.52–0.75). The subgroup of case-control studies (summary OR = 0.54; 95% CI = 0.4–0.72; $I^2=64.2\%$; $P=0.001$) and the subgroup of

cohort studies (summary OR = 0.76; 95% CI = 0.62–0.93; $I^2=0\%$; $P=1$) showed a respective 46% and 24% statistically significant reduction in risk of esophageal cancer associated with citrus fruits intake (Figure 2). In subgroup analyses defined by study type, cancer subtype, geographical location, source of controls, research quality, and adjusted confounders, citrus intake was inversely associated with risk of esophageal cancer in most subgroups, with no evidence of significant heterogeneity between subgroups with meta-regression analyses. (Table 4).

Publication Bias

No publication bias was observed in the selected studies. Visualization of Begg funnel plot was symmetrical (Figure 3). Formal testing using the Egger method supports the notion that there was no publication bias (intercept = -0.79, $P=0.288$); however, the result of Begg test suggested an obvious publication bias ($P=0.046$). And the outcome of trim and fill analysis demonstrated that there was no publication bias.

DISCUSSION

The overall summary OR in our study presents an inverse association between citrus fruits and esophageal cancer (summary OR = 0.63; 95% CI = 0.52–0.75; $P=0$). The result is supported by the strengths of our review, which includes a systematic literature search, strict selection criteria, comprehensive data abstraction, and rigorous statistical analysis. Additionally, the results of similar reviews about the association between citrus fruits and other cancers^{26–29} are encouraging.

TABLE 2. Summary of Case–Control Studies Included in the Meta-Analysis

Study/Years of Publication	Country	No. of Case/Control	Sources of Controls	Types of Cancer	Exposure	Adjusted Factors	Comparison of Exposure Level	Adjusted OR/RR (95% CI)	P value	NOS Scale
Tuyns 1983 ³²	France	1246/1976	Population	EC	Citrus and juices	Age, alcohol consumption, tobacco smoking, and urban or rural residence	Yes vs no	0.62[0.50–0.77]	—	5
Brown et al 1988 ³³	United States	207/422	Hospital	EC	citrus and juices	use of cigarettes and alcohol	Q3 vs Q1	0.5[0.29–0.87]	≤0.01	5
Cheng et al 1992 ¹⁰	Hong Kong	400/1598	Population	EC	Citrus fruits	Age, educational attainment, place of birth, hot drinks or soups, green leaf vegetables, pickled vegetables, tobacco, alcohol, whether had meals or eating out	Q6 vs Q1	0.089[0.02–0.35]	—	7
Castelletto et al 1994 ³⁴	Argentina	131/262	Hospital	SCC	Citrus fruits	Design variables, age, sex, hospital, education, average number of cigarettes/day, alcohol consumption	Q3 vs Q1	1.6[0.8–3.1]	—	6
Brown et al 1995 ³⁵	United States	174/750	Population	EAC	Citrus fruits	barbecued meat, potatoes, raw vegetables, cooked vegetables	Q4 vs Q1	0.7[0.06–7.65]	0.77	7
Zhang et al 1997 ³⁶	United States	95/132	Hospital	EAC	Citrus fruits	Age, area, smoking, liquor use, income, calories from food, and BMI	Q4 vs Q1	0.9[0.61–1.32]	0.532	8
Brown et al 1998 ¹⁸	United States	114/681	Population	SCC	Citrus fruits	Age, sex, race, education, total dietary intake of calories, pack-years of smoking, alcohol use, and BMI	Q4 vs Q1	0.8[0.02–38.32]	0.91	7
Launoy et al 1998 ³⁷	France	208/399	Hospital	SCC	Citrus fruits	Age, area, smoking, alcohol, and food calories	Q4 vs Q1	0.54[0.33–0.89]	<0.05	6
Bosetti et al 2000 ³⁸	Italy	304/743	Hospital	SCC	Citrus fruits	Age, interviewer, smoking, beer, amised aperitifs, hot Calvados, whisky, total alcohol, and total energy intake	Q5 vs Q1	0.42[0.25–0.71]	—	7
Levi et al 2000 ¹⁷	Switzerland	101/327	Hospital	EC	Citrus fruits	Age, sex, area of residence, education, tobacco smoking, alcohol, drinking, and nonalcohol energy	Q3 vs Q1	0.22[0.09–0.54]	—	7
Chen et al 2002 ³⁹	United States	124/449	Population	EAC	Citrus fruits	Age, sex, Education, Smoking, alcohol, and nonalcohol total energy intake	Q4 vs Q1	0.48[0.21–1.10]	0.03	7
De Stefani et al 2005 ⁴⁰	Uruguay	200/400	Hospital	SCC	citrus fruits	Age, sex, energy intake, respondent type, BMI, alcohol use, tobacco use, education	Q4 vs Q1	0.28[0.15–0.54]	0.0001	6
Sapkota et al 2008 ⁹	Japan	116/	Hospital	SCC	Citrus fruits	Age, sex, residence, urban/rural status, birthplace, education, smoking status, number of cigarettes smoked per day, years since quit, alcohol drinking, mate temperature, stewed meat, and total energy intake	Q3 vs Q1	0.78[0.48–1.26]	0.21	6

BMI = body mass index, CI = confidence interval, EAC = esophageal adenocarcinoma, EC = esophageal carcinoma, OR = odds ratio, RR = relative risk, SCC = squamous cell carcinoma.

TABLE 3. Summary of Cohort Studies Included in the Meta-Analysis

Study/Years of Publication	Country	No. of Case/person-years	Sources of Controls	Subtype of Cancer	Exposure	Adjusted Factors	Comparison of Exposure Level	Adjusted OR/RR (95% CI)	P value	NOS Scale
Boeing et al 2006 ⁴¹	European	352/2,182,560	Population	SCC	Citrus fruits	Age, sex, center, BMI, energy from fat sources, energy from nonfat sources, education, smoking status categories	Q5 vs Q1	0.76[0.51–1.13]	0.129	8
Gonzalez et al 2006 ⁴²	European	65/3,110,034	Population	EAC	Citrus fruits	Sex, height, weight, education level, tobacco smoking, cigarette smoking intensity, work and leisure, physical activity, alcohol intake, energy intake, red meat intake, and processed meat intake	Q3 vs Q1	0.73[0.39–1.37]	0.22	8
Freedman et al 2007 ³¹	United States	103/2,193,751	Population	SCC	Citrus fruits	Sex, age at entry into cohort, BMI, education, alcohol intake, cigarette-smoke-dose, vigorous physical activity, usual activity throughout the day, and total energy	Q3 vs Q1	0.58[0.34–0.99]	0.046	7
Freedman et al 2007 ³¹	United States	233/2,193,751	Population	EAC	Citrus fruits	Sex, age at entry into cohort, BMI, education, alcohol intake, cigarette-smoke-dose, vigorous physical activity, usual activity throughout the day, and total energy	Q3 vs Q1	0.96[0.69–1.35]	>0.05	7
Yamaji et al 2008 ¹⁶ Li et al 2010 ⁴³	European Japan	116/297,651 151/329,985	Population population	SCC EC	Citrus fruits Citrus	Age, alcohol, smoking (continuous variable), sex (for total participants), job status, year of education, BMI, physical activity, energy intake, cigarette, alcohol	Q3 vs Q1 Q3 vs Q1	0.78[0.48–1.25] 0.71[0.43–1.16]	0.21 0.178	9 9
Steevens et al 2011 ¹⁵	Netherlands	101/50,785	Population	SCC	Citrus fruits	Age (years), sex, cigarette smoking (current smoking [yes/no]), frequency (number of cigarettes per day), duration (number of years), alcohol consumption (g ethanol/day), total vegetable intake and all other fruits, consumption of red meat (g/day), consumption of fish (g/day)	Q5 vs Q1	0.54[0.27–1.07]	0.38	9
Steevens et al 2011 ¹⁵	Netherlands	144/50,785	Population	EAC	Citrus fruits	Age, sex, cigarette smoking, frequency, duration, alcohol consumption, total vegetable intake and all other fruits, consumption of red meat, consumption of fish	Q5 vs Q1	0.97[0.90–1.04]	0.37	9

BMI = body mass index, CI = confidence interval, EAC = esophageal adenocarcinoma, EC = esophageal carcinoma, OR = odds ratio, RR = relative risk, SCC = squamous cell carcinoma of esophagus.




FIGURE 2. Summary estimates of the association between citrus intake and esophageal cancer risks sorted by effect estimate. CI = confidence interval; df = degree of freedom; χ^2 = chi-square statistic; I^2 = the percentage of total variation across studies that is due to heterogeneity rather than change; fixed = using fixed-effect model.

Nevertheless, some limitations of our study should be taken into consideration. First, the considerable variables within observational studies made the outcome more likely to be suspicious.⁵⁰ Furthermore, the included studies were evaluated to be low quality using the Grades of Recommendation, Assessment, Development and Evaluation system.⁵¹ These inherent drawbacks of observational study make outcomes more conservative.⁵² In addition, most of the included studies were not originally designed to evaluate citrus fruits and esophageal cancer, which possibly turns the pooled result into a simple summary.⁵⁰ Measurement errors resulting from citrus intake should also be considered because of diversity of consumption patterns. Therefore, in the process of food intake measurement, various confounding factors made it difficult to obtain accurate dietary exposure information.²⁷ Most case-control studies on diet are based on recent estimates of dietary intake,¹⁵ whereas the development of cancer after exposure to even a potent risk factor takes several decades.^{53,54}

We found heterogeneity among the included studies, which decreases the quality of evidence to very low quality.⁵¹ Although rigorous criteria would make selective studies homogeneous, these could give rise to an inclusion bias.⁵⁵ We excluded 2 studies^{48,49} sharing the same population, because the inclusion of duplicated data may lead to overestimation of exposure effects.^{29,56} Cancer deaths reflect failure of treatment as well as the occurrence of the cancer.⁵⁷ Therefore, incidence rates are preferable as an early indicator of the impact of a risk

factor. After careful screening of the eligible studies without CI or original data,^{18,35} we calculated the corresponding SE by the ORs and exact *P* values.⁵⁸ The most appropriate way of handling the selection of studies is to perform sensitivity analyses with regard to the different possible entry criteria.⁵⁵ Considering that the wide confidence interval of studies^{18,35} may obviously affect our outcome, we conducted sensitivity analysis.²⁹ The analytic result showed that the studies have no apparent impact on the overall outcome. We also omitted 1 study⁵⁹ that provided no citrus intake measurement. Methodology is a significant source of heterogeneity,⁵⁰ so we performed subgroup analysis to verify the effect of study designs on heterogeneity. Figure 2 shows that study design causes heterogeneity. In the 6 cohort studies, the I^2 of summary OR was 0%, because prospective studies can avoid recall and selection biases. In the 13 case-control studies, the I^2 of summary OR was 64.2%. Both study designs demonstrate that citrus intake could reduce the incidence of esophageal cancer with summary OR 0.57 (CI 0.4–0.72), 0.76 (CI 0.62–0.93) for the case-control study and cohort study, respectively. The discrepancies between study results can be explained by recall and selection biases in case-control studies and by imprecise dietary measurements and limited variability of dietary intake in cohort studies.^{27,60} To further explore the source of heterogeneity, we conducted subgroups analyses and meta-regression analyses by many factors such as cancer subtypes, geographical location, source of controls, research quality, and adjusted confounders.

TABLE 4. Subgroup Analyses of Citrus Intake and Risk of Esophageal Cancer, Sensitivity Analysis, Meta-Regression Analysis

Subgroups	No. of studies	OR (95% CI)	I ² (%)	P Heterogeneity	
				*	†
Study					
Case-control	13	0.538 (0.405–0.715)	64.2	0.001	0.162
Cohort	6	0.761 (0.623–0.929)	0	1	
Cancer subtype					
SCC	10	0.63 (0.48–0.82)	51	0.031	0.366
EAC	6	0.81(0.66–1)	0	0.458	
Geographical location					
Europe	8	0.62 (0.5–0.75)	33.9	0.158	0.592
USA	6	0.72 (0.56–0.93)	0	0.549	
Asia	3	0.48 (0.21–1.08)	76.9	0.013	
Latin America	2	0.67 (0.12–3.68)	92.5	0	
Controls					
Population	10	0.65 (0.54–0.78)	21	0.249	0.866
Hospital	9	0.6 (0.41–0.86)	69.3	0.001	
Research quality					
High	6	0.61 (0.44–0.86)	66.1	0.011	0.945
Middle	13	0.63 (0.5–0.8)	45.5	0.037	
Adjustment for confounders					
Alcohol					
Yes	17	0.61 (0.5–0.75)	56.5	0.002	0.615
No	2	0.76 (0.51–1.12)	0	0.947	
BMI					
Yes	6	0.57 (0.44–0.73)	62.1	0.002	0.263
No	13	0.77 (0.62–0.95)	0	0.005	
Sensitivity analysis					
All studies except Cheng et al (1992) ¹⁰	18	0.65 (0.55–0.77)	41.9	0.032	

BMI = body mass index, CI = confidence interval, EAC = esophageal adenocarcinoma, OR = odds ratio.

* Within each subgroup.

† Between subgroups with meta-regression analysis.

The pathogenesis and risk factors for different types of esophageal cancer^{18,21} vary widely, so exploring the impacts of citrus intake on these cancers is essential. Four of the included studies did not describe the specific cancer subtypes or included




FIGURE 3. Funnel plot of studies evaluating the association between citrus fruit intake and esophageal cancer risks. Dot lines are 95% pseudo-confidence intervals. SE = standard error; OR = odds ratio.

both subtypes. Table 4 shows the association between citrus and SCC (summary OR 0.63; CI 0.48–0.82) and EAC (summary OR 0.81; CI 0.66–1). The lack of overlapping confidence interval could partially explain the study heterogeneity. The forest plot (Figure 2) demonstrates that there is no overlap in CIs between 3 studies¹⁰ and the summary OR. Repeat meta-analysis of a new model excluded the study¹⁰ from all 19 selected articles was conducted.⁶¹ The level of heterogeneity decreased from high ($I^2 = 52%$) to low ($I^2 = 0$).

Citrus fruits include many bioactive components.^{25,62} Dietary antioxidants are emerging as potentially modifiable risk factors for EAC. High intake of beta-carotene may be associated with decreased risk of dysplastic Barrett esophagus, which is regarded as the precursor of EAC.⁶³ Some studies^{64,65} showed that carotenoids may be responsible nutritional factors (as nutritional scavengers) in the development of different malignant diseases including esophageal cancer.^{66,67} Carotenoids may intervene in cancer-related molecular pathways and the expression proteins involved in cell proliferation, differentiation, apoptosis and angiogenesis, carcinogen detoxification, DNA damage, and repair.⁶⁸ A related study indicates⁶⁹ that a high intake of vitamin C is associated with a reduced risk of EAC and reflux esophagitis. Antioxidants may also play a role in the pathogenesis of reflux esophagitis and EAC and may be more important in terms of progression rather than initiation of the disease process⁶⁹; however, low intake of vitamin C and E

correlates significantly with the development of SCC as well as EAC in males.^{70,71} Regarding the mechanism, researchers think that vitamin C could enhance the EGCG- and TF3-induced apoptosis in SPC-A-1 and Eca-109 cells via MAPK pathways.⁷² Additionally, folate and other dietary methyl group factors are implicated in the etiology of EAC and its precursors. Folate is implicated in carcinogenesis via effects on DNA synthesis, repair, and methylation.^{21,22} Some studies indicate that flavanone intake is inversely associated with SCC risk and may account for the protective effect of fruit, especially citrus fruits, on esophageal cancer.^{73,74} Because citrus fruits account for 90% of flavanone intake, the findings of Rossi et al⁷³ suggest that flavanones may play a role in the protective effect of citrus fruits on esophageal cancer. Therefore, the basic research of mechanisms flavanones protect against esophageal cancer are worth studying. Although the results are exciting, we have to taken in account the interaction between medicines and fruits. The research by Bailey et al⁷⁵ suggested that there exist adverse reactions when grape is combined with some drugs.

Our review demonstrates that citrus fruit intake could reduce the incidence of esophageal cancer by 37% based on published results of epidemiologic studies. The trends are consistent between case-control studies and cohort studies; however, considering the drawbacks mentioned above, our conclusions should be taken cautiously. There are no relevant studies that provide explicit evidence for the inconsistency between SCC and EAC. The low quantity of EAC cases and the limitations of meta-analysis are responsible for the results. Therefore, larger studies with rigorous and prospective methodology should be considered to validate the association between citrus fruits and esophageal cancer. It is still unknown which components in citrus fruits have an effect on esophageal cancer prevention. Our conclusion may encourage researchers to further explore the protective elements and potential mechanisms, which may contribute to reducing the esophageal cancer risk. We hope further research will explore this issue.

REFERENCES

1. Esophageal cancer: epidemiology, pathogenesis and, prevention. *Nat Clin Pract Gastroenterol Hepatol*. 2008;5:517–526doi: 10.1038/ncpgasthep1223.
2. Zhang Y. Epidemiology of esophageal cancer. *World J Gastroenterol*. 2013;19:5598–5606doi: 10.3748/wjg.v19.i34.5598.
3. Pennathur A, Gibson MK, Jobe BA, et al. Oesophageal carcinoma. *Lancet*. 2013;381:400–412doi: 10.1016/s0140-6736(12)60643-6.
4. Chen Y, Yu C, Li Y. Physical activity and risks of esophageal and gastric cancers: a meta-analysis. *PLoS One*. 2014;9:e88082doi: 10.1371/journal.pone.0088082.
5. Napier KJ, Scheerer M, Misra S. Esophageal cancer: a review of epidemiology, pathogenesis, staging workup and treatment modalities. *World J Gastrointest Oncol*. 2014;6:112–120doi: 10.4251/wjgo.v6.i5.112.
6. Herszenyi L, Tulassay Z. Epidemiology of gastrointestinal and liver tumors. *Eur Rev Med Pharmacol Sci*. 2010;14:249–258.
7. Lindblad M, Rodriguez LA, Lagergren J. Body mass, tobacco and alcohol and risk of esophageal, gastric cardia, and gastric non-cardia adenocarcinoma among men and women in a nested case-control study. *Cancer Causes Control*. 2005;16:285–294doi: 10.1007/s10552-004-3485-7.
8. Choi Y, Song S, Song Y, et al. Consumption of red and processed meat and esophageal cancer risk: meta-analysis. *World J Gastroenterol*. 2013;19:1020–1029doi: 10.3748/wjg.v19.i7.1020.
9. Sapkota A, Hsu CC, Zaridze D, et al. Dietary risk factors for squamous cell carcinoma of the upper aerodigestive tract in central and eastern Europe. *Cancer Causes Control*. 2008;19:1161–1170doi: 10.1007/s10552-008-9183-0.
10. Cheng KK, Day NE, Duffy SW, et al. Pickled vegetables in the aetiology of oesophageal cancer in Hong Kong Chinese. *Lancet*. 1992;339:1314–1318.
11. Carr JS, Zafar SF, Saba N, et al. Risk factors for rising incidence of esophageal and gastric cardia adenocarcinoma. *J Gastrointest Cancer*. 2013;44:143–151doi: 10.1007/s12029-013-9480-z.
12. Lepage C, Drouillard A, Jouve JL, et al. Epidemiology and risk factors for oesophageal adenocarcinoma. *Dig Liver Dis*. 2013;45:625–629doi: 10.1016/j.dld.2012.12.020.
13. Salehi M, Moradi-Lakeh M, Salehi MH, et al. Meat, fish, and esophageal cancer risk: a systematic review and dose-response meta-analysis. *Nutr Rev*. 2013;71:257–267doi: 10.1111/nure.12028.
14. Spechler SJ. Barrett esophagus and risk of esophageal cancer: a clinical review. *JAMA*. 2013;310:627–636doi: 10.1001/jama.2013.226450.
15. Steevens J, Schouten LJ, Goldbohm RA, et al. Vegetables and fruits consumption and risk of esophageal and gastric cancer subtypes in the Netherlands Cohort Study. *Int J Cancer*. 2011;129:2681–2693doi: 10.1002/ijc.25928.
16. Yamaji T, Inoue M, Sasazuki S, et al. Fruit and vegetable consumption and squamous cell carcinoma of the esophagus in Japan: the JPHC study. *Int J Cancer*. 2008;123:1935–1940doi: 10.1002/ijc.23744.
17. Levi F, Pasche C, Lucchini F, et al. Food groups and oesophageal cancer risk in Vaud, Switzerland. *Eur J Cancer Prev*. 2000;9:257–263.
18. Brown LM, Swanson CA, Gridley G, et al. Dietary factors and the risk of squamous cell esophageal cancer among black and white men in the United States. *Cancer Causes Control*. 1998;9:467–474.
19. Coleman HG, Murray LJ, Hicks B, et al. Dietary fiber and the risk of precancerous lesions and cancer of the esophagus: a systematic review and meta-analysis. *Nutr Rev*. 2013;71:474–482.
20. Zhu HC, Yang X, Xu LP, et al. Meat consumption is associated with esophageal cancer risk in a meat- and cancer-histological-type dependent manner. *Dig Dis Sci*. 2014;59:664–673.
21. Sharp L, Carsin AE, Cantwell MM, et al. Intakes of dietary folate and other B vitamins are associated with risks of esophageal adenocarcinoma, Barrett's esophagus, and reflux esophagitis. *J Nutr*. 2013;143:1966–1973doi: 10.3945/jn.113.174664.
22. Jiao L, Kramer JR, Rugge M, et al. Dietary intake of vegetables, folate, and antioxidants and the risk of Barrett's esophagus. *Cancer Causes Control*. 2013;24:1005–1014doi: 10.1007/s10552-013-0175-3.
23. Ge XX, Xing MY, Yu LF, et al. Carotenoid intake and esophageal cancer risk: a meta-analysis. *Asian Pac J Cancer Prev*. 2013;14:1911–1918.
24. Abnet CC, Qiao YL, Dawsey SM, et al. Prospective study of serum retinol, beta-carotene, beta-cryptoxanthin, and lutein/zeaxanthin and esophageal and gastric cancers in China. *Cancer Causes Control*. 2003;14:645–655.
25. Silalahi J. Anticancer and health protective properties of citrus fruit components. *Asia Pac J Clin Nutr*. 2002;11:79–84.
26. Fountoulakis A, Martin IG, White KL, et al. Plasma and esophageal mucosal levels of vitamin C: role in the pathogenesis and neoplastic progression of Barrett's esophagus. *Dig Dis Sci*. 2004;49:914–919.
27. Bae JM, Lee EJ, Guyatt G. Citrus fruit intake and pancreatic cancer risk: a quantitative systematic review. *Pancreas*. 2009;38:168–174doi: 10.1097/MPA.0b013e318188c497.

28. Bae JM, Lee EJ, Guyatt G. Citrus fruits intake and prostate cancer risk: a quantitative systematic review. *J Prev Med Public Health*. 2008;41:159–164.
29. Bae JM, Lee EJ, Guyatt G. Citrus fruit intake and stomach cancer risk: a quantitative systematic review. *Gastric Cancer*. 2008;11:23–32doi: 10.1007/s10120-007-0447-2.
30. Deeks J, Altman DG, Bradburn MJ. Statistical methods for examining heterogeneity and combining results from several studies in meta-analysis. In: Egger M, Smith DG, Altman DG, eds. *Systematic Reviews in Health Care*. 2nd ed London, England: BMJ Books; 2007:285–312.
31. Freedman ND, Park Y, Subar AF, et al. Fruit and vegetable intake and esophageal cancer in a large prospective cohort study. *Int J Cancer*. 2007;121:2753–2760doi: 10.1002/ijc.22993.
32. Tuyns AJ. Protective effect of citrus fruit on esophageal cancer. *Nutr Cancer*. 1983;5:195–200doi: 10.1080/01635588309513796.
33. Brown LM, Blot WJ, Schuman SH, et al. Environmental factors and high risk of esophageal cancer among men in coastal South Carolina. *J Natl Cancer Inst*. 1988;80:1620–1625.
34. Castelletto R, Castellsague X, Munoz N, et al. Alcohol, tobacco, diet, mate drinking, and esophageal cancer in Argentina. *Cancer Epidemiol Biomarkers Prev*. 1994;3:557–564.
35. Brown LM, Swanson CA, Gridley G, et al. Adenocarcinoma of the esophagus: role of obesity and diet. *J Natl Cancer Inst*. 1995;87:104–109.
36. Zhang ZF, Kurtz RC, Yu GP, et al. Adenocarcinomas of the esophagus and gastric cardia: the role of diet. *Nutr Cancer*. 1997;27:298–309doi: 10.1080/01635589709514541.
37. Launoy G, Milan C, Day NE, et al. Diet and squamous-cell cancer of the oesophagus: a French multicentre case-control study. *Int J Cancer*. 1998;76:7–12.
38. Bosetti C, La Vecchia C, Talamini R, et al. Food groups and risk of squamous cell esophageal cancer in northern Italy. *Int J Cancer*. 2000;87:289–294.
39. Chen H, Ward MH, Graubard BI, et al. Dietary patterns and adenocarcinoma of the esophagus and distal stomach. *Am J Clin Nutr*. 2002;75:137–144.
40. De Stefani E, Boffetta P, Deneo-Pellegrini H, et al. The role of vegetable and fruit consumption in the aetiology of squamous cell carcinoma of the oesophagus: a case-control study in Uruguay. *Int J Cancer*. 2005;116:130–135doi: 10.1002/ijc.20950.
41. Boeing H, Dietrich T, Hoffmann K, et al. Intake of fruits and vegetables and risk of cancer of the upper aero-digestive tract: the prospective EPIC-study. *Cancer Causes Control*. 2006;17:957–969doi: 10.1007/s10552-006-0036-4.
42. Gonzalez CA, Pera G, Agudo A, et al. Fruit and vegetable intake and the risk of stomach and oesophagus adenocarcinoma in the European Prospective Investigation into Cancer and Nutrition (EPIC-EURGAST). *Int J Cancer*. 2006;118:2559–2566doi: 10.1002/ijc.21678.
43. Li WQ, Kuriyama S, Li Q, et al. Citrus consumption and cancer incidence: the Ohsaki cohort study. *Int J Cancer*. 2010;127:1913–1922doi: 10.1002/ijc.25203.
44. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21:1539–1558.
45. Fleiss JL, Gross AJ. Meta-analysis in epidemiology, with special reference to studies of the association between exposure to environmental tobacco smoke and lung cancer: a critique. *J Clin Epidemiol*. 1991;44:127–139.
46. The Cochrane Collaboration. Review Manager Version 4.2 for Windows. Copenhagen, Denmark: The Nordic Cochrane Centre; 2003.
47. Stata Corporation. Stata Statistical Software. Special Edition. 8.2 for Windows. College Station, TX: Stata Corporation; 2004.
48. Tuyns AJ, Riboli E, Doornbos G, et al. Diet and esophageal cancer in Calvados (France). *Nutr Cancer*. 1987;9:81–92doi: 10.1080/01635588709513915.
49. De Stefani E, Deneo-Pellegrini H, Ronco AL, et al. Food groups and risk of squamous cell carcinoma of the oesophagus: a case-control study in Uruguay. *Br J Cancer*. 2003;89:1209–1214doi: 10.1038/sj.bjc.6601239.
50. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283:2008–2012.
51. Schunemann HJ, Jaeschke R, Cook DJ, et al. An official ATS statement: grading the quality of evidence and strength of recommendations in ATS guidelines and recommendations. *Am J Respir Crit Care Med*. 2006;174:605–614doi: 10.1164/rccm.200602-197ST.
52. Easterbrook PJ, Berlin JA, Gopalan R, et al. Publication bias in clinical research. *Lancet*. 1991;337:867–872.
53. Pou SA, Niclis C, Aballay LR, et al. Cancer and its association with dietary patterns in Cordoba (Argentina). *Nutr Hosp*. 2014;29:618–628doi: 10.3305/nh.2014.29.3.7192.
54. Michaud DS, Skinner HG, Wu K, et al. Dietary patterns and pancreatic cancer risk in men and women. *J Natl Cancer Inst*. 2005;97:518–524doi: 10.1093/jnci/dji094.
55. Egger M, Smith GD. Bias in location and selection of studies. *BMJ*. 1998;316:61–66.
56. Ioannidis JP, Trikalinos TA, Zintzaras E. Extreme between-study homogeneity in meta-analyses could offer useful insights. *J Clin Epidemiol*. 2006;59:1023–1032doi: 10.1016/j.jclinepi.2006.02.013.
57. Ezzat A. Book review: national cancer control programmes, policies and managerial guidelines. *Ann Saudi Med*. 1996;16:358.
58. Greenland S. Quantitative methods in the review of epidemiologic literature. *Epidemiol Rev*. 1987;9:1–30.
59. Navarro Silvera SA, Mayne ST, Risch H, et al. Food group intake and risk of subtypes of esophageal and gastric cancer. *Int J Cancer*. 2008;123:852–860doi: 10.1002/ijc.23544.
60. Bai Y, Yuan H, Li J, et al. Relationship between bladder cancer and total fluid intake: a meta-analysis of epidemiological evidence. *World J Surg Oncol*. 2014;12:223doi: 10.1186/1477-7819-12-223.
61. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557–560doi: 10.1136/bmj.327.7414.557.
62. Parmar HS, Dixit Y, Kar A. Fruit and vegetable peels: paving the way towards the development of new generation therapeutics. *Drug Discov Ther*. 2010;4:314–325.
63. Ibiebele TI, Hughes MC, Nagle CM, et al. Dietary antioxidants and risk of Barrett's esophagus and adenocarcinoma of the esophagus in an Australian population. *Int J Cancer*. 2013;133:214–224doi: 10.1002/ijc.28016.
64. Stice CP, Wang XD. Carotenoids and alcoholic liver disease. *Hepatobiliary Surg Nutr*. 2013;2:244–247doi: 10.3978/j.issn.2304-3881.2013.10.01.
65. Peng HC, Chen YL, Yang SY, et al. The antiapoptotic effects of different doses of beta-carotene in chronic ethanol-fed rats. *Hepatobiliary Surg Nutr*. 2013;2:132–141doi: 10.3978/j.issn.2304-3881.2013.06.08.
66. Rumi G Jr, Matus Z, Toth G, et al. Changes of serum carotenoids in patients with esophageal, gastric, hepatocellular, pancreatic and colorectal cancer. *J Physiol, Paris*. 2001;95:239–242.

67. Hammerich L, Tacke F. Eat more carrots? Dampening cell death in ethanol-induced liver fibrosis by beta-carotene. *Hepatobiliary Surg Nutr*. 2013;2:248–251doi: 10.3978/j.issn.2304-3881.2013.10.03.
68. Palozza P. Carotenoids and modulation of cancer: molecular targets. *Curr Pharmacogenomics*. 2004;2:35–45.
69. Murphy SJ, Anderson LA, Ferguson HR, et al. Dietary antioxidant and mineral intake in humans is associated with reduced risk of esophageal adenocarcinoma but not reflux esophagitis or Barrett's esophagus. *J Nutr*. 2010;140:1757–1763doi: 10.3945/jn.110.124362.
70. Bollschweiler E, Wolfgarten E, Nowroth T, et al. Vitamin intake and risk of subtypes of esophageal cancer in Germany. *J Cancer Res Clin Oncol*. 2002;128:575–580doi: 10.1007/s00432-002-0380-z.
71. Malekshah AF, Kimiagar M, Pourshams A, et al. Vitamin deficiency in Golestan Province, northern Iran: a high-risk area for esophageal cancer. *Arch Iran Med*. 2010;13:391–394doi: 010135/aim.005.
72. Gao Y, Li W, Jia L, et al. Enhancement of (-)-epigallocatechin-3-gallate and theaflavin-3-3'-digallate induced apoptosis by ascorbic acid in human lung adenocarcinoma SPC-A-1 cells and esophageal carcinoma Eca-109 cells via MAPK pathways. *Biochem Biophys Res Commun*. 2013;438:370–374doi: 10.1016/j.bbrc.2013.07.078.
73. Rossi M, Garavello W, Talamini R, et al. Flavonoids and risk of squamous cell esophageal cancer. *Int J Cancer*. 2007;120:1560–1564doi: 10.1002/ijc.22499.
74. Woo HD, Kim J. Dietary flavonoid intake and smoking-related cancer risk: a meta-analysis. *PLoS One*. 2013;8:e75604doi: 10.1371/journal.pone.0075604.
75. Bailey DG, Dresser G, Arnold JM. Grapefruit-medication interactions: forbidden fruit or avoidable consequences? *CMAJ*. 2013;185:309–316.