



Mullerian inhibiting substance preferentially inhibits stem/progenitors in human ovarian cancer cell lines compared with chemotherapeutics

Citation

Wei, X., D. Dombkowski, K. Meirelles, R. Pieretti-Vanmarcke, P. P. Szotek, H. L. Chang, F. I. Preffer, et al. 2010. "Mullerian Inhibiting Substance Preferentially Inhibits Stem/progenitors in Human Ovarian Cancer Cell Lines Compared with Chemotherapeutics." Proceedings of the National Academy of Sciences 107, no. 44: 18874–18879. doi:10.1073/pnas.1012667107.

Published Version

doi:10.1073/pnas.1012667107

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:27527253>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Müllerian inhibiting substance preferentially inhibits stem/progenitors in human ovarian cancer cell lines compared with chemotherapeutics

Xiaolong Wei^{a,1}, David Dombkowski^b, Katia Meirelles^a, Rafael Pieretti-Vanmarcke^a, Paul P. Szotek^a, Henry L. Chang^a, Frederic I. Preffer^b, Peter R. Mueller^c, Jose Teixeira^d, David T. MacLaughlin^a, and Patricia K. Donahoe^{a,1}

^aPediatric Surgical Research Laboratories, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114; ^bFlow Cytometry Laboratory, Department of Pathology and Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA 02114; and ^cAbdominal Imaging and Intervention Division and ^dVincent Center for Reproductive Biology, Massachusetts General Hospital, Boston, MA 02114

Contributed by Patricia K. Donahoe, September 2, 2010 (sent for review July 2, 2010)

Cancer stem cells are proposed to be tumor-initiating cells capable of tumorigenesis, recurrence, metastasis, and drug resistance, and, like somatic stem cells, are thought to be capable of unlimited self-renewal and, when stimulated, proliferation and differentiation. Here we select cells by expression of a panel of markers to enrich for a population with stem cell-like characteristics. A panel of eight was initially selected from 95 human cell surface antigens as each was shared among human ovarian primary cancers, ovarian cancer cell lines, and normal fimbria. A total of 150 combinations of markers were reduced to a panel of three—CD44, CD24, and Epcam—which selected, in three ovarian cancer cell lines, those cells which best formed colonies. Cells expressing CD44, CD24, and Epcam exhibited stem cell characteristics of shorter tumor-free intervals *in vivo* after limiting dilution, and enhanced migration in invasion assays *in vitro*. Also, doxorubicin, cisplatin, and paclitaxel increased this enriched population which, conversely, was significantly inhibited by Müllerian inhibiting substance (MIS) or the MIS mimetic SP600125. These findings demonstrate that flow cytometry can be used to detect a population which shows differential drug sensitivity, and imply that treatment of patients can be individualized to target both stem/progenitor cell enriched and nonenriched subpopulations. The findings also suggest that this population, amenable to isolation by flow cytometry, can be used to screen for novel treatment paradigms, including biologic agents such as MIS, which will improve outcomes for patients with ovarian cancer.

anthrapyrazolone | chemotherapy resistance | stem/progenitor cell-enriched populations

As evidence is accumulating to indicate that cancer could be a stem cell disease (1–4), it is becoming increasingly important to be able to identify cancer stem/progenitor cells and to develop treatment modalities that specifically target the stem cell enriched population, coupled with treatments effective against the larger population not enriched for stem cells. The concept of cancer stem cells has opened new areas of research in carcinogenesis, but has the more immediate translational potential of uncovering new treatment targets.

We previously identified somatic label-retaining cells with stem cell features in ovarian surface epithelium (5), and with others (6, 7), postulate that somatic stem cells or their immediate progenitors can revert to cancer stem cells (8). It is also possible that the stem cells may remain the same, but that signals which control the stem/progenitor cell activity may change. Several recent studies have demonstrated that cancer stem cells may confer chemotherapeutic resistant ovarian tumor growth and metastasis (2, 9).

Ovarian cancer is diagnosed in approximately 25,000 new cases per year in the United States and is associated with a 50% mortality rate (10, 11); more than 90% of cases are epithelial in origin (12, 13). Epithelial ovarian cancers fall into four main subtypes: mucinous, endometrioid, clear cell, and serous (14). Serous ovarian carcinomas (15, 16) account for the overall high ovarian cancer-related mortality. Fewer than 25% are detected

at an early stage; thus, there is an urgent need for better predictive molecular markers that characterize early oncologic transformation (17) to permit earlier detection, to uncover additional therapeutic targets, and to change therapeutic protocols.

“Side population” (SP) cells identified by Hoechst 33342 dye exclusion in a wide range of cancers were found to be enriched for cancer stem cells (1, 18, 19). We found that SP cells form larger tumors and have higher tumorigenic propensity than do non-SP (NSP) cells, but did so in mouse ovarian cancer cell lines (MOV-CAR7). These stem/progenitor cells were inhibited by Müllerian inhibiting substance (MIS), whereas the lipophilic chemotherapeutic agent doxorubicin more significantly inhibited the NSP cells (4). These findings predict that chemotherapeutic agents and MIS may differentially affect populations in human ovarian cancer that are relatively chemoresistant and demonstrate stem cell characteristics.

In the present study, we performed flow cytometry and FACS to screen for stem cell markers and identified stem/progenitor cell enriched populations as defined by functional assays of colony formation and invasion *in vitro*, and by limiting dilution implantation *in vivo*, in a number of human ovarian cancer cell lines. In an attempt to define other markers specific to enriched stem/progenitor cell populations in these epithelial cancers for later use in screening human patients, 130 markers were screened and those amenable to flow cytometry and FACS ($n = 95$) (Fig. 1A), and conserved between patient primary ovarian cancer cells in ascites, human ovarian cancer cell lines, and the epithelium of normal human fimbria, were selected for further study (Fig. 1A). Eight surface markers conserved in all three sources were tested in various combinations and, of these, CD44⁺CD24⁺Epcam⁺ most consistently enriched for a population capable of colony growth.

These “triple-positive” (3+) cells also had a shorter tumor-free interval *in vivo* when xenotransplanted by limiting dilution, and migrated better in invasion assays *in vitro* than did triple-negative (3–) cells. Further, we assessed the effects of chemotherapeutic agents and MIS on survival of these ovarian cancer cell populations enriched or not enriched for stem-like cells. From these studies, we demonstrated that, although chemotherapeutic agents significantly inhibited viable nonenriched populations, they consistently enhanced stem/progenitor cell enriched populations. By contrast, MIS and its anthrapyrazolone small molecule agonist, SP600125, preferentially inhibited the stem/progenitor cell-enriched population. These differential results suggest that chemotherapeutic drugs and MIS should in the future be studied to determine if they can function in rationally selected combinations

Author contributions: X.W., F.I.P., J.T., D.T.M., and P.K.D. designed research; X.W., D.D., K.M., R.P.-V., P.P.S., and H.L.C. performed research; D.D., F.I.P., P.R.M., and J.T. contributed new reagents/analytic tools; X.W., D.D., K.M., R.P.-V., P.P.S., H.L.C., D.T.M., and P.K.D. analyzed data; and X.W. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence may be addressed. E-mail: xwei2@partners.org or pdonahoe@partners.org.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1012667107/-DCSupplemental.

to assure that both populations are effectively targeted (20, 21). Identification of a readily definable stem cell like population that is amenable to isolation by flow cytometry will aid in the development of future therapeutic strategies.

Results

In Fig. 1*B*, the left arm indicates that unseparated cells were cultured and treated with chemotherapeutic agents, MIS, or vehicle alone for 3 d. Flow cytometry analyzed 50,000 cells for viability and for percent of cells in side population relative to NSP or expressing Epcam⁺, CD44⁺, and CD24⁺ relative to those not expressing these three markers.

The right arm reflects cells separated by FACS and analyzed after using 150 combinations of eight markers. Combinations with the most robust colony growth in all cell lines were chosen for functional testing in vitro by invasion and proliferation assays, and by in vivo assays to determine tumor-free interval after injection of cells subjected to limiting dilutions.

SP of Ovarian Cancer Cell Line Is Enhanced After Treatment with Chemotherapeutic Agents. The verapamil-sensitive SP of human ovarian cancer cell lines was selected by exclusion of the DNA-binding Hoechst 33342 dye (Fig. S1). SKOV-3, OVCAR-5, and IGROV-1 cells, treated for 3 d with doxorubicin, showed an increased percentage of SP cells (Fig. S2*A* and *C*) and a decreased percentage of NSP cells. OVCAR-5 and IGROV-1 cells treated with cisplatin also significantly increased the percentage of SP cells in OVCAR-5 (Fig. S2*B* and *D*) and in IGROV-1 while decreasing that of NSP cells. Paclitaxel also caused an increase in the percentage of SP cells in OVCAR-5 and SKOV-3 cells (Fig. S2*B* and *D*). Chemotherapeutic agents dose-dependently reduced the total number of cells after 3 d of treatment (Table S1).

MIS and Its Anthracycline Agonist, SP600125, Diminish SP Cells. The receptor-mediated ligand MIS, compared with vehicle alone, decreased the percentage of SP cells (Fig. S3*A* and *C*) in the human ovarian cancer cell lines known to be responsive to MIS in vitro or in vivo, namely OVCAR-5 (20) and IGROV-1 (21, 22). SP600125, an anthracycline, discovered in a small molecule screen to act as an MIS agonist in MIS type II receptor (23, 24), but not in BMP type II receptor transfected cells, also significantly reduced the percentage of SP cells in OVCAR-5 and IGROV-1 (Fig. S3*B* and *D* and Table S1). SP600125 also decreased the NSP of IGROV-1 cells, but at higher doses (Fig. S3*D*). Thus, different from chemotherapeutic agents, MIS and its agonist SP600125,

after 3 d of treatment in culture, significantly reduce the ratio of viable SP cells of a number of human ovarian cancer cell lines.

Enrichment of Cancer Stem Cell Populations in Human Ovarian Cancer Cell Lines by Selection with a Marker Panel Compatible with Flow Cytometry. Primary ascites from three patients with ovarian cancer were screened in flow cytometry for relative binding of 95 fluorescently labeled surface markers. After negative selection for CD45 and CD31 cells, the panel was narrowed to 24 cell surface markers reconfirmed in five ovarian cancer cell lines (Fig. 1*A*). A panel of eight surface markers consistently found in both primary cancers and cancer cell lines, as well as normal Fallopian tube infundibular cells (Fig. 1*A*), was further tested in more than 50 different combinations (Table S2) for their ability to form colonies (25) in each of three cell lines, OVCAR-5, IGROV-1, or SKOV-3 cells (Fig. 1*B*, right arm). The CD44⁺CD24⁺Epcam⁺ population formed more large colonies after 14 d than did other combinations (Fig. 2*A* and *B* and Table S2; each combination represents $n = 3-5$ experiments). Moreover, we analyzed the association of SP and the CD44⁺CD24⁺Epcam⁺ population, and found that the 3+ cell percentage is increased in the SP cells (Fig. S4) and that SP cells are dramatically increased in the 3+ cells of OVCAR-5, indicating that there is an overlap between SP and CD44⁺CD24⁺Epcam⁺ cells.

Triple-Positive Cells Invade Matrigel More Effectively than Do 3-Cells. CD44⁺CD24⁺Epcam⁺ cells in serum-free media invade through an extracellular matrix-coated membrane insert after incubation for an additional 72 h in serum containing media. When stained and then quantitated by ImageJ software (National Institutes of Health), 3+ cells showed significantly more invasion through the matrix than did 3- cells ($P < 0.05$; $n = 3$; Fig. 2*B*, *Right*).

Triple Positive Cells Injected into the Right Flank of NOD/SCID Mice Grew Tumors Earlier than Did Triple Negative Left Flank Cells. CD44⁺CD24⁺Epcam⁺ cells grew tumors earlier at comparable cell dilutions (10^3 and 10^2) than did 3- cells. Tumor-free intervals were then subjected to Kaplan-Meier analysis and log-rank (Mantel-Cox) and Gehan-Breslow-Wilcoxon tests for significance. The tumor free interval was shorter ($P = 0.015$ for 10^3 cells, $P = 0.049$ for 10^2 cells; $n = 5$ animals for each) for the 3+ cells compared with 3- cells (Fig. 2*C*).

Marker Panel-Selected Cells Are Diminished by MIS and SP600125. After treatment with chemotherapeutic agents or MIS, remaining viable populations of OVCAR-5 cells were selected by flow




Fig. 1. (A) Selection of human ovarian cancer cell surface markers conserved between primary ascites from ovarian cancer patients, ovarian cancer cell lines, and normal human Fallopian tube fimbria. Combinations of eight conserved markers were tested, as shown in a flowchart of experimental design (B).




Fig. 2. Characterization of populations enriched for stem/progenitor cells in human ovarian cancer cell lines by colony growth and invasion in vitro and by tumor free interval in vivo. (A and B) Colony growth of OVCAR-5 (A) and SKOV-3 (B) in six-well plates. CD44/CD24/Epcam 3+ (Upper) gave optimal colony growth compared with 3- (Lower), as quantified on the right. (B) The 3+ cells isolated from SKOV-3 and plated on a membrane insert coated with ECM invaded to the undersurface of the membrane better than did 3- cells (* $P < 0.05$; $n = 3$). (C) The 3+ cells sorted from OVCAR-5 and serially diluted and injected s.c. into 5-wk-old female NOD/SCID mice ($n = 5$ mice for each group) grew tumors more readily than did 3- cells. (Bars indicate SD.)

cytometry for 3+ markers (Fig. 1B, left arm), and the remaining cell numbers determined and compared with those cells that were 3-. There was a dramatic expansion of the ratio of 3+ OVCAR-5 cells (Table S4, right column) when treated with doxorubicin (4.52–39.9%) or cisplatin (2.03–10.41%) (Fig. 3A and B and Table S4) whereas the total number of viable cells slightly decreased. By comparison, both the total numbers of viable unseparated cells and 3+ separated cells decreased significantly when treated with MIS or SP600125 (Fig. 3C and D and Table S4).

Effects of Chemotherapeutic Agents and MIS and Its Agonist on Cell Survival of 3+ and 3- Cells. We tested the effect of chemotherapeutic agents and MIS on the survival of 3+ and 3- populations in 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assays. Separated OVCAR-5 cells plated at 2,000 cells per well were treated with different doses of doxorubicin for 3 d. Doxorubicin slightly inhibited the survival of 3+ cells ($P < 0.05$), but more strongly inhibited 3- cells ($P < 0.01$) (Fig. 4A, Upper), as did cisplatin (Fig. 4A, Lower). Similar results were seen on cells separated by FACS into NSP (Fig. S5A), which were more inhibited than SP by doxorubicin and cisplatin. These results indicate that chemotherapeutic agents preferentially inhibit 3- and NSP (Fig. S5A), and that 3+ and SP cells show relative resistance to chemotherapeutic agents.

The 3+ and 3- cells from OVCAR-5 cells were treated with different doses of MIS for 3 d. In contrast, MIS significantly inhibits 3+ cells at lower doses than those required to inhibit 3- cells, although both were significant (Fig. 4B, Upper, for 3 d; Fig. S5C, after 7 d). SP600125 also significantly inhibited growth of 3+ cells ($P < 0.01$), but required higher doses to inhibit 3- cells, although both were significant (Fig. 4B, Lower, and Fig. S5D). Survival of SP cells was more inhibited by MIS and SP600125 compared with NSP cells (Fig. S5B).

Discussion

It is essential to identify ovarian cancer populations enriched for tumor-initiating cells and, at the same time, to elucidate the molecular mechanisms that regulate normal self-renewal or differentiation, and to determine how such mechanisms can be co-opted or changed to contribute to transformation and drug resistance and whether modulation of these mechanisms will affect patient outcome. SP cells have been described as being enriched for cancer stem cells and characterized in several mammalian tissues or cancers by efflux of lipophilic substrates, including the cell-permeable DNA-specific bisbenzimidazole dye Hoechst 33342 and some chemotherapeutic agents (1, 4, 5, 19, 26–28). Having identified SP cells in mouse ovarian cancer cell lines (4) and shown that they were more tumorigenic than NSP cells, we now show




Fig. 3. MIS and its agonist SP600125 decrease the percentage of marker-enriched stem cell populations in human ovarian cancer cell lines. OVCAR-5 cells were treated with increasing doses of doxorubicin (A), cisplatin (B), MIS (C), or SP600125 (D) for 3 d and stained with a combination of CD44, CD24, and Epcam antibodies. A total of 50,000 cells were tested by flow analysis for viability. Percentages of marker panel selected populations were determined and fold changes calculated ($n = 3$ separate experiments performed in triplicate at each dose for each drug).

that SP populations existing in human ovarian cancer cell lines (OVCAR-5, IGROV-1, and SKOV-3) form more colonies than NSP cells and are resistant to chemotherapeutic agents (doxorubicin, cisplatin, and paclitaxel), but more sensitive to MIS and its agonist, SP600126.

To consider other therapeutic agents that can target SP cells and NSP cells, we turned to extracellularly active receptor mediated agents and tested the effects of MIS, as mouse ovarian surface epithelium expresses MIS RII receptors (21, 29, 30), as do human ovarian cancer cell lines and primary ovarian cancer ascites cells (20). We found that MIS and its small molecule MISRII receptor-dependent agonist, SP600125 (23), significantly decreased SP cell percentages in OVCAR-5, IGROV-1, and SKOV-3.

As little is known about markers more specific for the stem cell population of human ovarian cancer, we performed a screen of 95 flow cytometry-compatible cell surface markers that were conserved in human primary ovarian cancer ascites and in human ovarian cancer cell lines (OVCAR-3, OVCAR-5, OVCAR-8, SKOV-3, and IGROV-1; Fig. 14). Eight of these (Epcam, CD24, CD44, CD90, CD105, CD133, E-cadherin, and SP) were preserved in normal human fimbria from 16 patients undergoing excision for benign disease, as fimbria removed prophylactically with ovaries in patients with familial breast cancer were found to be the site of occult tumors bearing characteristics of serous cystadenocarcinoma of the ovary (31–33). Normal infundibulum as a surrogate for ovarian surface epithelium contained a significant SP after enzymatic digestion of the human normal fimbria, and therefore was used to screen cancer stem cell surface markers that could be subsequently used in human patient ascites.

MTT assay after treatment of sorted OVCAR-5 cells




Fig. 4. Chemotherapeutic agents and MIS differentially affect survival of marker-enriched stem cell populations. CD44/CD24/Epcam 3+ and 3− cells were sorted from OVCAR-5 cells by FACS and tested in MTT assays. (A) Doxorubicin treatment (10, 30, 60 nM) and cisplatin treatment (0.5, 1, 2 μM) inhibited the proliferation of 3− cells compared with 3+ cells ($*P < 0.05$; $**P < 0.01$). (B) MIS treatment (37.5, 112.5, 225 nM) and SP600125 treatment (5, 10, 16 μM), by contrast, inhibited proliferation of the 3+ population significantly more than the 3− population ($*P < 0.05$; $**P < 0.01$). (Bars indicate SD.)

Several of these markers have been identified in human cancer (34–41). For example, Epcam (CD326) is a glycosylated type I membrane adhesion molecule expressed in a variety of human epithelial tissues, cancers, and stem cells (39). CD44 is the receptor for the extracellular matrix component hyaluronic acid (34, 37). Epcam/CD44 expression profiles are significantly higher in colorectal primary tumors than in normal colonic tissues, and tumors originating from Epcam⁺/CD44⁺ cells generated the full morphologic and phenotypic heterogeneity of their parental lesions (36). CD133 (prominin-1) originally used to isolate hematopoietic and endothelial progenitor cells, was subsequently found to be a marker of tumor-initiating cells in a number of other human cancers (9, 35, 42–44). CD24, a mucin-like adhesion molecule, is highly expressed in a large variety of human cancers and contributes to tumor growth and metastases by binding to platelet-selectin (45, 46). Like CD44⁺ cells (47), CD24⁺ cells (48) from human ovarian tumors were recently reported to be enriched in cancer stem cells and to form tumor xenografts in nude mice, whereas CD24⁻ cells were nontumorigenic. Stem cells from different cancers may have different phenotypes, i.e., breast cancer stem cells are CD44⁺CD24^{low} and colorectal cancer-derived cell lines are CD44⁺CD24⁺ (49). Our findings demonstrate that a phenotype of CD44⁺CD24⁺Epcam⁺ is enriched for stem/progenitor cells in human ovarian cancer cell lines.

E-cadherin (CD324), a ligand for integrin α -E/ β -7, encodes a glycoprotein with five extracellular cadherin repeats. Loss of E-cadherin is associated with epithelial-to-mesenchymal transformation and cancer progression by increasing proliferation, invasion, and/or metastasis (50, 51).

After detecting verapamil-sensitive SP cells in OVCAR-3, SKOV-3, and IGROV-1, in multiple replicates, we used SP for comparison while developing a panel of surface markers that would allow us to recover live cells in which to examine molecular mechanisms and to observe the response of separated cells to drugs, for later use in human patients.

Colony formation assays used to screen the eight marker panel on three human ovarian cancer cell lines showed that the combination of Epcam⁺, CD24⁺, and CD44⁺ formed more colonies than other marker combinations. It was necessary to use this 3+ panel in combination, as each marker alone was not sufficiently selective (Table S3).

As chemotherapeutic agents have been shown in other cancers to contribute to resistance (37, 52) and to enrich for stem cell characteristics (37), we examined the effects of chemotherapeutic agents and MIS and demonstrated that chemotherapeutic agents preferentially inhibited the population not enriched for tumor initiating cells; in contrast, receptor-mediated MIS and its agonist SP600125 inhibited significantly the populations enriched for tumor-initiating cells. The fact that these enriched populations showed resistance to chemotherapeutic agents, but maintained sensitivity to MIS and to SP600125, indicates that there is experimental rationale to use this combination of markers to study primary ascites or tumor from patients and to screen for additional selective inhibitors of human ovarian cancer populations enriched for stem-like cells. Furthermore, there is a need to design individualized treatment which specifically addresses the patient-specific population enriched for tumor initiating cells. This approach will hopefully lead to improved outcomes for patients with ovarian cancer. In addition, the panel-selected epithelial cells can now be used as an experimental tool to examine for variations (53, 54) that highlight mechanistic differences that could serve as future therapeutic targets. Furthermore, this marker panel is immediately translatable to the clinic to select patient-specific treatment strategies.

Materials and Methods

Cell Lines, Chemotherapeutic Agents, and MIS. Human ovarian cancer cell lines—OVCAR-5 (55, 56), SKOV-3 (57, 58), and IGROV-1 (59) cells—were maintained in the pediatric surgical research laboratories as previously described (4, 21). Cells were treated with doxorubicin, cisplatin, paclitaxel (all from NovaPlus), MIS, or SP600125 (Sigma) (23). MIS was purified in

our laboratory, and its bioactivity assessed in embryonic Müllerian duct regression assays (60, 61). The treatment doses were selected (21) around the IC₅₀ for each drug for each cell line (10–60 nM for doxorubicin, 0.5–2 μ M for cisplatin, 0.5–4 nM for paclitaxel, 37.5–225 nM for MIS, and 5–16 μ M for SP600125).

Harvesting of Primary Human Ovarian Cancer Ascites. Primary ascites were removed therapeutically from patients with ovarian cancer at the Massachusetts General Hospital [institutional review board (IRB) approval 2007-P-001918], centrifuged at 450 \times g for 20 min, resuspended in DMEM/F-12 medium, filtered using a sterile cell strainer (70- μ m nylon mesh; Fisher Scientific), recentrifuged at 225 \times g for 5 min, resuspended with ammonium chloride solution to lyse red blood cells, diluted with DMEM/F-12 medium, and recentrifuged. Cells were then washed twice in the same media, stained with antibodies for 20 min at 4 $^{\circ}$ C, washed again, and resuspended in PBS solution for immediate flow analysis.

Digestion of Normal Fimbria Epithelium. Normal human Fallopian tube fimbria removed from sixteen patients undergoing surgery for benign uterine disease at the Massachusetts General Hospital (IRB approval 2007-P-001918/4) or at Dana Farber Cancer Institute (IRB Legacy 02–0251) were enzymatically digested at 37 $^{\circ}$ C in 0.2% (wt/vol) with collagenase type II (type II in DMEM; Gibco-BRL) for 30 to 45 min, followed by dilution with DMEM/F-12 medium. Normal epithelial cells, released by scraping the fimbria, were collected, centrifuged at 225 \times g for 5 min, resuspended in DMEM/F-12 medium, filtered through a sterile cell strainer (70 μ m nylon mesh), and treated in the same manner as the ascites cells before flow cytometric analysis.

Flow Cytometry and Cell Number After Treatment of Cell Lines. Cells were plated in T75 flasks at 10% to 15% confluency (Fig. 1B, left arm), then treated with increasing doses of doxorubicin, cisplatin, paclitaxel, MIS, SP600125, or vehicle alone for 3 d. Flow cytometry was performed in the Massachusetts General Hospital Department of Pathology Flow Cytometry Laboratory using a custom designed high-resolution 7-laser LSR (BD Science) (4, 62). Cells were stained with Hoechst dye 33342 (Invitrogen) for 90 min at 37 $^{\circ}$ C, or antibodies to the markers for 30 min at 4 $^{\circ}$ C. A total of 50,000 cells were tested by flow analysis for viability using 7-AAD (Sigma) and for verapamil-sensitive (50–100 μ g/mL; Invitrogen) exclusion of Hoechst dye or for binding of optimal marker combinations (as detailed later).

Colony Formation to Select Optimal Combination of Surface Markers. Single cell suspensions were selected using a 5-laser FACSDiVa cell sorter (BD Science) and stained with a variety of combinations of antibodies to 95 markers to choose eight markers conserved in human ovarian cancer cell lines, primary ascites, and primary human fimbria epithelial cells and also detected on other cancer stem cells: anti-human CD24-PE (eBioscience), anti-mouse/human CD44-APC/Cy7 or CD44-Alexa Fluor-700 (BioLegend), anti-human CD90-APC (BioLegend), anti-human CD105-PE (Invitrogen), anti-human CD133-APC (BioLegend), anti-human Epcam-Alexa Fluor-647 or Epcam-APC (BioLegend), and anti-human E-cadherin-FITC (CD324; BioLegend; Fig. 1B, right arm). (For additional antibodies tested, see link to Stem Progenitor Cell Marker Panel at www.massgeneral.org/children/research/researchers/wei_xiaolong.aspx.) Cells sorted from the three cell lines were separated by various markers and selected for viability, plated on six-well plates (500, 1,000, or 2,000 cells/well), and incubated for 14 d at 37 $^{\circ}$ C, with a media change at day 7. At day 14, cells were washed, fixed with methanol, stained with Giemsa, and analyzed with ImageJ (National Institutes of Health).

Effects of Treatment on Cell Survival. The SP and NSP populations and the 3+ and 3- populations, after sorting, were tested in a cell survival tetrazole MTT assay as previously described (21) at 2,000 cells per well in 200 μ L of media per well after treatment for 3 d with different doses of chemotherapeutic agents (doxorubicin, cisplatin, paclitaxel), MIS, and the MIS agonist SP600125, then compared with an appropriate vehicle as control.

Cell Invasion Assay. In vitro cell invasion assay was performed according to the manufacturer's protocols of an ECM invasion assay kit (Millipore), comparing 3+ versus 3- populations.

Tumor-Free Interval or Time to Appearance of Xenotransplanted Tumors After Limiting Dilution. The 3+ and 3- cell populations were isolated from OVCAR-5 cell lines by FACS, serially diluted (10^5 , 10^3 , 10^2 cells), resuspended in 1:1 PBS/Matrigel (BD Biosciences) at 4 $^{\circ}$ C in 200 μ L, and injected s.c. into the right flank and 3- into the left flank of 4- to 6-wk-old female NOD.CB17-Prkdc/

Scid/J mice (Jackson Laboratory) under a protocol approved by the Massachusetts General Hospital Institutional Animal Care and Use Committee (IRB approval 2009 N000033/1). Mice were monitored weekly for tumor formation, and time of appearance was recorded after euthanasia by CO₂ inhalation. Tumors were measured and then fixed or frozen for further study.

Statistical Analysis. Univariate two-tailed *t* tests were used (4, 21) to compare MIS and drugs with vehicle as a control. Kaplan-Meier and log-rank (Mantel-Cox) and Gehan-Breslow-Wilcoxon analyses were used to compare

differences of time to tumor appearance. Analyses were performed using Prism 5 software for Mac OSX (version 5.0a; GraphPad Software).

ACKNOWLEDGMENTS. We thank Caroline Coletti for excellent editorial support. This work was supported by a Massachusetts General Hospital Executive Committee on Research Award for Medical Discovery (to X.W.), National Institutes of Health Grant R01 CA17393 (to P.K.D. and D.T.M.), Harvard Stem Cell Institute (P.K.D.), Ovarian Cancer Foundation (D.T.M.), and gifts from Commons Development, the McBride Family Foundation, the Austen Foundation, and the Surdna-Gar Foundation.

- Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF (2003) Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 100:3983–3988.
- Bapat SA, Mali AM, Koppikar CB, Kurrey NK (2005) Stem and progenitor-like cells contribute to the aggressive behavior of human epithelial ovarian cancer. *Cancer Res* 65:3025–3029.
- Li C, et al. (2007) Identification of pancreatic cancer stem cells. *Cancer Res* 67:1030–1037.
- Szotek PP, et al. (2006) Ovarian cancer side population defines cells with stem cell-like characteristics and Mullerian inhibiting substance responsiveness. *Proc Natl Acad Sci USA* 103:11154–11159.
- Szotek PP, et al. (2008) Normal ovarian surface epithelial label-retaining cells exhibit stem/progenitor cell characteristics. *Proc Natl Acad Sci USA* 105:12469–12473.
- Knoblich JA (2008) Mechanisms of asymmetric stem cell division. *Cell* 132:583–597.
- Morrison SJ, Spradling AC (2008) Stem cells and niches: mechanisms that promote stem cell maintenance throughout life. *Cell* 132:598–611.
- Gupta PB, Chaffer CL, Weinberg RA (2009) Cancer stem cells: Mirage or reality? *Nat Med* 15:1010–1012.
- Baba T, et al. (2009) Epigenetic regulation of CD133 and tumorigenicity of CD133+ ovarian cancer cells. *Oncogene* 28:209–218.
- Cannistra SA, et al. (2003) Progress in the management of gynecologic cancer: consensus summary statement. *J Clin Oncol* 21(suppl 10):129s–132s.
- Jemal A, et al. (2009) Cancer statistics, 2009. *CA Cancer J Clin* 59:225–249.
- Alvarez AA, Krigman HR, Whitaker RS, Dodge RK, Rodriguez GC (1999) The prognostic significance of angiogenesis in epithelial ovarian carcinoma. *Clin Cancer Res* 5:587–591.
- Quirk JT, Natarajan N, Mettlin CJ (2005) Age-specific ovarian cancer incidence rate patterns in the United States. *Gynecol Oncol* 99:248–250.
- Ho SM (2003) Estrogen, progesterone and epithelial ovarian cancer. *Reprod Biol Endocrinol* 1:73.
- Körner M, Burckhardt E, Mazzucchelli L (2005) Different proportions of aneuploid cells in ovarian inclusion cysts associated with serous borderline tumours and serous high-grade carcinomas support different pathogenetic pathways. *J Pathol* 207:20–26.
- Landen CN, Jr, Birrer MJ, Sood AK (2008) Early events in the pathogenesis of epithelial ovarian cancer. *J Clin Oncol* 26:995–1005.
- Shih IeM, Kurman RJ (2005) Molecular pathogenesis of ovarian borderline tumors: new insights and old challenges. *Clin Cancer Res* 11:7273–7279.
- Goodell MA, Brose K, Paradis G, Conner AS, Mulligan RC (1996) Isolation and functional properties of murine hematopoietic stem cells that are replicating in vivo. *J Exp Med* 183:1797–1806.
- Kondo T, Setoguchi T, Taga T (2004) Persistence of a small subpopulation of cancer stem-like cells in the C6 glioma cell line. *Proc Natl Acad Sci USA* 101:781–786.
- Masiakos PT, et al. (1999) Human ovarian cancer, cell lines, and primary ascites cells express the human Mullerian inhibiting substance (MIS) type II receptor, bind, and are responsive to MIS. *Clin Cancer Res* 5:3488–3499.
- Pieretti-Vanmarcke R, et al. (2006) Mullerian inhibiting substance enhances subclinical doses of chemotherapeutic agents to inhibit human and mouse ovarian cancer. *Proc Natl Acad Sci USA* 103:17426–17431.
- Stephen AE, et al. (2002) Highly purified müllerian inhibiting substance inhibits human ovarian cancer in vivo. *Clin Cancer Res* 8:2640–2646.
- Renlund N, et al. (2008) c-Jun N-terminal kinase inhibitor II (SP600125) activates Mullerian inhibiting substance type II receptor-mediated signal transduction. *Endocrinology* 149:108–115.
- Teixeira J, et al. (1996) Developmental expression of a candidate müllerian inhibiting substance type II receptor. *Endocrinology* 137:160–165.
- Fuller AF, Jr, Guy S, Budzik GP, Donahoe PK (1982) Mullerian inhibiting substance inhibits colony growth of a human ovarian carcinoma cell line. *J Clin Endocrinol Metab* 54:1051–1055.
- Haraguchi N, et al. (2006) Characterization of a side population of cancer cells from human gastrointestinal system. *Stem Cells* 24:506–513.
- Moserle L, Ghisi M, Amadori A, Indraccolo S (2010) Side population and cancer stem cells: therapeutic implications. *Cancer Lett* 288:1–9.
- Wulf GG, et al. (2001) A leukemic stem cell with intrinsic drug efflux capacity in acute myeloid leukemia. *Blood* 98:1166–1173.
- Connolly DC, et al. (2003) Female mice chimeric for expression of the simian virus 40 TAG under control of the MISIR promoter develop epithelial ovarian cancer. *Cancer Res* 63:1389–1397.
- Rodina AV, et al. (2008) Localization of Mullerian inhibiting substance receptors in various human cancer cell lines. *Biochemistry (Moscow)* 73:797–805.
- Callahan MJ, et al. (2007) Primary fallopian tube malignancies in BRCA-positive women undergoing surgery for ovarian cancer risk reduction. *J Clin Oncol* 25:3985–3990.
- Crum CP, et al. (2007) Lessons from BRCA: the tubal fimbria emerges as an origin for pelvic serous cancer. *Clin Med Res* 5:35–44.
- Crum CP, et al. (2007) The distal fallopian tube: A new model for pelvic serous carcinogenesis. *Curr Opin Obstet Gynecol* 19:3–9.
- Alvero AB, et al. (2009) Molecular phenotyping of human ovarian cancer stem cells unravels the mechanisms for repair and chemoresistance. *Cell Cycle* 8:158–166.
- Choi D, et al. (2009) Cancer stem cell markers CD133 and CD24 correlate with invasiveness and differentiation in colorectal adenocarcinoma. *World J Gastroenterol* 15:2258–2264.
- Dalerba P, et al. (2007) Phenotypic characterization of human colorectal cancer stem cells. *Proc Natl Acad Sci USA* 104:10158–10163.
- Hong SP, Wen J, Bang S, Park S, Song SY (2009) CD44-positive cells are responsible for gemcitabine resistance in pancreatic cancer cells. *Int J Cancer* 125:2323–2331.
- Horst D, Kriegl L, Engel J, Kirchner T, Jung A (2009) Prognostic significance of the cancer stem cell markers CD133, CD44, and CD166 in colorectal cancer. *Cancer Invest* 27:844–850.
- Marhaba R, et al. (2008) CD44 and EpCAM: Cancer-initiating cell markers. *Curr Mol Med* 8:784–804.
- Munz M, Baeuerle PA, Gires O (2009) The emerging role of EpCAM in cancer and stem cell signaling. *Cancer Res* 69:5627–5629.
- Takaishi S, et al. (2009) Identification of gastric cancer stem cells using the cell surface marker CD44. *Stem Cells* 27:1006–1020.
- Curley MD, et al. (2009) CD133 expression defines a tumor initiating cell population in primary human ovarian cancer. *Stem Cells* 27:2875–2883.
- Mizrak D, Brittan M, Alison MR (2008) CD133: Molecule of the moment. *J Pathol* 214:3–9.
- Shmelkov SV, St Clair R, Lyden D, Rafii S (2005) AC133/CD133/Prominin-1. *Int J Biochem Cell Biol* 37:715–719.
- Lim SC (2005) CD24 and human carcinoma: Tumor biological aspects. *Biomed Pharmacother* 59(suppl 2):S351–S354.
- Sagiv E, Arber N (2008) The novel oncogene CD24 and its arising role in the carcinogenesis of the GI tract: From research to therapy. *Expert Rev Gastroenterol Hepatol* 2:125–133.
- Zhang S, et al. (2008) Identification and characterization of ovarian cancer-initiating cells from primary human tumors. *Cancer Res* 68:4311–4320.
- Gao MQ, Choi YP, Kang S, Youn JH, Cho NH (2010) CD24+ cells from hierarchically organized ovarian cancer are enriched in cancer stem cells. *Oncogene* 29:2672–2680.
- Yeung TM, Gandhi SC, Wilding JL, Muschel K, Bodmer WF (2010) Cancer stem cells from colorectal cancer-derived cell lines. *Proc Natl Acad Sci USA* 107:3722–3727.
- Geiger B, Ayalon O (1992) Cadherins. *Annu Rev Cell Biol* 8:307–332.
- Mareel M, Vlemminckx K, Vermeulen S, Bracke M, Van Roy F (1992) E-cadherin expression: A counterbalance for cancer cell invasion. *Bull Cancer* 79:347–355.
- Dave B, Chang J (2009) Treatment resistance in stem cells and breast cancer. *J Mammary Gland Biol Neoplasia* 14:79–82.
- Corney DC, et al. (2010) Frequent downregulation of miR-34 family in human ovarian cancers. *Clin Cancer Res* 16:1119–1128.
- Corney DC, Nikitin AY (2008) MicroRNA and ovarian cancer. *Histol Histopathol* 23:1161–1169.
- Johnson SW, Laub PB, Beesley JS, Ozols RF, Hamilton TC (1997) Increased platinum-DNA damage tolerance is associated with cisplatin resistance and cross-resistance to various chemotherapeutic agents in unrelated human ovarian cancer cell lines. *Cancer Res* 57:850–856.
- Schilder RJ, et al. (1990) Metallothionein gene expression and resistance to cisplatin in human ovarian cancer. *Int J Cancer* 45:416–422.
- Fogh J, Wright WC, Loveless JD (1977) Absence of HeLa cell contamination in 169 cell lines derived from human tumors. *J Natl Cancer Inst* 58:209–214.
- Hirschmann-Jax C, et al. (2004) A distinct “side population” of cells with high drug efflux capacity in human tumor cells. *Proc Natl Acad Sci USA* 101:14228–14233.
- Bénard J, et al. (1985) Characterization of a human ovarian adenocarcinoma line, IGOV1, in tissue culture and in nude mice. *Cancer Res* 45:4970–4979.
- Lorenzo HK, et al. (2002) New approaches for high-yield purification of Müllerian inhibiting substance improve its bioactivity. *J Chromatogr B Analyt Technol Biomed Life Sci* 766:89–98.
- Ragin RC, Donahoe PK, Kenneally MK, Ahmad MF, MacLaughlin DT (1992) Human müllerian inhibiting substance: Enhanced purification imparts biochemical stability and restores antiproliferative effects. *Protein Expr Purif* 3:236–245.
- Pfeffer F, Dombkowski D (2009) Advances in complex multiparameter flow cytometry technology: Applications in stem cell research. *Cytometry B Clin Cytom* 76:295–314.