



First-in-man assessment of plaque rupture by polarization-sensitive optical frequency domain imaging in vivo

Citation

van der Sijde, Johannes N., Antonios Karanasos, Martin Villiger, Brett E. Bouma, and Evelyn Regar. 2016. "First-in-man assessment of plaque rupture by polarization-sensitive optical frequency domain imaging in vivo." *European Heart Journal* 37 (24): 1932. doi:10.1093/eurheartj/ehw179. <http://dx.doi.org/10.1093/eurheartj/ehw179>.

Published Version

doi:10.1093/eurheartj/ehw179

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:27662122>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehw179

Online publish-ahead-of-print 12 May 2016

First-in-man assessment of plaque rupture by polarization-sensitive optical frequency domain imaging *in vivo*

Johannes N. van der Sijde¹, Antonios Karanasos¹, Martin Villiger², Brett E. Bouma², and Evelyn Regar^{1*}

¹Department of Interventional Cardiology, Thoraxcenter, BA-585, Erasmus Medical Center, Gravendijkwal 230, Rotterdam 3015 CE, The Netherlands; and ²Harvard Medical School and Massachusetts General Hospital, Wellman Center for Photomedicine, Boston, USA

*Corresponding author. Tel: +31 10 7035232, Fax: +31 10 7035046, Email: e.regar@erasmusmc.nl

A 50-year-old man underwent primary percutaneous coronary intervention for anterior ST-segment elevation myocardial infarction 3 h after symptom onset. After thrombectomy, polarization-sensitive optical frequency domain imaging (PS-OFDI) was performed at the culprit left anterior descending artery lesion using a prototype imaging system (Wellman Center for Photomedicine). In addition to the conventional image of plaque structure, PS-OFDI provides a measure of tissue birefringence. This optical tissue property is elevated in tissue with regularly organized architecture, such as thick collagen fibres or smooth muscle cells.

Structure images showed a thin-cap fibroatheroma (TCFA) (Panel B1; arrow) with plaque rupture (Panel C1; white arrow) and thrombus (Panels C1–D1). Inspection of birefringence images revealed the presence of low birefringence (yellow colour) in the fibrous cap region both at the rupture site and proximally, suggesting the absence of thick collagen fibres (Panels B2–C2; white arrow). The intracoronary thrombus had predominantly low birefringence,

consistent with the unorganized architecture of acute thrombus (Panel D2), but also featured areas of intermediate-high birefringence, suggesting a higher degree of organization (Panel C2; black arrow). The lesion was then successfully treated with bioresorbable scaffold implantation.

Thrombotic coronary occlusion following plaque rupture is the most common pathomechanism of myocardial infarction. In our case, we witnessed an acute coronary syndrome, due to rupture of a TCFA, which presumably lacked thick collagen fibres in the fibrous cap. It is yet unknown how the collagen composition of the fibrous cap impacts plaque stability and whether thrombus healing impacts clinical event manifestation. Considering that invasive detection of TCFA predicts future culprit lesions poorly, a tool allowing an enhanced structural assessment such as PS-OFDI might improve invasive risk stratification in the future.

Funding

This research was supported in part by the National Institutes of Health (grants P41EB-015903 and R01HL-119065) and by Terumo Corporation. Funding to pay the Open Access publication charges for this article was provided by Wellman Center for Photomedicine.

Supplementary material is available at *European Heart Journal* online.

© The Author 2016. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

