



Antibody Secreting Cell Responses following Vaccination with Bivalent Oral Cholera Vaccine among Haitian Adults

The Harvard community has made this article openly available. [Please share](#) how this access benefits you. Your story matters

Citation	Matias, W. R., B. Falkard, R. C. Charles, L. M. Mayo-Smith, J. E. Teng, P. Xu, P. Kováč, et al. 2016. "Antibody Secreting Cell Responses following Vaccination with Bivalent Oral Cholera Vaccine among Haitian Adults." PLoS Neglected Tropical Diseases 10 (6): e0004753. doi:10.1371/journal.pntd.0004753. http://dx.doi.org/10.1371/journal.pntd.0004753 .
Published Version	doi:10.1371/journal.pntd.0004753
Citable link	http://nrs.harvard.edu/urn-3:HUL.InstRepos:27822377
Terms of Use	This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

RESEARCH ARTICLE

Antibody Secreting Cell Responses following Vaccination with Bivalent Oral Cholera Vaccine among Haitian Adults

Wilfredo R. Matias^{1,2}✉, Brie Falkard³✉, Richelle C. Charles^{3,4}, Leslie M. Mayo-Smith³, Jessica E. Teng⁵, Peng Xu⁶, Pavol Kováč⁶, Edward T. Ryan³, Firdausi Qadri⁷, Molly F. Franke¹, Louise C. Ivers^{1,2,5}, Jason B. Harris^{3,8*}

1 Department of Global Health & Social Medicine, Harvard Medical School, Boston, Massachusetts, United States of America, **2** Partners In Health, Boston, Massachusetts, United States of America, **3** Division of Infectious Diseases, Massachusetts General Hospital, Boston, Massachusetts, United States of America, **4** Harvard Medical School, Boston, Massachusetts, United States of America, **5** Division of Global Health Equity, Brigham & Women's Hospital, Boston, Massachusetts, United States of America, **6** NIDDK, LBC, National Institutes of Health, Bethesda, Maryland, United States of America, **7** International Centre for Diarrhoeal Disease Research Bangladesh (icddr,b), Dhaka, Bangladesh, **8** Department of Pediatrics, Harvard Medical School, Boston, Massachusetts, United States of America

✉ These authors contributed equally to this work.

* jbharris@mgd.harvard.edu



OPEN ACCESS

Citation: Matias WR, Falkard B, Charles RC, Mayo-Smith LM, Teng JE, Xu P, et al. (2016) Antibody Secreting Cell Responses following Vaccination with Bivalent Oral Cholera Vaccine among Haitian Adults. *PLoS Negl Trop Dis* 10(6): e0004753. doi:10.1371/journal.pntd.0004753

Editor: Joseph M. Vinetz, University of California San Diego School of Medicine, UNITED STATES

Received: March 10, 2016

Accepted: May 11, 2016

Published: June 16, 2016

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This research was supported by core grants to the icddr,b, the Doris Duke Charitable Foundation through a grant supporting the Doris Duke International Clinical Research Fellows Program at Harvard Medical School (WRM), through programs funded by the National Institutes of Health, including the National Institute of Allergy and Infectious Diseases (AI099243 [JBH and LCI], AI103055 [JBH], AI106878 [ETR and FQ], and K08 AI089721 [RCC]), and the Intramural Research

Abstract

Background

The bivalent whole-cell (BivWC) oral cholera vaccine (Shanchol) is effective in preventing cholera. However, evaluations of immune responses following vaccination with BivWC have been limited. To determine whether BivWC induces significant mucosal immune responses, we measured *V. cholerae* O1 antigen-specific antibody secreting cell (ASC) responses following vaccination.

Methodology/Principal Findings

We enrolled 24 Haitian adults in this study, and administered doses of oral BivWC vaccine 14 days apart (day 0 and day 14). We drew blood at baseline, and 7 days following each vaccine dose (day 7 and 21). Peripheral blood mononuclear cells (PBMCs) were isolated, and ASCs were enumerated using an ELISPOT assay. Significant increases in Ogawa (6.9 cells per million PBMCs) and Inaba (9.5 cells per million PBMCs) OSP-specific IgA ASCs were detected 7 days following the first dose ($P < 0.001$), but not the second dose. The magnitude of *V. cholerae*-specific ASC responses did not appear to be associated with recent exposure to cholera. ASC responses measured against the whole lipopolysaccharide (LPS) antigen and the OSP moiety of LPS were equivalent, suggesting that all or nearly all of the LPS response targets the OSP moiety.

Conclusions/Significance

Immunization with the BivWC oral cholera vaccine induced ASC responses among a cohort of healthy adults in Haiti after a single dose. The second dose of vaccine resulted in minimal

Program of the NIH and NIDDK (PX and PK), the Robert Wood Johnson Foundation Harold Amos Medical Faculty Development Program [RCC], the Abundance foundation [LCI]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

ASC responses over baseline, suggesting that the current dosing schedule may not be optimal for boosting mucosal immune responses to *V. cholerae* antigens for adults in a cholera-endemic area.

Author Summary

The bivalent whole-cell (BivWC) oral cholera vaccine (Shanchol) is effective in preventing cholera. Despite its increasing use as part of comprehensive cholera prevention and control efforts, evaluations of immune responses following vaccination with BivWC have been limited. In this study, we measured the development of cholera-specific antibody secreting cells, markers of mucosal immunity, following vaccination with BivWC among a population of adults in Haiti, where cholera is now endemic. BivWC induced development of robust immune responses following the first dose of vaccine, but similar ASC responses were not detected following the second dose, suggesting that the currently recommended 14-day interval between doses may not be optimal for boosting mucosal immune responses among adults in cholera endemic regions. These findings suggest that additional evaluation of the optimal dosing schedule for oral cholera vaccines is warranted with the goal of improving long-term immunity.

Introduction

Cholera, a diarrheal disease with epidemic potential caused by the bacterium *Vibrio cholerae*, remains a significant cause of morbidity and mortality, accounting for approximately 1.4 to 4.3 million cases of acute diarrhea and 28,000 to 142,000 deaths annually.[1] *V. cholerae* serogroup O1 is the current predominant cause of cholera, while serogroup O139 was a major cause of cholera in the 1990s and early 2000s.[2] The O1 serogroup is further classified into Ogawa and Inaba serotypes.[3] These differ by the presence of a 2-O-methyl group present in the terminal carbohydrate of the O-specific polysaccharide (OSP) in the lipopolysaccharide (LPS) of the Ogawa serotype.[4–6] Protection against cholera is serogroup specific, and serogroup specificity is defined by the OSP moiety of LPS.[7–9]

Recent years have seen the frequency and severity of cholera outbreaks increase worldwide.[10] For example, the cholera epidemic that began in Haiti in October 2010 has accounted for 750,752 reported cases of acute diarrhea and 9,031 deaths,[11] making it the largest outbreak of cholera in recent history. While the development of robust water and sanitation infrastructure is essential in regions affected by cholera, vaccination offers another important and complementary tool for cholera prevention and control.[12]

There are currently three World Health Organization (WHO) pre-qualified, commercially available oral cholera vaccines (OCV): Dukoral (Crucell, Sweden), which consists of whole-cell killed *V. cholerae* O1 Inaba and Ogawa and recombinant cholera toxin B subunit (WC-rBS); and Shanchol (Shantha Biotechnics, India) and Euvichol (EuBiologics, Korea), which are similar bivalent whole-cell (BivWC) vaccines containing multiple biotypes of *V. cholerae* O1 and O139 which lack the cholera toxin B subunit (CtxB) found in Dukoral. Following Shanchol's prequalification by the WHO in 2011, multiple programs in diverse settings including Haiti have demonstrated the feasibility and efficacy of integrating vaccination with BivWC into comprehensive cholera control strategies.[13,14] In 2013, the WHO created an OCV stockpile to respond to cholera outbreaks worldwide.[15]

Despite the rapidly growing use of oral cholera vaccination with BivWC, there remain significant gaps in our understanding of immune responses following vaccination with BivWC. Since vaccine immunogenicity is a surrogate for protection, such studies are especially important in settings where large scale evaluations of direct effectiveness are not practical, such as in comparing vaccine formulations, dosing schedules, and in estimating vaccine efficacy across specific populations. Trials to assess safety and immunogenicity have demonstrated development of significant vibriocidal antibody and OSP-specific plasma immunoglobulin A (IgA) responses among adults and children following vaccination [16–20], including comparable vibriocidal antibody responses following a 14-day versus a 28-day dosing interval in a cholera-endemic area of Kolkata, India. [18] Previously, our group demonstrated significant vibriocidal responses and OSP-specific IgA responses following vaccination with BivWC among adults, children, and HIV-infected adults in Haiti. [21,22] However, compared to studies characterizing immune responses following natural cholera infection [23–28] and vaccination with WC-rBS [29–33], studies evaluating immune responses following vaccination with BivWC are limited.

Among evaluations of immunogenicity, the detection of antibody secreting cells (ASCs) in blood is a standard measure of the mucosal immune response. Following stimulation in the gastrointestinal tract by live *V. cholerae* or by OCVs, these cells transiently migrate into the systemic circulation where they can be detected by standard Enzyme-Linked ImmunoSpot (ELISPOT) assays. [34] This migration peaks approximately at day 7 following primary infection or vaccination. Many of these circulating ASCs express gut-homing markers and return to mucosal tissue. [35] Detection of LPS-specific ASCs in the peripheral blood correlates with the development of *V. cholerae* antigen-specific antibodies in the small intestinal lamina propria up to 6 months following infection. [36] As such, ASCs provide a critical and early window into subsequent immunologic memory at the mucosal surface.

To determine whether vaccination with BivWC induces significant mucosal immune responses, we measured levels of ASC responses against *V. cholerae* LPS and OSP at multiple time points following vaccination with BivWC among healthy participants in Haiti.

Methods

Study Population and Study Design

Healthy adults, ages 18–60 years old attending the Saint Nicolas Hospital outpatient clinic in Saint Marc, Haiti were invited to participate in May 2015, and were deemed eligible if they lived in the catchment area. Individuals with an active gastrointestinal disorder, pregnancy (as determined by urine pregnancy test), those that had previously received OCV, and those unable to give informed consent were excluded from the study. The BivWC vaccine was stored at 2°C–8°C prior to administration. Two doses of vaccine were given 14 days apart, in accordance with the package insert. Venous blood from each participant was obtained prior to immunization and 7 days after each dose of vaccine (on days 0, 7, and 21). A stool specimen was collected on day 7 to test for enteric pathogens, and a questionnaire on social and nutritional factors was conducted on day 14. The participants enrolled in this study were part of a larger ongoing study to measure long-term immune responses among adult cholera vaccine recipients.

Ethics Statement

Written informed consent was obtained from all potential adult participants prior to enrolling in the study. The study protocol was reviewed and approved by the institutional review board

of Partners HealthCare in Boston, Massachusetts, and the Haitian National Bioethics Committee in Port-au-Prince, Haiti.

Sample Collection and Isolation of PBMCs

We collected venous blood from each participant in Vacutainer blood collection tubes (Becton Dickinson), containing sodium heparin anticoagulant. Samples were obtained at day 0 (before vaccination), day 7 (7 days after the first dose of vaccine), and day 21 (1 week after the second dose of vaccine). Following sample collection, blood was diluted in PBS (pH 7.4) and processed for separation of plasma and peripheral blood mononuclear cells (PBMCs). PBMCs were prepared by differential centrifugation in Leucosep tubes (Greiner Bio-One Ltd) filled with Ficoll-Isopaque (Pharmacia, Piscataway, NJ), and after two subsequent PBS washes, resuspended at a concentration of 5×10^6 cells/mL in RPMI complete medium (Gibco, Carlsbad, CA) supplemented with 10% heat-inactivated fetal bovine serum (FBS) (HyClone, Logan, UT) and 1% Penicillin-Streptomycin (Sigma-Aldrich). These cells were used immediately as described below for measuring ASC responses. Plasma was shipped in a liquid nitrogen dry vapor unit and stored at -80°C prior to analysis in Boston.

Antibody Responses in Plasma

Vibriocidal antibody responses in plasma were measured using a standard protocol described previously.[\[24\]](#) Briefly, *V. cholerae* O1 El Tor Ogawa (X25049) and Inaba (T19479) were used as the target organism. The vibriocidal titer was defined as the reciprocal of the highest plasma dilutions resulting in a greater than 50% optical density reduction associated with *V. cholerae* O1 growth when compared to control wells without plasma. An increase of titer by 4-fold was considered to represent seroconversion, by convention. Control plasma, a pool of day 7 plasma from culture-confirmed *V. cholerae* infected individuals in Dhaka, Bangladesh, was used to monitor intra-assay variability between plates.

OSP was purified and conjugated to bovine serum albumin (BSA) as previously described;[\[37\]](#) source strains of OSP Inaba and Ogawa were *V. cholerae* O1 El Tor PIC018 and PIC058, respectively.[\[38\]](#) Anti-OSP IgA responses in plasma were measured using a previously described ELISA protocol.[\[24\]](#) In brief, 96-well polystyrene plates (Nunc, low affinity plates) were coated with *V. cholerae* O1 Ogawa OSP conjugated to BSA, or Inaba OSP conjugated to BSA (1 $\mu\text{g}/\text{mL}$), dissolved in 50mM carbonate buffer (pH 9.6). We added 100 μL of plasma diluted 1:25 in 0.1% BSA in PBS-0.05% Tween. Plates were incubated at 37°C for 60 minutes while shaking (100 rpm) and then washed. We detected OSP-specific antibodies using 100 μL per well of horseradish peroxidase-conjugated goat anti-human IgA at a dilution of 1:1,000 (Jackson ImmunoResearch, West Grove, PA) in 0.1% BSA in PBS-0.05% Tween. The plates were incubated at 37°C for 1 hour while shaking (100rpm), washed three times with PBS-0.05% Tween, and developed with ABTS/ H_2O_2 (Sigma-Aldrich, St. Louis, MO) in 0.1M citrate-phosphate buffer (pH 4.5) and 0.03% hydrogen peroxide. Measurements were made using a kinetic reading at 405 nm wavelength using the SoftMax Pro software.

Antibody Secreting Cell Responses to OSP and LPS

ASC responses were measured by enzyme-linked immunosorbent spot (ELISPOT) as described previously.[\[34\]](#) To measure the total number of circulating ASCs, nitrocellulose-bottom plates (Millipore, Bedford, MA) were coated with 100 μL of affinity-purified goat anti-human IgG F(ab)₂ (Jackson ImmunoResearch, West Grove, PA) at a concentration of 5 $\mu\text{g}/\text{mL}$ in PBS (pH 7.4). Nitrocellulose plates were also coated with 100 μL of the following antigens, to detect *V. cholerae*-specific antigen responses: *V. cholerae* O1 Ogawa and Inaba OSP:BSA (10 $\mu\text{g}/\text{mL}$),

Ogawa and Inaba-LPS (25 µg/mL), CtxB (2.5 µg/mL; Sigma-Aldrich), and keyhole limpet hemocyanin (KLH, Pierce Biotechnology, Rockford, IL, 2.5 µg/mL). While KLH is a standard control antigen, CtxB was also used to control for ASC responses from possible exposure to natural *V. cholerae* O1 infection. The BivWC vaccine lacks recombinant CtxB, therefore should not elicit a detectable CtxB-specific response.

Plates were incubated with antigen overnight at 4°C, and then blocked for 1 hour at 37°C with 200 µL of 2% milk-RPMI. A total of 1×10^5 PBMCs were added to the total Ig-coated wells and serially diluted two times by a factor of 10. A total of 5×10^5 PBMCs were added to the OSP, LPS, CtxB and KLH-coated wells. Plates were then incubated at 37°C for 3 hours and then washed. To detect IgG and IgA ASCs, plates were incubated overnight at 4°C with horseradish peroxidase-conjugated mouse anti-human IgA (Hybridoma Reagent Laboratory, Baltimore, MD) and alkaline phosphatase-conjugated IgG (Southern Biotech, Birmingham, AL), each diluted 1:500 in sterile filtered PBS-0.1%-BSA-tween-0.05%. Following overnight incubation at 4°C, plates were developed with 5-bromo-4-chloro-3-indolylphosphate-nitroblue tetrazolium (BCIP/NBT, Sigma-Aldrich) and 3-amino-9-ethylcarbazole (AEC pre-mix solution, Sigma-Aldrich). Two individuals using a stereomicroscope independently quantified ASC numbers. The number of antigen-specific IgG and IgA ASCs were expressed per 10^6 total PBMCs and as a percentage of total IgG or IgA expressing cells.

Statistical Analyses

Statistical analyses were performed using STATA Version 14 (StataCorp, LP, College Station, TX) and Graph-Pad Prism (Graph Pad Software, Inc., La Jolla, CA). Vibriocidal and antigen-specific plasma antibody responses were expressed as geometric mean titers (GMT) with 95% confidence intervals. We used the Wilcoxon signed-rank test to compare within-person antigen-specific ASC responses, vibriocidal antibody responses, and antigen-specific plasma antibody responses across different time points.

We used Spearman's correlation to quantify associations between antigen-specific ASC, vibriocidal and antigen-specific antibody responses. Comparisons between groups previously exposed and unexposed to *V. cholerae* were performed using the Mann Whitney-U test. We tested for equivalence between anti-LPS and anti-OSP ASC responses using two-sided 90% confidence intervals and an equivalence bound of ± 5 ASCs per 10^6 PBMCs difference. Confidence intervals within this bound were considered equivalent. All P-values were two-tailed, with a $P \leq 0.05$ defined as the threshold for statistical significance.

Our sample size was based on the number of participants needed to detect a difference between the baseline and post-vaccination frequency of LPS-antigen specific ASCs. Extrapolating from other studies, [32,34,39] we estimated that 19 participants would be needed to detect an increase in both IgG and IgA ASC responses, with at least 90% power following vaccination.

Results

Study Population

[Table 1](#) describes the demographic characteristics of the 24 individuals that enrolled in the study. One participant dropped out following the first blood draw on day 0, and another dropped out following the second blood draw on day 7. A third participant was excluded from the analysis of ASC responses because of excessive red blood cell contamination in PBMCs at all time points. Thus, 22 participants were included in the analysis of responses following first dose of vaccine; Twenty-one individuals were included in the analysis following the second dose of vaccine. Baseline vibriocidal titers among the study population were consistent with widespread exposure to cholera in a substantial portion of the cohort. 8/22 (36%) participants

had a vibriocidal ≥ 80 for the Ogawa serotype, and the same proportion of participants (36%) had a vibriocidal ≥ 80 for the Inaba serotype. A total of 11/22 (50%) had a vibriocidal ≥ 80 for either the Ogawa or Inaba serotype.

Plasma Antibody Responses to Vaccination

A summary of immune responses to BivWC at each time point is listed in [Table 2](#). Vibriocidal antibody responses are also depicted in [Fig 1](#). Robust vibriocidal antibody responses to Ogawa and Inaba were detected by day 7 after a single dose of vaccine ($p < 0.0001$) with a geometric mean fold rise (GMF) of 7.3 to *V. cholerae* O1 Ogawa and 9.1 to Inaba. Vibriocidal titers at day 21 were slightly lower compared to day 7 responses, although these differences were not significant. However, vibriocidal GMT at day 21 remained significantly increased compared to baseline values for Ogawa ($p < 0.01$) and Inaba ($p < 0.001$). Vibriocidal antibody seroconversion rates are depicted in [Table 2](#). The majority of participants seroconverted following the first dose of BivWC; Ogawa (64%) and Inaba (73%). The seroconversion rate increased to 76% and 81% for Ogawa and Inaba, respectively, following two doses of vaccine. In addition to vibriocidal antibody responses, we observed significant OSP-specific plasma IgA responses to both Ogawa and Inaba serotypes at days 7 and 21 ([Table 2](#)).

Antibody Secreting Cell (ASC) Responses to Vaccination

We did not observe an overall increase in the number of total IgA and IgG antibody secreting cells either in total or relative to the total number of peripheral blood mononuclear cells following vaccination with BivWC ([Table 2](#)). Among antigen-specific responses, there was no response to either of the control antigens, CtxB or KLH, at any time point. Some artefactual spots (typically < 3) are typical of ELISPOT assays. We observed a single KLH spot in only three of the 130 ELISPOT samples analyzed (2.3%), and no more than a single KLH spot was observed in any of the samples. The low number of KLH spots suggests that there was negligible artefactual background in the other antigen-specific ASC measurements. Similarly, in four of the 130 samples (3.1%) analyzed as part of the study, we observed a single CtxB spot, and again, in no samples, were more than a single CtxB spot observed. As such, we defined the presence of ≥ 2 ASCs per 10^6 PBMCs as a positive ASC response. The lack of CtxB responses suggests that responses detected to the OSP or LPS antigens were not likely the result of intercurrent infection with or exposure to *V. cholerae*, since natural infection is associated with significant responses to the CtxB antigen.

Table 1. Demographic characteristics of study participants.

Characteristics	Total
	N = 22
Mean age, years (S.E.)*	27 (19–50, 10.08)
Gender (%)	
Females	8 (36)
Males	15 (64)
Blood Type (%)	
A	7 (32)
B	4 (18)
O	10 (45)
AB	1 (5)

*S.E. Standard Error

doi:10.1371/journal.pntd.0004753.t001

Table 2. Immune responses to *V. cholerae* (O1 Ogawa and O1 Inaba) following vaccination with BivWC.

	Total ASC	O1 Ogawa	O1 Inaba
Day 0 (n = 22)			
Baseline Vibriocidal GMT ^a (95% CI)		35.3 (15.1–82.2)	29.2 (14.7–58.1)
Baseline Serum OSP IgA GMT ^b (95% CI)		65.8 (52.6–82.4)	60.7 (49.6–74.3)
Baseline Total IgA ASCs	334.5 (237.4–431.6)		
Baseline OSP IgA ASCs (per million)		0.02 (-0.02–0.07)	0.07 (-0.01–0.15)
Baseline Total IgG ASCs	920.4 (680.6–1160.1)		
Baseline OSP IgG ASCs (mean, per million)		0.045 (-0.020–0.11)	0.045 (-0.049–0.14)
Day 7 (n = 22)			
Vibriocidal GMT ^a (95% CI)		256.7 (146.0–451.2)	264.9 (150.9–464.9)
Vibriocidal GMF ^b (95% CI)		7.3 (3.4–15.6)	9.1 (4.9–16.9)
Vibriocidal Seroconversion ^c (#, %)		14 (64)	16 (73)
Serum OSP IgA GMT ^a (95% CI)		93.6 (66.1–132.7)	86.6 (64.0–117.3)
Serum OSP IgA GMF ^b (95% CI)		1.4 (1.0–1.9)	1.4 (1.1–1.9)
Serum OSP IgA Seroconversion ^d (#, %)		7 (32)	4 (18)
Total IgA ASCs	290 (233.5–346.5)		
OSP IgA ASCs (mean, per million)		6.9 (1.9–12.0)	
OSP IgA ASCs Responders ^e (#, %)		13 (59)	
Total IgG ASCs	1017 (666.7–1367.3)		
OSP IgG ASCs (mean, per million)		1.2 (0.58–1.87)	0.36 (-0.0059–0.73)
OSP IgG ASCs Responders ^e (#, %)		5 (22.7)	3 (13.6)
Day 21 (n = 21)			
Vibriocidal GMT ^a (95% CI)		165.4 (81.0–333.6)	245.7 (141.7–426.3)
Vibriocidal GMF ^b (95% CI)		4.9 (2.5–9.6)	9.1 (4.6–18.2)
Vibriocidal Seroconversion ^c (#, %)		16 (76)	17 (81)
Serum OSP IgA GMT ^a (95% CI)		86.4 (66.0–113.1)	83.5 (65.6–106.5)
Serum OSP IgA GMF ^b (95% CI)		1.4 (1.1–1.9)	1.5 (1.2–1.8)
Serum OSP IgA Seroconversion ^d (#, %)		8 (38)	6 (29)
Total IgA ASCs	408.3 (306.9–509.7)		
OSP IgA ASCs (mean, per million)		1 (0.1–1.9)	0.8 (0.1–1.5)
OSP IgA ASCs Responders ^e (#, %)		5 (23.8)	5 (23.8)
Total IgG ASCs	1035.1 (786.3–1283.9)		
OSP IgG ASCs (mean, per million)		0.41 (-0.083–0.90)	0.2 (-0.091–0.50)
OSP IgG ASCs Responders ^e (#, %)		1 (4.8)	1 (4.8)

^a Geometric mean titer.

^b Geometric mean-fold rise from baseline to 7 days after 1st dose or from baseline to 7 days after 2nd dose.

^c # with ≥ 4-fold rise in titers from baseline to 7 days after 1st dose or from baseline to 7 days after 2nd dose.

^d 3 with ≥ 2-fold rise in titers from baseline to 7 days after 1st dose or from baseline to 7 days after 2nd dose.

^e # with ≥ 2 ASCs detected

Number of participants

doi:10.1371/journal.pntd.0004753.t002

OSP-specific ASC responses are shown in Fig 2. Notably, OSP-specific IgA ASC responses to Ogawa and Inaba peaked on day 7 following the first dose of vaccine, and were significantly increased from baseline ($P < 0.001$). Notably, day 7 OSP-specific IgA antibody responses detected in plasma were strongly correlated with day 7 OSP-specific IgA ASC responses to Ogawa (Spearman $r = 0.6877$, $P = 0.0004$) and Inaba (Spearman $r = 0.6077$, $P = 0.0027$)

An analysis of ASC response based on stratification by a pre-vaccination vibriocidal titer of ≥ 80 , shows no significant difference in mean ASC responses. OSP-specific IgA responses at




Fig 1. Vibriocidal Responses. Geometric mean titer (+95% CI) of vibriocidal response to *V. cholerae* O1 Ogawa (A) and Inaba (B) at baseline (day 0) and 7 days after each immunization (day 7 and 21). Statistically significant differences relative to baseline are indicated (** = $P < 0.01$, *** = $P < 0.001$, **** = $P < 0.0001$).

doi:10.1371/journal.pntd.0004753.g001

day 7 to the Ogawa serotype were 5.2 ASCs per million PBMCs and 7.9 ASCs per million PBMCs among those exposed and unexposed ($P = 0.8894$), respectively, whereas OSP-specific IgA responses to the Inaba serotype were 10.4 ASCs per million PBMCs and 9 ASCs per million PBMCs ($P = 0.285$), respectively. Additionally, no significant differences were seen when comparing OSP-specific IgA ASC responses at day 7 to the Ogawa and Inaba serotype between individuals with previous exposure to either Ogawa or Inaba ($P = 0.6905$ for Inaba OSP response, $P = 0.7889$ for Ogawa response). However, our sample size is limited and not sufficiently powered to definitively assess whether prior exposure in this setting impacts mucosal immune responses.

OSP-specific IgA ASC responses after the second dose of the vaccine decreased nearly to baseline levels by day 21. OSP-specific IgG antibody responses were proportionally lower than IgA responses, although OSP-specific IgG ASC responses nonetheless demonstrated a statistically significant increase from baseline for the Ogawa ($p < 0.001$), but not the Inaba serotype ($p = 0.099$) (Fig 3).




Fig 2. Ogawa and Inaba O-specific polysaccharide (OSP) Immunoglobulin A (IgA) Antibody secreting cell (ASC) responses in vaccinees. Individual and mean circulating antigen-specific IgA responses to Ogawa and Inaba OSP with standard error bars. P values are indicated where statistically significant differences from baseline (Day 0) are present.

doi:10.1371/journal.pntd.0004753.g002




Fig 3. Ogawa and Inaba O-specific polysaccharide (OSP) Immunoglobulin G (IgG) Antibody secreting cell (ASC) responses in vaccinees. Individual and mean circulating antigen-specific IgG responses to Ogawa and Inaba OSP with standard error bars. P values are indicated where statistically significant differences from baseline (Day 0) are present.

doi:10.1371/journal.pntd.0004753.g003

We noted the OSP and LPS responses were not only highly correlated, but appeared nearly equivalent. (Fig 4) Testing for equivalence of IgA ASC responses to LPS and OSP demonstrated a mean difference of 0.70 ASCs per million PBMCs (90% CI: -0.799–2.21) and 0.73 ASCs per 10⁶ PBMCs (90% CI: -3.86–2.41) for Ogawa and Inaba, respectively. The confidence intervals for these mean cell differences lie within the predetermined equivalence range of +/- 5 ASCs per 10⁶ PBMCs, suggesting equivalence or near equivalence between anti-LPS and anti-OSP ASC responses at a significance level of 0.05.

Discussion

To our knowledge, this study is the first report of ASC responses to the BivWC oral cholera vaccine. We demonstrate that when administered according to the recommended two dose schedule in adults living in Haiti, BivWC induces a robust mucosal response to the *V. cholerae*




Fig 4. Correlation between Day 7 anti-OSP and anti-LPS ASC Responses. Ogawa: Spearman's $r = 0.7706$, $p < 0.0001$. Inaba: Spearman's $r = 0.9030$, $p < 0.0001$.

doi:10.1371/journal.pntd.0004753.g004

O1 antigen following the first vaccine dose. However, a second dose of the vaccine administered 14 days later did not result in a robust mucosal response to the *V. cholerae* O1 antigen.

V. cholerae is a non-invasive pathogen, and it is thought that immunity against cholera is most likely mediated by *V. cholerae* O1-antigen specific IgA produced at the mucosal surface. For this reason, IgA-secreting ASCs, which transiently appear in the circulating following vaccination and ultimately differentiate into mucosal plasma cells provide an important window into mucosal immunity. In this context, our finding that a second dose of the OCV administered 14 days after the first dose does not result in significant ASC responses in Haitian adults raises important questions about the appropriate dosing interval for the BivWC vaccine in cholera-endemic areas. While the 14-day interval may allow for the delivery of two vaccine doses to people during an outbreak as quickly as possible, it is not clear whether this is a sufficient time period to generate sufficient immunologic memory in order to achieve a prime-boost effect with the second vaccine dose. Given that our findings mirror previous studies conducted in cholera-endemic areas of South Asia following vaccination with the WC-rBS vaccine, [32,34,40] we suspect that the current 14-day dosing schedule is too short for the generation of memory B cells that can respond to secondary antigenic stimulation upon re-exposure to *V. cholerae* O1 OSP.

To address this question regarding the timing of the second vaccine dose definitively would require additional studies. Specifically, evaluating the effect of re-vaccination with oral cholera vaccine at different intervals and determining whether repeated vaccination results in increasing levels of class switching and affinity maturation among the resulting *V. cholerae*-specific ASCs, would determine whether *bona fide* functional and long lasting B cell memory can be boosted following repeated doses of oral cholera vaccination administered at longer intervals. These results could then provide mechanistic context for larger vaccine effectiveness studies comparing both single and multi-dose OCV regimens.

Another notable finding in this report was derived from measuring ASC responses separately to both *V. cholerae* O1 LPS and the O-antigen-specific polysaccharide (OSP) moiety of LPS. These responses were not only strongly correlated, but also nearly equivalent, suggesting that immune responses primarily target the OSP portion of the antigen, rather than the O-antigen core, lipid A, or other component of the LPS preparation. These findings have implications both for subsequent evaluations of immune responses following cholera infection or vaccination, and the development of more immunogenic vaccines for cholera.

Another finding in our study, was the significant correlation we observed between day 7 antigen-specific plasma IgA responses and day 7 antigen-specific ASC responses. This correlation has been previously described following infection with *V. cholerae* or vaccination, [8,34] and suggests that plasma IgA responses on day 7 following vaccination does indeed serve as a potential proxy for the mucosal ASC response in some circumstances.

Our study has a number of limitations. Because cholera is now endemic in Haiti, our cohort had high pre-existing levels of vibriocidal antibodies before vaccination with 50% of participants having a vibriocidal ≥ 80 for either the Ogawa or Inaba. These findings are in line with a serosurvey conducted in Haiti in 2011 demonstrating that 64% of individuals had a vibriocidal antibody titer ≥ 80 for either *V. cholerae* Inaba or Ogawa, suggesting high levels of prior infection. [41] This is important because ASC responses to vaccination are likely to differ between populations based on the level of previous antigen exposure. For example, in contrast to our results in Haiti where evidence of cholera exposure was common, a previous study conducted among Swedish volunteers demonstrated significant CT-specific ASC responses to a WC-rBS vaccine following the second dose of vaccine as part of a similar vaccination schedule spaced 14 days apart, suggesting differences between responses in endemic and non-endemic settings. [39] Levels of previous exposure also effect the kinetics of the ASC response, and ASC

responses peak earlier among primed populations.[42] Because we only measured ASC responses at a single time point after each vaccination (7-days) we could not evaluate the kinetic aspects of the response. While this is a limitation of the study, we do not think it changes the major findings, because based on previous studies, it is very unlikely that such priming would shift the kinetics of ASC responses to the extent they would become undetectable by day 7.[42] Lastly, while our results should be generalizable to adults living in other cholera endemic areas, additional studies are also needed in children and other subpopulations in cholera-affected regions, who may differ in their mucosal response to the BivWC vaccine.

In conclusion, vaccination with the BivWC oral cholera vaccine induces a significant mucosal ASC response targeting the OSP portion of *V. cholerae* LPS in Haitian adults, while a second dose of vaccine does not appear to stimulate antibody secreting cell responses in this population. These findings are similar to responses with the WC-rBS oral cholera vaccine in other cholera-endemic regions. More detailed evaluations of whether alternate vaccine dosing intervals can stimulate more robust mucosal immune responses, and contribute to improved duration of memory B cell targeting the *V. cholerae* O-antigen should be a priority for future studies.

Supporting Information

S1 Dataset. Table containing all relevant study data.
(XLS)

Acknowledgments

We are grateful to the study participants that made this work possible, and the Zanmi Lasante staff in Haiti that supported the study.

Author Contributions

Conceived and designed the experiments: WRM BF RCC LMMS JET ETR FQ MFF LCI JBH. Performed the experiments: WRM BF RCC LMMS. Analyzed the data: WRM BF RCC MFF LCI JBH. Contributed reagents/materials/analysis tools: PX PK FQ MFF. Wrote the paper: WRM BF RCC LMMS JET PX PK ETR FQ MFF LCI JBH.

References

1. Ali M, Lopez AL, You YA, Kim YE, Sah B, Maskery B, et al. The global burden of cholera. Bull World Health Organ [Internet]. 2012; 90(3):209–18A. Available from: <http://www.who.int/bulletin/volumes/90/3/11-093427.pdf> doi: [10.2471/BLT.11.093427](https://doi.org/10.2471/BLT.11.093427) PMID: [22461716](https://pubmed.ncbi.nlm.nih.gov/22461716/)
2. Harris JB, LaRocque RC, Qadri F, Ryan ET, Calderwood SB. Cholera. Lancet [Internet]. 2012; 379(9835):2466–76. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3761070&tool=pmcentrez&rendertype=abstract> doi: [10.1016/S0140-6736\(12\)60436-X](https://doi.org/10.1016/S0140-6736(12)60436-X) PMID: [22748592](https://pubmed.ncbi.nlm.nih.gov/22748592/)
3. Stroehrer UH, Karageorgos LE, Morona R, Manning P a. Serotype conversion in *Vibrio cholerae* O1. Proc Natl Acad Sci U S A [Internet]. 1992; 89(7):2566–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1372980> PMID: [1372980](https://pubmed.ncbi.nlm.nih.gov/1372980/)
4. Hisatsune K, Kondo S, Isshiki Y, Iguchi T, Haishima Y. Occurrence of 2-O-methyl-N-(3-deoxy-L-glycero-tetronyl)-D-perosamine (4-amino-4,6-dideoxy-D-manno-pyranose) in lipopolysaccharide from Ogawa but not from Inaba O forms of O1 *Vibrio cholerae*. Biochem Biophys Res Commun [Internet]. 1993; 190(1):302–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8422256> PMID: [8422256](https://pubmed.ncbi.nlm.nih.gov/8422256/)
5. Wang J, Villeneuve S, Zhang J, Lei PS, Miller CE, Lafaye P, et al. On the antigenic determinants of the lipopolysaccharides of *Vibrio cholerae* O:1, serotypes Ogawa and Inaba. J Biol Chem [Internet]. 1998; 273(5):2777–83. Available from: http://apps.webofknowledge.com/accesdistant.upmc.fr/full_record.do?product=WOS&search_mode=Refine&qid=65&SID=T13oOmbkP175DeAApmG&page=2&doc=12 PMID: [9446585](https://pubmed.ncbi.nlm.nih.gov/9446585/)

6. Ito T, Higuchi T, Hirobe M, Hiramatsu K, Yokota T. Identification of a novel sugar, 4-amino-4,6-dideoxy-2-O-methylmannose in the lipopolysaccharide of *Vibrio cholerae* O1 serotype Ogawa. *Carbohydr Res* [Internet]. 1994; 256(1):113–28. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8194067> PMID: [8194067](http://www.ncbi.nlm.nih.gov/pubmed/8194067)
7. Albert MJ, Alam K, Ansaruzzaman M, Qadri F, Sack RB. Lack of cross-protection against diarrhea due to *Vibrio cholerae* O139 (Bengal strain) after oral immunization of rabbits with *V. cholerae* O1 vaccine strain CVD103-HgR. *J Infect Dis* [Internet]. 1994 Jan; 169(1):230–1. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8277193> PMID: [8277193](http://www.ncbi.nlm.nih.gov/pubmed/8277193)
8. Qadri F, Wennerås C, Albert MJ, Hossain J, Mannoor K, Begum YA, et al. Comparison of immune responses in patients infected with *Vibrio cholerae* O139 and O1. *Infect Immun* [Internet]. 1997 Sep; 65(9):3571–6. Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9284121 PMID: [9284121](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=9284121)
9. Waldor MK, Colwell R, Mekalanos JJ. The *Vibrio cholerae* O139 serogroup antigen includes an O-antigen capsule and lipopolysaccharide virulence determinants. *Proc Natl Acad Sci U S A* [Internet]. 1994 Nov 22; 91(24):11388–92. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7972070> PMID: [7972070](http://www.ncbi.nlm.nih.gov/pubmed/7972070)
10. Centers for Disease Control and Prevention (CDC). Update: cholera outbreak—Haiti, 2010. *MMWR. Morbidity and mortality weekly report*. 2010. 1473–1479 p.
11. Ministère de la Santé Publique et de la Population. Rapport Journaliers du MSPP sur l'évolution du choléra en Haiti. 30 Septembre. [Internet]. 2015. Available from: http://mspp.gouv.ht/site/downloads/Rapport_Web_30.09.2015_Avec_Courbes_departementales.pdf
12. Martin S, Lopez AL, Bellos A, Deen J, Ali M, Alberti K, et al. Post-licensure deployment of oral cholera vaccines: a systematic review. *Bull World Health Organ*. 2014; 92(12):881–93. doi: [10.2471/BLT.14.139949](https://doi.org/10.2471/BLT.14.139949) PMID: [25552772](https://pubmed.ncbi.nlm.nih.gov/25552772/)
13. Ivers LC, Teng JE, Lascher J, Raymond M, Weigel J, Victor N, et al. Use of oral cholera vaccine in haiti: A rural demonstration project. *Am J Trop Med Hyg*. 2013; 89(4):617–24. doi: [10.4269/ajtmh.13-0183](https://doi.org/10.4269/ajtmh.13-0183) PMID: [24106187](https://pubmed.ncbi.nlm.nih.gov/24106187/)
14. Luquero FJ, Grout L, Ciglenecki I, Sakoba K, Traore B, Heile M, et al. First outbreak response using an oral cholera vaccine in Africa: vaccine coverage, acceptability and surveillance of adverse events, Guinea, 2012. *PLoS Negl Trop Dis*. 2013; 7(10):e2465. doi: [10.1371/journal.pntd.0002465](https://doi.org/10.1371/journal.pntd.0002465) PMID: [24147164](https://pubmed.ncbi.nlm.nih.gov/24147164/)
15. Martin S, Costa A, Perea W. Stockpiling oral cholera vaccine. *Bull World Health Organ* [Internet]. 2012; 90(10):714. Available from: <http://www.who.int/bulletin/volumes/90/10/12-112433.pdf> doi: [10.2471/BLT.12.112433](https://doi.org/10.2471/BLT.12.112433) PMID: [23109735](https://pubmed.ncbi.nlm.nih.gov/23109735/)
16. Anh DD, Canh DG, Lopez AL, Thiem VD, Long PT, Son NH, et al. Safety and immunogenicity of a reformulated Vietnamese bivalent killed, whole-cell, oral cholera vaccine in adults. *Vaccine* [Internet]. 2007; 25(6):1149–55. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0264410X06010498> PMID: [17055622](https://pubmed.ncbi.nlm.nih.gov/17055622/)
17. Mahalanabis D, Lopez AL, Sur D, Deen J, Manna B, Kanungo S, et al. A randomized, placebo-controlled trial of the bivalent killed, whole-cell, oral cholera vaccine in adults and children in a cholera endemic area in Kolkata, India. *PLoS One*. 2008; 3(6).
18. Kanungo S, Paisley A, Lopez AL, Bhattacharya M, Manna B, Kim DR, et al. Immune responses following one and two doses of the reformulated, bivalent, killed, whole-cell, oral cholera vaccine among adults and children in Kolkata, India: A randomized, placebo-controlled trial. *Vaccine*. 2009; 27(49):6887–93. doi: [10.1016/j.vaccine.2009.09.008](https://doi.org/10.1016/j.vaccine.2009.09.008) PMID: [19761838](https://pubmed.ncbi.nlm.nih.gov/19761838/)
19. Saha A, Chowdhury MI, Khanam F, Bhuiyan MS, Chowdhury F, Khan AI, et al. Safety and immunogenicity study of a killed bivalent (O1 and O139) whole-cell oral cholera vaccine Shanchol, in Bangladeshi adults and children as young as 1 year of age. *Vaccine*. 2011; 29(46):8285–92. doi: [10.1016/j.vaccine.2011.08.108](https://doi.org/10.1016/j.vaccine.2011.08.108) PMID: [21907255](https://pubmed.ncbi.nlm.nih.gov/21907255/)
20. Baik YO, Choi SK, Olveda RM, Espos R a., Ligsay AD, Montellano MB, et al. A randomized, non-inferiority trial comparing two bivalent killed, whole cell, oral cholera vaccines (Euvichol vs Shanchol) in the Philippines. *Vaccine* [Internet]. Elsevier Ltd; 2015;6–11. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0264410X15012281>
21. Charles RC, Hilaire IJ, Mayo-Smith LM, Teng JE, Jerome JG, Franke MF, et al. Immunogenicity of a killed bivalent (O1 and O139) whole cell oral cholera vaccine, Shanchol, in Haiti. *PLoS Negl Trop Dis* [Internet]. 2014; 8(5):e2828. Available from: <http://dx.plos.org/10.1371/journal.pntd.0002828> doi: [10.1371/journal.pntd.0002828](https://doi.org/10.1371/journal.pntd.0002828) PMID: [24786645](https://pubmed.ncbi.nlm.nih.gov/24786645/)
22. Ivers LC, Charles RC, Hilaire IJ, Mayo-Smith LM, Teng JE, Jerome JG, et al. Immunogenicity of the Bivalent Oral Cholera Vaccine Shanchol in Haitian Adults With HIV Infection. *J Infect Dis* [Internet].

- 2015; 212(5):779–83. Available from: <http://jid.oxfordjournals.org/lookup/doi/10.1093/infdis/jiv108> doi: [10.1093/infdis/jiv108](https://doi.org/10.1093/infdis/jiv108) PMID: [25722294](https://pubmed.ncbi.nlm.nih.gov/25722294/)
23. Jayasekera CR, Harris JB, Bhuiyan S, Chowdhury F, Khan AI, Faruque ASG, et al. Cholera toxin-specific memory B cell responses are induced in patients with dehydrating diarrhea caused by *Vibrio cholerae* O1. *J Infect Dis* [Internet]. 2008; 198(7):1055–61. Available from: <http://jid.oxfordjournals.org/content/198/7/1055.full> doi: [10.1086/591500](https://doi.org/10.1086/591500) PMID: [18729782](https://pubmed.ncbi.nlm.nih.gov/18729782/)
 24. Harris AM, Bhuiyan MS, Chowdhury F, Khan AI, Hossain A, Kendall E a., et al. Antigen-specific memory B-cell responses to *Vibrio cholerae* O1 infection in Bangladesh. *Infect Immun*. 2009; 77(9):3850–6. doi: [10.1128/IAI.00369-09](https://doi.org/10.1128/IAI.00369-09) PMID: [19528207](https://pubmed.ncbi.nlm.nih.gov/19528207/)
 25. Kuchta A, Rahman T, Sennott EL, Bhuyian TR, Uddin T, Rashu R, et al. *Vibrio cholerae* O1 infection induces proinflammatory CD4 + T-cell responses in blood and intestinal mucosa of infected humans. *Clin Vaccine Immunol*. 2011; 18(8):1371–7. doi: [10.1128/CVI.05088-11](https://doi.org/10.1128/CVI.05088-11) PMID: [21697339](https://pubmed.ncbi.nlm.nih.gov/21697339/)
 26. Leung DT, Rahman MA, Mohasin M, Riyadh MA, Patel SM, Alam MM, et al. Comparison of memory B cell, antibody-secreting cell, and plasma antibody responses in young children, older children, and adults with infection caused by *Vibrio cholerae* O1 El Tor Ogawa in Bangladesh. *Clin Vaccine Immunol* [Internet]. 2011; 18(8):1317–25. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.05124-11> doi: [10.1128/CVI.05124-11](https://doi.org/10.1128/CVI.05124-11) PMID: [21697337](https://pubmed.ncbi.nlm.nih.gov/21697337/)
 27. Leung DT, Chowdhury F, Calderwood SB, Qadri F, Ryan ET. Immune responses to cholera in children. *Expert Rev Anti Infect Ther* [Internet]. 2012; 10(4):435–44. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3378723&tool=pmcentrez&rendertype=abstract> doi: [10.1586/eri.12.23](https://doi.org/10.1586/eri.12.23) PMID: [22512753](https://pubmed.ncbi.nlm.nih.gov/22512753/)
 28. Johnson RA, Uddin T, Aktar A, Mohasin M, Alam MM, Chowdhury F, et al. Comparison of immune responses to the O-specific polysaccharide and lipopolysaccharide of *Vibrio cholerae* O1 in Bangladeshi adult patients with cholera. *Clin Vaccine Immunol* [Internet]. 2012; 19(11):1712–21. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.00321-12> doi: [10.1128/CVI.00321-12](https://doi.org/10.1128/CVI.00321-12) PMID: [22993410](https://pubmed.ncbi.nlm.nih.gov/22993410/)
 29. Alam MM, Riyadh MA, Fatema K, Rahman MA, Akhtar N, Ahmed T, et al. Antigen-specific memory B-cell responses in Bangladeshi adults after one- or two-dose oral killed cholera vaccination and comparison with responses in patients with naturally acquired cholera. *Clin Vaccine Immunol*. 2011; 18(5):844–50. doi: [10.1128/CVI.00562-10](https://doi.org/10.1128/CVI.00562-10) PMID: [21346055](https://pubmed.ncbi.nlm.nih.gov/21346055/)
 30. Arifuzzaman M, Rashu R, Leung DT, Hosen MI, Bhuiyan TR, Bhuiyan MS, et al. Antigen-specific memory T cell responses after vaccination with an oral killed cholera vaccine in Bangladeshi children and comparison to responses in patients with naturally acquired cholera. *Clin Vaccine Immunol*. 2012; 19(8):1304–11. doi: [10.1128/CVI.00196-12](https://doi.org/10.1128/CVI.00196-12) PMID: [22739692](https://pubmed.ncbi.nlm.nih.gov/22739692/)
 31. Leung DT, Rahman MA, Mohasin M, Patel SM, Aktar A, Khanam F, et al. Memory B cell and other immune responses in children receiving two doses of an oral killed cholera vaccine compared to responses following natural cholera infection in Bangladesh. *Clin Vaccine Immunol* [Internet]. 2012; 19(5):690–8. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.05615-11> doi: [10.1128/CVI.05615-11](https://doi.org/10.1128/CVI.05615-11) PMID: [22441386](https://pubmed.ncbi.nlm.nih.gov/22441386/)
 32. Uddin T, Aktar A, Xu P, Johnson R a., Rahman MA, Leung DT, et al. Immune responses to O-specific polysaccharide and lipopolysaccharide of *Vibrio cholerae* O1 Ogawa in adult Bangladeshi recipients of an oral killed cholera vaccine and comparison to responses in patients with cholera. *Am J Trop Med Hyg* [Internet]. 2014; 90(5):873–81. Available from: <http://www.ajtmh.org/cgi/doi/10.4269/ajtmh.13-0498> doi: [10.4269/ajtmh.13-0498](https://doi.org/10.4269/ajtmh.13-0498) PMID: [24686738](https://pubmed.ncbi.nlm.nih.gov/24686738/)
 33. Leung DT, Uddin T, Xu P, Aktar A, Johnson RA, Rahman MA, et al. Immune responses to the O-specific polysaccharide antigen in children who received a killed oral cholera vaccine compared to responses following natural cholera infection in Bangladesh. *Clin Vaccine Immunol* [Internet]. 2013; 20(6):780–8. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.00035-13> doi: [10.1128/CVI.00035-13](https://doi.org/10.1128/CVI.00035-13) PMID: [23515016](https://pubmed.ncbi.nlm.nih.gov/23515016/)
 34. Rahman A, Rashu R, Bhuiyan TR, Chowdhury F, Khan AI, Islam K, et al. Antibody-secreting cell responses after *Vibrio cholerae* O1 infection and oral cholera vaccination in adults in Bangladesh. *Clin Vaccine Immunol* [Internet]. 2013; 20(10):1592–8. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.00347-13> doi: [10.1128/CVI.00347-13](https://doi.org/10.1128/CVI.00347-13) PMID: [23945156](https://pubmed.ncbi.nlm.nih.gov/23945156/)
 35. Qadri F, Mäkelä PH, Holmgren J, Albert MJ, Mannoor K, Kantele A, et al. Enteric infections in an endemic area induce a circulating antibody-secreting cell response with homing potentials to both mucosal and systemic tissues. *J Infect Dis*. 1998; 177(6):1594–9. PMID: [9607838](https://pubmed.ncbi.nlm.nih.gov/9607838/)
 36. Uddin T, Harris JB, Bhuiyan TR, Shirin T, Uddin MI, Khan AI, et al. Mucosal immunologic responses in cholera patients in Bangladesh. *Clin Vaccine Immunol* [Internet]. 2011; 18(3):506–12. Available from: <http://cvi.asm.org/cgi/doi/10.1128/CVI.00481-10> doi: [10.1128/CVI.00481-10](https://doi.org/10.1128/CVI.00481-10) PMID: [21248157](https://pubmed.ncbi.nlm.nih.gov/21248157/)
 37. Xu P, Alam MM, Kalsy A, Charles RC, Calderwood SB, Qadri F, et al. Simple, direct conjugation of bacterial O-SP-core antigens to proteins: development of cholera conjugate vaccines. *Bioconjug Chem*

- [Internet]. 2011 Oct 19; 22(10):2179–85. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21899371> doi: [10.1021/bc2001984](https://doi.org/10.1021/bc2001984) PMID: [21899371](https://pubmed.ncbi.nlm.nih.gov/21899371/)
38. Sayeed MA, Bufano MK, Xu P, Eckhoff G, Charles RC, Alam MM, et al. A Cholera Conjugate Vaccine Containing O-specific Polysaccharide (OSP) of *V. cholerae* O1 Inaba and Recombinant Fragment of Tetanus Toxin Heavy Chain (OSP:rTTHc) Induces Serum, Memory and Lamina Proprial Responses against OSP and Is Protective in Mice. *PLoS Negl Trop Dis* [Internet]. 2015 Jul; 9(7):e0003881. Available from: <http://dx.plos.org/10.1371/journal.pntd.0003881> doi: [10.1371/journal.pntd.0003881](https://doi.org/10.1371/journal.pntd.0003881) PMID: [26154421](https://pubmed.ncbi.nlm.nih.gov/26154421/)
 39. Czerkinsky C, Svennerholm A-M, Quiding M, Jonsson R, Holmgren J. Antibody-producing cells in peripheral blood and salivary glands after oral cholera vaccination of humans. *Infect Immun*. 1991; 59(3):996–1001. PMID: [1997444](https://pubmed.ncbi.nlm.nih.gov/1997444/)
 40. Shamsuzzaman S, Ahmed T, Mannoor K, Begum YA, Bardhan PK, Sack RB, et al. Robust gut associated vaccine-specific antibody-secreting cell responses are detected at the mucosal surface of Bangladeshi subjects after immunization with an oral killed bivalent *V. cholerae* O1/O139 whole cell cholera vaccine: comparison with other muco. *Vaccine* [Internet]. 2009 Feb 25; 27(9):1386–92. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19146897> doi: [10.1016/j.vaccine.2008.12.041](https://doi.org/10.1016/j.vaccine.2008.12.041) PMID: [19146897](https://pubmed.ncbi.nlm.nih.gov/19146897/)
 41. Jackson BR, Talkington DF, Pruckler JM, Fouché MDB, Lafosse E, Nygren B, et al. Seroepidemiologic survey of epidemic cholera in Haiti to assess spectrum of illness and risk factors for severe disease. *Am J Trop Med Hyg*. 2013; 89(4):654–64. doi: [10.4269/ajtmh.13-0208](https://doi.org/10.4269/ajtmh.13-0208) PMID: [24106192](https://pubmed.ncbi.nlm.nih.gov/24106192/)
 42. Leach S, Lundgren A, Svennerholm A-M. Different kinetics of circulating antibody-secreting cell responses after primary and booster oral immunizations: a tool for assessing immunological memory. *Vaccine* [Internet]. Elsevier Ltd; 2013 Jun 26; 31(30):3035–8. Available from: <http://linkinghub.elsevier.com/retrieve/pii/S0264410X13005355> doi: [10.1016/j.vaccine.2013.04.066](https://doi.org/10.1016/j.vaccine.2013.04.066) PMID: [23664997](https://pubmed.ncbi.nlm.nih.gov/23664997/)