



# miR-17-3p Exacerbates Oxidative Damage in Human Retinal Pigment Epithelial Cells

## Citation

Tian, Bo, Daniel E. Maidana, Bernard Dib, John B. Miller, Peggy Bouzika, Joan W. Miller, Demetrios G. Vavvas, and Haijiang Lin. 2016. "miR-17-3p Exacerbates Oxidative Damage in Human Retinal Pigment Epithelial Cells." PLoS ONE 11 (8): e0160887. doi:10.1371/journal.pone.0160887. <http://dx.doi.org/10.1371/journal.pone.0160887>.

## Published Version

doi:10.1371/journal.pone.0160887

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:29002607>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

RESEARCH ARTICLE

# miR-17-3p Exacerbates Oxidative Damage in Human Retinal Pigment Epithelial Cells

Bo Tian<sup>1,2</sup>, Daniel E. Maidana<sup>1</sup>, Bernard Dib<sup>1</sup>, John B. Miller<sup>1</sup>, Peggy Bouzika<sup>1</sup>, Joan W. Miller<sup>1</sup>, Demetrios G. Vavvas<sup>1\*</sup>, Haijiang Lin<sup>1\*</sup>

**1** Retina Service, Angiogenesis Laboratory, Department of Ophthalmology, Massachusetts Eye and Ear Infirmary, Harvard Medical School, Boston, MA, United States of America, **2** Department of Ophthalmology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan, P.R. China

\* [Haijiang\\_Lin@meei.harvard.edu](mailto:Haijiang_Lin@meei.harvard.edu) (HL); [Demetrios\\_Vavvas@meei.harvard.edu](mailto:Demetrios_Vavvas@meei.harvard.edu) (DGV)



**OPEN ACCESS**

**Citation:** Tian B, Maidana DE, Dib B, Miller JB, Bouzika P, Miller JW, et al. (2016) miR-17-3p Exacerbates Oxidative Damage in Human Retinal Pigment Epithelial Cells. PLoS ONE 11(8): e0160887. doi:10.1371/journal.pone.0160887

**Editor:** Zsolt Ablonczy, Medical University of South Carolina, UNITED STATES

**Received:** April 17, 2016

**Accepted:** July 26, 2016

**Published:** August 9, 2016

**Copyright:** © 2016 Tian et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data are available from Dryad in the following temporary url: <http://datadryad.org/review?doi=doi:10.5061/dryad.cf860>

**Funding:** This work was supported by: NEI R21EY023079-01A1 (DGV); the Yeatts Family Foundation (DGV, JWM); the Loefflers Family Fund (DGV, JWM); the 2013 Macula Society Research Grant award (DGV); a Physician Scientist Award (DGV), an unrestricted grant (JWM) from the research to prevent blindness foundation; NEI grant EY014104 (Meei Core Grant); and Bayer Healthcare Global Ophthalmology Award (DEM). The funders had no role in study design, data collection and

## Abstract

Oxidative stress has been shown to contribute to the development of age-related macular degeneration (AMD). MicroRNAs (miRNA) are small non-coding RNA molecules that function in RNA silencing and post-transcriptional regulation of gene expression. We showed miR-17-3p to be elevated in macular RPE cells from AMD patients and in ARPE-19 cells under oxidative stress. Transfection of miR-17-3p mimic in ARPE-19 induced cell death and exacerbated oxidative lethality that was alleviated by miR-17-3p inhibitor. The expression of antioxidant enzymes manganese superoxide dismutase (MnSOD) and thioredoxin reductase-2 (TrxR<sub>2</sub>) were suppressed by miR-17-3p mimic and reversed by miR-17-3p inhibitor. These results suggest miR-17-3p aggravates oxidative damage-induced cell death in human RPE cells, while miR-17-3p inhibitor acts as a potential protector against oxidative stress by regulating the expression of antioxidant enzymes.

## Introduction

Age-related macular degeneration is the leading cause of irreversible blindness among the elderly in developed countries. Although therapeutic strategies for the less prevalent neovascular form exist, therapies for the more prevalent non-neovascular (“dry”) form are lacking. A better understanding its pathogenesis may guide researchers towards the development of novel therapies. It has been postulated that oxidative stress [1–4] and inflammation [5–13] play a critical role in AMD pathogenesis. Retinal pigment epithelium (RPE) cells are extremely important to photoreceptor function, being responsible for the recycling of the visual pigments and in phagocytosis of photoreceptor outer segments [14,15]. Due to its location in a highly oxygenated and lit environment, RPE is at high risk for oxidative injury that could lead to cellular dysfunction, inflammation, and eventually cell death [2–4,16–20].

There are several enzymes responsible for oxidative stress management, such as superoxide dismutase (SOD) and thioredoxin reductase. Among the three SOD isoforms, MnSOD is the essential mitochondrial antioxidant enzyme. It has been demonstrated that RPE cells deriving from MnSOD-deficient mice are more susceptible to oxidative stress than wild-type RPE cells [21]. Three well characterized isoenzymes of thioredoxin reductases, namely cytosolic TrxR<sub>1</sub>,

analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have no conflicts of interest to disclose.

mitochondrial TrxR<sub>2</sub>, and testicular TrxR<sub>3</sub> are flavoproteins that reduce thioredoxin, a major protein involved in the reduction of cellular oxidative stress [22,23]. TrxR<sub>2</sub> controls H<sub>2</sub>O<sub>2</sub> emission by maintaining the level of active thioredoxin [24], which has been shown to efficiently decrease RPE cell death caused by oxidative stress [25]. Overall, dysfunction of antioxidant enzymes can cause ROS accumulation, and ROS defense systems have been shown to be important in RPE [26–36]. Consequently, the regulation of antioxidant enzymes is important in the study of AMD, as efficient antioxidant defense systems are needed to protect the RPE cells.

MicroRNAs (miRNAs) are small non-coding RNAs that play an essential role in regulating gene expression, either by degrading messenger RNA (mRNA) or stalling translation [37]. They can be generated from either the 5p or 3p or both arms of pre-miRNA, and depending on which arm they are generated from, they are notated as: miR-#-5p or miR-#-3p [38]. These molecules have been proven to be involved in extensive pathological processes, including angiogenesis, oxidative stress, immune response and inflammation [39–42], all of which are critical processes in age-related macular degeneration (AMD). MicroRNA-30b has previously been shown to impair oxidative stress mechanisms in ARPE-19 [43], whereas miR-9 has been shown to be upregulated by a retinoic acid analogue in the same cells [44]. Using human iPSC-derived RPE cells under Paraquat stress, Garcia et al. showed upregulation of miR-146a and miR-29a, downregulation of miR-144, miR-200a and miR-21, whereas a biphasic response was seen on miR-27b [45]. We previously reported that miR-23 enhances RPE cell resistance to oxidative stress damage and is downregulated in macular RPE cells from AMD patients [46]. miR-17-3p is a member of miR-17/92 cluster, originally found to be involved in tumorigenesis, but more recently, members of this cluster have been shown to be involved in many aging disorders [47]. Although most of the work regarding miR-17-3p has focused on regulation of cell proliferation pathways, a study using prostate cancer cell lines demonstrated that miR-17-3p is also involved in regulating antioxidant enzymes [48]. In this study, we aim to explore the role of miR-17-3p in ARPE-19 cell viability and antioxidant enzyme production under oxidative stress, a major factor in AMD pathogenesis.

## Materials and Methods

### Cell culture

Human donor eyes from AMD patients (70–90 years old) and age-matched controls were acquired from the Minnesota Lions Eye Bank (Saint Paul, MN), in accordance with the provisions of the Declaration of Helsinki for research involving human tissues. Macular RPE cells were isolated as explained previously [49]. Briefly, cornea, anterior segment, vitreous, and neural retina were carefully removed without disturbing the RPE layer. The dissection was performed by an 8-mm sterile trephine punch through the RPE cell layer, Bruch's membrane, and choroid, centered on the macula. RPE cells from this region were collected as macular RPE cells. Subsequently, the RPE cells were dissociated after trypsin digestion (30 minutes at 37°C) in pre-warmed medium (DMEM/F12; Cat#11330–057, Gibco, Grand Island, NY). Centrifugation of cells for 5 minutes was carried at 168 × g at 4°C. Supernatant was carefully aspirated and the cell pellet was re-suspended in DMEM medium. To guarantee the purity of RPE cells during the isolating procedure, the isolated cells were analyzed morphologically and stained with cytokeratin-18 antibody, which confirmed that the percentage of RPE cells was more than 97%.

Human retinal pigment epithelium and ARPE-19 cells (CRL-2302, ATCC, Manassas, VA) were cultured in medium (DMEM/F-12; Cat#11330–057, Gibco, Grand Island, NY) supplemented with 10% fetal bovine serum (FBS; Cat#10438–026, Gibco, Grand Island, NY) and

penicillin-streptomycin (100 U/mL-100 $\mu$ g/ml; Cat#15140, Gibco, Grand Island, NY) at 37°C in a humidified 5% CO<sub>2</sub> incubator.

### miRNA transfection

ARPE-19 cells were cultured in DMEM/F12 medium to reach 70% confluence, and transfection was performed for functional analysis. miR-17-3p mimic (Cat#4464066, Life technologies), antisense miR-17-3p mimic (inhibitor, Cat#4464084, Life technologies), scrambled mimic (sc-mimic, Cat#4464058, Life technologies), and scrambled inhibitor (sc-inhibitor, Cat#4464076, Life technologies) were transfected into ARPE-19 cells with Lipofectamine<sup>®</sup> RNAiMAX (Cat#13778030, Invitrogen, Carlsbad, CA) according to the manufacturer's manual, respectively. The final concentration of miRNAs was 20 nM. Cells were then incubated for 72 h in a humidified incubator with 5% CO<sub>2</sub> at 37°C until utilized for experiments.

### H<sub>2</sub>O<sub>2</sub> and TBH treatment

ARPE-19 cells were seeded to a density of 3.5 × 10<sup>5</sup>/ml in 6-well plates, and cultured in medium supplemented with 10% fetal bovine serum and penicillin-streptomycin. Twenty-four hours later, cells at 90~100% confluence were exposed to H<sub>2</sub>O<sub>2</sub> (25, 50, 100, or 200 $\mu$ M) (Cat#H-1009, Sigma-Aldrich, St.Louis, MO) and tert-Butyl hydrogen peroxide solution, TBH (Cat#458139, 37.5, 75, or 150 $\mu$ M) (Sigma-Aldrich, St. Louis, MO) diluted in cell culture medium (DMEM/F12; Cat#11330-057, Gibco, Grand Island, NY) without fetal bovine serum or phenol red.

### Cell viability assay

Cell viability was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, Cat#M6494, Molecular Probes, Eugene, OR) assay. Specifically, after the incubation, the cell medium was aspirated from the 96-well plate. Cells were then washed with Dulbecco's Phosphate Buffer Saline (DPBS; Cat#17-515Q, Lonza, Walkersville, MD) twice and treated with 0.5mg/ml MTT. Following 3 hours of incubation at 37°C, the MTT solution was aspirated and 100 $\mu$ l of hydrochloride-isopropanol (0.04N) (Cat#258148, Cat#109827, Sigma-Aldrich, St. Louis, MO) was added to each well. Plates were incubated for 15 minutes on a shaker and absorbance was read at 590nm using a SpectraMax 190 microplate reader (Molecular Devices, Sunnyvale, CA).

### Determination of miR-17-3p expression level

Primary cultured macular RPE or ARPE-19 cells were seeded at 2 × 10<sup>5</sup>/ well in 6-well plates and incubated to 80% confluence. Primary cultured RPE cells were then harvested at passage 3. Total RNA was extracted from RPE cells (miRNeasy Mini kit; Cat#217004, QIAGEN, Germany). Complementary DNA (cDNA) was synthesized using predesigned TaqMan miRNA Assays (Applied Biosystems, Foster City, CA) and the Reverse Transcription Kit (Cat#4368814, Applied Biosystems, Foster City, CA) for miRNA after total RNA extraction according to the manufacturer's instructions. Briefly, the reaction contained a volume of 5 $\mu$ l RNA extract, 3 $\mu$ l of reverse transcription (RT) primer, and 7 $\mu$ l master mix. RT reactions were performed on a Thermal cycler (Bio-Rad, Hercules, CA) under the following conditions: 16°C for 30 min, 42°C for 30 min, 85°C for 5 min, and 4°C on hold. All TaqMan assays were run in triplicate on an AB Step One Plus real-time PCR system using TaqMan Universal PCR Master Mix II without UNG (Cat#4440040, Applied Biosystems, Foster City, CA). Real-time PCR cycling conditions consisted of 95°C for 10 min, followed by 40 cycles of 95°C for 15 s and 60°C for 1 min.

## Western blot

Total protein was extracted from cultured ARPE-19 cells using NP40 cell lysis buffer (Cat#FNN0021, Invitrogen, Frederick, MD), and the concentration was calculated by DC protein assay (Cat#500-0016, Bio-Rad, Philadelphia, PA). Equivalent amounts of extracted protein were loaded into each well and electrophoresed on a Nupage 4–12% Bis-Tris gel (Cat#NP0335BOX, Novex, Carlsbad, CA), then transferred to a 0.2 $\mu$ m PVDF membrane (Cat#ISEQ00010, Millipore, Billerica, MA), blocked with non-fat dry milk (Cat#9999, Cell signaling, Danvers, MA) and incubated with primary antibodies against MnSOD (Cat#16956, Abcam, Cambridge, MA) at 1:2000, TrxR<sub>2</sub> (Cat#sc-46278, Santa Cruz Biotechnology, Dallas, TX) at 1:250, and GAPDH (Cat#5174, Cell Signaling Technology, Danvers, MA) at 1:1000, for 2 hours. The membrane was washed 3 times for 10 minutes each time and incubated with secondary antibodies at 1:10000 for 1.5 hours. After washing, membranes were incubated with the chemiluminescent substrate (Cat#RPN2235, ECL Select western blotting detection reagents, GE Healthcare Life Sciences, Piscataway, NJ) for 5 minutes. Band signals were detected by an image-scanning densitometer (ChemIDoc imaging system; Bio-Rad) and quantitated by Image J 2.0.

## TUNEL (terminal deoxynucleotidyl transferase-mediated dUTP nick-end labelling) staining

ARPE-19 cells were washed with PBS once and then fixed in 4% paraformaldehyde in PBS for 10 min at room temperature (25°C). After two washes in PBS for 5 minutes each, cells were post-fixed in cooled ethanol-acetic acid (v:v, 2:1) and stained using an In Situ Apoptosis Detection Kit (Cat#S7110, Millipore, Temecula, CA) according to the manufacturer's protocol. The cell nuclei were counterstained with DAPI (4', 6-diamidino-2-phenylindole, Cat#62248, Life technologies). The number of TUNEL-positive cells was counted and processed for statistical analysis.

## Statistical analysis

Statistical analysis was performed on GraphPad Prism 7.0a. Statistically significant differences between two groups were assessed using Student's t-test. One-way ANOVA followed by Tukey's post-hoc test was applied to multiple group comparisons. Differences with  $p < 0.05$  were considered statistically significant.


## Results

### miR-17-3p expression increases in macular RPE cells from AMD patients

A microarray assay previously performed by our group showed that miR-17-3p expression level increases in macular RPE cells of AMD patients' donor eyes compared to age-matched normal control donor eyes. To confirm this data, total RNA was isolated from macular RPE cells of each AMD donor eye ( $n = 5$ ) and age-matched normal control eye ( $n = 5$ ). miR-17-3p expression level was analyzed by qRT-PCR. [Fig 1](#) illustrates that miR-17-3p was significantly upregulated ( $p < 0.05$ ) in macular RPE cells from AMD donor eyes compared to controls (1.4-fold). It suggests that increased miR-17-3p expression may be associated with AMD.

### miR-17-3p causes ARPE-19 cell death

Due to the fact that miR-17-3p was upregulated in macular RPE cells from AMD patients, we hypothesized that it may be implicated in AMD pathogenesis. Since RPE cell atrophy and






**Fig 1. Differential miR-17-3p expression in macular RPE cells from AMD patients and normal controls.** Expression levels of miR-17-3p in macular RPE cells from AMD and age-matched normal control donor eyes were analyzed by qRT-PCR. The expression levels are shown relative to control values in five independent experiments. Results are displayed as mean  $\pm$  SD. \* $p < 0.05$ .

doi:10.1371/journal.pone.0160887.g001

death are features of AMD, we assessed cell viability by MTT assay following miR-17-3p transfection into ARPE-19 cells. miR-17-3p mimic sequence transfection caused significant reduction in cell viability (63.7%) compared to scrambled sequence (95.4%), and this effect was partially reversed by co-transfection with miR-17-3p-specific inhibitor (87.5%), but not a scrambled inhibitor (Fig 2A). Moreover, the ratio of TUNEL-positive cells/total cells in the miR-17-3p transfected group was 30.5% higher compared to the scrambled sequence group, while co-transfection with the inhibitor abrogated miR-17-3p cell death (Fig 2B and 2C). These results indicate that miR-17-3p induces ARPE-19 cell death.

### Increased expression of miR-17-3p in ARPE-19 under oxidative stress

Since oxidative stress has been linked to RPE senescence and AMD [50], we sought to evaluate miR-17-3p expression in RPE cells under oxidative stress. It was found that miR-17-3p was raised 1.3-, 1.5- and 2.3-fold in ARPE-19 cells after treatment with increasing concentrations of  $H_2O_2$  (25, 50, 100 $\mu$ M) compared with the untreated control respectively (Fig 3A). Meanwhile, treatment with a similar oxidant stimulus, TBH, exhibited an analogous increase of miR-17-3p expression ranging from 1.2- to 2-fold compared with the untreated control (Fig 3B). However, doses higher than 100 $\mu$ M of  $H_2O_2$  and 75 $\mu$ M of TBH failed to increase miR-17-3p expression, most likely due to cytotoxicity. These results indicate that miR-17-3p can be up regulated in ARPE-19 cells in the presence of oxidizing stimuli.




**Fig 2. Effect of miR-17-3p on ARPE-19 cell viability.** (A) ARPE-19 cells were transfected with miR-17-3p mimic (mimic), scrambled miR-17-3p mimic (sc-mimic), miR-17-3p antisense (inhibitor) or scrambled miR-17-3p antisense (sc-inhibitor) as indicated for 72 hours. Cell viability was assessed by MTT assay. The mimic sequence transfection reduced cell viability compared to scrambled sequence (sc-mimic); however, decreased cell viability was reversed by co-transfection with miR-17-3p inhibitor. Data shown as mean  $\pm$  SD. (n = 8) (B) After 72 hours of transfection, cell death was evaluated by TUNEL staining. Representative images were chosen from three independent experiments. (C) The average number of TUNEL positive cells from different transfection groups was counted from six random microscope fields. Results are displayed as mean  $\pm$  SD. \* $p$ <0.05.

doi:10.1371/journal.pone.0160887.g002


### miR-17-3p increases vulnerability of RPE cells to oxidative stress

Since miR-17-3p was shown to cause cell death in ARPE-19 cells, and its expression was upregulated under oxidative stress, we speculated that miR-17-3p could potentially augment RPE cell death under oxidative stress. ARPE-19 cells were differentially transfected with miR-17-3p (mimic) or co-transfected with miR-17-3p antisense (inhibitor) for 72 hours, and then treated with H<sub>2</sub>O<sub>2</sub> (100 $\mu$ M) or TBH (75 $\mu$ M) for 9 hours. Cell viability was assessed by MTT assay, which showed that cell death increased to 26.3% and 19.6% after H<sub>2</sub>O<sub>2</sub> and TBH treatment, respectively, compared with the untreated group. Transfection with miR-17-3p mimic further increased this value to 58.9% and 51.8%, respectively (Fig 4A and 4B), whereas transfection with miR-17-3p inhibitor but not scrambled inhibitor was able to reverse H<sub>2</sub>O<sub>2</sub> and TBH toxicity (37.7% and 31.5% of cell death by co-transfection with inhibitor in the H<sub>2</sub>O<sub>2</sub> and TBH treated groups, respectively). Moreover, transfection with miR-17-3p antisense (INH) alone but not scrambled inhibitor (sc-INH) alone was able to partially reverse H<sub>2</sub>O<sub>2</sub> or TBH toxicity (Fig 4). These data suggest that miR-17-3p can increase ARPE-19 cell vulnerability to oxidative stress and miR-17-3p antisense plays a protective role under oxidative stress.



**Fig 3. Expression of miR-17-3p in ARPE-19 cells under increasing amounts of oxidative stress.** ARPE-19 cells were treated with different concentrations of H<sub>2</sub>O<sub>2</sub> (0, 25, 50, 100, 200 $\mu$ M) or TBH (0, 37.5, 75, 150 $\mu$ M) for 6 hours. miR-17-3p expression levels were determined by qRT-PCR in both H<sub>2</sub>O<sub>2</sub> (A) and TBH (B) treated group. Results are displayed as mean  $\pm$  SD. \* $p$ <0.05.

doi:10.1371/journal.pone.0160887.g003



**Fig 4. miR-17-3p increases vulnerability of RPE cells to oxidative stress.** ARPE-19 cells were transfected with miR-17-3p (mimic) or miR-17-3p antisense (inhibitor, INH) as indicated for 72 hours, and then treated with (A) H<sub>2</sub>O<sub>2</sub> (100µM) or (B) TBH (75µM) for 9 hours. Cell viability was then determined by MTT assay. Data are shown as mean ± SD. (n = 8) \*p<0.05.

doi:10.1371/journal.pone.0160887.g004

### miR-17-3p reduced the expression of MnSOD and TrxR<sub>2</sub> in ARPE-19 cells under oxidative stress

To explore the potential targets of miR-17-3p, which are responsible for the increased vulnerability under oxidative stress, we performed a bioinformatics sequence analysis and found that two antioxidant enzymes, MnSOD and TrxR<sub>2</sub>, possess binding sites of miR-17-3p at the 3' untranslated regions (3' UTR) of mRNA (Fig 5A), which is validated by Xu et al. using luciferase assay [48]. miR-17-3p mimic transfection reduced MnSOD and TrxR<sub>2</sub> protein expression levels to 39.6% and 56.9% respectively compared to a transfection control. Furthermore, antisense-miR-17-3p was able to reverse MnSOD and TrxR<sub>2</sub> downregulations to 78.7% and 81.0%, respectively (Fig 5B and 5C). Collectively, these results demonstrate that miR-17-3p is a negative regulator of two important antioxidant enzymes in ARPE-19 cells.

### Discussion

Our study provides evidence that miR-17-3p is elevated in macular RPE cells of AMD patients and is cytotoxic to ARPE-19 cells. This cytotoxic effect can be partially reversed by the inhibitor of miR-17-3p. Furthermore, the expression level of miR-17-3p increases under oxidative stress, and it increases ARPE-19 vulnerability to oxidative stress by downregulating the mitochondrial antioxidant enzymes MnSOD and TrxR<sub>2</sub>. This effect can be abolished by the inhibitor of miR-17-3p.

There are only a few studies examining the role of miRNA in dry AMD. Our 2011 study [46] was the first to show that miR-23 is downregulated in human AMD eyes and its putative protective effect is mediated by downregulation of the Fas ligand. A year later, Lukiw et al. [51] showed that several miRNAs, including miRNA-9, miRNA-125b, miRNA-146a and miRNA-155, were upregulated in both Alzheimer's disease and AMD, and they were common to the pathogenetic mechanism of complement factor H (CFH) deficiency which drives inflammatory neurodegeneration. In 2014, Murad et al. [52] reported that miR-184 was significantly downregulated in primary RPE cells isolated from AMD donors, and its downregulation resulted in



apoptosis [46,55,56]. TBH is a short-chain organic hydro peroxide, also used as an oxidant for investigating mechanisms of cellular alterations resulting from oxidative stress in cells and tissues, such as the RPE [57]. We found that miR-17-3p was elevated in ARPE-19 cells after treatment with increasing concentrations of oxidizing reagents H<sub>2</sub>O<sub>2</sub> and TBH. Multiple oxidation-sensitive genes and factors are induced in RPE cells exposed to ROS [58,59]. Alterations in gene expression in response to oxidative stress have been extensively studied at transcriptional levels [60,61], however it is accepted that post-transcriptional regulation of gene expression is also very important [62]. MicroRNAs serve as significant post-transcriptional regulators of gene expression.

Evidence stemming from prior studies in prostate cancer cells as well as from our bioinformatic analysis suggests that antioxidant enzymes MnSOD and TrxR<sub>2</sub> act as targets of miR-17-3p [48]. MnSOD, a ROS scavenger, plays a vital role in promoting cell survival under stressful conditions, and several studies have confirmed that the decrease in cell viability observed under oxidative stress is correlated with MnSOD downregulation [21,63]. Moreover, some exogenous antioxidants, such as estrogen and progesterone, exhibit their antioxidant function by increasing the expression of mitochondrial-localized gene products, such as MnSOD and Bcl-2, which display antioxidant activities [64–66]. Verline et al. found that MnSOD knockdown in mice leads to oxidative damage of RPE cells, and more importantly, the appearance of certain key features of AMD [67]. The mitochondrial enzyme TrxR<sub>2</sub> preserves thioredoxin in a reduced state, thereby playing a key role in maintaining the cellular redox environment [68]. Inhibition of TrxR<sub>2</sub> in mitochondria isolated from guinea pigs, as well as in mouse cardiomyocytes, resulted in increased H<sub>2</sub>O<sub>2</sub> production [24]. This correlates with our findings, as cell viability was reduced with miR-17-3p mimic transfection under oxidative stress, in parallel with MnSOD and TrxR<sub>2</sub> downregulation.

Due to increased miR-17-3p levels in AMD eyes in combination with the fact that miR-17-3p can specifically downregulate MnSOD and TrxR<sub>2</sub> expression, its role as a mediator for RPE susceptibility to oxidative stress in AMD should not be overlooked. Interestingly, another recent study by Haque et al. [43] showed that in ARPE-19 cells, a sublethal dose of H<sub>2</sub>O<sub>2</sub> upregulated miR-30b, which inhibited catalase, another anti-oxidant enzyme. Furthermore, a miR-30b antagonist protected RPE cells from oxidative stress, concurring with our findings after using miR-17-3p.

miR-17-3p is a member of miR-17/92 cluster which is extensively studied in tumorigenesis [69]. Most of the work on miR17-3p has thus been focused on studying the regulation of cell proliferation pathways. More recently though, studies have shown more varied effects of miR-17-3p, such as involvement in development and spinal progenitor cells [70], induction by TNF alpha and LPS in HeLa cells [71], downregulating flk-1 in endothelial cells [72], suppressing epithelial to mesenchymal transformation of ovarian epithelial cells [73], inhibiting fibroblast senescence [74] and being downregulated in the plasma of Alzheimer patients [75]. More relevant to our work, it has been shown that miR-17-3p downregulates important anti-oxidant enzymes, such as manganese superoxide dismutase, glutathione peroxidase-2 and thioredoxin reductase-2 (TrxR<sub>2</sub>) in prostate cancer cell lines [48] and blood mononuclear cells [76]. To confirm the previous studies, our bioinformatic sequence analysis found that there is a putative binding site for the miR-17-3p in the mRNA 3' untranslated regions of MnSOD and TrxR<sub>2</sub> (Fig 5A) that is verified by in vitro experiments (Fig 5B and 5C). The mechanism by which oxidative stress regulates miR-17-3p levels remains elusive. It is interesting to note that downregulation of c-Myc suppresses pri-miR-17/92 cluster expression [77] and that c-Myc is downregulated by peroxiredoxin 1, which is involved under oxidative stress [78]. Although this mechanism has not been studied in RPE cells or in AMD, it is a promising target for future investigation.

In conclusion, this study demonstrates increased miR-17-3p expression in human AMD eyes. It also shows that miR-17-3p expression is increased by oxidative stimuli and that it has detrimental effects on RPE antioxidant defense mechanisms by downregulating the anti-oxidant enzymes MnSOD and TrxR<sub>2</sub>. Therefore, inhibition of miR-17-3p could be a novel therapeutic approach to protect RPE cells from oxidative stress.

## Acknowledgments

This work was supported by: NEI R21EY023079-01A1 (DGV); the Yeatts Family Foundation (DGV, JWM); the Loefflers Family Fund (DGV, JWM); the 2013 Macula Society Research Grant award (DGV); a Physician Scientist Award (DGV), an unrestricted grant (JWM) from the research to prevent blindness foundation; NEI grant EY014104 (Meei Core Grant); and Bayer Healthcare Global Ophthalmology Award (DEM). The authors would like to thank Kasandra Kosmidou for proofreading this manuscript.

## Author Contributions

**Conceived and designed the experiments:** HL DGV JWM.

**Performed the experiments:** BT.

**Analyzed the data:** HL BT.

**Contributed reagents/materials/analysis tools:** DEM BD PB JBM DGV JWM.

**Wrote the paper:** HL DGV BT DEM PB.

## References

1. Solberg Y, Rosner M, Belkin M. The association between cigarette smoking and ocular diseases. *Surv Ophthalmol* n.d.; 42:535–47.
2. Hollyfield JG, Bonilha VL, Rayborn ME, Yang X, Shadrach KG, Lu L, et al. Oxidative damage-induced inflammation initiates age-related macular degeneration. *Nat Med* 2008; 14:194–8. doi: [10.1038/nm1709](https://doi.org/10.1038/nm1709) PMID: [18223656](https://pubmed.ncbi.nlm.nih.gov/18223656/)
3. Hollyfield JG. Age-related macular degeneration: the molecular link between oxidative damage, tissue-specific inflammation and outer retinal disease: the Proctor lecture. *Invest Ophthalmol Vis Sci* 2010; 51:1275–81. doi: [10.1167/iovs.09-4478](https://doi.org/10.1167/iovs.09-4478)
4. Beatty S, Koh H-H, Phil M, Henson D, Boulton M. The Role of Oxidative Stress in the Pathogenesis of Age-Related Macular Degeneration. *Surv Ophthalmol* 2000; 45:115–34. doi: [10.1016/S0039-6257\(00\)00140-5](https://doi.org/10.1016/S0039-6257(00)00140-5) PMID: [11033038](https://pubmed.ncbi.nlm.nih.gov/11033038/)
5. Whitmore SS, Sohn EH, Chirco KR, Drack A V, Stone EM, Tucker BA, et al. Complement activation and choriocapillaris loss in early AMD: implications for pathophysiology and therapy. *Prog Retin Eye Res* 2015; 45:1–29. doi: [10.1016/j.preteyeres.2014.11.005](https://doi.org/10.1016/j.preteyeres.2014.11.005) PMID: [25486088](https://pubmed.ncbi.nlm.nih.gov/25486088/)
6. Klein RJ, Zeiss C, Chew EY, Tsai J-Y, Sackler RS, Haynes C, et al. Complement factor H polymorphism in age-related macular degeneration. *Science* 2005; 308:385–9. doi: [10.1126/science.1109557](https://doi.org/10.1126/science.1109557) PMID: [15761122](https://pubmed.ncbi.nlm.nih.gov/15761122/)
7. Hageman GS, Anderson DH, Johnson L V, Hancox LS, Taiber AJ, Hardisty LI, et al. A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc Natl Acad Sci U S A* 2005; 102:7227–32. doi: [10.1073/pnas.0501536102](https://doi.org/10.1073/pnas.0501536102) PMID: [15870199](https://pubmed.ncbi.nlm.nih.gov/15870199/)
8. Johnson L V, Leitner WP, Rivest AJ, Staples MK, Radeke MJ, Anderson DH. The Alzheimer's A beta-peptide is deposited at sites of complement activation in pathologic deposits associated with aging and age-related macular degeneration. *Proc Natl Acad Sci U S A* 2002; 99:11830–5. doi: [10.1073/pnas.192203399](https://doi.org/10.1073/pnas.192203399) PMID: [12189211](https://pubmed.ncbi.nlm.nih.gov/12189211/)
9. Wang JCC, Cao S, Wang A, To E, Law G, Gao J, et al. CFH Y402H polymorphism is associated with elevated vitreal GM-CSF and choroidal macrophages in the postmortem human eye. *Mol Vis* 2015; 21:264–72. PMID: [25814824](https://pubmed.ncbi.nlm.nih.gov/25814824/)

10. Lad EM, Cousins SW, Van Arnam JS, Proia AD. Abundance of infiltrating CD163+ cells in the retina of postmortem eyes with dry and neovascular age-related macular degeneration. *Graefes Arch Clin Exp Ophthalmol* 2015; 253:1941–5. doi: [10.1007/s00417-015-3094-z](https://doi.org/10.1007/s00417-015-3094-z) PMID: [26148801](https://pubmed.ncbi.nlm.nih.gov/26148801/)
11. Ambati J, Atkinson JP, Gelfand BD. Immunology of age-related macular degeneration. *Nat Rev Immunol* 2013; 13:438–51. doi: [10.1038/nri3459](https://doi.org/10.1038/nri3459) PMID: [23702979](https://pubmed.ncbi.nlm.nih.gov/23702979/)
12. Penfold PL, Madigan MC, Gillies MC, Provis JM. Immunological and aetiological aspects of macular degeneration. *Prog Retin Eye Res* 2001; 20:385–414. PMID: [11286898](https://pubmed.ncbi.nlm.nih.gov/11286898/)
13. Dib B, Lin H, Maidana DE, Tian B, Miller JB, Bouzika P, et al. Mitochondrial DNA has a pro-inflammatory role in AMD. *Biochim Biophys Acta—Mol Cell Res* 2015; 1853:2897–906. doi: [10.1016/j.bbamcr.2015.08.012](https://doi.org/10.1016/j.bbamcr.2015.08.012)
14. Kevany BM, Palczewski K. Phagocytosis of retinal rod and cone photoreceptors. *Physiology (Bethesda)* 2010; 25:8–15. doi: [10.1152/physiol.00038.2009](https://doi.org/10.1152/physiol.00038.2009)
15. Sun M, Finnemann SC, Febbraio M, Shan L, Annangudi SP, Podrez EA, et al. Light-induced oxidation of photoreceptor outer segment phospholipids generates ligands for CD36-mediated phagocytosis by retinal pigment epithelium: a potential mechanism for modulating outer segment phagocytosis under oxidant stress conditions. *J Biol Chem* 2006; 281:4222–30. doi: [10.1074/jbc.M509769200](https://doi.org/10.1074/jbc.M509769200) PMID: [16354659](https://pubmed.ncbi.nlm.nih.gov/16354659/)
16. Brunk UT, Wihlmark U, Wrigstad A, Roberg K, Nilsson SE. Accumulation of lipofuscin within retinal pigment epithelial cells results in enhanced sensitivity to photo-oxidation. *Gerontology* 1995; 41 Suppl 2:201–12. PMID: [8821332](https://pubmed.ncbi.nlm.nih.gov/8821332/)
17. Becquet F, Goureau O, Soubrane G, Coscas G, Courtois Y, Hicks D. Superoxide inhibits proliferation and phagocytic internalization of photoreceptor outer segments by bovine retinal pigment epithelium in vitro. *Exp Cell Res* 1994; 212:374–82. doi: [10.1006/excr.1994.1157](https://doi.org/10.1006/excr.1994.1157) PMID: [8187832](https://pubmed.ncbi.nlm.nih.gov/8187832/)
18. Dorey CK, Delori FC, Akeo K. Growth of cultured RPE and endothelial cells is inhibited by blue light but not green or red light. *Curr Eye Res* 1990; 9:549–59. PMID: [2117518](https://pubmed.ncbi.nlm.nih.gov/2117518/)
19. He Y, Ge J, Burke JM, Myers RL, Dong ZZ, Tombran-Tink J. Mitochondria impairment correlates with increased sensitivity of aging RPE cells to oxidative stress. *J Ocul Biol Dis Infor* 2010; 3:92–108. doi: [10.1007/s12177-011-9061-y](https://doi.org/10.1007/s12177-011-9061-y) PMID: [22833778](https://pubmed.ncbi.nlm.nih.gov/22833778/)
20. Dong A, Xie B, Shen J, Yoshida T, Yokoi K, Hackett SF, et al. Oxidative stress promotes ocular neovascularization. *J Cell Physiol* 2009; 219:544–52. doi: [10.1002/jcp.21698](https://doi.org/10.1002/jcp.21698) PMID: [19142872](https://pubmed.ncbi.nlm.nih.gov/19142872/)
21. Kasahara E, Lin L-R, Ho Y-S, Reddy VN. SOD2 protects against oxidation-induced apoptosis in mouse retinal pigment epithelium: implications for age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2005; 46:3426–34. doi: [10.1167/iovs.05-0344](https://doi.org/10.1167/iovs.05-0344) PMID: [16123448](https://pubmed.ncbi.nlm.nih.gov/16123448/)
22. Flores LC, Ortiz M, Dube S, Hubbard GB, Lee S, Salmon A, et al. Thioredoxin, oxidative stress, cancer and aging. *Longev Heal* 2012; 1:4. doi: [10.1186/2046-2395-1-4](https://doi.org/10.1186/2046-2395-1-4)
23. Nishinaka Y, Masutani H, Nakamura H, Yodoi J. Regulatory roles of thioredoxin in oxidative stress-induced cellular responses. *Redox Rep* 2001; 6:289–95. doi: [10.1179/135100001101536427](https://doi.org/10.1179/135100001101536427) PMID: [11778846](https://pubmed.ncbi.nlm.nih.gov/11778846/)
24. Stanley BA, Sivakumaran V, Shi S, McDonald I, Lloyd D, Watson WH, et al. Thioredoxin reductase-2 is essential for keeping low levels of H(2)O(2) emission from isolated heart mitochondria. *J Biol Chem* 2011; 286:33669–77. doi: [10.1074/jbc.M111.284612](https://doi.org/10.1074/jbc.M111.284612) PMID: [21832082](https://pubmed.ncbi.nlm.nih.gov/21832082/)
25. Sugano E, Isago H, Murayama N, Tamai M, Tomita H. Different Anti-Oxidant Effects of Thioredoxin 1 and Thioredoxin 2 in Retinal Epithelial Cells. *CELL Struct Funct* 2013; 38:81–8. PMID: [23485938](https://pubmed.ncbi.nlm.nih.gov/23485938/)
26. Akeo K, Curran SA, Dorey CK. Superoxide dismutase activity and growth of retinal pigment epithelial cells are suppressed by 20% oxygen in vitro. *Curr Eye Res* 1988; 7:961–7. PMID: [3229124](https://pubmed.ncbi.nlm.nih.gov/3229124/)
27. Dorey CK, Khouri GG, Syniuta LA, Curran SA, Weiter JJ. Superoxide production by porcine retinal pigment epithelium in vitro. *Invest Ophthalmol Vis Sci* 1989; 30:1047–54. PMID: [2543650](https://pubmed.ncbi.nlm.nih.gov/2543650/)
28. Newsome DA, Dobard EP, Liles MR, Oliver PD. Human retinal pigment epithelium contains two distinct species of superoxide dismutase. *Invest Ophthalmol Vis Sci* 1990; 31:2508–13. PMID: [2265990](https://pubmed.ncbi.nlm.nih.gov/2265990/)
29. Del Vecchio PJ, Shaffer JB. Regulation of antioxidant enzyme expression in LPS-treated bovine retinal pigment epithelial and corneal endothelial cells. *Curr Eye Res* 1991; 10:919–25. PMID: [1720368](https://pubmed.ncbi.nlm.nih.gov/1720368/)
30. Oliver PD, Newsome DA. Mitochondrial superoxide dismutase in mature and developing human retinal pigment epithelium. *Invest Ophthalmol Vis Sci* 1992; 33:1909–18. PMID: [1582797](https://pubmed.ncbi.nlm.nih.gov/1582797/)
31. Wang M, Lam TT, Fu J, Tso MO. TEMPOL, a superoxide dismutase mimic, ameliorates light-induced retinal degeneration. *Res Commun Mol Pathol Pharmacol* 1995; 89:291–305. PMID: [8680798](https://pubmed.ncbi.nlm.nih.gov/8680798/)
32. Tanito M, Masutani H, Kim Y-C, Nishikawa M, Ohira A, Yodoi J. Sulforaphane induces thioredoxin through the antioxidant-responsive element and attenuates retinal light damage in mice. *Invest Ophthalmol Vis Sci* 2005; 46:979–87. doi: [10.1167/iovs.04-1120](https://doi.org/10.1167/iovs.04-1120) PMID: [15728556](https://pubmed.ncbi.nlm.nih.gov/15728556/)

33. Yokoyama T, Yamane K, Minamoto A, Tsukamoto H, Yamashita H, Izumi S, et al. High glucose concentration induces elevated expression of anti-oxidant and proteolytic enzymes in cultured human retinal pigment epithelial cells. *Exp Eye Res* 2006; 83:602–9. doi: [10.1016/j.exer.2006.02.016](https://doi.org/10.1016/j.exer.2006.02.016) PMID: [16697369](https://pubmed.ncbi.nlm.nih.gov/16697369/)
34. Sugano E, Isago H, Murayama N, Tamai M, Tomita H. Different anti-oxidant effects of thioredoxin 1 and thioredoxin 2 in retinal epithelial cells. *Cell Struct Funct* 2013; 38:81–8. PMID: [23485938](https://pubmed.ncbi.nlm.nih.gov/23485938/)
35. Sugano E, Murayama N, Takahashi M, Tabata K, Tamai M, Tomita H. Essential role of thioredoxin 2 in mitigating oxidative stress in retinal epithelial cells. *J Ophthalmol* 2013; 2013:185825. doi: [10.1155/2013/185825](https://doi.org/10.1155/2013/185825) PMID: [24319591](https://pubmed.ncbi.nlm.nih.gov/24319591/)
36. Kasahara E, Lin L-R, Ho Y-S, Reddy VN. SOD2 protects against oxidation-induced apoptosis in mouse retinal pigment epithelium: implications for age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2005; 46:3426–34. doi: [10.1167/iops.05-0344](https://doi.org/10.1167/iops.05-0344) PMID: [16123448](https://pubmed.ncbi.nlm.nih.gov/16123448/)
37. Wu L, Fan J, Belasco JG. MicroRNAs direct rapid deadenylation of mRNA. *Proc Natl Acad Sci U S A* 2006; 103:4034–9. doi: [10.1073/pnas.0510928103](https://doi.org/10.1073/pnas.0510928103) PMID: [16495412](https://pubmed.ncbi.nlm.nih.gov/16495412/)
38. Guo L, Yu J, Yu H, Zhao Y, Chen S, Xu C, et al. Evolutionary and Expression Analysis of miR-#-5p and miR-#-3p at the miRNAs/isomiRs Levels. *Biomed Res Int* 2015; 2015. doi: [10.1155/2015/168358](https://doi.org/10.1155/2015/168358)
39. Raisch J, Darfeuille-Michaud A, Nguyen HTT. Role of microRNAs in the immune system, inflammation and cancer. *World J Gastroenterol* 2013; 19:2985–96. doi: [10.3748/wjg.v19.i20.2985](https://doi.org/10.3748/wjg.v19.i20.2985) PMID: [23716978](https://pubmed.ncbi.nlm.nih.gov/23716978/)
40. Yildirim SS, Akman D, Catalucci D, Turan B. Relationship between downregulation of miRNAs and increase of oxidative stress in the development of diabetic cardiac dysfunction: junctin as a target protein of miR-1. *Cell Biochem Biophys* 2013; 67:1397–408. doi: [10.1007/s12013-013-9672-y](https://doi.org/10.1007/s12013-013-9672-y) PMID: [23723006](https://pubmed.ncbi.nlm.nih.gov/23723006/)
41. Suárez Y, Sessa WC. MicroRNAs as novel regulators of angiogenesis. *Circ Res* 2009; 104:442–54. doi: [10.1161/CIRCRESAHA.108.191270](https://doi.org/10.1161/CIRCRESAHA.108.191270) PMID: [19246688](https://pubmed.ncbi.nlm.nih.gov/19246688/)
42. Li Q-J, Chau J, Ebert PJR, Sylvester G, Min H, Liu G, et al. miR-181a is an intrinsic modulator of T cell sensitivity and selection. *Cell* 2007; 129:147–61. doi: [10.1016/j.cell.2007.03.008](https://doi.org/10.1016/j.cell.2007.03.008) PMID: [17382377](https://pubmed.ncbi.nlm.nih.gov/17382377/)
43. Haque R, Chun E, Howell JC, Sengupta T, Chen D, Kim H. MicroRNA-30b-mediated regulation of catalase expression in human ARPE-19 cells. *PLoS One* 2012; 7:e42542. doi: [10.1371/journal.pone.0042542](https://doi.org/10.1371/journal.pone.0042542) PMID: [22880027](https://pubmed.ncbi.nlm.nih.gov/22880027/)
44. Kutty RK, Samuel W, Jaworski C, Duncan T, Nagineni CN, Raghavachari N, et al. MicroRNA expression in human retinal pigment epithelial (ARPE-19) cells: increased expression of microRNA-9 by N-(4-hydroxyphenyl)retinamide. *Mol Vis* 2010; 16:1475–86. PMID: [20806079](https://pubmed.ncbi.nlm.nih.gov/20806079/)
45. Garcia TY, Gutierrez M, Reynolds J, Lamba DA. Modeling the Dynamic AMD-Associated Chronic Oxidative Stress Changes in Human ESC and iPSC-Derived RPE Cells. *Invest Ophthalmol Vis Sci* 2015; 56:7480–8. doi: [10.1167/iops.15-17251](https://doi.org/10.1167/iops.15-17251) PMID: [26595608](https://pubmed.ncbi.nlm.nih.gov/26595608/)
46. Lin H, Qian J, Castillo AC, Long B, Keyes KT, Chen G, et al. Effect of miR-23 on Oxidant-Induced Injury in Human Retinal Pigment Epithelial Cells. *Invest Ophthalmol Vis Sci* 2011; 52:6308–14. doi: [10.1167/iops.10-6632](https://doi.org/10.1167/iops.10-6632) PMID: [21693609](https://pubmed.ncbi.nlm.nih.gov/21693609/)
47. Mogilyansky E, Rigoutsos I. The miR-17/92 cluster: a comprehensive update on its genomics, genetics, functions and increasingly important and numerous roles in health and disease. *Cell Death Differ* 2013; 20:1603–14. doi: [10.1038/cdd.2013.125](https://doi.org/10.1038/cdd.2013.125) PMID: [24212931](https://pubmed.ncbi.nlm.nih.gov/24212931/)
48. Xu Y, Fang F, Zhang J, Josson S, Clair WHS, Clair DKS. Mir-17\* suppresses tumorigenicity of prostate cancer by inhibiting mitochondrial antioxidant enzymes. *PLoS One* 2010; 5:1–8. doi: [10.1371/journal.pone.0014356](https://doi.org/10.1371/journal.pone.0014356)
49. Lin H, Xu H, Liang F-Q, Liang H, Gupta P, Havey AN, et al. Mitochondrial DNA damage and repair in RPE associated with aging and age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2011; 52:3521–9. doi: [10.1167/iops.10-6163](https://doi.org/10.1167/iops.10-6163) PMID: [21273542](https://pubmed.ncbi.nlm.nih.gov/21273542/)
50. Marazita MC, Dugour A, Marquioni-Ramella MD, Figueroa JM, Suburo AM. Oxidative stress-induced premature senescence dysregulates VEGF and CFH expression in retinal pigment epithelial cells: Implications for Age-related Macular Degeneration. *Redox Biol* 2016; 7:78–87. doi: [10.1016/j.redox.2015.11.011](https://doi.org/10.1016/j.redox.2015.11.011) PMID: [26654980](https://pubmed.ncbi.nlm.nih.gov/26654980/)
51. Lukiw WJ, Surjyadipta B, Dua P, Alexandrov PN. Common micro RNAs (miRNAs) target complement factor H (CFH) regulation in Alzheimer's disease (AD) and in age-related macular degeneration (AMD). *Int J Biochem Mol Biol* 2012; 3:105–16. PMID: [22509485](https://pubmed.ncbi.nlm.nih.gov/22509485/)
52. Murad N, Kokkinaki M, Gunawardena N, Gunawan MS, Hathout Y, Janczura KJ, et al. miR-184 regulates ezrin, LAMP-1 expression, affects phagocytosis in human retinal pigment epithelium and is down-regulated in age-related macular degeneration. *FEBS J* 2014; 281:5251–64. doi: [10.1111/febs.13066](https://doi.org/10.1111/febs.13066) PMID: [25251993](https://pubmed.ncbi.nlm.nih.gov/25251993/)

53. Szemraj J. Serum MicroRNAs as Potential Biomarkers of AMD. *Med Sci Monit* 2015; 21:2734–42. doi: [10.12659/MSM.893697](https://doi.org/10.12659/MSM.893697) PMID: [26366973](https://pubmed.ncbi.nlm.nih.gov/26366973/)
54. Blasiak J, Petrovski G, Veréb Z, Facsó A, Kaarniranta K. Oxidative stress, hypoxia, and autophagy in the neovascular processes of age-related macular degeneration. *Biomed Res Int* 2014; 2014:768026. doi: [10.1155/2014/768026](https://doi.org/10.1155/2014/768026) PMID: [24707498](https://pubmed.ncbi.nlm.nih.gov/24707498/)
55. Kim M-H, Chung J, Yang J, Chung S-M, Kwag N-H, Yoo J-S. Hydrogen peroxide-induced cell death in a human retinal pigment epithelial cell line, ARPE-19. *Korean J Ophthalmol* 2003; 17:19–28. doi: [10.3341/kjo.2003.17.1.19](https://doi.org/10.3341/kjo.2003.17.1.19) PMID: [12882504](https://pubmed.ncbi.nlm.nih.gov/12882504/)
56. Lu L, Hackett SF, Mincey A, Lai H, Campochiaro PA. Effects of different types of oxidative stress in RPE cells. *J Cell Physiol* 2006; 206:119–25. doi: [10.1002/jcp.20439](https://doi.org/10.1002/jcp.20439) PMID: [15965958](https://pubmed.ncbi.nlm.nih.gov/15965958/)
57. Kučera O, Endlicher R, Roušar T, Lotková H, Gamol T, Drahotka Z, et al. The Effect of tert-Butyl Hydroperoxide-Induced Oxidative Stress on Lean and Steatotic Rat Hepatocytes In Vitro. *Oxid Med Cell Longev* 2014; 2014:1–12. doi: [10.1155/2014/752506](https://doi.org/10.1155/2014/752506)
58. Weigel AL, Ida H, Boylan SA, Hjelmeland LM. Acute hyperoxia-induced transcriptional response in the mouse RPE/choroid. *Free Radic Biol Med* 2003; 35:465–74. doi: [10.1016/S0891-5849\(03\)00320-4](https://doi.org/10.1016/S0891-5849(03)00320-4) PMID: [12927596](https://pubmed.ncbi.nlm.nih.gov/12927596/)
59. Alizadeh M, Wada M, Gelfman CM, Handa JT, Hjelmeland LM. Downregulation of Differentiation Specific Gene Expression by Oxidative Stress in ARPE-19 Cells. *Invest Ophthalmol Vis Sci* 2001; 42:2706–13. PMID: [11581219](https://pubmed.ncbi.nlm.nih.gov/11581219/)
60. Weigel AL, Handa JT, Hjelmeland LM. Microarray analysis of H2O2-, HNE-, or tBH-treated ARPE-19 cells. *Free Radic Biol Med* 2002; 33:1419–32. doi: [10.1016/S0891-5849\(02\)01082-1](https://doi.org/10.1016/S0891-5849(02)01082-1) PMID: [12419474](https://pubmed.ncbi.nlm.nih.gov/12419474/)
61. Vandenbroucke K, Robbens S, Vandepoele K, Inzé D, Van de Peer Y, Van Breusegem F. Hydrogen peroxide-induced gene expression across kingdoms: a comparative analysis. *Mol Biol Evol* 2008; 25:507–16. doi: [10.1093/molbev/msm276](https://doi.org/10.1093/molbev/msm276) PMID: [18187560](https://pubmed.ncbi.nlm.nih.gov/18187560/)
62. Halbeisen RE, Galgano A, Scherrer T, Gerber AP. Post-transcriptional gene regulation: from genome-wide studies to principles. *Cell Mol Life Sci* 2008; 65:798–813. doi: [10.1007/s00018-007-7447-6](https://doi.org/10.1007/s00018-007-7447-6) PMID: [18043867](https://pubmed.ncbi.nlm.nih.gov/18043867/)
63. Fukui M, Zhu BT. Mitochondrial superoxide dismutase SOD2, but not cytosolic SOD1, plays a critical role in protection against glutamate-induced oxidative stress and cell death in HT22 neuronal cells. *Free Radic Biol Med* 2010; 48:821–30. doi: [10.1016/j.freeradbiomed.2009.12.024](https://doi.org/10.1016/j.freeradbiomed.2009.12.024) PMID: [20060889](https://pubmed.ncbi.nlm.nih.gov/20060889/)
64. Matsuoka A, Kizuka F, Lee L, Tamura I, Taniguchi K, Asada H, et al. Progesterone increases manganese superoxide dismutase expression via a cAMP-dependent signaling mediated by noncanonical Wnt5a pathway in human endometrial stromal cells. *J Clin Endocrinol Metab* 2010; 95:E291–9. doi: [10.1210/jc.2010-0619](https://doi.org/10.1210/jc.2010-0619) PMID: [20685861](https://pubmed.ncbi.nlm.nih.gov/20685861/)
65. Tripanichkul W, Sripanichkulchai K, Duce JA, Finkelstein DI. 17Beta-estradiol reduces nitrotyrosine immunoreactivity and increases SOD1 and SOD2 immunoreactivity in nigral neurons in male mice following MPTP insult. *Brain Res* 2007; 1164:24–31. doi: [10.1016/j.brainres.2007.05.076](https://doi.org/10.1016/j.brainres.2007.05.076) PMID: [17640623](https://pubmed.ncbi.nlm.nih.gov/17640623/)
66. Kowaltowski AJ, Fenton RG, Fiskum G. Bcl-2 family proteins regulate mitochondrial reactive oxygen production and protect against oxidative stress. *Free Radic Biol Med* 2004; 37:1845–53. doi: [10.1016/j.freeradbiomed.2004.09.005](https://doi.org/10.1016/j.freeradbiomed.2004.09.005) PMID: [15528043](https://pubmed.ncbi.nlm.nih.gov/15528043/)
67. Justilien V, Pang JJ, Renganathan K, Zhan X, Crabb JW, So RK, et al. SOD2 knockdown mouse model of early AMD. *Investig Ophthalmol Vis Sci* 2007; 48:4407–20. doi: [10.1167/iovs.07-0432](https://doi.org/10.1167/iovs.07-0432)
68. Myers CR, Myers JM. The effects of acrolein on peroxiredoxins, thioredoxins, and thioredoxin reductase in human bronchial epithelial cells. *Toxicology* 2009; 257:95–104. doi: [10.1016/j.tox.2008.12.013](https://doi.org/10.1016/j.tox.2008.12.013) PMID: [19135121](https://pubmed.ncbi.nlm.nih.gov/19135121/)
69. Mogilyansky E, Rigoutsos I. The miR-17/92 cluster: a comprehensive update on its genomics, genetics, functions and increasingly important and numerous roles in health and disease. *Cell Death Differ* 2013; 20:1603–14. doi: [10.1038/cdd.2013.125](https://doi.org/10.1038/cdd.2013.125) PMID: [24212931](https://pubmed.ncbi.nlm.nih.gov/24212931/)
70. Chen J-A, Huang Y-P, Mazzoni EO, Tan GC, Zavadil J, Wichterle H. Mir-17-3p Controls Spinal Neural Progenitor Patterning by Regulating Olig2/Irx3 Cross-Repressive Loop. *Neuron* 2011; 69:721–35. doi: [10.1016/j.neuron.2011.01.014](https://doi.org/10.1016/j.neuron.2011.01.014) PMID: [21338882](https://pubmed.ncbi.nlm.nih.gov/21338882/)
71. Jiang X, Li N. Induction of MiR-17-3p and MiR-160a by TNFα and LPS. *Cell Biochem Funct* 2011; 29:164–70. doi: [10.1002/cbf.1728](https://doi.org/10.1002/cbf.1728) PMID: [21370248](https://pubmed.ncbi.nlm.nih.gov/21370248/)
72. Yin R, Wang R, Guo L, Zhang W, Lu Y. MiR-17-3p inhibits angiogenesis by downregulating flk-1 in the cell growth signal pathway. *J Vasc Res* 2013; 50:157–66. doi: [10.1159/000345697](https://doi.org/10.1159/000345697) PMID: [23258273](https://pubmed.ncbi.nlm.nih.gov/23258273/)
73. Liu T, Qin W, Huang Y, Zhao Y, Wang J. Induction of Estrogen-Sensitive Epithelial Cells Derived from Human-Induced Pluripotent Stem Cells to Repair Ovarian Function in a Chemotherapy-Induced Mouse

- Model of Premature Ovarian Failure. *DNA Cell Biol* 2013; 32:130914053323001. doi: [10.1089/dna.2013.2032](https://doi.org/10.1089/dna.2013.2032)
74. Du WW, Li X, Li T, Li H, Khorshidi A, Liu F, et al. The microRNA miR-17-3p inhibits mouse cardiac fibroblast senescence by targeting Par4. *J Cell Sci* 2015; 128:293–304. doi: [10.1242/jcs.158360](https://doi.org/10.1242/jcs.158360) PMID: [25472717](https://pubmed.ncbi.nlm.nih.gov/25472717/)
  75. Satoh JI, Kino Y, Niida S. MicroRNA-seq data analysis pipeline to identify blood biomarkers for Alzheimer's disease from public data. *Biomark Insights* 2015; 10:21–31. doi: [10.4137/BMIMI.S25132](https://doi.org/10.4137/BMIMI.S25132) PMID: [25922570](https://pubmed.ncbi.nlm.nih.gov/25922570/)
  76. Curti V, Capelli E, Boschi F, Nabavi SF, Bongiorno AI, Habtemariam S, et al. Modulation of human miR-17-3p expression by methyl 3-O-methyl gallate as explanation of its in vivo protective activities. *Mol Nutr Food Res* 2014:1–9. doi: [10.1002/mnfr.201400007](https://doi.org/10.1002/mnfr.201400007)
  77. Thomas M, Lange-Grünweller K, Hartmann D, Golde L, Schlereth J, Streng D, et al. Analysis of transcriptional regulation of the human miR-17-92 cluster; evidence for involvement of Pim-1. *Int J Mol Sci* 2013; 14:12273–96. doi: [10.3390/ijms140612273](https://doi.org/10.3390/ijms140612273) PMID: [23749113](https://pubmed.ncbi.nlm.nih.gov/23749113/)
  78. Graves JA, Metukuri M, Scott D, Rothermund K, Prochownik E V. Regulation of reactive oxygen species homeostasis by peroxiredoxins and c-Myc. *J Biol Chem* 2009; 284:6520–9. doi: [10.1074/jbc.M807564200](https://doi.org/10.1074/jbc.M807564200) PMID: [19098005](https://pubmed.ncbi.nlm.nih.gov/19098005/)