



PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS

Citation

Southey, M. C., D. E. Goldgar, R. Winqvist, K. Pyrkäs, F. Couch, M. Tischkowitz, W. D. Foulkes, et al. 2016. "PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS." *Journal of medical genetics* 53 (12): 800-811. doi:10.1136/jmedgenet-2016-103839. <http://dx.doi.org/10.1136/jmedgenet-2016-103839>.

Published Version

doi:10.1136/jmedgenet-2016-103839

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:29739204>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



Published in final edited form as:

J Med Genet. 2016 December ; 53(12): 800–811. doi:10.1136/jmedgenet-2016-103839.

PALB2, CHEK2 and ATM rare variants and cancer risk: data from COGS

A full list of authors and affiliations appears at the end of the article.

Abstract

Background—The rarity of mutations in *PALB2*, *CHEK2* and *ATM* make it difficult to estimate precisely associated cancer risks. Population-based family studies have provided evidence that at least some of these mutations are associated with breast cancer risk as high as those associated with rare *BRCA2* mutations. We aimed to estimate the relative risks associated with specific rare variants in *PALB2*, *CHEK2* and *ATM* via a multicentre case-control study.

Methods—We genotyped 10 rare mutations using the custom iCOGS array: *PALB2* c.1592delT, c.2816T>G and c.3113G>A, *CHEK2* c.349A>G, c.538C>T, c.715G>A, c.1036C>T, c.1312G>T, and c.1343T>G and *ATM* c.7271T>G. We assessed associations with breast cancer risk (42 671 cases and 42 164 controls), as well as prostate (22 301 cases and 22 320 controls) and ovarian (14 542 cases and 23 491 controls) cancer risk, for each variant.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

Correspondence to Professor Melissa C. Southey, Genetic Epidemiology Laboratory, Department of Pathology, The University of Melbourne, Melbourne, Victoria 3010, Australia; msouthey@unimelb.edu.au.

Contributors All authors provided DNA samples and/or data and have participated in the Breast Cancer Association Consortium through attendance at regular meetings and planning of the iCOGS experiment. Each author has made substantial contribution through designing and coordinating the studies listed in the supplemental material and therefore have made substantial contributions to the conception or design of this work. Many authors played multiple roles across these activities. Specifically, MCS conceived this study, worked to include the rare variants on the iCOGS and drafted the manuscript. RLM led the statistical analysis and drafted the paper. Members of the PALB2 interest group, MCS, DEG, RW, KP, FC, MT, WF, JD, KM, EJvR, TH, HN, JLH, TD, KC, JR-F, ZLT, PR, IC, PP, HT, FAO, JGD contributed to the inclusion of the PALB2 rare variants on iCOGS. DFE coordinated the BCAC project and contributed to statistical analysis along with DEG. SVT contributed to the selection of CHEK2 rare variants. GC-T contributed to the selection of rare variants in ATM. AD, CL and JD made significant contribution to the data quality related to the calling of the rare genetics variants on iCOGS. MKS, AB, FBH, SV, JC, CC, RJS, PAF, LH, ABE, MWB, JP, IDSS, OF, NJ, MKB, EJS, IT, MJK, NM, FM, BB, RY, PG, TT, FM, MS, SB, SFN, HF, JB, MPZ, JIAP, PM, HAC, SN, AZ, CCD, HB, VA, CS, HB, TB, YDK, TAM, KA, CB, NVB, NNA, AL, SM, AM, VK, V-MK, JMH, AS, EW, DS, BB, GF, JCC, AR, PS, DFJ, JEO, CV, VSP, CM, CAH, BEH, FS, LLM, VK, GGA, WZ, DJH, SL, SEH, PK, IA, JAK, GG, AMM, AJV, MG, SK, PD, RAEMT, CS, AH, MGC, JF, SJC, JL, KC, HD, ME, DME, SR, WJT, SMG, MJH, JWMM, JMC, MTL, PH, JL, JSB, KH, AC, MWRR, CL, CB, AD, UH, DT, HUU, TR, AJ, JL, KJ, KD, SS, AET, CBA, DY, AS, AA, NO, MJ, AGN, GP, MRA, NA, DH, DCT, DV, FB, JS, MD, PS, RE, KM, FW, HG, JS, MW, BGN, RCT, DN, JLD, FCH, KTK, JLS, WJB, ST, DJS, JLK, CM, ASK, CC, LCA, KB, JP, RK, JB, MRT, ZKJ, AAAO, SB, SPR, AH, AH, MR, DL, EVN, IV, SL, JAD, MAR, SN, UE, SWG, KO, LES, GF, GL, JLK, LRW, MTG, IR, PAH, LMP, RB, FM, RPE, RBN, KBM, ADB, FH, PH, SK, BYK, CW, JL, AJ, SKK, EH, BP, BB, LB, ELG, BLF, RAV, JMC, MCL, ZCF, KRK, DL, KHL, MATH, XW, DAL, FD, MB, AB, ESI, JRM, LA, DWC, KLT, ELP, MS, SST, EVB, IO, SHO, LB, HBS, AMVA, KKHA, LAK, LFAGM, TP, YB, ABW, LEK, LSC, NDL, BG, JG, JM, CKH, LL, LN, SAE, ED, JT, IC, IN, JP, NS, RG, ASW, JHR, VG, WS, HC, XOS, RTT, RS, JRM, SAN, CP, ANM, DF, HYL, JPW, TAS, TAC, YYT, ZC, AGM, SAG, SJR, UM, AHW, CLP, DVDB, MCP, ADM, JPH, JMS, JK, PP, HS, IW, GC-T, GGG, SVT, DFE, RLM provided DNA samples and/or phenotypical data. All authors read and approved the final manuscript.

Further information about the financial report received is outlined in the supplementary file online.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement This would vary for each study—each study is listed in the supplemental material.

Results—For European women, strong evidence of association with breast cancer risk was observed for *PALB2* c.1592delT OR 3.44 (95% CI 1.39 to 8.52, $p=7.1\times 10^{-5}$), *PALB2* c.3113G>A OR 4.21 (95% CI 1.84 to 9.60, $p=6.9\times 10^{-8}$) and *ATM* c.7271T>G OR 11.0 (95% CI 1.42 to 85.7, $p=0.0012$). We also found evidence of association with breast cancer risk for three variants in *CHEK2*, c.349A>G OR 2.26 (95% CI 1.29 to 3.95), c.1036C>T OR 5.06 (95% CI 1.09 to 23.5) and c.538C>T OR 1.33 (95% CI 1.05 to 1.67) ($p=0.017$). Evidence for prostate cancer risk was observed for *CHEK2* c.1343T>G OR 3.03 (95% CI 1.53 to 6.03, $p=0.0006$) for African men and *CHEK2* c.1312G>T OR 2.21 (95% CI 1.06 to 4.63, $p=0.030$) for European men. No evidence of association with ovarian cancer was found for any of these variants.

Conclusions—This report adds to accumulating evidence that at least some variants in these genes are associated with an increased risk of breast cancer that is clinically important.

INTRODUCTION

The rapid introduction of massive parallel sequencing (MPS) into clinical genetics services is enabling the screening of multiple breast cancer susceptibility genes in one assay at reduced cost for women who are at increased risk of breast (and other) cancer. These gene panels now typically include the so-called ‘moderate-risk’ breast cancer susceptibility genes, including *PALB2*, *CHEK2* and *ATM*.^{1–3} However, mutations in these genes are individually extremely rare and limited data are available with which to accurately estimate the risk of cancer associated with them.

Estimation of the age-specific cumulative risk (penetrance) of breast cancer associated with specific mutations in these three genes has been limited to those that have been observed more frequently, such as *PALB2* c.1592delT (a Finnish founder mutation), *PALB2* c.3113G>A and *ATM* c.7271T>G. These mutations have been estimated to be associated with a 40% (95% CI 17% to 77%), 91% (95% CI 44% to 100%) and 52% (95% CI 28% to 80%) cumulative risk of breast cancer to the age of 70 years, respectively.^{4–7} These findings, based on segregation analyses in families of population-based case series, indicate that at least some mutations in these ‘moderate-risk’ genes are associated with a breast cancer risk comparable to that of the average pathogenic mutation in *BRCA2*: 45% (95% CI 31% to 56%).⁸ However, such estimates are imprecise and, moreover, may be confounded by modifying genetic variants or other familial risk factors.

Case-control studies provide an alternative approach to estimating cancer risks associated with specific variants. This design can estimate the relative risk directly, without making assumptions about the modifying effects of other risk factors. However, because these variants are rare, such studies need to be extremely large to provide precise estimates.

The clearest evidence for association, and the most precise breast cancer risk estimates, for rare variants in *PALB2*, *CHEK2* and *ATM* relate to protein truncating and splice-junction variants.^{9,10} However, studies based on mutation screening in case-control studies, combined with stratification of variants by their evolutionary likelihood suggest that at least some evolutionarily unlikely missense substitutions are associated with a similar risk to those conferred by truncating mutations.^{11–13} For example, Tavtigian *et al*.¹² estimated an OR of 2.85 (95% CI 0.83 to 4.86) for evolutionarily unlikely missense substitutions in the 3′ third

of *ATM*, which is comparable to that for truncating variants. Specifically, *ATM* c.7271C>G has been associated with a more substantial breast cancer risk in several studies.^{7,13} Le Calvez-Kelm *et al*,¹¹ estimated that the ORs associated with rare mutations in *CHEK2* from similarly designed studies were 6.18 (95% CI 1.76 to 21.8) for rare protein-truncating and splice-junction variants and 8.75 (95% CI 1.06 to 72.2) for evolutionarily unlikely missense substitutions.¹¹

It is plausible that monoallelic mutations in *PALB2*, *CHEK2* and *ATM* could be associated with increased risk of cancers other than breast cancer, as has been observed for *BRCA1* and *BRCA2* and both ovarian and prostate cancers.^{14–17} However, with the exception of pancreatic cancer in *PALB2* carriers, there is little evidence to support or refute the existence of such associations, although a few individually striking pedigrees have been observed.^{4,8,18–20}

In this study we selected rare genetic variants on the basis that they had been observed in breast cancer candidate gene case-control screening projects involving *PALB2*, *CHEK2* or *ATM*. These included three rare variants in *PALB2*: the protein truncating variants c.1592delT (p.Leu531Cysfs)⁴ and c.3113 G>A (p.Trp1038*)⁶ and the missense variant c.2816T>G, (p. Leu939Trp), six rare missense variants in *CHEK2*: c.349A>G (p.Arg117Gly) and c.1036C>T (p.Arg346Cys) predicted to be deleterious on the basis of evolutionary conservation,¹¹ c.538C>T (p.Arg180Cys), c.715G>A (p.Glu239Lys), c.1312G>T (p.Asp438Tyr) and c.1343T>G (p.Ile448Ser) and *ATM* c.7271T>G (p.Val2424Gly).⁷ We assessed the association of these variants with breast, ovarian and prostate risk by case-control analyses in three large consortia participating in the Collaborative Oncological Gene-environment Study.^{21,22}

METHODS

Participants

Participants were drawn from studies participating in three consortia as follows:

The *Breast Cancer Association Consortium (BCAC)*, involving a total of 48 studies: 37 of women from populations with predominantly European ancestry (42 671 cases and 42 164 controls), 9 of Asian women (5795 cases and 6624 controls) and 2 of African-American women (1046 cases and 932 controls). All cases had invasive breast cancer. The majority of studies were population-based or hospital-based case-control studies, but some studies of European women oversampled cases with a family history or with bilateral disease (see online supplementary table S1). Overall, 79% of BCAC cases with known Estrogen Receptor (ER) status (23% missing) are ER-positive. The proportion of cases selected by family history that are ER-positive is 78% (38% missing).

The *Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome (PRACTICAL)* involving a total of 26 studies: 25 included men with European ancestry (22 301 cases and 22 320 controls) and 3 included African-American men (623 cases and 569 controls). The majority of studies were population-based or hospital-based case-control studies (see online supplementary table S2).

The *Ovarian Cancer Association Consortium (OCAC)*, involving a total of 46 studies. Some studies were case-only and their data were combined with case-control studies from the same geographical region (leaving 36 study groupings). Of these groupings, 33 included women from populations with predominantly European ancestry (16 287 cases (14 542 with invasive disease) and 23 491 controls), 25 included Asian women (813 cases (720 with invasive disease) and 1574 controls), 17 included African-American women (186 cases (150 with invasive disease) and 200 controls) and 29 included women of other ethnic origin (893 cases (709 with invasive disease) and 864 controls). The majority of studies were population-based or hospital-based case-control studies (see online supplementary table S3).

Details regarding sample quality control have been published previously.^{22,23} All study participants gave informed consent and all studies were approved by the corresponding local ethics committees (see online supplementary tables S1–S3).

Variant selection

We selected for genotyping 13 rare mutations that had been observed in population-based case-control mutation screening studies. These variants were *PALB2* (c.1592delT, p. Leu531Cysfs;⁴⁵¹⁰ c.2323C>T p.Gln775*;²⁰ c.2816T>G, p. Leu939Trp;²²⁰ c.3113G>A, p.Trp1038*;²⁶²⁰ c.3116delA, p. Asn1039Ilefs;²⁶²⁰ c.3549C>G, p.Tyr1183*²), *CHEK2* (c.349A>G, p.ArgR117Gly; c.538C>T, p.Arg180Cys; c.715G>A p.Glu239Lys; c.1036C>T, p.Arg346Cys; c.1312G>T, p.Asp438Tyr; c.1343T>G, p.Ile448Ser)¹¹ and *ATM* (c.7271T>G, p.Val2424Gly)⁷¹³²⁴ see table 1. A DNA sample carrying each of these variants was included in a plate of control DNAs that was distributed to each genotyping centre to assist with quality control and genotype calling.

Genotyping

Three *PALB2* variants c.2323C>T (p.Gln775*), c.3116delA (p.Asn1039Ilefs) and c.3549C>G (p.Tyr1183*) were unable to be designed for measurement on the custom Illumina iSelect genotyping array and were not considered further (table 1). Genotyping was conducted using a custom Illumina Infinium array (iCOGS) in four centres, as part of a multiconsortia collaboration as described previously.²² Genotypes were called using Illumina's proprietary GenCall algorithm and then, for the data generated from the rare variant probes, manually confirmed with reference to the positive control sample. Two per cent of samples were provided in duplicate by all studies and 270 HapMap2 samples were genotyped in all four genotyping centres. Subjects with an overall call rate <95% were excluded. Plates with call rates <90% were excluded on a variant-by-variant basis. Cluster plots generated for all of the 10 rare variants were manually checked to confirm automated calls (see online supplementary figure S1).

Statistical methods

The association of each variant with breast, prostate and ovarian cancer risk was assessed using unconditional logistic regression to estimate ORs for carriers versus non-carriers, adjusting for study (categorical). p Values were determined by the likelihood ratio test comparing models with and without carrier status as a covariate. We also applied conditional logistic regression, defining risk sets by study, and found that this made no difference to the

OR estimates, CIs or p values to two significant figures; since model convergence was a problem for this latter regression analysis, all subsequent analyses were based on unconditional logistic regression. For the main analyses of breast cancer risk in European women, we also included as covariates the first six principal components, together with a seventh component specific to one study (Leuven Multidisciplinary Breast Centre (LMBC)) for which there was substantial inflation not accounted for by the components derived from the analysis of all studies. Addition of further principal components did not reduce inflation further. Data from all breast cancer studies were included to assess statistical significance. Data from cases selected for inclusion based on personal or family history of breast cancer were excluded in order to obtain unbiased OR estimates for the general population of white European women (leaving 37 039 cases and 38 260 controls from 32 studies). Multiple testing was adjusted for using the Benjamini-Hochberg procedure to control the false discovery rate, with a significance threshold of 0.05.²⁵ Reported p values are unadjusted unless otherwise stated. Reported CIs are all nominal. We included two race-specific principal components in each of the main breast cancer analyses of Asian and African-American women. Similar analyses were conducted using the data from PRACTICAL and OCAC, consistent with those used previously.^{23,26} All analyses were carried out using Stata: Release V.10 (StataCorp, 2008).

RESULTS

PALB2

In BCAC, *PALB2* c.1592delT (Leu531Cysfs) was only observed in 35 cases and 6 controls, all from four studies from Sweden and Finland (Helsinki Breast Cancer Study (HEBCS), Kuopio Breast Cancer Project (KBCP), Oulu Breast Cancer Study (OBCS) and Karolinska Mammography Project for Risk Prediction Breast Cancer (pKARMA); see online supplementary table S1), giving strong evidence of association with breast cancer risk ($p=7.1\times 10^{-5}$); the OR estimate was 4.52 (95% CI 1.90 to 10.8) based on all studies and 3.44 (95% CI 1.39 to 8.52) based on unselected cases and controls (table 2). We also found evidence of heterogeneity by ER status ($p=0.0023$), the association being stronger for ER-negative disease (OR 6.49 (95% CI 2.17 to 19.4) versus 2.24 (95% CI 1.05 to 7.24) for ER-positive disease).

PALB2 c.3113G>A (p.Trp1038*) was identified in 44 cases and 8 controls from nine BCAC studies. Only one carrier of the variant was of non-European origin. Strong evidence of association with breast cancer risk was observed ($p=6.9\times 10^{-8}$), with an estimated OR of 5.93 (95% CI 2.77 to 12.7) based on all studies and 4.21 (95% CI 1.85 to 9.61) based on unselected cases and controls. There was no evidence of a differential association by ER status ($p=0.15$).

Based on unselected cases, the estimated OR associated with carrying either of these *PALB2* variants (c.1592delT or c.3113G>A) was 3.85 (95% CI 2.09 to 7.09).

PALB2 c.2816T>G (p.Leu939Trp) was identified in 150 cases and 145 controls and there was no evidence of association with risk of breast cancer. There was no evidence of

association with risk of prostate or ovarian cancer for any of the three *PALB2* variants (see tables 3 and 4).

CHEK2

CHEK2 c.349A>G (p.Arg117Gly) was identified in 44 cases and 18 controls in studies participating in BCAC; all of these women were of European origin. We found evidence of association with breast cancer (p=0.003), with little change in the OR after excluding selected cases (OR 2.03 (95% CI 1.10 to 3.73)).

CHEK2 c.538C>T (p.Arg180Cys) was identified in 158 breast cancer cases and 142 controls in studies of white Europeans. Evidence of association with breast cancer risk (p=0.016) was observed, with an unbiased OR estimate of 1.34 (95% CI 1.06 to 1.70). A consistent OR estimate was observed for Asian women, based on 45 case and 45 control carriers (OR 1.16 (95% CI 0.75 to 1.76)).

CHEK2 c.715G>A (p.Glu239Lys) mutations were identified in 15 cases and 9 controls, all European women participating in BCAC and no evidence of association with risk of breast cancer was observed (p=0.21).

CHEK2 c.1036C>T (p.Arg346Cys) was identified in nine cases from seven studies and two controls from two different studies in BCAC (neither control carrier was from a study that had case carriers), all of European origin. We found evidence of association with breast cancer risk (p=0.017) with reduced OR estimate of 3.39 (95% CI 0.68 to 16.9) after excluding selected cases.

None of the above four *CHEK2* variants (*CHEK2* c.349A>G (p.Arg117Gly); c.538C>T (p.Arg180Cys); c.715G>A (p.Glu239Lys) and c.1036C>T (p.Arg346Cys)) were found to be associated with an increased risk of prostate or ovarian cancer (tables 3 and 4). *CHEK2* variant c.1312G>T (p.Asp438Tyr) was not associated with risk of breast cancer for European women (p=0.91). Variant c.1343T>G (p.Ile448Ser) was not observed in any breast cancer cases of European or Asian origin. It was detected in 48 cases and 29 controls of African origin, giving weak evidence of association (OR 1.52 (95% CI 0.95 to 2.43, p=0.083)). *CHEK2* c.1312G>T (p.Asp438Tyr) was identified in 23 cases and 11 controls from PRACTICAL, all European, providing evidence of association with prostate cancer risk (OR 2.21 (95% CI 1.06 to 4.63, p=0.030)). *CHEK2* c.1343T>G (p.Ile448Ser) was observed in 35 cases and 11 controls, all African, participating in PRACTICAL and was also associated with an increased risk of prostate cancer (OR 3.03 (95% CI 1.53 to 6.03, p=0.00059)). There was no evidence that these *CHEK2* variants were associated with risk of ovarian cancer (table 4).

ATM

ATM c.7271T>G (p.Val2424Gly) was identified in 12 cases and 1 control in studies participating in BCAC, all of European origin, giving evidence of association with breast cancer risk (p=0.0012). The OR estimate based on unselected studies was 11.0 (95% CI 1.42 to 85.7). There was no evidence of association of this variant with prostate or ovarian cancer risk (see tables 3 and 4).

DISCUSSION

The present report adds to an accumulating body of evidence that at least some *rare variants* in so-called ‘moderate-risk’ genes are associated with an increased risk of breast cancer that is of clinical relevance.

These findings are presented at a time when detailed information about variants in these genes is becoming more readily available via the translation of diagnostic genetic testing from Sanger sequencing-based testing platforms to MPS platforms that test panels of genes in single assays.^{27–29} The vast majority of information about *PALB2*, *CHEK2* and *ATM*, variants generated from these new testing platforms is not being used in clinical genetics services due to lack of reliable estimates of the cancer risk associated with individual variants, or groups of variants, in each gene. Previous analyses have been largely based on selected families, relying on data on the segregation of the variant. The present study is by far the largest to take a case-control approach. Consistent with previous reports,^{5–7,911–13} *PALB2* c.3113G>A (p.Trp1038*), *PALB2* c.1592delT (p.Leu531Cysfs) and *ATM* c.7271T>G (p.Val2424Gly) were found to be associated with substantially increased risk of breast cancer all with associated relative risk estimates of 3.44 or greater.

The estimates for the two loss-of-function *PALB2* variants (c.1592delT and c.3113G<A) were consistent with each other and with estimates based on segregation analysis.⁵⁶⁹ We found no evidence of association with breast cancer for *PALB2* c.2816T>G (p.Leu939Trp), with an upper 95% confidence limit excluding an OR >1.5 which is notable given the Align-Grantham Variation Grantham Deviation (Align-GVGD) score and the observed impact on protein function.³⁰

The estimate for *ATM* c.7271T>G (p.Val2424Gly) was also consistent with that found by segregation analysis.⁷¹³ The substantial increased risk of breast cancer associated with *ATM* c.7271T>G (p.Val2424Gly) could be due to the reduction in kinase activity (with near-normal protein levels) observed for *ATM* p.Val2424Gly,³¹ thus this variant is likely to be acting as a dominant negative mutation.³²

In contrast, we found no evidence of an association with risk of prostate or ovarian cancer with any of these three variants: however, the confidence limits were wide; based on the upper 95% confidence limit we could exclude an OR of >1.4 for prostate cancer for the loss-of-function *PALB2* c.3113G>A and 1.9 for c.1592delT and c.3113G>A combined.

We analysed six rare missense variants in *CHEK2*. Two of these (*CHEK2* c.349A>G (p.Arg117Gly; rs28909982) and c.1036C>T (p.Arg346Cys)) had evidence of a significant impact on the protein based on in silico prediction. We proposed these variants for inclusion in the iCOGS design as they had been identified in 3/1242 cases and 1/1089 controls and 3/1242 cases and 0/1089 controls, respectively, in a population-based case-control mutation screening study of *CHEK2*.¹¹ In that study, Le Calvez-Kelm *et al*, estimated an OR of 8.75 (95% CI 1.06 to 72.2) for variants with an Align-GVGD score C65 (based on nine cases and one control). The current analysis provides confirmatory evidence of this association in a much larger sample (OR 2.18 (95% CI 1.23 to 3.85)) including 40 unselected case and 18 control carriers. The evidence that *CHEK2* is a breast cancer susceptibility gene is largely

based on studies of protein truncating variants, in particular *CHEK2* 1100delC.³³ Reports of the association of the missense variant I157T, (C15) and breast cancer risk have been conflicting but a large meta-analysis involving 15 985 breast cancer cases and 18 609 controls estimated a modest OR of 1.58 (95% CI 1.42 to 1.75).³⁴ We also found evidence ($p=0.015$) of an association for c.538C>T (Align-GVGD C25); OR 1.34 (95% CI 1.06 to 1.70), a risk comparable to I157T.

The p values reported above have not been adjusted for multiple testing. This was not considered appropriate for the associations with breast cancer risk of *PALB2* c.1592delT, c.3113G>A and *ATM* c.7271T>G because these associations had previously been reported; our aim was to more precisely estimate the associated relative risks. All three associations with breast cancer risk reported for *CHEK2* variants remained statistically significant after adjusting for the other tests conducted in relation to breast cancer risk, but not after correcting for all tests for all cancers. Nevertheless, the findings for *CHEK2* c.349A>G and c.1036C>T confirmed those reported previously, although collectively. The association observed with *CHEK2* c.538C>T requires independent replication.

Do this approach and new data have an impact on clinical recommendations for women and families carrying these rare genetic variants? Although age-specific cumulative risks for cancer are more informative for genetic counselling and clinical management of carriers, our study provides information that is relevant to clinical recommendations. As discussed in Easton *et al.*,³⁵ a relative risk of 4 will place a woman in a 'high-risk' category (in the absence of any other risk factor) and a relative risk between 2 and 4 will place a woman in this category if other risk factors are present. Thus, several of the variants included in this report (*PALB2* c.1592delT; c.3113G>A *ATM* c.7271T>G) would place the carrier in a high-risk group, especially if other risk factors, such as a family history, are present. The high level of breast cancer risk associated with *PALB2* c.1592delT and c.3113G>A reported here is consistent with the penetrance estimate reported for a group of loss-of-function mutations in *PALB2*⁹ and has an advantage in terms of clinical utility that the estimates in this study have been made at a mutation-specific level. Therefore, this work provides important information for risk reduction recommendations (such as prophylactic mastectomy and potentially salpingo-oophorectomy) for carriers of these variants. However, further prospective research is required to characterise these risks and to understand the potential of other risk-reducing strategies such as salpingo-oophorectomy and chemoprevention.

The consistency of the relative risk estimates with those derived through family based studies supports the hypothesis that these variants combine multiplicatively with other genetic loci and familial risk factors; this information is critical for deriving comprehensive risk models. Even with very large sample sizes such as those studied here, however, it is still only possible to derive individual risk estimates for a limited set of variants, and even for these variants the estimates are still imprecise. This internationally collaborative approach also has limited capacity to improve risk estimates for rare variants that are only observed in specific populations. Inevitably, therefore, risk models will depend on combining data across multiple variants, using improved in silico predictions and potentially biochemical/functional evidence to synthesise these estimates efficiently. It will also be necessary

develop counselling and patient management strategies that can accommodate a multifactorial approach to variant classification.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Melissa C Southey¹, David E Goldgar², Robert Winqvist³, Katri Pylkäs³, Fergus Couch⁴, Marc Tischkowitz⁵, William D Foulkes⁶, Joe Dennis⁷, Kyriaki Michailidou⁷, Elizabeth J van Rensburg⁸, Tuomas Heikkinen⁹, Heli Nevanlinna⁹, John L Hopper¹⁰, Thilo Dörk¹¹, Kathleen BM Claes¹², Jorge Reis-Filho¹³, Zhi Ling Teo¹, Paolo Radice¹⁴, Irene Catucci¹⁵, Paolo Peterlongo¹⁵, Helen Tsimiklis¹, Fabrice A Odefrey¹, James G Dowty¹⁰, Marjanka K Schmidt¹⁶, Annegien Broeks¹⁶, Frans B Hogervorst¹⁶, Senno Verhoef¹⁶, Jane Carpenter¹⁷, Christine Clarke¹⁸, Rodney J Scott¹⁹, Peter A Fasching^{20,21}, Lothar Haeberle^{20,22}, Arif B Ekici²³, Matthias W Beckmann²⁰, Julian Peto²⁴, Isabel dos-Santos-Silva²⁴, Olivia Fletcher²⁵, Nichola Johnson²⁵, Manjeet K Bolla⁷, Elinor J Sawyer²⁶, Ian Tomlinson²⁷, Michael J Kerin²⁸, Nicola Miller²⁸, Federik Marme^{29,30}, Barbara Burwinkel^{29,31}, Rongxi Yang^{29,31}, Pascal Guénel^{32,33}, Thérèse Truong^{32,33}, Florence Menegaux^{32,33}, Marie Sanchez^{32,33}, Stig Bojesen^{34,35}, Sune F Nielsen^{34,35}, Henrik Flyger³⁶, Javier Benitez^{37,38}, M Pilar Zamora³⁹, Jose Ignacio Arias Perez⁴⁰, Primitiva Menéndez⁴¹, Hoda Anton-Culver⁴², Susan Neuhausen⁴³, Argyrios Ziogas⁴⁴, Christina A Clarke⁴⁵, Hermann Brenner^{46,47,48}, Volker Arndt⁴⁶, Christa Stegmaier⁴⁹, Hiltrud Brauch^{48,50,51}, Thomas Brüning⁵², Yon-Dschun Ko⁵³, Taru A Muranen⁵⁴, Kristiina Aittomäki⁵⁵, Carl Blomqvist⁵⁶, Natalia V Bogdanova^{11,57}, Natalia N Antonenkova⁵⁸, Annika Lindblom⁵⁹, Sara Margolin⁶⁰, Arto Mannermaa^{61,62}, Vesa Kataja^{63,64}, Veli-Matti Kosma^{61,62}, Jaana M Hartikainen^{61,62}, Amanda B Spurdle⁶⁵, kConFab Investigators⁶⁶, Australian Ovarian Cancer Study Group^{65,66}, Els Wauters^{67,68}, Dominiek Smeets^{67,68}, Benoit Beuselinck⁶⁹, Giuseppe Floris⁶⁹, Jenny Chang-Claude⁷⁰, Anja Rudolph⁷⁰, Petra Seibold⁷⁰, Dieter Flesch-Janys⁷¹, Janet E Olson⁷², Celine Vachon⁷², Vernon S Pankratz⁷², Catriona McLean⁷³, Christopher A Haiman⁷⁴, Brian E Henderson⁷⁴, Fredrick Schumacher⁷⁴, Loic Le Marchand⁷⁵, Vessela Kristensen^{76,77}, Grethe Grenaker Alnæs⁷⁶, Wei Zheng⁷⁸, David J Hunter^{79,80}, Sara Lindstrom^{79,80}, Susan E Hankinson^{80,81}, Peter Kraft^{79,80}, Irene Andrulic^{82,83}, Julia A Knight^{84,85}, Gord Glendon⁸², Anna Marie Mulligan^{86,87}, Arja Jukkola-Vuorinen⁸⁸, Mervi Grip⁸⁹, Salla Kauppila⁹⁰, Peter Devilee⁹¹, Robert A E M Tollenaar⁹¹, Caroline Seynaeve^{92,98}, Antoinette Hollestelle^{92,98}, Montserrat Garcia-Closas⁹³, Jonine Figueroa⁹⁴, Stephen J Chanock⁹⁴, Jolanta Lissowska⁹⁵, Kamila Czene⁹⁶, Hatef Darabi⁹⁶, Mikael Eriksson⁹⁶, Diana M Eccles⁹⁷, Sajjad Rafiq⁹⁷, William J Tapper⁹⁷, Sue M Gerty⁹⁷, Maartje J Hooning⁹⁸, John W M Martens⁹⁸, J Margriet Collée⁹⁹, Madeleine Tilanus-Linthorst¹⁰⁰, Per Hall¹⁰¹, Jingmei Li¹⁰², Judith S Brand¹⁰¹, Keith Humphreys¹⁰¹, Angela Cox¹⁰³, Malcolm W R Reed¹⁰³, Craig Luccarini¹⁰⁴, Caroline Baynes¹⁰⁴, Alison M Dunning¹⁰⁴, Ute Hamann¹⁰⁵, Diana Torres^{105,106}, Hans Ulrich Ulmer¹⁰⁷, Thomas Rüdiger¹⁰⁸, Anna Jakubowska¹⁰⁹, Jan

Lubinski¹⁰⁹, Katarzyna Jaworska^{109,110}, Katarzyna Durda¹⁰⁹, Susan Slager⁷², Amanda E Toland¹¹¹, Christine B Ambrosone¹¹², Drakoulis Yannoukakos¹¹³, Anthony Swerdlow^{114,115}, Alan Ashworth⁹³, Nick Orr⁹³, Michael Jones¹¹⁴, Anna González-Neira³⁷, Guillermo Pita³⁷, M Rosario Alonso³⁷, Nuria Álvarez³⁷, Daniel Herrero³⁷, Daniel C Tessier¹¹⁶, Daniel Vincent¹¹⁷, Francois Bacot¹¹⁷, Jacques Simard¹¹⁸, Martine Dumont¹¹⁸, Penny Soucy¹¹⁸, Rosalind Eeles^{119,120}, Kenneth Muir¹²¹, Fredrik Wiklund¹²², Henrik Gronberg¹²², Johanna Schleutker^{123,124}, Børge G Nordestgaard¹²⁵, Maren Weischer¹²⁶, Ruth C Travis¹²⁷, David Neal¹²⁸, Jenny L Donovan¹²⁹, Freddie C Hamdy¹³⁰, Kay-Tee Khaw¹³¹, Janet L Stanford^{132,133}, William J Blot¹³⁴, Stephen Thibodeau⁴, Daniel J Schaid⁷², Joseph L Kelley¹³⁵, Christiane Maier^{136,137}, Adam S Kibel^{138,139}, Cezary Cybulski¹⁴⁰, Lisa Cannon-Albright¹⁴¹, Katja Butterbach⁴⁶, Jong Park¹⁴², Radka Kaneva¹⁴³, Jyotsna Batra¹⁴⁴, Manuel R Teixeira¹⁴⁵, Zsafia Kote-Jarai¹¹⁹, Ali Amin Al Olama⁷, Sara Benlloch⁷, Stefan P Renner¹⁴⁶, Arndt Hartmann¹⁴⁷, Alexander Hein¹⁴⁶, Matthias Ruebner¹⁴⁶, Diether Lambrechts^{148,149}, Els Van Nieuwenhuysen¹⁵⁰, Ignace Vergote¹⁵⁰, Sandrina Lambrechts¹⁵⁰, Jennifer A Doherty¹⁵¹, Mary Anne Rossing^{152,153}, Stefan Nickels¹⁵⁴, Ursula Eilber¹⁵⁴, Shan Wang-Gohrke¹⁵⁵, Kunle Odunsi¹⁵⁶, Lara E Sucheston-Campbell¹⁵⁶, Grace Friel¹⁵⁶, Galina Lurie¹⁵⁷, Jeffrey L Killeen¹⁵⁸, Lynne R Wilkens¹⁵⁷, Marc T Goodman^{159,160}, Ingo Runnebaum¹⁶¹, Peter A Hillemanns¹⁶², Liisa M Pelttari⁹, Ralf Butzow¹⁶³, Francesmary Modugno^{164,165}, Robert P Edwards¹³⁵, Roberta B Ness¹⁶⁶, Kirsten B Moysich¹⁶⁷, Andreas du Bois^{168,169}, Florian Heitz^{168,169}, Philipp Harter^{168,169}, Stefan Kommoss^{169,170}, Beth Y Karlan¹⁷¹, Christine Walsh¹⁷¹, Jenny Lester¹⁷¹, Allan Jensen¹⁷², Susanne Krüger Kjaer^{172,173}, Estrid Høgdall^{172,174}, Bernard Peissel¹⁷⁵, Bernardo Bonanni¹⁷⁶, Loris Bernard¹⁷⁷, Ellen L Goode⁷², Brooke L Fridley¹⁷⁸, Robert A Vierkant⁷², Julie M Cunningham⁴, Melissa C Larson⁷², Zachary C Fogarty⁷², Kimberly R Kalli¹⁷⁹, Dong Liang¹⁸⁰, Karen H Lu¹⁸¹, Michelle A T Hildebrandt¹⁸², Xifeng Wu¹⁸², Douglas A Levine¹⁸³, Fanny Dao¹⁸³, Maria Bisogna¹⁸³, Andrew Berchuck¹⁸⁴, Edwin S Iversen¹⁸⁵, Jeffrey R Marks¹⁸⁶, Lucy Akushevich¹⁸⁷, Daniel W Cramer¹⁸⁸, Joellen Schildkraut¹⁸⁷, Kathryn L Terry¹⁸⁸, Elizabeth M Poole^{189,190}, Meir Stampfer^{80,189}, Shelley S Tworoger^{189,190}, Elisa V Bandera¹⁹¹, Irene Orlow¹⁹², Sara H Olson¹⁹², Line Bjorge^{193,194}, Helga B Salvesen^{193,194}, Anne M van Altena¹⁹⁵, Katja K H Aben^{196,197,198}, Lambertus A Kiemeny¹⁹⁶, Leon F A G Massuger¹⁹⁵, Tanja Pejovic¹⁹⁹, Yukie Bean¹⁹⁹, Angela Brooks-Wilson^{200,201}, Linda E Kelemen^{202,203}, Linda S Cook²⁰⁴, Nhu D Le²⁰⁵, Bohdan Górski²⁰⁶, Jacek Gronwald²⁰⁶, Janusz Menkiszak²⁰⁷, Claus K Høgdall¹⁷³, Lene Lundvall²⁰⁸, Lotte Nedergaard²⁰⁹, Svend Aage Engelholm²¹⁰, Ed Dicks²¹¹, Jonathan Tyrer²¹¹, Ian Campbell²¹², Iain McNeish²¹³, James Paul²¹⁴, Nadeem Siddiqui²¹⁵, Rosalind Glasspool²¹⁵, Alice S Whittemore²¹⁶, Joseph H Rothstein²¹⁶, Valerie McGuire²¹⁶, Weiva Sieh²¹⁶, Hui Cai⁷⁸, Xiao-Ou Shu⁷⁸, Rachel T Teten²¹⁷, Rebecca Sutphen²¹⁷, John R McLaughlin²¹⁸, Steven A Narod²¹⁹, Catherine M Phelan²²⁰, Alvaro N Monteiro²²⁰, David Fenstermacher²²¹, Hui-Yi Lin²²¹, Jennifer B Permuth²²⁰, Thomas A Sellers²²⁰, Y Ann Chen²²¹, Ya-Yu Tsai²²⁰, Zhihua Chen²²¹, Aleksandra Gentry-Maharaj²²², Simon A Gayther²²³, Susan J Ramus²²³, Usha Menon²²², Anna H Wu²²³, Celeste L Pearce²²³, David Van Den Berg²²³, Malcolm C Pike^{223,224},

Agnieszka Dansonka-Mieszkowska²²⁵, Joanna Plisiecka-Halasa²²⁵, Joanna Moes-Sosnowska²²⁵, Jolanta Kupryjanczyk²²⁵, Paul DP Pharoah²¹¹, Honglin Song²¹¹, Ingrid Winship^{226,227}, Georgia Chenevix-Trench⁶⁵, Graham G Giles^{10,228}, Sean V Tavtigian², Doug F Easton⁷, and Roger L Milne^{10,228}

Affiliations

¹Genetic Epidemiology Laboratory, Department of Pathology, The University of Melbourne, Melbourne, Australia ²Huntsman Cancer Institute, Salt Lake City, UT, USA ³Laboratory of Cancer Genetics and Tumor Biology, Cancer and Translational Medicine Research Unit and Biocenter Oulu, University of Oulu, Nordlab Oulu, Oulu, Finland ⁴Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA ⁵Department of Medical Genetics and National Institute for Health Research Cambridge Biomedical Research Centre, University of Cambridge, and the Department of Clinical Genetics, East Anglian Regional Genetics Service, Addenbrooke's Hospital ⁶Program in Cancer Genetics, Department of Human Genetics and Oncology, Lady Davis Institute, and Research Institute, McGill University Health Centre, McGill University, Montreal, Canada ⁷Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Strangeways Laboratory, Worts Causeway, Cambridge, UK ⁸Department of Genetics, University of Pretoria, South Africa ⁹Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland ¹⁰Centre for Epidemiology and Biostatistics, School of Population and Global Health, The University of Melbourne, Melbourne, Australia ¹¹Gynaecology Research Unit, Hannover Medical School, Hannover, Germany ¹²Center for Medical Genetics, Ghent University Hospital, De Pintelaan 185, 9000, Ghent, Belgium ¹³Department of Pathology and Human Oncology and Pathogenesis Program, Memorial Sloan-Kettering Cancer Center, New York, New York, USA ¹⁴Unit of Molecular Bases of Genetic Risk and Genetic Testing, Department of Preventive and Predictive Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori (INT), Milan, Italy ¹⁵IFOM, the FIRC Institute of Molecular Oncology, Milan, Italy ¹⁶Netherlands Cancer Institute, Antoni van Leeuwenhoek hospital, Amsterdam, The Netherlands ¹⁷Australian Breast Cancer Tissue Bank, University of Sydney at the Westmead Institute for Medical Research, NSW, Australia ¹⁸Centre for Cancer Research, University of Sydney at the Westmead Institute for Medical Research, NSW, Australia ¹⁹Division of Molecular Medicine, Pathology North, Newcastle and University of Newcastle, NSW, Australia ²⁰University Breast Center Franconia, Department of Gynecology and Obstetrics, University Hospital Erlangen, Friedrich-Alexander University Erlangen-Nuremberg, Comprehensive Cancer Center Erlangen-EMN, Erlangen, Germany ²¹David Geffen School of Medicine, Department of Medicine Division of Hematology and Oncology, University of California at Los Angeles, CA, USA ²²Unit of Biostatistics, Department of Gynecology and Obstetrics, University Hospital Erlangen, Friedrich-Alexander University Erlangen-Nuremberg, Erlangen, Germany ²³Institute of Human Genetics, University Hospital Erlangen, Friedrich Alexander University Erlangen-Nuremberg, Erlangen, Germany ²⁴Non-communicable Disease Epidemiology Department,

London School of Hygiene and Tropical Medicine, London, UK ²⁵Breakthrough Breast Cancer Research Centre, The Institute of Cancer Research, London, UK ²⁶Division of Cancer Studies, NIHR Comprehensive Biomedical Research Centre, Guy's & St. Thomas' NHS Foundation Trust in partnership with King's College London, London, UK ²⁷Wellcome Trust Centre for Human Genetics and Oxford Biomedical Research Centre, University of Oxford, UK and Oxford NIHR Biomedical Research Centre, Headington, OX3 7LE ²⁸Surgery, Lambe Institute for Translational Science, NUI Galway, University Hospital Galway, Galway, Ireland ²⁹Department of Obstetrics and Gynecology, University of Heidelberg, Heidelberg, Germany ³⁰National Center for Tumor Diseases, University of Heidelberg, Heidelberg, Germany ³¹Molecular Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany ³²Inserm (National Institute of Health and Medical Research), CESP (Center for Research in Epidemiology and Population Health), U1018, Environmental Epidemiology of Cancer, Villejuif, France ³³University Paris-Sud, UMRS 1018, Villejuif, France ³⁴Copenhagen General Population Study, Herlev Hospital, Copenhagen University Hospital, University of Copenhagen, Copenhagen, Denmark ³⁵Department of Clinical Biochemistry, Herlev Hospital, Copenhagen University Hospital, University of Copenhagen, Copenhagen, Denmark ³⁶Department of Breast Surgery, Herlev Hospital, Copenhagen University Hospital, Copenhagen, Denmark ³⁷Human Genetics Group, Human Cancer Genetics Program, Spanish National Cancer Research Centre (CNIO), Madrid, Spain ³⁸Centro de Investigación en Red de Enfermedades Raras (CIBERER), Valencia, Spain ³⁹Servicio de Oncología Médica, Hospital Universitario La Paz, Madrid, Spain ⁴⁰Servicio de Cirugía General y Especialidades, Hospital Monte Naranco, Oviedo, Spain ⁴¹Servicio de Anatomía Patológica, Hospital Monte Naranco, Oviedo, Spain ⁴²Department of Epidemiology, University of California Irvine, Irvine, California, USA ⁴³Beckman Research Institute of City of Hope, Duarte, California, USA ⁴⁴Department of Epidemiology, University of California Irvine, Irvine, California, USA ⁴⁵Cancer Prevention Institute of California, Fremont, California, USA ⁴⁶Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany ⁴⁷Division of Preventive Oncology, German Cancer Research Center (DKFZ), Heidelberg, Germany ⁴⁸German Cancer Consortium (DKTK), German Cancer Research Center (DKFZ), Heidelberg, Germany ⁴⁹Saarland Cancer Registry, Saarbrücken, Germany ⁵⁰Dr. Margarete Fischer-Bosch-Institute of Clinical Pharmacology, Stuttgart ⁵¹University of Tübingen, Tübingen, Germany ⁵²Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr University, Bochum (IPA), Germany ⁵³Department of Internal Medicine, Evangelische Kliniken Bonn gGmbH, Johanniter Krankenhaus, Bonn, Germany ⁵⁴Department of Obstetrics and Gynecology, University of Helsinki and Helsinki University Central Hospital, Helsinki, Finland ⁵⁵Department of Clinical Genetics, Helsinki University Central Hospital, Helsinki, Finland ⁵⁶Department of Oncology, Helsinki University Central Hospital, Helsinki, Finland ⁵⁷Department of Radiation Oncology, Hannover Medical School, Hannover, Germany ⁵⁸N.N. Alexandrov Research Institute of Oncology and Medical

Radiology, Minsk, Belarus ⁵⁹Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden ⁶⁰Department of Oncology – Pathology, Karolinska Institutet, Stockholm, Sweden ⁶¹School of Medicine, Institute of Clinical Medicine, Pathology and Forensic Medicine, and Cancer Center of Eastern Finland, University of Eastern Finland, Kuopio, Finland ⁶²Imaging Center, Department of Clinical Pathology, Kuopio University Hospital, Kuopio, Finland ⁶³School of Medicine, Institute of Clinical Medicine, Oncology, University of Eastern Finland, Kuopio, Finland ⁶⁴Biocenter Kuopio, Cancer Center of Eastern Finland, Kuopio University Hospital, Kuopio, Finland ⁶⁵QIMR Berghofer Medical Research Institute, Brisbane, Australia ⁶⁶Research Department, Peter MacCallum Cancer Centre and The Sir Peter MacCallum Department of Oncology, University of Melbourne, Victoria, Australia ⁶⁷Vesalius Research Center (VRC), VIB, Leuven, Belgium ⁶⁸Laboratory for Translational Genetics, Department of Oncology, University of Leuven, Leuven, Belgium ⁶⁹University Hospital Gasthuisberg, Leuven, Belgium ⁷⁰Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany ⁷¹Department of Cancer Epidemiology/Clinical Cancer Registry and Institute for Medical Biometrics and Epidemiology, University Clinic Hamburg-Eppendorf, Hamburg, Germany ⁷²Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA ⁷³Anatomical Pathology, The Alfred Hospital, Melbourne, Australia ⁷⁴Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA ⁷⁵Epidemiology Program, Cancer Research Center, University of Hawaii, Honolulu, HI, USA ⁷⁶Department of Genetics, Institute for Cancer Research, Oslo University Hospital, Radiumhospitalet, Oslo, Norway ⁷⁷Faculty of Medicine (Faculty Division Ahus), University of Oslo (UiO), Norway ⁷⁸Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN, USA ⁷⁹Program in Molecular and Genetic Epidemiology, Harvard School of Public Health, Boston, MA, USA ⁸⁰Department of Epidemiology, Harvard School of Public Health, Boston, MA, USA ⁸¹Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA ⁸²Ontario Cancer Genetics Network, Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Toronto, Ontario, Canada ⁸³Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada ⁸⁴Prosserman Centre for Health Research, Lunenfeld-Tanenbaum Research Institute of Mount Sinai Hospital, Toronto, Ontario, Canada ⁸⁵Division of Epidemiology, Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada ⁸⁶Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada ⁸⁷Laboratory Medicine Program, University Health Network, Toronto, Ontario; Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada ⁸⁸Department of Oncology, Oulu University Hospital, University of Oulu, Oulu, Finland ⁸⁹Department of Surgery, Oulu University Hospital, University of Oulu, Oulu, Finland ⁹⁰Department of Pathology, Oulu University Hospital, University of Oulu, Oulu, Finland ⁹¹Department of Surgical Oncology, Leiden University Medical Center, 2300 RC Leiden, The Netherlands

⁹²Family Cancer Clinic, Department of Medical Oncology, Erasmus MC-Daniel den Hoed Cancer Centre, Rotterdam, The Netherlands ⁹³The Breast Cancer Now Toby Robins Research Centre, The Institute of Cancer Research, London, SW3 6JB, UK ⁹⁴Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland, USA ⁹⁵Department of Cancer Epidemiology and Prevention, M. Sklodowska-Curie Memorial Cancer Center & Institute of Oncology, Warsaw, Poland ⁹⁶Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 17177, Sweden ⁹⁷Faculty of Medicine, University of Southampton (UoS), Southampton UK ⁹⁸Department of Medical Oncology, Family Cancer Clinic, Erasmus MC Cancer Institute, Rotterdam, The Netherlands ⁹⁹Department of Clinical Genetics, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands ¹⁰⁰Department of Surgical Oncology, Family Cancer Clinic, Erasmus University Medical Center, Rotterdam, The Netherlands ¹⁰¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm 17177, Sweden ¹⁰²Human Genetics Division, Genome Institute of Singapore, Singapore 138672, Singapore ¹⁰³Sheffield Cancer Research, Department of Oncology, University of Sheffield, Sheffield, UK ¹⁰⁴Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK ¹⁰⁵Molecular Genetics of Breast Cancer, German Cancer Research Center (DKFZ), Heidelberg, Germany ¹⁰⁶Institute of Human Genetics, Pontificia Universidad Javeriana, Bogota, Colombia ¹⁰⁷Frauenklinik der Stadtklinik Baden-Baden, Baden-Baden, Germany ¹⁰⁸Institute of Pathology, Städtisches Klinikum Karlsruhe, Karlsruhe, Germany ¹⁰⁹Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland ¹¹⁰Postgraduate School of Molecular Medicine, Warsaw Medical University, Warsaw, Poland ¹¹¹Department of Molecular Virology, Immunology and Medical Genetics, Comprehensive Cancer Center, The Ohio State University, Columbus, OH, USA ¹¹²Roswell Park Cancer Institute, Buffalo, New York, USA ¹¹³Molecular Diagnostics Laboratory, IRRP, National Centre for Scientific Research "Demokritos", Aghia Paraskevi Attikis, Athens, Greece ¹¹⁴Division of Genetics and Epidemiology, Institute of Cancer Research, London, UK ¹¹⁵Division of Breast Cancer Research, Institute of Cancer Research, London, UK ¹¹⁶Centre d'innovation Genome Quebec et University McGill Montreal Quebec, Canada ¹¹⁷McGill University, Montreal, Quebec, Canada ¹¹⁸Cancer Genomics Laboratory, Centre Hospitalier Universitaire de Quebec Research Center. Laval University, Quebec, Canada ¹¹⁹The Institute of Cancer Research, London, SM2 5NG, UK ¹²⁰Royal Marsden NHS Foundation Trust, Fulham, London, SW3 6JJ, UK ¹²¹University of Warwick, Coventry, UK ¹²²Department of Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden ¹²³Department of Medical Biochemistry and Genetics, University of Turku, and Tyks Microbiology and Genetics, Department of Medical Genetics, Turku University Hospital, Turku, Finland ¹²⁴Institute of Biomedical Technology/BioMediTech, University of Tampere, Tampere, Finland ¹²⁵Department of Clinical Biochemistry, Herlev Hospital, Copenhagen University Hospital, Herlev Ringvej 75, DK-2730 Herlev, Denmark ¹²⁶Department of Human Genetics University of Utah, Salt Lake City, UT, USA and Department of Clinical Biochemistry, Herlev

Hospital, Copenhagen University Hospital, University of Copenhagen, Copenhagen, Denmark ¹²⁷Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK ¹²⁸Surgical Oncology (Uro-Oncology: S4), University of Cambridge, Box 279, Addenbrooke's Hospital, Hills Road, Cambridge, UK and Cancer Research UK Cambridge Research Institute, Li Ka Shing Centre, Cambridge, UK ¹²⁹Professor of Social Medicine, University of Bristol, Canynge Hall, 39 Whatley Road, Bristol BS8 2PS ¹³⁰Nuffield Department of Surgical Sciences, Old Road Campus Research Building (off Roosevelt Drive), University of Oxford, Headington, Oxford, OX3 7DQ ¹³¹Cambridge Institute of Public Health, University of Cambridge, Forvie Site, Robinson Way, Cambridge CB2 0SR ¹³²Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA ¹³³Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington, USA ¹³⁴International Epidemiology Institute, 1455 Research Blvd., Suite 550, Rockville, MD 20850 ¹³⁵Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA ¹³⁶Department of Urology, University Hospital Ulm, Germany ¹³⁷Institute of Human Genetics University Hospital Ulm, Germany ¹³⁸Brigham and Women's Hospital/Dana-Farber Cancer Institute, 45 Francis Street-ASB II-3, Boston, MA 02115 ¹³⁹Washington University, St Louis, Missouri ¹⁴⁰International Hereditary Cancer Center, Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland ¹⁴¹Division of Genetic Epidemiology, Department of Medicine, University of Utah School of Medicine ¹⁴²Division of Cancer Prevention and Control, H. Lee Moffitt Cancer Center, 12902 Magnolia Dr., Tampa, Florida, USA ¹⁴³Molecular Medicine Center and Department of Medical Chemistry and Biochemistry, Medical University – Sofia, 2 Zdrave St, 1431, Sofia, Bulgaria ¹⁴⁴Australian Prostate Cancer Research Centre-Qld, Institute of Health and Biomedical Innovation and Schools of Life Science and Public Health, Queensland University of Technology, Brisbane, Australia ¹⁴⁵Department of Genetics, Portuguese Oncology Institute, Porto, Portugal and Biomedical Sciences Institute (ICBAS), Porto University, Porto, Portugal ¹⁴⁶University Hospital Erlangen, Department of Gynecology and Obstetrics, Friedrich-Alexander-University Erlangen-Nuremberg, Comprehensive Cancer Center Erlangen-EMN, Universitaetsstrasse 21-23, 91054 Erlangen, Germany ¹⁴⁷University Hospital Erlangen, Institute of Pathology, Friedrich-Alexander-University Erlangen-Nuremberg, Comprehensive Cancer Center Erlangen-EMN, Universitaetsstrasse 21-23, 91054 Erlangen, German ¹⁴⁸Vesalius Research Center, VIB, Leuven, Belgium ¹⁴⁹Laboratory for Translational Genetics, Department of Oncology, University of Leuven, Belgium ¹⁵⁰Department of Epidemiology, The Geisel School of Medicine at Dartmouth, Lebanon, NH, USA ¹⁵¹Department of Epidemiology, The Geisel School of Medicine at Dartmouth, Hannover, NH, USA ¹⁵²Program in Epidemiology, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA ¹⁵³Department of Epidemiology, University of Washington, Seattle, WA, USA ¹⁵⁴German Cancer Research Center, Division of Cancer Epidemiology, Heidelberg, Germany ¹⁵⁵Department of Obstetrics and

Gynecology, University of Ulm, Ulm, Germany ¹⁵⁶Department of Gynecological Oncology, Roswell Park Cancer Institute, Buffalo, NY ¹⁵⁷Cancer Epidemiology Program, University of Hawaii Cancer Center, Hawaii, USA ¹⁵⁸Department of Pathology, Kapiolani Medical Center for Women and Children, John A. Burns School of Medicine, University of Hawaii, Honolulu, Hawaii 96826, USA ¹⁵⁹Cancer Prevention and Control, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California, USA ¹⁶⁰Community and Population Health Research Institute, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, California, USA ¹⁶¹Department of Gynecology and Obstetrics, Friedrich Schiller University, Jena University Hospital, Jena, Germany ¹⁶²Clinics of Obstetrics and Gynaecology, Hannover Medical School, Hannover, Germany ¹⁶³Department of Pathology, Helsinki University Central Hospital, Helsinki, 00029 HUS, Finland ¹⁶⁴University of Pittsburgh Department of Obstetrics, Gynecology and Reproductive Sciences and Ovarian Cancer Center of Excellence Pittsburgh PA USA ¹⁶⁵University of Pittsburgh Department of Epidemiology, University of Pittsburgh Graduate School of Public Health and Womens Cancer Research Program, Magee-Womens Research Institute and University of Pittsburgh Cancer Institute Pittsburgh PA USA ¹⁶⁶The University of Texas School of Public Health, Houston, TX, USA ¹⁶⁷Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY ¹⁶⁸Department of Gynecology and Gynecologic Oncology, Kliniken Essen-Mitte/ Evang. Huysens-Stiftung/ Knappschaft GmbH, Essen, Germany ¹⁶⁹Department of Gynecology and Gynecologic Oncology, Dr. Horst Schmidt Kliniken Wiesbaden, Wiesbaden, Germany ¹⁷⁰Tuebingen University Hospital, Department of Women's Health, Tuebingen, Germany ¹⁷¹Women's Cancer Program at the Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California ¹⁷²Department of Virus, Lifestyle and Genes, Danish Cancer Society Research Center, Copenhagen, Denmark ¹⁷³Department of Obstetrics and Gynecology, Rigshospitalet, Copenhagen, Denmark ¹⁷⁴Molecular Unit, Department of Pathology, Herlev Hospital, University of Copenhagen, Copenhagen, Denmark ¹⁷⁵Unit of Medical Genetics, Department of Preventive and Predictive Medicine, Fondazione IRCCS Istituto Nazionale dei Tumori (INT), Milan, Italy ¹⁷⁶Division of Cancer Prevention and Genetics, Istituto Europeo di Oncologia (IEO), Milan, Italy ¹⁷⁷Department of Experimental Oncology, Istituto Europeo di Oncologia (IEO), Milan, Italy and Cogentech Cancer Genetic Test Laboratory, Milan, Italy ¹⁷⁸University of Kansas Medical Center, Kansas City, KS, USA ¹⁷⁹Department of Medical Oncology, Mayo Clinic, Rochester, Minnesota, USA ¹⁸⁰College of Pharmacy and Health Sciences, Texas Southern University, Houston, Texas, USA ¹⁸¹Department of Gynecologic Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA ¹⁸²Department of Epidemiology, The University of Texas MD Anderson Cancer Center, Houston, Texas, USA ¹⁸³Gynecology Service, Department of Surgery, Memorial Sloan-Kettering Cancer Center, New York, NY, USA ¹⁸⁴Department of Obstetrics and Gynecology, Duke University Medical Center, Durham, North Carolina, USA ¹⁸⁵Department of

Statistical Science, Duke University, Durham, North Carolina, USA ¹⁸⁶Department of Surgery, Duke University Medical Center, Durham, North Carolina, USA ¹⁸⁷Cancer Prevention, Detection & Control Research Program, Duke Cancer Institute, Durham, North Carolina, USA ¹⁸⁸Obstetrics and Gynecology Epidemiology Center, Brigham and Women's Hospital, Boston, Massachusetts, USA ¹⁸⁹Channing Division of Network Medicine, Brigham and Women's Hospital and Harvard Medical School ¹⁹⁰Department of Epidemiology, Harvard TH Chan School of Public Health, Boston, Massachusetts, USA ¹⁹¹Cancer Prevention and Control Program, Rutgers Cancer Institute of New Jersey, The State University of New Jersey, New Brunswick, NJ, USA ¹⁹²Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, USA ¹⁹³Department of Gynecology and Obstetrics, Haukeland University Hospital, Bergen, Norway ¹⁹⁴Centre for Cancer Biomarkers, Department of Clinical Sciences, University of Bergen, Bergen, Norway ¹⁹⁵Radboud university medical center, Department of Gynaecology, Nijmegen, Netherlands ¹⁹⁶Radboud university medical centre, Radboud Institute for Health Sciences, Nijmegen, Netherlands ¹⁹⁷Netherlands Comprehensive Cancer Organisation, Utrecht, Netherlands ¹⁹⁸Department of Obstetrics & Gynecology, Oregon Health & Science University ¹⁹⁹Knight Cancer Institute, Oregon Health & Science University, Portland, Oregon, USA ²⁰⁰Canada's Michael Smith Genome Sciences Centre, BC Cancer Agency, Vancouver, BC, Canada ²⁰¹Department of Biomedical Physiology and Kinesiology, Simon Fraser University, Burnaby, BC Canada ²⁰²Department of Public Health Sciences, College of Medicine, Medical University of South Carolina, SC, USA ²⁰³Hollings Cancer Center, Medical University of South Carolina, SC, USA ²⁰⁴Division of Epidemiology and Biostatistics, Department of Internal Medicine, University of New Mexico, Albuquerque, New Mexico, USA ²⁰⁵Cancer Control Research, BC Cancer Agency, Vancouver, BC, Canada ²⁰⁶International Hereditary Cancer Center, Department of Genetics and Pathology, Pomeranian Medical University, Szczecin, Poland ²⁰⁷Department of Gynecological Surgery and Gynecological Oncology of Adults and Adolescents, Pomeranian Medical University, Szczecin, Poland ²⁰⁸Gyn Clinic, Rigshospitalet, University of Copenhagen, Denmark ²⁰⁹Department of Pathology, Rigshospitalet, University of Copenhagen, Denmark ²¹⁰Department of Oncology, Rigshospitalet, University of Copenhagen, Denmark ²¹¹Department of Oncology, University of Cambridge, Strangeways Research laboratory, Cambridge, UK ²¹²Cancer Genetics Laboratory, Research Division, Peter MacCallum Cancer Centre, St Andrews Place, East Melbourne ²¹³Institute of Cancer Sciences, University of Glasgow, Wolfson Wohl Cancer Research Centre, Beatson Institute for Cancer Research, Glasgow, UK ²¹⁴The Cancer Research UK Clinical Trials Unit, Beatson West of Scotland Cancer Centre, 1053 Great Western Road, Glasgow, G12 0YN ²¹⁵Department of Gynaecological Oncology, Glasgow Royal Infirmary ²¹⁶Department of Health Research and Policy - Epidemiology, Stanford University School of Medicine, Stanford CA, USA ²¹⁷Epidemiology Center, College of Medicine, University of South Florida, Tampa, Florida, USA ²¹⁸Public Health Ontario, Toronto, Canada ²¹⁹Women's College Research Institute, University of Toronto, Toronto, Ontario, Canada ²²⁰Department of Cancer Epidemiology, Moffitt

Cancer Center, Tampa, FL, USA ²²¹Department of Biostatistics and Bioinformatics, Moffitt Cancer Center, Tampa, FL, USA ²²²Women's Cancer, Institute for Women's Health, UCL, London, United Kingdom ²²³Department of Preventive Medicine, Keck School of Medicine, University of Southern California Norris Comprehensive Cancer Center, Los Angeles, California, USA ²²⁴Department of Epidemiology and Biostatistics, Memorial Sloan-Kettering Cancer Center, New York, New York, USA ²²⁵Department of Pathology and Laboratory Diagnostics, The Maria Sklodowska-Curie Memorial Cancer Center and Institute of Oncology, Warsaw, Poland ²²⁶Department of Medicine, The University of Melbourne Health, Australia ²²⁷The Royal Melbourne Hospital, Victoria 3050, Australia ²²⁸Cancer Epidemiology Centre, Cancer Council Victoria, Victoria, Australia

Acknowledgments

The authors thank the following for their contributions to this study: Qin Wang (BCAC), Lesley McGuffog, and Ken Offit (CIMBA), Andrew Lee, and Ed Dicks and the staff of the Centre for Genetic Epidemiology Laboratory, staff of the CNIO genotyping unit, Sylvie LaBoissière and Frederic Robidoux and the staff of the McGill University and Génome Québec Innovation Centre, staff of the Copenhagen DNA laboratory, and Sharon A Windebank, Christopher A Hilker, Jeffrey Meyer and the staff of Mayo Clinic Genotyping Core Facility. The authors also thank UM1 CA167552 (Willett) Cancer Epidemiology Cohort in Male Health Professionals, P01 CA87969 (Stampfer) Dietary and Hormonal Determinants of Cancer in Women, UM1 CA186107 (Stampfer) Long term multidisciplinary study of cancer in women: The Nurses' Health Study, R01CA141298 (Stampfer) Growth Factors and Lethal Prostate Cancer Signature, NCI: K07-CA80668 DAMD17-02-1-0669 NIH/National Center for Research Resources/General Clinical Research Center grant MO1-RR000056, R01CA095023, P50-CA159981, NCI CCSG award P30-CA008748 (Memorial Sloan Kettering Cancer Center), Department of Defense (W81XWH-07-0449), National Health and Medical Research Council of Australia APP1029974 and APP1061177, Deutsche Krebshilfe (Maier), Wellcome Trust Centre for Human Genetics from the Wellcome Trust (090532/Z/09/Z). The authors thank the National Institute of Health Research for their support to The Institute of Cancer Research and Royal Marsden NHS Foundation Trust, London, UK.

Funding Funding for the iCOGS infrastructure came from: the European Community's Seventh Framework Programme under grant agreement n° 223175 (HEALTH-F2-2009-223175) (COGS), Cancer Research UK (C1287/A10118, C1287/A 10710, C12292/A11174, C5047/A8384, C5047/A15007, C5047/A10692, CRUK C8197/A10123), the National Institutes of Health (CA128978) and Post-Cancer GWAS initiative (No. 1 U19 CA 148537—the GAME-ON initiative), the Department of Defense (W81XWH-10-1-0341), the Canadian Institutes of Health Research (CIHR) for the CIHR Team in Familial Risks of Breast Cancer, Komen Foundation for the Cure, the Breast Cancer Research Foundation, the Ovarian Cancer Research Fund and Susan G Komen (WF).

REFERENCES

1. Stratton MR, Rahman N. The emerging landscape of breast cancer susceptibility. *Nat Genet.* 2008; 40:17–22. [PubMed: 18163131]
2. Rahman N, Seal S, Thompson D, Kelly P, Renwick A, Elliott A, Reid S, Spanova K, Barfoot R, Chagtai T, Jayatilake H, McGuffog L, Hanks S, Evans DG, Eccles D. Breast Cancer Susceptibility Collaboration (UK), Easton DF, Stratton MR. PALB2, which encodes a BRCA2-interacting protein, is a breast cancer susceptibility gene. *Nat Genet.* 2007; 39:165–167. [PubMed: 17200668]
3. Southey MC, Teo ZL, Winship I. PALB2 and breast cancer: ready for clinical translation! *Appl Clin Genet.* 2013; 6:43–52. [PubMed: 23935381]
4. Erkkö H, Xia B, Nikkilä J, Schleutker J, Syrjäkoski K, Mannermaa A, Kallioniemi A, Pykäs K, Karpainen SM, Rapakko K, Miron A, Sheng Q, Li G, Mattila H, Bell DW, Haber DA, Grip M, Reiman M, Jukkola-Vuorinen A, Mustonen A, Kere J, Aaltonen LA, Kosma VM, Kataja V, Soini Y, Drapkin RI, Livingston DM, Winqvist R. A recurrent mutation in PALB2 in Finnish cancer families. *Nature.* 2007; 446:316–319. [PubMed: 17287723]

5. Erkko H, Dowty JG, Nikkilä J, Syrjäkoski K, Mannermaa A, Pylkäs K, Southey MC, Holli K, Kallioniemi A, Jukkola-Vuorinen A, Kataja V, Kosma VM, Xia B, Livingston DM, Winqvist R, Hopper JL. Penetrance analysis of the PALB2 c.1592delT founder mutation. *Clin Cancer Res.* 2008; 14:4667–4671. [PubMed: 18628482]
6. Southey MC, Teo ZL, Dowty JG, Odefrey FA, Park DJ, Tischkowitz M, Sabbaghian N, Apicella C, Byrnes GB, Winship I, Baglietto L, Giles GG, Goldgar DE, Foulkes WD, Hopper JL, kConFab for the Beast Cancer Family Registry. A PALB2 mutation associated with high risk of breast cancer. *Breast Cancer Res.* 2010; 12:R109. [PubMed: 21182766]
7. Bernstein JL, Teraoka S, Southey MC, Jenkins MA, Andrulis IL, Knight JA, John EM, Lapinski R, Wolitzer AL, Whittemore AS, West D, Seminara D, Olson ER, Spurdle AB, Chenevix-Trench G, Giles GG, Hopper JL, Concannon P. Population-based estimates of breast cancer risks associated with ATM gene variants c.7271T>G and c.1066-6T>G (IVS10-6T>G) from the Breast Cancer Family Registry. *Hum Mutat.* 2006; 27:1122–1128. [PubMed: 16958054]
8. Antoniou AC, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL, Olsson H, Johannsson O, Borg A, Pasini B, Radice P, Manoukian S, Eccles DM, Tang N, Olah E, Anton-Culver H, Warner E, Lubinski J, Gronwald J, Gorski B, Tulinius H, Thorlacius S, Eerola H, Nevanlinna H, Syrjäkoski K, Kallioniemi OP, Thompson D, Evans C, Peto J, Lalloo F, Evans DG, Easton DF. Average risks of breast and ovarian cancer associated with BRCA1 or BRCA2 mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet.* 2003; 72:1117–1130. [PubMed: 12677558]
9. Antoniou AC, Casadei S, Heikkinen T, Barrowdale D, Pylkäs K, Roberts J, Lee A, Subramanian D, De Leeneer K, Fostira F, Tomiak E, Neuhausen SL, Teo ZL, Khan S, Aittomäki K, Moilanen JS, Turnbull C, Seal S, Mannermaa A, Kallioniemi A, Lindeman GJ, Buys SS, Andrulis IL, Radice P, Tondini C, Manoukian S, Toland AE, Miron P, Weitzel JN, Domchek SM, Poppe B, Claes KB, Yannoukakos D, Concannon P, Bernstein JL, James PA, Easton DF, Goldgar DE, Hopper JL, Rahman N, Peterlongo P, Nevanlinna H, King MC, Couch FJ, Southey MC, Winqvist R, Foulkes WD, Tischkowitz M. Breast-cancer risk in families with mutations in PALB2. *N Engl J Med.* 2014; 371:497–506. [PubMed: 25099575]
10. Heikkinen T, Kärkkäinen H, Aaltonen K, Milne RL, Heikkilä P, Aittomäki K, Blomqvist C, Nevanlinna H. The breast cancer susceptibility mutation PALB2 1592delT is associated with an aggressive tumor phenotype. *Clin Cancer Res.* 2009; 15:3214–3222. [PubMed: 19383810]
11. Le Calvez-Kelm F, Lesueur F, Damiola F, Vallée M, Voegelé C, Babikyan D, Durand G, Forey N, McKay-Chopin S, Robinot N, Nguyen-Dumont T, Thomas A, Byrnes GB, Breast Cancer Family Registry. Hopper JL, Southey MC, Andrulis IL, John EM, Tavtigian SV. Rare, evolutionarily unlikely missense substitutions in CHEK2 contribute to breast cancer susceptibility: results from a breast cancer family registry case-control mutation-screening study. *Breast Cancer Res.* 2011; 13:R6. [PubMed: 21244692]
12. Tavtigian SV, Oefner PJ, Babikyan D, Hartmann A, Healey S, Le Calvez-Kelm F, Lesueur F, Byrnes GB, Chuang SC, Forey N, Feuchtinger C, Gioia L, Hall J, Hashibe M, Herte B, McKay-Chopin S, Thomas A, Vallée MP, Voegelé C, Webb PM, Whiteman DC, Australian Cancer Study; Breast Cancer Family Registries (BCFR); Kathleen Cuningham Foundation Consortium for Research into Familial Aspects of Breast Cancer (kConFab). Sangrajrang S, Hopper JL, Southey MC, Andrulis IL, John EM, Chenevix-Trench G. Rare, evolutionarily unlikely missense substitutions in ATM confer increased risk of breast cancer. *Am J Hum Genet.* 2009; 85:427–446. [PubMed: 19781682]
13. Goldgar DE, Healey S, Dowty JG, Da Silva L, Chen X, Spurdle AB, Terry MB, Daly MJ, Buys SM, Southey MC, Andrulis I, John EM, BCFR; kConFab. Khanna KK, Hopper JL, Oefner PJ, Lakhani S, Chenevix-Trench G. Rare variants in the ATM gene and risk of breast cancer. *Breast Cancer Res.* 2011; 13:R73. [PubMed: 21787400]
14. Thompson D, Easton DF. Breast Cancer Linkage Consortium. Cancer incidence in BRCA1 mutation carriers. *J Natl Cancer Inst.* 2002; 94:1358–1365. [PubMed: 12237281]
15. Edwards SM, Kote-Jarai Z, Meitz J, Hamoudi R, Hope Q, Osin P, Jackson R, Southgate C, Singh R, Falconer A, Dearnaley DP, Ardern-Jones A, Murkin A, Dowe A, Kelly J, Williams S, Oram R, Stevens M, Teare DM, Ponder BA, Gayther SA, Easton DF, Eeles RA. Cancer Research UK/ British Prostate Group UK Familial Prostate Cancer Study Collaborators; British Association of Urological Surgeons Section of Oncology. Two percent of men with early-onset prostate cancer

- harbor germline mutations in the BRCA2 gene. *Am J Hum Genet.* 2003; 72:1–12. [PubMed: 12474142]
16. Breast Cancer Linkage Consortium. Cancer risks in BRCA2 mutation carriers. *J Natl Cancer Inst.* 1999; 91:1310–1316. [PubMed: 10433620]
 17. Leongamornlert D, Mahmud N, Tymrakiewicz M, Saunders E, Dadaev T, Castro E, Goh C, Govindasami K, Guy M, O'Brien L, Sawyer E, Hall A, Wilkinson R, Easton D, UKGPCS, Collaborators Goldgar D, Eeles R, Kote-Jarai Z. Germline BRCA1 mutations increase prostate cancer risk. *Br J Cancer.* 2012; 106:1697–1701. [PubMed: 22516946]
 18. Tischkowitz M, Sabbaghian N, Ray AM, Lange EM, Foulkes WD, Cooney KA. Analysis of the gene coding for the BRCA2-interacting protein PALB2 in hereditary prostate cancer. *Prostate.* 2008; 68:675–678. [PubMed: 18288683]
 19. Dansonka-Mieszkowska A, Kluska A, Moes J, Dabrowska M, Nowakowska D, Niwinska A, Derlatka P, Cendrowski K, Kupryjanczyk J. A novel germline PALB2 deletion in Polish breast and ovarian cancer patients. *BMC Med Genet.* 2010; 11:20. [PubMed: 20122277]
 20. Teo ZL, Park DJ, Provenzano E, Chatfield CA, Odefrey FA, Nguyen-Dumont T, kConFab. Dowty JG, Hopper JL, Winship I, Goldgar DE, Southey MC. Prevalence of PALB2 mutations in Australasian multiple-case breast cancer families. *Breast Cancer Res.* 2013; 15:R17. [PubMed: 23448497]
 21. Sakoda LC, Jorgenson E, Witte JS. Turning of COGS moves forward findings for hormonally mediated cancers. *Nat Genet.* 2013; 45:345–348. [PubMed: 23535722]
 22. Michailidou K, Hall P, Gonzalez-Neira A, Ghoussaini M, Dennis J, Milne RL, Schmidt MK, Chang-Claude J, Bojesen SE, Bolla MK, Wang Q, Dicks E, Lee A, Turnbull C, Rahman N, Breast and Ovarian Cancer Susceptibility Collaboration; Fletcher O, Peto J, Gibson L, Dos Santos Silva I, Nevanlinna H, Muranen TA, Aittomäki K, Blomqvist C, Czene K, Irwanto A, Liu J, Waisfisz Q, Meijers-Heijboer H, Adank M, Hereditary Breast and Ovarian Cancer Research Group Netherlands (HEBON); van der Luijt RB, Hein R, Dahmen N, Beckman L, Meindl A, Schmutzler RK, Müller-Myhsok B, Lichtner P, Hopper JL, Southey MC, Makalic E, Schmidt DF, Uitterlinden AG, Hofman A, Hunter DJ, Chanock SJ, Vincent D, Bacot F, Tessier DC, Canisius S, Wessels LF, Haiman CA, Shah M, Luben R, Brown J, Luccarini C, Schoof N, Humphreys K, Li J, Nordestgaard BG, Nielsen SF, Flyger H, Couch FJ, Wang X, Vachon C, Stevens KN, Lambrechts D, Moisse M, Paridaens R, Christiaens MR, Rudolph A, Nickels S, Flesch-Janys D, Johnson N, Aitken Z, Aaltonen K, Heikkinen T, Broeks A, Veer LJ, van der Schoot CE, Guénel P, Truong T, Laurent-Puig P, Menegaux F, Marme F, Schneeweiss A, Sohn C, Burwinkel B, Zamora MP, Perez JI, Pita G, Alonso MR, Cox A, Brock IW, Cross SS, Reed MW, Sawyer EJ, Tomlinson I, Kerin MJ, Miller N, Henderson BE, Schumacher F, Le Marchand L, Andrulis IL, Knight JA, Glendon G, Mulligan AM, kConFab Investigators, Australian Ovarian Cancer Study Group; Lindblom A, Margolin S, Hooning MJ, Hollestelle A, van den Ouweland AM, Jager A, Bui QM, Stone J, Dite GS, Apicella C, Tsimiklis H, Giles GG, Severi G, Baglietto L, Fasching PA, Haeberle L, Ekici AB, Beckmann MW, Brenner H, Müller H, Arndt V, Stegmaier C, Swerdlow A, Ashworth A, Orr N, Jones M, Figueroa J, Lissowska J, Brinton L, Goldberg MS, Labrèche F, Dumont M, Winqvist R, Pylkäs K, Jukkola-Vuorinen A, Grip M, Brauch H, Hamann U, Brüning T, GENICA (Gene Environment Interaction and Breast Cancer in Germany) Network/. Radice P, Peterlongo P, Manoukian S, Bonanni B, Devilee P, Tollenaar RA, Seynaeve C, van Asperen CJ, Jakubowska A, Lubinski J, Jaworska K, Durda K, Mannermaa A, Kataja V, Kosma VM, Hartikainen JM, Bogdanova NV, Antonenkova NN, Dörk T, Kristensen VN, Anton-Culver H, Slager S, Toland AE, Edge S, Fostira F, Kang D, Yoo KY, Noh DY, Matsuo K, Ito H, Iwata H, Sueta A, Wu AH, Tseng CC, Van Den Berg D, Stram DO, Shu XO, Lu W, Gao YT, Cai H, Teo SH, Yip CH, Phuah SY, Cornes BK, Hartman M, Miao H, Lim WY, Sng JH, Muir K, Lophatananon A, Stewart-Brown S, Siriwanarangsana P, Shen CY, Hsiung CN, Wu PE, Ding SL, Sangrajrang S, Gaborieau V, Brennan P, McKay J, Blot WJ, Signorello LB, Cai Q, Zheng W, Deming-Halverson S, Shrubsole M, Long J, Simard J, Garcia-Closas M, Pharoah PD, Chenevix-Trench G, Dunning AM, Benitez J, Easton DF. Large-scale genotyping identifies 41 new loci associated with breast cancer risk. *Nat Genet.* 2013; 45:353–361. [PubMed: 23535729]
 23. Pharoah PD, Tsai YY, Ramus SJ, Phelan CM, Goode EL, Lawrenson K, Buckley M, Fridley BL, Tyrer JP, Shen H, Weber R, Karevan R, Larson MC, Song H, Tessier DC, Bacot F, Vincent D, Cunningham JM, Dennis J, Dicks E, Australian Cancer Study, Australian Ovarian Cancer Study

- Group. Aben KK, Anton-Culver H, Antonenkova N, Armasu SM, Baglietto L, Bandera EV, Beckmann MW, Birrer MJ, Bloom G, Bogdanova N, Brenton JD, Brinton LA, Brooks-Wilson A, Brown R, Butzow R, Campbell I, Carney ME, Carvalho RS, Chang-Claude J, Chen YA, Chen Z, Chow WH, Cicek MS, Coetzee G, Cook LS, Cramer DW, Cybulski C, Dansonka-Mieszkowska A, Despierre E, Doherty JA, Dörk T, du Bois A, Dürst M, Eccles D, Edwards R, Ekici AB, Fasching PA, Fenstermacher D, Flanagan J, Gao YT, Garcia-Closas M, Gentry-Maharaj A, Giles G, Gjyshi A, Gore M, Gronwald J, Guo Q, Halle MK, Harter P, Hein A, Heitz F, Hillemanns P, Hoatlin M, Høgdall E, Høgdall CK, Hosono S, Jakubowska A, Jensen A, Kalli KR, Karlan BY, Kelemen LE, Kiemeny LA, Kjaer SK, Konecny GE, Krakstad C, Kupryjanczyk J, Lambrechts D, Lambrechts S, Le ND, Lee N, Lee J, Leminen A, Lim BK, Lissowska J, Lubinski J, Lundvall L, Lurie G, Massuger LF, Matsuo K, McGuire V, McLaughlin JR, Menon U, Modugno F, Moysich KB, Nakanishi T, Narod SA, Ness RB, Nevanlinna H, Nickels S, Noushmehr H, Odunsi K, Olson S, Orlow I, Paul J, Pejovic T, Pelttari LM, Permut-Wey J, Pike MC, Poole EM, Qu X, Risch HA, Rodriguez-Rodriguez L, Rossing MA, Rudolph A, Runnebaum I, Rzepecka IK, Salvesen HB, Schwaab I, Severi G, Shen H, Shridhar V, Shu XO, Sieh W, Southey MC, Spellman P, Tajima K, Teo SH, Terry KL, Thompson PJ, Timorek A, Tworoger SS, van Altena AM, van den Berg D, Vergote I, Vierkant RA, Vitonis AF, Wang-Gohrke S, Wentzensen N, Whittemore AS, Wik E, Winterhoff B, Woo YL, Wu AH, Yang HP, Zheng W, Ziogas A, Zulkifli F, Goodman MT, Hall P, Easton DF, Pearce CL, Berchuck A, Chenevix-Trench G, Iversen E, Monteiro AN, Gayther SA, Schildkraut JM, Sellers TA. GWAS meta-analysis and replication identifies three new susceptibility loci for ovarian cancer. *Nat Genet.* 2013; 45:362–370. [PubMed: 23535730]
24. Chenevix-Trench G, Spurdle AB, Gatei M, Kelly H, Marsh A, Chen X, Donn K, Cummings M, Nyholt D, Jenkins MA, Scott C, Pupo GM, Dörk T, Bendix R, Kirk J, Tucker K, McCredie MR, Hopper JL, Sambrook J, Mann GJ, Khanna KK. Dominant negative ATM mutations in breast cancer families. *J Natl Cancer Inst.* 2002; 94:205–215. [PubMed: 11830610]
25. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B.* 1995; 57:289–300.
26. Eeles RA, Olama AA, Benlloch S, Saunders EJ, Leongamornlert DA, Tymrakiewicz M, Ghoussaini M, Luccarini C, Dennis J, Jugurnauth-Little S, Dadaev T, Neal DE, Hamdy FC, Donovan JL, Muir K, Giles GG, Severi G, Wiklund F, Gronberg H, Haiman CA, Schumacher F, Henderson BE, Le Marchand L, Lindstrom S, Kraft P, Hunter DJ, Gapstur S, Chanock SJ, Berndt SI, Albanes D, Andriole G, Schleutker J, Weischer M, Canzian F, Riboli E, Key TJ, Travis RC, Campa D, Ingles SA, John EM, Hayes RB, Pharoah PD, Pashayan N, Khaw KT, Stanford JL, Ostrander EA, Signorello LB, Thibodeau SN, Schaid D, Maier C, Vogel W, Kibel AS, Cybulski C, Lubinski J, Cannon-Albright L, Brenner H, Park JY, Kaneva R, Batra J, Spurdle AB, Clements JA, Teixeira MR, Dicks E, Lee A, Dunning AM, Baynes C, Conroy D, Maranian MJ, Ahmed S, Govindasami K, Guy M, Wilkinson RA, Sawyer EJ, Morgan A, Dearnaley DP, Horwich A, Huddart RA, Khoo VS, Parker CC, Van As NJ, Woodhouse CJ, Thompson A, Dudderidge T, Ogden C, Cooper CS, Lophatananon A, Cox A, Southey MC, Hopper JL, English DR, Aly M, Adolfsson J, Xu J, Zheng SL, Yeager M, Kaaks R, Diver WR, Gaudet MM, Stern MC, Corral R, Joshi AD, Shahabi A, Wahlfors T, Tammela TL, Auvinen A, Virtamo J, Klarskov P, Nordestgaard BG, Røder MA, Nielsen SF, Bojesen SE, Siddiq A, Fitzgerald LM, Kolb S, Kwon EM, Karyadi DM, Blot WJ, Zheng W, Cai Q, McDonnell SK, Rinckleb AE, Drake B, Colditz G, Wokolorczyk D, Stephenson RA, Teerlink C, Muller H, Rothenbacher D, Sellers TA, Lin HY, Slavov C, Mitev V, Lose F, Srinivasan S, Maia S, Paulo P, Lange E, Cooney KA, Antoniou AC, Vincent D, Bacot F, Tessier DC. COGS–Cancer Research UK GWAS–ELLIPSE (part of GAME-ON) Initiative, Australian Prostate Cancer Bioresource, UK Genetic Prostate Cancer Study Collaborators/British Association of Urological Surgeons’ Section of Oncology, UK ProtecT (Prostate testing for cancer and Treatment) Study Collaborators, PRACTICAL (Prostate Cancer Association Group to Investigate Cancer-Associated Alterations in the Genome) Consortium, Kote-Jarai Z, Easton DF. Identification of 23 new prostate cancer susceptibility loci using the iCOGS custom genotyping array. *Nat Genet.* 2013; 45:385–391. [PubMed: 23535732]
27. Glusman G. Clinical applications of sequencing take center stage. *Genome Biol.* 2013; 14:303. [PubMed: 23651925]
28. Sikkema-Raddatz B, Johansson LF, de Boer EN, Almomani R, Boven LG, van den Berg MP, van Spaendonck-Zwarts KY, van Tintelen JP, Sijmons RH, Jongbloed JD, Sinke RJ. Targeted Next-

- Generation Sequencing can Replace Sanger Sequencing in Clinical Diagnostics. *Hum Mutat.* 2013; 34:1035–1042. [PubMed: 23568810]
29. Rattenberry E, Vialard L, Yeung A, Bair H, McKay K, Jafri M, Canham N, Cole TR, Denes J, Hodgson SV, Irving R, Izatt L, Korbonits M, Kumar AV, Lalloo F, Morrison PJ, Woodward ER, Macdonald F, Wallis Y, Maher ER. A comprehensive next generation sequencing based genetic testing strategy to improve diagnosis of inherited pheochromocytoma and paraganglioma. *J Clin Endocrinol Metab.* 2013; 98:E1248–E1256. [PubMed: 23666964]
 30. Park JY, Singh TR, Nassar N, Zhang F, Freund M, Hanenberg H, Meetei AR, Andreassen PR. Breast cancer-associated missense mutants of the PALB2 WD40 domain, which directly binds RAD51C, RAD51 and BRCA2, disrupt DNA repair. *Oncogene.* 2014; 33:4803–4812. [PubMed: 24141787]
 31. Taylor AM, Lam Z, Last JI, Byrd PJ. Ataxia telangiectasia: more variation at clinical and cellular levels. *Clin Genet.* 2015 Mar.87:199–208. [PubMed: 25040471]
 32. Waddell N, Jonnalagadda J, Marsh A, Grist S, Jenkins M, Hobson K, Taylor M, Lindeman GJ, Tavtigian SV, Suthers G, Goldgar D, Oefner PJ, kConFab Investigators. Taylor D, Grimmond S, Khanna KK, Chenevix-Trench G. Characterization of the breast cancer associated ATM 7271T<G (V2424G) mutation by gene expression profiling. *Genes Chromosomes Cancer.* 2006; 45:1169–1181. [PubMed: 17001622]
 33. Weischer M, Bojesen SE, Ellervik C, Tybjaerg-Hansen A, Nordestgaard BG. CHEK2*1100delC genotyping for clinical assessment of breast cancer risk: meta-analyses of 26,000 patient cases and 27,000 controls. *J Clin Oncol.* 2008; 26:542–548. [PubMed: 18172190]
 34. Han FF, Guo CL, Liu LH. The effect of CHEK2 variant I157T on cancer susceptibility: evidence from a meta-analysis. *DNA Cell Biol.* 2013; 32:329–335. [PubMed: 23713947]
 35. Easton DF, Pharoah PD, Antoniou AC, Tischkowitz M, Tavtigian SV, Nathanson KL, Devilee P, Meindl A, Couch FJ, Southey M, Goldgar DE, Evans DG, Chenevix-Trench G, Rahman N, Robson M, Domchek SM, Foulkes WD. Gene-panel sequencing and the prediction of breast-cancer risk. *N Engl J Med.* 2015; 372:2243–2257. [PubMed: 26014596]

Table 1

Rare genetic variants included in the iCOGS array.

Breast cancer risk estimates									
Gene	Variant*	Amino acid*	dbSNP rs	OR (95% CI)	Penetrance [†] (95% CI)	Align-GVGD	Reference(s)	Designed [‡]	Genotyped
PALB2	c.1592delT	p.Leu531Cysfs	rs180177102	3.94 (1.5–12.1), [§]	40% (17–77)	na	4, 5, 10	Yes	Yes
	c.2323C>T	p.Gln775*	rs180177111			na	25, 26	No	No
	c.2816T>G	p.Leu939Trp	rs45478192			C55	20	Yes	Yes
	c.3113G>A	p.Trp1038*	rs180177132		95% (44–100)	na	2, 6, 20	Yes	Yes
	c.3116delA	p.Asn1039Ilefs	rs180177133			na	2	No	No
CHEK2	c.3549C>G	p.Tyr1183*	rs118203998			na	2	No	No
	c.349A>G	p.Arg117Gly	rs28909982	8.75 (1.06–72.2), [¶]		C65	11	Yes	Yes
	c.538C>T	p.Arg180Cys	rs77130927	2.47 (0.45–13.49), ^{**}		C25	11	Yes	Yes
	c.715G>A	p.Glu239Lys	rs121908702	1.82 (0.62–5.34), ^{††}		C15	11	Yes	Yes
	c.1036C>T	p.Arg346Cys	na	8.75 (1.06–72.2), [¶]		C65	11	Yes	Yes
	c.1312G>T	p.Asp438Tyr	na	2.47 (0.45–13.49), ^{**}		C25	11	Yes	Yes
	c.1343T>G	p.Ile448Ser	rs17886163	1.82 (0.62–5.34), ^{††}		C15	11	Yes	Yes
	c.7271T>G	p.Val2424Gly	rs28904921		52% (28–80)	C65	7, 13, 23, 27	Yes	Yes

* Human Genome Variation Society (HGVS); reference sequences PALB2, NM_024675.3, NP_078951.2; CHEK2, NM_007194.3, NP_009125.1; ATM, NM_000051.3, NP_000042.3.

[†] Age-specific cumulative risk of breast cancer to age 70 years.^{5–7}

[‡] Able to be designed for measurement on the custom Illumina iSelect genotyping array.^{21,22}

[§] Breast cancer cases unselected for family history of breast cancer.⁴

[¶] OR estimated in a combined group of C65 CHEK2 variants.¹¹

^{*} OR estimated in a combined group of C25 CHEK2 variants.¹¹

^{††} OR estimated in a combined group of C15 CHEK2 variants.¹¹

na, not available.

Summary results from Breast Cancer Association Consortium studies of white Europeans (42 671 invasive breast cancer cases and 42 164 controls)

Table 2

Variant	Frequency* Controls	Frequency* Cases	OR (95% CI)	LRT p Value	OR [†] (95% CI)	LRT p Value [‡]
<i>PALB2</i> [§]						
c.1592delT (p.Leu531Cysfs)	0.00014	0.00082	4.52 (1.90 to 10.8)	7.1×10 ⁻⁵	3.44 (1.39 to 8.52)	0.003
c.2816T>G (p.Leu939Trp)	0.00342	0.00352	1.05 (0.83 to 1.32)	0.70	1.03 (0.80 to 1.32)	0.82
c.3113G>A (p.Trp1038*)	0.00019	0.00101	5.93 (2.77 to 12.7)	6.9×10 ⁻⁸	4.21 (1.84 to 9.60)	1.2×10 ⁻⁴
<i>CHEK2</i>						
c.349A>G (p.Arg117Gly)	0.00043	0.00103	2.26 (1.29 to 3.95)	0.003	2.03 (1.10 to 3.73)	0.020
c.538C>T (p.Arg180Cys)	0.00337	0.00370	1.33 (1.05 to 1.67)	0.016	1.34 (1.06 to 1.70)	0.015
c.715G>A (p.Glu239Lys)	0.00021	0.00035	1.70 (0.73 to 3.93)	0.210	1.47 (0.60 to 3.64)	0.40
c.1036C>T (p.Arg346Cys)	0.00005	0.00021	5.06 (1.09 to 23.5)	0.017	3.39 (0.68 to 16.9)	0.11
c.1312G>T (p.Asp438Tyr)	0.00078	0.00082	1.03 (0.62 to 1.71)	0.910	0.87 (0.49 to 1.52)	0.62
c.1343T>G (p.Ile448Ser) [‡]	0.00002	0	–	–	–	–
<i>ATM</i>						
c.727T>G (p.Val2424Gly)	0.00002	0.00028	11.6 (1.50 to 89.9)	0.0012	11.0 (1.42 to 85.7)	0.0019

* Proportion of subjects carrying the variant.

[†] Excluding women from five studies that selected all cases based on family history or bilateral disease and the subset of selected cases from other studies (based on 34 488 unselected cases and 34 059 controls).

[‡] *CHEK2* c.1343T>G (p.Ile448Ser) was only observed in one control and no cases of white European origin.

[§] *PALB2* c.3113G>A (p.Trp1038*) only observed in the UK, Australia, the USA and Canada. *PALB2* c.1592delT (p.Leu531Cysfs) only observed in Finland and Sweden.

LRT, likelihood ratio test; OR, OR for carriers of the variant versus common-allele homozygotes, adjusted for study and seven principal components.

Table 3

Summary results from the Prostate Cancer Association Group to Investigate Cancer Associated Alterations in the Genome studies for white European men* (22 301 prostate cancer cases and 22 320 controls)

Variant	Frequency [†] Controls	Frequency [†] Cases	OR (95% CI)	LRT p Value
<i>PALB2</i>				
c.1592delT (p.Leu531Cysfs)	0.00018	0.00031	2.06 (0.59 to 7.11)	0.24
c.2816T>G (p.Leu939Trp)	0.00354	0.00381	0.95 (0.69 to 1.29)	0.73
c.3113G>A (p.Trp1038*)	0.00045	0.00027	0.49 (0.18 to 1.36)	0.16
<i>CHEK2</i> [‡]				
c.349A>G (p.Arg117Gly)	0.00063	0.00081	1.46 (0.71 to 3.02)	0.30
c.538C>T (p.Arg180Cys)	0.00341	0.00296	1.02 (0.73 to 1.44)	0.90
c.715G>A (p.Glu239Lys)	0.00018	0.00027	1.47 (0.41 to 5.35)	0.55
c.1036C>T (p.Arg346Cys)	0.00018	0.00022	1.07 (0.28 to 4.07)	0.93
c.1312G>T (p.Asp438Tyr)	0.00049	0.00103	2.21 (1.06 to 4.63)	0.03
c.1343T>G (p.Ile448Ser)	0	0.00009	–	–
c.1343T>G (Africans [§])	0.019	0.057	3.03 (1.53 to 6.03)	0.001
ATM				
c.7271T>G (p.Val2424Gly)	0.00004	0.00027	4.37 (0.52 to 36.4)	0.17

* For white European men, unless otherwise indicated.

[†] Proportion of subjects carrying the variant.

[‡] *CHEK2* c.1343T>G (p.Ile448Ser) was the only *CHEK2* variant observed in African men and was identified in two cases and no controls of white European origin.

[§] Based on data from 623 and 569 African-American cases and controls, respectively.

LRT, likelihood ratio test; OR, OR for carriers of the variant versus common-allele homozygotes, adjusted for study and seven principal components.

Table 4

Summary results from the Ovarian Cancer Association Consortium studies for white European women (14 542 invasive ovarian cancer cases and 23 491 controls)

Variant	Frequency* Controls	Frequency* Cases	OR (95% CI)	LRT p Value
<i>PALB2</i>				
c.1592delT (p.Leu531Cysfs)	0.00004	0.00012	2.50 (0.21 to 29.1)	0.45
c.2816T>G (p.Leu939Trp)	0.00413	0.00399	0.96 (0.69 to 1.34)	0.81
c.3113G>A (p.Trp1038*)	0.00034	0.00031	1.34 (0.36 to 4.97)	0.66
<i>CHEK2</i>				
c.349A>G (p.Arg117Gly)	0.00038	0.00031	1.07 (0.32 to 3.60)	0.92
c.538C>T (p.Arg180Cys)	0.00128	0.00160	1.49 (0.83 to 2.67)	0.18
c.715G>A (p.Glu239Lys)	0.00021	0.00037	1.47 (0.42 to 5.22)	0.54
c.1036C>T (p.Arg346Cys) [‡]	0	0	–	–
c.1312G>T (p.Asp438Tyr)	0.00081	0.00074	0.92 (0.42 to 1.99)	0.83
c.1343T>G (p.Ile448Ser)	0.00009	0	–	–
<i>ATM</i>				
c.7271T>G (p.Val2424Gly)	0	0.00012	–	–

* Proportion of subjects carrying the variant.

[‡]c.1036C>T (p.Arg346Cys) was not observed in any sample.

LRT, likelihood ratio test; OR, OR for carriers of the variant versus common-allele homozygotes, adjusted for study and seven principal components.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript