



SHP-2 complex formation with the SHP-2 substrate-1 during C2C12 myogenesis

Citation

Kontaridis, Maria, Xiangdong Liu, Lei Zhang, and Anton M. Bennett. 2001. SHP-2 complex formation with the SHP-2 substrate-1 during C2C12 myogenesis. *Journal of Cell Science* 114: 2187-2198.

Published Version

<http://jcs.biologists.org/content/joces/114/11/2187.full.pdf>

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:30203536>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

SHP-2 complex formation with the SHP-2 substrate-1 during C2C12 myogenesis

Maria I. Kontaridis, Xiangdong Liu, Lei Zhang and Anton M. Bennett*

Department of Pharmacology, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06520-8066, USA

*Author for correspondence (e-mail: anton.bennett@yale.edu)

Accepted 13 March 2001

Journal of Cell Science 114, 2187-2198 (2001) © The Company of Biologists Ltd

SUMMARY

Myogenesis is a highly ordered process that involves the expression of muscle-specific genes, cell-cell recognition and multinucleated myotube formation. Although protein tyrosine kinases have figured prominently in myogenesis, the involvement of tyrosine phosphatases in this process is unknown. SHP-2 is an SH2 domain-containing tyrosine phosphatase, which positively regulates growth and differentiation. We show that in C2C12 myoblasts, SHP-2 becomes upregulated early on during myogenesis and associates with a 120 kDa tyrosyl-phosphorylated complex. We have identified that the 120 kDa complex consists of the SHP-2 substrate-1 (SHPS-1) and the Grb2-associated binder-1 (Gab-1). SHPS-1, but not Gab-1, undergoes tyrosyl phosphorylation and association with SHP-2 during myogenesis, the kinetics of which correlate with the

expression of MyoD. Either constitutive expression or inducible activation of MyoD in 10T $\frac{1}{2}$ fibroblasts promotes SHPS-1 tyrosyl phosphorylation and its association with SHP-2. It has been shown that p38 mitogen-activated protein kinase (MAPK) activity is required for the expression/activation of MyoD and MyoD-responsive genes. Inhibition of p38 MAPK by SB203580 in differentiating C2C12 myoblasts blocks MyoD expression, SHPS-1 tyrosyl phosphorylation and the association of SHPS-1 with SHP-2. These data suggest that SHPS-1/SHP-2 complex formation is an integral signaling component of skeletal muscle differentiation.

Key words: SHP-2, Tyrosine phosphatase, Myogenesis, Signalling, Myoblast, SHPS-1

INTRODUCTION

Skeletal muscle differentiation is a multi-step process that involves the coordinated expression of muscle-specific genes, cell cycle withdrawal and fusion of myoblasts into multinucleated myotubes. The early stages of myogenesis involve the activation of muscle-specific transcription factors of the E-box family, such as MyoD, myogenin, Myf-5 and MRF4 (Dias et al., 1994; Lassar et al., 1994; Olson and Klein, 1994; Weintraub, 1993). Although previous studies have defined extensively the mechanisms of myogenesis at the transcriptional level, the intracellular signaling mechanisms involved are still poorly understood. Work from several laboratories has shown that intracellular signaling cascades regulated by tyrosyl phosphorylation are critical components of myogenic progression. For example, the mitogen-activated protein kinases (MAPKs), specifically the extracellular-regulated protein kinases (Erks 1 and 2), have been implicated as critical for early myogenic progression (Bennett and Tonks, 1997; Coolican et al., 1997; Gredinger et al., 1998; Weyman and Wolfman, 1998; Wu et al., 2000). The p38 MAPK and Erk5/BMK function to activate MEF2C, another muscle-specific gene (Kato et al., 1997; Zetser et al., 1999; Zhao et al., 1999b). In addition, SAPK3/Erk6/p38 γ is induced during myogenesis and promotes myotube formation (Lechner et al., 1996). Other signaling molecules involved in myogenesis and regulated by protein tyrosine kinases (PTKs) include phosphatidylinositol 3'-kinase (PI-3K) (Coolican et al., 1997; Kaliman et al., 1996), its downstream serine/

threonine kinase target PKB/Akt (Fujio et al., 1999; Jiang et al., 1999), and the integrins (Sastry et al., 1996). Given that tyrosyl phosphorylation is a balance between the opposing actions of PTKs and protein tyrosine phosphatases (PTPs), it is likely that PTPs will also play a critical role in myogenic progression.

We have focused on determining whether SHP-2, an Src homology 2 (SH2) domain-containing PTP, which is expressed ubiquitously and is highly abundant in skeletal muscle, is involved in myogenesis (Adachi et al., 1992; Ahmad et al., 1993; Feng et al., 1993; Freeman et al., 1992; Vogel et al., 1993). Studies in *Drosophila* (Perkins et al., 1992), *Xenopus* (Tang et al., 1995), *C. elegans* (Gutch et al., 1998) and mice (Saxton et al., 1997) demonstrate that SHP-2 is required for development. The PTP activity of SHP-2 mediates positive signaling (growth and differentiation promoting) and functions either upstream of, and/or parallel to, Ras in the Raf/MEK/Erk pathway (Neel, 1993; Neel and Tonks, 1997; Tonks and Neel, 1996). In addition to signaling to the Erks, SHP-2 plays an important role in cell migration/motility, morphogenesis and cell spreading via integrin-dependent mechanisms (O'Reilly et al., 2000; Oh et al., 1999; Tsuda et al., 1998; Yu et al., 1998).

In most cases, if not all, SHP-2 propagates its downstream signal(s) by first interacting via its SH2 domains with tyrosyl-phosphorylated proteins. This event serves both to localize and to activate SHP-2, and ensures that the target of SHP-2 is appropriately activated in a coordinated and specific manner. To begin defining the role of SHP-2 in skeletal muscle

differentiation, we have attempted to identify proteins with which it may interact during myogenesis. Such interacting proteins will likely dictate how and when SHP-2 is activated during myogenesis. The SH2 domains of SHP-2 mediate its direct interaction with the activated platelet-derived growth factor (PDGF) receptor β (Lechleider et al., 1993) as well as multi-substrate adapter proteins such as IRS-1 (Kuhné et al., 1993), Gab-1 (Holado-Madruga et al., 1997) Gab-2 (Gu et al., 1998; Zhao et al., 1999a), FRS-2/SNT-1, SNT-2 (Kouhara et al., 1997; Xu et al., 1998) and DOS (Herbst et al., 1996; Raabe et al., 1996). SHP-2 also interacts with transmembrane glycoproteins, such as the signal regulatory proteins (SIRPs) (Kharitonov et al., 1997) (also known as SHP-2 substrate-1 (SHPS-1) (Fujioka et al., 1996), brain-immunoglobulin like molecule (BIT) (Sano et al., 1997), p84 (Comu et al., 1997) and macrophage fusion receptor (MFR) (Saginario et al., 1998) and the platelet/endothelial cell adhesion molecule (PECAM) (Jackson et al., 1997; Sagawa et al., 1997). The basis for how and when SHP-2 participates in cell-type and growth factor-specific signaling is likely regulated by the mutually exclusive complement of protein-protein interactions that SHP-2 forms. We have identified that SHP-2 interacts with Grb-2-associated binder-1 (Gab-1) and SHPS-1 in the C2C12 myoblast cell line. We show that SHPS-1/SHP-2 complex formation is an integral signaling event during C2C12 myoblast differentiation, suggesting a functional link between SHP-2 and myogenic progression.

MATERIALS AND METHODS

Cell lines

C2C12 myoblasts were purchased from the American Tissue Culture Collection (ATCC, Rockville, MD, USA) and cultured at 37°C and 5% CO₂ in growth medium (GM) containing DMEM (Gibco, Grand Island, NY, USA) supplemented with 10% FBS (Sigma, St Louis, MO, USA), 1% sodium pyruvate and 1% penicillin/streptomycin. C3H10T $\frac{1}{2}$ (10T $\frac{1}{2}$) and C3H10T $\frac{1}{2}$ -MyoD (10T $\frac{1}{2}$ -MyoD) fibroblasts were provided by Dr Andrew Lassar (Harvard Medical School, Boston, MA, USA) and maintained in DMEM supplemented by 10% FBS, 1% sodium pyruvate and 1% penicillin/streptomycin. Cells were induced to differentiate by transfer to differentiation medium (DM) containing DMEM supplemented with 0.1% FBS, 1% sodium pyruvate, 1% penicillin/streptomycin and 5 μ g/ml each of insulin (Sigma) and transferrin (Sigma). The L8 rat myogenic cell line was obtained from ATCC and cultured as described for C2C12 myoblasts. C2C12 myoblasts were seeded at 5 \times 10⁵ cells per 10 cm dish for 24 hours prior to the initiation of differentiation. C2C12 myoblasts at various stages during myogenesis were visualized with a Zeiss Axiovert 100 inverted microscope and photographed using a SPOT CCD camera (Diagnostic Instruments, Sterling Heights, MI, USA). The MyoD-ER 10T $\frac{1}{2}$ (clone 9-5) inducible cell line was a gift from Stephen Tapscott (Fred Hutchinson Cancer Center, Seattle, WA, USA) and Stephen Konieczny (Purdue University, Purdue, IN, USA) (Hollenberg et al., 1993). MyoD-ER 10T $\frac{1}{2}$ cell lines were maintained in DMEM plus 10% FCS (Hyclone) on gelatin-coated plates. Experiments with the MyoD-ER 10T $\frac{1}{2}$ inducible cell lines were performed on non-coated plates and MyoD activation was induced by the addition of 0.1 μ M 17 β -estradiol for 24 hours.

Immunoprecipitation and immunoblotting

SHPS-1 and Gab-1 polyclonal antibodies were purchased from Upstate Biotechnology (Lake Placid, NY, USA). SHPS-1 polyclonal

antibodies were also provided by Dr Benjamin Neel (Beth Israel Deaconess Hospital, Boston, MA, USA). Anti-SIRP- α 1/MFR (10C4) monoclonal antibodies were provided by Dr Agnes Vignery (Yale University School of Medicine, New Haven, CT, USA) (Saginario et al., 1995). SHP-2 polyclonal antibodies used for immunoprecipitation experiments were obtained from Gen-Shen Feng (Burnham Institute, La Jolla, CA, USA) or were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). SHP-2 immunoblots were performed using a monoclonal antibody purchased from Transduction Laboratories (Lexington, KY, USA). Dr Peter Houghton (St Jude Children's Research Hospital, Memphis, TN, USA) provided antibodies to MyoD. Anti-phosphotyrosine immunoblots were carried out using the 4G10 antibody purchased from Upstate Biotechnology and myosin heavy chain antibodies (MF20) were purchased from the Developmental Studies Hybridoma Bank (University of Iowa, Iowa City, IA, USA). Phospho-specific antibodies to Akt and MAPK were obtained from New England Biolabs (Beverly, MA, USA). Anti-Akt antibodies were purchased from Santa Cruz Biotechnology and anti-MAPK (C1) antibodies were a kind gift from John Blenis (Harvard Medical School, Boston, MA, USA). Insulin-like growth factor-I and II, and SB203580 were purchased from Calbiochem (La Jolla, CA, USA). All other chemicals were purchased from Sigma.

Immunoprecipitation and immunoblotting experiments were performed by lysing cells on ice in 1 ml of NP-40 lysis buffer containing 1.0% NP-40, 5 μ g/ml leupeptin, 5 μ g/ml aprotinin, 1 μ g/ml pepstatin A, 1 mM phenylmethylsulfonyl fluoride, 1 mM benzamidine and 0.2 mM sodium orthovanadate. Lysates were centrifuged at 14,000 r.p.m. for 15 minutes at 4°C and supernatants were precleared for 15 minutes with 2 μ l Pansorbin (Calbiochem-Novabiochem, La Jolla, CA, USA). Protein concentration was determined using the Bradford Assay (Pierce, Rockford, IL, USA). Approximately 0.5-1.0 mg of total cell lysates was used for each immunoprecipitation and approximately 50 μ g of total cell lysates were used for immunoblotting. Immunoprecipitation experiments were carried out overnight at 4°C. For immunoprecipitation experiments, 6 μ g of SHP-2 antibodies, 10 μ g SHPS-1 antibodies, 15 μ g of Gab-1 antibodies were used to immunoprecipitate the respective proteins. Immune complexes were collected on Protein A-sepharose beads, which were washed three times with lysis buffer containing 0.1 mM sodium orthovanadate, with a final wash in STE (150 mM NaCl, 20 mM Tris, pH 7.4, and 5 mM EDTA) plus 0.2 mM sodium orthovanadate. Immune complexes were resolved by SDS-PAGE and transferred onto Immobilon-P membranes (Millipore, Bedford, MA, USA). Immobilon-P membranes were washed three times for 10 minutes in TBST (20 mM Tris-HCl, pH 8.0, 150 mM NaCl and 0.05% Tween-20) and were blocked for at least 1 hour with 5% non-fat dry milk in TBST at 4°C. Immunoblots that were incubated with anti-phosphotyrosine 4G10 antibodies (pTyr) were blocked for at least 1 hour in 5% BSA (Boehringer-Mannheim, Indianapolis, IN, USA) in TBST. After blocking, membranes were washed three times for 10 minutes in TBST. Membranes were then incubated with primary antibodies (anti-SHP-2 at 1:1,000; anti-SHPS-1 at 1:5,000; anti-myosin heavy chain antibody at 1:12; anti-Gab-1 at 1:1,000; anti-MyoD at 1:1,000; anti-Akt at 1:1,000; and anti-myogenin at 1:12) diluted in 2.5% non-fat dry milk, for 2-4 hours at 4°C. Anti-phosphotyrosine 4G10 antibodies were diluted to 1:2,000 in 2.5% BSA/TBST and anti-phospho-MAPK and anti-phospho-Akt antibodies were diluted to 1:1,000 in 2.5% nonfat dry milk/TBST. Primary antibodies were incubated with membranes for either 3 hours at room temperature or overnight at 4°C. Following incubation with primary antibodies, membranes were washed three times for 10 minutes in TBST and primary antibodies were detected with either anti-rabbit or anti-mouse horseradish peroxidase-conjugated antibodies at a 1:5,000 dilution in TBST for 30 minutes. Membranes were washed and primary antibodies visualized using enhanced chemiluminescence detection (Amersham Pharmacia Biotech, Piscataway, NJ, USA).




Fig. 1. Tyrosyl phosphorylation and SHP-2 protein expression during myogenic differentiation in the murine C2C12 cell line. C2C12 myoblasts were cultured either in the presence of growth medium (GM) or in differentiation medium (DM) for the indicated times as described in Materials and Methods. (A) (Top) Total cell lysates were prepared, separated by SDS-PAGE and immunoblotted with 4G10 anti-phosphotyrosine antibodies (pTyr). The arrows to the right indicate either specific proteins or a molecular mass range in which total tyrosyl-phosphorylation changes are observed during differentiation. The molecular mass markers (Gibco-BRL) are depicted on the left. (Bottom) Representative photomicrographs of C2C12 myoblasts, fixed in methanol and visualized with Wright Giemsa Stain, at each stage of differentiation corresponding to the total tyrosyl-phosphorylation analysis in top panel. (B) C2C12 cell lysates (50 µg) prepared from A were immunoblotted with the indicated antibodies. The positions of molecular mass markers are shown.

RESULTS

Changes in protein tyrosyl phosphorylation and induction of SHP-2 protein expression during C2C12 myogenesis

To characterize the involvement of SHP-2 in myogenesis, we analyzed the expression of SHP-2 protein and total protein tyrosyl phosphorylation during differentiation of the C2C12 myoblast cell line. Upon mitogen withdrawal, C2C12 myoblasts exit the cell cycle and express muscle-specific genes. At this stage of myogenesis (12-24 hours), we observe an increase in tyrosyl phosphorylation of several proteins, notably a 180 kDa protein and proteins within the 100-130 kDa range (Fig. 1A). In contrast, a 35 and a 50 kDa protein undergo tyrosyl dephosphorylation (Fig. 1A). During the early periods of myogenesis (12-24 hours), the levels of SHP-2 protein expression induced are concomitant with the induction of MyoD and myogenin (Fig. 1B). SHP-2 protein expression remains elevated throughout the later stages of myogenesis (48-72 hours), as C2C12 myoblasts undergo terminal differentiation, express myosin heavy chain (Fig. 1B) and fuse to form multinucleated myotubes (Fig. 1A). These data

indicate that during C2C12 myogenesis, both tyrosyl phosphorylation and dephosphorylation events occur, and that the expression of SHP-2 is induced concomitant with that of early muscle-specific gene expression.

SHP-2 interacts with a 120 kDa tyrosyl-phosphorylated protein during C2C12 myogenesis

We hypothesized that if SHP-2 is involved in myogenic progression it would interact with tyrosyl-phosphorylated protein(s), via its SH2 domains, during differentiation. To test this possibility, SHP-2 was immunoprecipitated from undifferentiated and differentiating C2C12 myoblasts. The recovered SHP-2 immune complexes were resolved and immunoblotted with anti-phosphotyrosine antibodies. These experiments reveal that SHP-2 complexes with a 120 kDa (p120) tyrosyl-phosphorylated protein in undifferentiated myoblasts (Fig. 2A). In myoblasts undergoing differentiation, there is a transient increase in the mass of this p120 complex between 24 and 72 hours of differentiation (Fig. 2A). SHP-2 itself is tyrosyl-phosphorylated in undifferentiated and in differentiating myoblasts, but undergoes tyrosyl dephosphorylation as terminally differentiated multinucleated




Fig. 2. SHP-2 forms a complex with a p120 tyrosyl-phosphorylated protein during C2C12 myogenesis. C2C12 myoblasts cultured at the indicated times were subjected to (A) immunoprecipitation (IP) with either preimmune (PI) antisera or anti-SHP-2 antibodies. Immune complexes were resolved by SDS-PAGE, transferred to Immobilon and immunoblotted with anti-phosphotyrosine (pTyr) antibodies. The arrows to the right indicate the positions of the p120 and p180 tyrosyl-phosphorylated proteins that associate with SHP-2 during differentiation. The immunoblot was reprobred with anti-SHP-2 antibodies (bottom). (B) Cell lysates from C2C12 myoblasts cultured as indicated were immunoprecipitated with anti-SHP-2 antibodies and immunoblotted with pTyr antibodies. As a control, this immunoblot was reprobred with anti-SHP-2 antibodies (bottom). The positions of molecular mass markers are shown.

myotubes form (Fig. 2B). In addition, we observe the presence of a 180 kDa (p180) tyrosyl-phosphorylated protein in these SHP-2 immune complexes (Fig. 2A). This p180 tyrosyl-phosphorylated protein associates with SHP-2 early on during myogenesis, and then undergoes either tyrosyl dephosphorylation and/or dissociation with SHP-2 between 48 to 72 hours as multinucleated myotubes form (Fig. 2A). These observations demonstrate that during C2C12 myogenic differentiation, SHP-2 forms a complex with at least two tyrosyl-phosphorylated proteins of 120 kDa and 180 kDa.

SHP-2-associated p120 tyrosyl-phosphorylated protein in C2C12 myoblasts is SHPS-1

SHP-2 interacts with a family of tyrosyl-phosphorylated transmembrane glycoproteins known as the SIRPs (Fujioka et al., 1996; Kharitonov et al., 1997; Saginario et al., 1998). Due to the glycosylated characteristics of the SIRPs, this family of proteins migrates with a molecular mass ranging from 100 to 150 kDa (Kharitonov et al., 1997). To test the possibility that the p120 was a SIRP family member, lysates were prepared from undifferentiated C2C12 myoblasts, and were subjected to immunoprecipitation with anti-SHP-2 antibodies. The immune complexes were immunoblotted using an antibody generated to the conserved cytoplasmic domain of SHPS-1/SIRP- α 1, which detects all SIRP-related family members. This SHPS-1/SIRP- α 1 antibody cross-reacts in an immune-specific manner with a protein of identical apparent molecular mass as the tyrosyl-phosphorylated p120 protein found in SHP-2 immune complexes (Fig. 3A, left). SHP-2 was recovered in an immune-specific manner, as confirmed by reprobred these immunoblots with anti-SHP-2 antibodies (Fig. 3A, bottom). These data suggest that the p120 SHP-2-interacting protein is a SIRP family member.

The antibodies to SHPS-1 used in the previous experiment are directed to the conserved cytoplasmic domain of the SIRPs, raising the possibility that SHPS-1 may not constitute the specific SIRP isoform bound by SHP-2 in myoblasts (Fig. 3A,

left). To determine whether the SHPS-1 isoform binds to SHP-2 in myoblasts, we asked whether SHP-2 could be coimmunoprecipitated using a monoclonal antibody (10C4) that recognizes specifically the first hypervariable Ig (V1) domain of SHPS-1/MFR (Han et al., 2000). For these experiments, we utilized L8 rat myoblasts because this SHPS-1 monoclonal antibody (10C4) crossreacts only with rat SHPS-1 (data not shown). Lysates prepared from undifferentiated L8 myoblasts were subjected to immunoprecipitation with SHPS-1-specific antibodies and immune complexes were immunoblotted either with SHPS-1 antibodies that recognize all SIRP isoforms, or with anti-SHP-2 antibodies. Fig. 3A (right) demonstrates that SHP-2 is present in immune complexes derived from SHPS-1-specific antibodies. Moreover, these immune complexes were also recognized by antibodies directed to the conserved cytoplasmic domain of the SIRPs (Fig. 3A, right). These data indicate that SHP-2 associates specifically with SHPS-1 in C2C12 myoblasts. We also performed RT-PCR analysis for SIRP family members in C2C12 myoblasts, to determine which isoforms are expressed. A single PCR product was generated using degenerate V1 loop primers from cDNA prepared from terminally differentiated C2C12 cultures. Of the ten individual clones isolated, all contained the SHPS-1-specific V1 loop sequence (data not shown). These data, in conjunction with those presented in Fig. 3, indicate that SHPS-1 is the predominant, if not the sole, SIRP family member that forms a complex specifically with SHP-2 in C2C12 myoblasts.

Induction of SHPS-1 protein expression, tyrosyl phosphorylation and association with SHP-2 during C2C12 myogenesis

We next examined the protein expression and tyrosyl phosphorylation of SHPS-1 during myogenic differentiation. Early on during C2C12 myogenesis, the expression of SHPS-1 increases and plateaus by 48 hours into differentiation (Fig. 3B). In order to determine the kinetics of SHPS-1 tyrosyl phosphorylation, SHPS-1 was immunoprecipitated at various




Fig. 3. The SHP-2-interacting p120 tyrosyl-phosphorylated protein is SHPS-1. (A) (Left) Proliferating C2C12 myoblasts cultured in GM were lysed and immunoprecipitated with either pre-

immune (PI) or anti-SHP-2 antibodies. Immune complexes were resolved by SDS-PAGE and immunoblotted with anti-SHPS-1 (epitope directed to the conserved cytoplasmic domain of SIRP family members). (Right) C2C12 myoblast lysates prepared as described above were subjected to immunoprecipitation with anti-SHPS-1/MFR (10C4) antibodies, which recognize the V1 hypervariable extracellular region of SHPS-1. Immune complexes were immunoblotted with anti-SHPS-1 antibodies. (Bottom) The above immunoblots were reprobed with anti-SHP-2 antibodies. (B) Lysates were prepared from C2C12 myoblasts cultured in either GM or DM for the indicated times and were immunoblotted with anti-SHPS-1 antibodies. (C,D) Tyrosyl phosphorylation of SHPS-1 and association with SHP-2 during C2C12 myogenesis. Lysates prepared from C2C12 myoblasts cultured in either GM or DM for the indicated times were subjected to immunoprecipitation with either anti-SHPS-1 antibodies, followed by pTyrosine immunoblotting (C) or immunoprecipitation with anti-SHP-2 antibodies followed by anti-SHPS-1 immunoblotting (D). The lower panel in C represents an equal fraction of the lysates from the experiment above, which was immunoprecipitated for SHPS-1 and immunoblotted with anti-SHPS-1 antibodies as a control. The lower panel in D is a reprobe of the above experiment with anti-SHP-2 antibodies. The positions of molecular mass markers are shown.

times during myogenic differentiation. These immune complexes were immunoblotted with anti-phosphotyrosine antibodies. In addition, SHP-2 was immunoprecipitated from these lysates and immune complexes were immunoblotted with anti-SHPS-1 antibodies. We found that SHPS-1 is basally tyrosyl-phosphorylated, and forms a complex with SHP-2 in undifferentiated myoblasts (Fig. 3C,D, top). Upon initiation of myogenesis, SHPS-1 becomes tyrosyl-phosphorylated between 12 and 24 hours into differentiation (Fig. 3C, top), which correlates with an increase in the level of SHPS-1 in SHP-2 immune complexes (Fig. 3D, top). The level of tyrosyl phosphorylation on SHPS-1 peaks at 48 hours and declines slightly by 72 hours as myoblasts undergo fusion to form multinucleated myotubes (Fig. 3C, top). These data demonstrate that SHPS-1 undergoes an increase in protein expression, becomes tyrosyl-phosphorylated and associates with SHP-2 during C2C12 myogenesis.

SHP-2 interacting p120 tyrosyl-phosphorylated protein is a complex that also contains Gab-1

SHPS-1 is a glycoprotein containing 15 N-glycosylation sites. Upon treatment of SHP-2 immune complexes with endoglycosidase F, SHPS-1 can be deglycosylated, resulting in the generation of a core protein of an approximate molecular

mass 60 kDa (Fujioka et al., 1996; Kharitonov et al., 1997; Saginario et al., 1995). Our experiments indicate that although SHPS-1 could be deglycosylated to its 60 kDa core protein, we were unable to reduce completely the mass of the p120 complex to a lower core molecular mass (data not shown). These results suggest that other non-glycosylated tyrosyl-phosphorylated protein(s) might also constitute a component of the p120 complex that associates with SHP-2. Gab-1 is a tyrosyl-phosphorylated multi-substrate docking protein of 120 kDa that binds SH2 domain-containing proteins such as PI-3 kinase, Grb2 and SHP-2. Therefore, we tested whether the p120 complex also contains Gab-1. In Fig. 4A, we show that when lysates from undifferentiated C2C12 myoblasts are subjected to immunoprecipitation with anti-SHP-2 antibodies, the immune complexes contain a protein that is recognized in an immune-specific manner by antibodies to Gab-1. These data indicate that in undifferentiated C2C12 myoblasts, SHP-2 also forms a complex with Gab-1.

Tyrosyl dephosphorylation and constitutive association of Gab-1 with SHP-2 during C2C12 myogenesis

Analysis of Gab-1 during C2C12 differentiation reveals that Gab-1 protein expression remains constant throughout




Fig. 4. SHP-2 associates with Gab-1 in C2C12 myoblasts. (A) Lysates prepared from C2C12 myoblasts cultured in GM were immunoprecipitated with either preimmune (PI) antisera or anti-SHP-2 antibodies. Immune complexes were immunoblotted with anti-Gab-1 antibodies. The bottom panel is a reprobe with anti-SHP-2 antibodies from the above experiment.

(B,C,D) Tyrosyl dephosphorylation of Gab-1 and constitutive association with SHP-2 during C2C12 myogenesis. Lysates prepared from C2C12 myoblasts were cultured in either GM or DM for the indicated times. (B) Total cell lysates were immunoblotted with anti-Gab-1 antibodies. Lysates prepared from C2C12 myoblasts under the indicated conditions were also subjected to immunoprecipitation with either (C) anti-Gab-1 antibodies followed by pTyr immunoblotting or (D) immunoprecipitation with anti-SHP-2 antibodies followed by anti-Gab-1 immunoblotting. The bottom panel in C represents an equal fraction of the lysates from the experiment above that was immunoprecipitated for Gab-1 and immunoblotted with anti-Gab-1 antibodies as a control. Bottom panel in D is a reprobe with anti-SHP-2 antibodies of the above experiment. The positions of molecular mass markers are shown.


myogenic differentiation (Fig. 4B), in contrast to SHPS-1 (Fig. 3B). To determine the kinetics of Gab-1 tyrosyl phosphorylation during myogenesis, lysates prepared from C2C12 myoblasts at various stages during differentiation were subjected to immunoprecipitation with anti-Gab-1 antibodies. These immune complexes were immunoblotted with antibodies to detect for phosphotyrosine on Gab-1. In undifferentiated myoblasts, Gab-1 is heavily tyrosyl-phosphorylated (Fig. 4C, top). However, later on during myogenesis, Gab-1 undergoes a dramatic decrease in tyrosyl phosphorylation between 48 and 72 hours (Fig. 4C), when multinucleated myotubes form. Surprisingly, when SHP-2 was immunoprecipitated from these same cell lysates, and immune complexes were immunoblotted with antibodies to Gab-1, we found that Gab-1 associates with SHP-2 throughout the entire course of differentiation, even as Gab-1 undergoes tyrosyl dephosphorylation (Fig. 4D). Unlike SHPS-1 (Fig. 3C,D) complex formation between Gab-1 and SHP-2 does not correlate with myogenic progression. Nevertheless, these results show that Gab-1 is a component of the SHP-2-associated p120 complex in differentiating myoblasts.

MyoD induces the association of SHPS-1 and SHP-2 in $10T\frac{1}{2}$ fibroblasts

The data presented in Fig. 3 indicate that the induction of

SHPS-1 tyrosyl phosphorylation and association with SHP-2 correlates with the progression of myogenesis. In contrast, Gab-1 tyrosyl phosphorylation and association with SHP-2 does not (Fig. 4). We hypothesized that the initiation and subsequent progression of myogenesis in C2C12 myoblasts induces tyrosyl phosphorylation of SHPS-1 and subsequently its association with SHP-2. To test this, we compared SHP-2 complex formation with SHPS-1 in $10T\frac{1}{2}$ fibroblasts and $10T\frac{1}{2}$ fibroblasts that were converted to the myogenic lineage by stable expression of MyoD ($10T\frac{1}{2}$ -MyoD). In Fig. 5A, SHP-2 was immunoprecipitated from $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD fibroblasts; the resultant immune complexes were immunoblotted with anti-phosphotyrosine antibodies. In $10T\frac{1}{2}$ -MyoD fibroblasts, we observed a dramatic increase in the amount of both p120 and p180 tyrosyl-phosphorylated proteins associated with SHP-2, as well as an increase in the levels of tyrosyl-phosphorylated SHP-2, as compared to $10T\frac{1}{2}$ fibroblasts (Fig. 5A). Importantly, the levels of SHP-2 that were immunoprecipitated between $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD fibroblasts were equivalent (Fig. 5A, bottom panel), as was the total amount of SHPS-1 protein between these two cell lines (Fig. 5B). In addition, SHP-2 immune complexes also contained higher levels of SHPS-1 in $10T\frac{1}{2}$ -MyoD as compared to $10T\frac{1}{2}$ fibroblasts (Fig. 5C). This was consistent with the observation that SHPS-1 itself was hypertyrosyl-

Fig. 5. Induction of SHPS-1 tyrosyl phosphorylation by constitutive expression of MyoD in $10T\frac{1}{2}$ fibroblasts. (A) $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD fibroblasts cultured in GM were subjected to immunoprecipitation with either pre-immune (PI) antisera or anti-SHP-2 specific antibodies. Immune complexes and one-tenth of the total cell lysates (TCL) were resolved and immunoblotted with pTyr antibodies. The bottom panel represents a reprobe of the above immunoblot with anti-SHP-2 antibodies. (B) Equal amounts (50 μ g) of total cell lysates from $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD cells were immunoblotted with anti-SHPS-1 antibodies. (C) $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD cells were immunoprecipitated with either pre-immune (PI) or anti-SHP-2 antibodies and immunoblotted for SHPS-1. The bottom panel shows the immunoblot reprobated for SHP-2. (D) Lysates prepared from $10T\frac{1}{2}$ and $10T\frac{1}{2}$ -MyoD cells were immunoprecipitated with anti-SHPS-1 antibodies and immune complexes were immunoblotted with pTyr antibodies. (Bottom) SHPS-1 was also immunoprecipitated from a fraction of these same cell lysates and immune complexes were immunoblotted with anti-SHPS-1 antibodies as a control. The positions of molecular mass markers are shown.





phosphorylated in $10T\frac{1}{2}$ -MyoD fibroblasts as compared to $10T\frac{1}{2}$ fibroblasts (Fig. 5D). These data demonstrate that the initiation of the muscle differentiation program in fibroblasts, by the expression of MyoD, is sufficient to induce SHPS-1/SHP-2 complex formation.

To substantiate further our observation that C2C12 myogenesis is sufficient to promote SHPS-1 tyrosyl phosphorylation and association with SHP-2, we utilized a $10T\frac{1}{2}$ fibroblast cell line that stably expresses an estradiol-inducible MyoD chimera (MyoD-ER $10T\frac{1}{2}$). These MyoD-ER $10T\frac{1}{2}$ cells can be induced to activate MyoD upon treatment with estradiol. Fig. 6A shows that MyoD-ER $10T\frac{1}{2}$ cells treated with estradiol activate MyoD, as reflected by the induction of MHC expression (Fig. 6A). Activation of MyoD in these MyoD-ER $10T\frac{1}{2}$ cells results in an increase in SHPS-1, but not Gab-1, tyrosyl phosphorylation (Fig. 6B). In addition, an increase in the association of tyrosyl-phosphorylated SHPS-1 with SHP-2 was observed when MyoD-ER $10T\frac{1}{2}$ cells were induced to activate MyoD (Fig. 6C). Collectively, the data presented in Figs 5 and 6 demonstrate that SHPS-1 tyrosyl phosphorylation and its association with SHP-2 occurs as a direct response to the initiation of myogenesis.

p38 MAPK is required for SHPS-1 tyrosyl phosphorylation and association with SHP-2 during myogenesis

In culture, the p38 MAPK inhibitor, SB203580, blocks

muscle-specific gene expression and subsequently myogenesis (Kato et al., 1997; Zetser et al., 1999; Zhao et al., 1999b). If SHPS-1 tyrosyl phosphorylation and its association with SHP-2 occur as integral signaling events of myogenic progression, then inhibition of p38 MAPK should block SHPS-1/SHP-2 complex assembly in differentiating C2C12 myoblasts treated with the p38 MAPK inhibitor. Treatment of C2C12 myoblasts prior to the initiation of differentiation with SB203580 (30 μ M) blocks myogenesis completely, as indicated by the absence of MyoD induction and multinucleated myotubes (Fig. 7A,B). Despite the fact that MyoD induction and myotube formation are prevented in differentiating C2C12 myoblasts treated with SB203580, the early induction of both SHP-2 and SHPS-1 protein expression is not. This observation suggests that both SHP-2 and SHPS-1 are induced early on during myogenesis independently of the p38 MAPK pathway. When SHPS-1 is immunoprecipitated from C2C12 myoblasts treated with SB203580 the levels of SHPS-1 tyrosyl phosphorylation are inhibited, as compared to control differentiating C2C12 myoblasts (Fig. 7C). Significantly, SHP-2 fails to associate with SHPS-1 in differentiating myoblasts treated with SB203580 (Fig. 7D). Immunoprecipitation of SHPS-1 (Fig. 7C, bottom panel) and immunoblotting with SHPS-1 antibodies or reprobating of the SHP-2 immunoprecipitates with SHP-2 antibodies (Fig. 7D, bottom panel) confirmed that comparable levels of these proteins between control and SB203580-treated cultures were analyzed. Although a



residual level of SHPS-1 tyrosyl phosphorylation is observed at 24 hours in SB203580 treated cultures (Fig. 7C), SHP-2 is not complexed with SHPS-1 at this time point. A likely explanation for these results is that SB203580 inhibits tyrosyl

phosphorylation on SHPS-1 at a specific residue(s) that mediates complex formation with SHP-2. These experiments demonstrate that p38 MAPK promotes the association of SHPS-1 with SHP-2 in C2C12 myoblasts, and suggest




Fig. 8. Insulin, IGF-I and IGF-II fail to induce SHPS-1 tyrosyl phosphorylation. C2C12 myoblasts were either left unstimulated, or were stimulated with insulin, IGF-I or IGF-II following serum deprivation for 6 hours. Cell lysates prepared from C2C12 myoblasts following stimulation with the indicated growth factor were immunoblotted with anti-phospho MAPK and reprobred with anti-MAPK antibodies (top). These cell lysates were also immunoblotted with anti-phospho-Akt antibodies and reprobred with anti-Akt antibodies (middle). These immunoblots were also reprobred with anti-Akt antibodies. SHPS-1 immunoprecipitated from the cell lysates shown in the top panels were immunoblotted with either anti-pTyr or anti-SHPS-1 antibodies (bottom).

that SHPS-1/SHP-2 interactions are linked to the p38 MAPK/MyoD pathway in myoblasts.

Insulin and the insulin-like growth factors fail to induce SHPS-1 tyrosyl phosphorylation in C2C12 myoblasts

The insulin-like growth factors (IGFs) are potent mediators of muscle differentiation, and expression of the IGFs is regulated in an autocrine manner by muscle-specific genes (Florini et al., 1996; Florini et al., 1993; Kou and Rotwein, 1993; Rosen et al., 1993; Stewart and Rotwein, 1996). The IGFs are therefore attractive candidates for mediating SHPS-1 tyrosyl phosphorylation during C2C12 differentiation. To test this, serum-starved (6 hours) C2C12 myoblasts were stimulated with either insulin, IGF-I or IGF-II for 15 minutes. Insulin and the IGFs transduce their downstream signals via both the Ras/Raf/MAPK and the PI 3-K/Akt pathways (Coolican et al., 1997; Fujio et al., 1999; Kaliman et al., 1998; Pinset et al., 1997; Rommel et al., 1999). As expected, stimulation of C2C12 myoblasts with either insulin, IGF-I or IGF-II led to the activation of both MAPK and Akt, as indicated by the detection of Akt and MAPK phosphospecific immunoreactivity (Fig. 8, top and middle). Anti-phosphotyrosine immunoblotting of SHPS-1 immune complexes recovered from these same cells did not, however, result in any detectable SHPS-1 tyrosyl phosphorylation following insulin, IGF-I or IGF-II stimulation (Fig. 8, bottom). These data indicate that in myoblasts, SHPS-1 is not tyrosyl-phosphorylated in response to either insulin, IGF-I or IGF-II.

DISCUSSION

In this report, we present data that lays the foundation towards

elucidating the role of SHP-2 in muscle function. We show that SHP-2, an SH2 domain-containing tyrosine phosphatase, interacts with SHPS-1, Gab-1 and a p180 protein in C2C12 myoblasts. Our results further demonstrate that the interaction of SHP-2 with SHPS-1 is a direct consequence of the myogenic process in C2C12 myoblasts. Together, these data suggest a role for SHP-2 and/or SHPS-1 in skeletal muscle differentiation.

SHP-2 is highly abundant in skeletal muscle and may play an important role in muscle development and/or post-developmental muscle function. Our observation that the expression of SHP-2 is induced during the initial periods of differentiation, concomitant with that of MyoD and myogenin induction (Fig. 1), is consistent with data from Mei et al. (1996). This group showed that SHP-2 is induced in differentiating myoblasts prepared from both embryonic mouse limbs and in developing rat muscle *in vivo* (Mei et al., 1996). In addition to SHP-2 becoming upregulated, the levels of SHPS-1 protein expression and tyrosyl phosphorylation increase during C2C12 differentiation (Figs 3 and 7). This correlates with an increase in the level of SHPS-1 that associates with SHP-2. Originally identified as a hypertyrosyl-phosphorylated protein in fibroblasts overexpressing a catalytically inactive mutant of SHP-2, SHPS-1 has been shown to bind and serve as a substrate for SHP-2 (Fujioka et al., 1996; Timms et al., 1998). The extracellular domain of SHPS-1 contains three Ig domains and a cytoplasmic domain comprising four tyrosyl-phosphorylation sites. Two of these tyrosyl residues are in the sequence context that describes an immunoreceptor tyrosine-based inhibitory motif (ITIM; I/V-X-pY-X-X-L/V/I). These ITIMs bind the SH2 domains of both SHP-2, and its related hematopoietic counterpart SHP-1 (Fujioka et al., 1996; Kharitonov et al., 1997; Saginario et al., 1998). SHPS-1 becomes tyrosyl-phosphorylated in response to a variety of growth factors and by integrin-mediated cell adhesion (Fujioka et al., 1996; Kharitonov et al., 1997; Ochi et al., 1997; Oh et al., 1999; Takeda et al., 1998; Tsuda et al., 1998). SHPS-1 is suggested to be involved in positive (Fujioka et al., 1996) and negative (Inagaki et al., 2000; Kharitonov et al., 1997) growth factor signaling, integrin-mediated signaling (Fujioka et al., 1996; Oh et al., 1999; Tsuda et al., 1998), neurite outgrowth (Abosch and Lagenaur, 1993), inhibition of IgE-induced mast cell activation (Lienard et al., 1999) and macrophage multinucleation (Saginario et al., 1995; Saginario et al., 1998). The functional consequences of SHPS-1 induction in expression, and tyrosyl phosphorylation during differentiation of C2C12 myoblasts, remain to be determined.

In addition to the p180 tyrosyl-phosphorylated protein (Fig. 2), which remains to be identified, we found that SHP-2 forms a complex with Gab-1 in C2C12 myoblasts (Fig. 4). Gab-1 tyrosyl-phosphorylation levels do not change when myoblasts are induced to differentiate (Fig. 3), or if MyoD is activated in the MyoD-ER 10T $\frac{1}{2}$ cells (Fig. 6). However, as myotubes form, Gab-1 undergoes tyrosyl dephosphorylation (Fig. 3), concomitant with that of tyrosyl dephosphorylation of SHP-2 (Fig. 2). Interestingly, we found that the amount of Gab-1 in SHP-2 immune complexes remains unchanged even when Gab-1 becomes dephosphorylated in terminally differentiated multinucleated myotubes (Fig. 4). Three possibilities can be proposed to explain this result. First, it is possible that SHP-2

interacts with Gab-1 in a non-phosphotyrosyl-dependent manner. Second, in myoblasts, SHP-2 could interact indirectly with Gab-1 via a third component. Finally, SHP-2 may interact directly with Gab-1 (Lehr et al., 1999), while SHP-2 and/or other PTPs dephosphorylate Gab-1 at sites other than those with which SHP-2 interacts. Since the level of Gab-1 protein during myogenesis remains constant (Fig. 4), the reduced levels of tyrosyl phosphorylation on Gab-1 during myogenesis are likely catalyzed through the actions of a PTP. There is both genetic and biochemical evidence implicating Gab-1 as an SHP-2 substrate (Herbst et al., 1996; Nishida et al., 1999; Raabe et al., 1996). Gab-1 mediates branching tubulogenesis in epithelial cells (Maroun et al., 1999; Weidner et al., 1996), and may also be involved in similar processes such as branching and/or multinucleation in myoblasts.

The most striking observation of this study is the result that tyrosyl phosphorylation of SHPS-1 and subsequently SHPS-1/SHP-2 complex formation are integral processes of the differentiation of C2C12 myoblasts (Figs 5-7). Three lines of evidence support this conclusion: (1) SHPS-1/SHP-2 association correlates with the induction of MyoD during differentiation (Figs 1 and 3); (2) constitutive expression and inducible activation of MyoD in fibroblasts result in SHPS-1 tyrosyl phosphorylation and association with SHP-2 (Figs 5 and 6) and (3) inhibition of p38 MAPK activity by SB203580 in C2C12 myoblasts blocks SHPS-1 tyrosyl phosphorylation and its association with SHP-2 (Fig. 7). Previous studies from several groups have shown that p38 MAPK is required for skeletal muscle differentiation. (Cuenda and Cohen, 1999; Han et al., 1997; Zetser et al., 1999; Zhao et al., 1999b). Our results support these observations, and further demonstrate that the interaction between SHPS-1 and SHP-2 is also dependent upon p38 MAPK activity in C2C12 myoblasts. As SHP-2 is recruited to SHPS-1 during differentiation, it may subsequently participate in multiple signaling pathways during myogenesis. For example, SHP-2 has been implicated in regulating cell adhesion and cell motility, which are important components of the myogenic process. Indeed, if SHPS-1/SHP-2 complex formation functions downstream of MyoD, then it is reasonable to propose that this complex initiates signaling events which occur in concert with muscle-specific gene expression during myogenesis. It is clear that a substantial amount of work is needed in order to uncover the precise role(s) that SHP-2 plays in skeletal muscle differentiation.

How does SHPS-1 become tyrosyl-phosphorylated during C2C12 differentiation? Several candidate autocrine growth factors that promote myogenesis have been identified. These include primarily the insulin-like growth factors (IGFs) (Florini et al., 1996; Florini et al., 1993). Moreover, a positive relationship between muscle-specific gene expression and IGF production during myogenesis has been reported (Florini et al., 1991; Kou and Rotwein, 1993; Rosen et al., 1993; Stewart and Rotwein, 1996). Interestingly, neither insulin, IGF-I nor IGF-II induce tyrosyl phosphorylation of SHPS-1 in C2C12 myoblasts, despite the activation of both the Erk and PI-3K/Akt pathways (Fig. 8). These observations suggest that growth factors other than insulin/IGFs, and/or other signaling mechanisms regulate SHPS-1 tyrosyl phosphorylation in myoblasts. Consistent with the inability of the IGFs to stimulate SHPS-1 tyrosyl phosphorylation, IGF-I also fails to induce p38 MAPK activation in C2C12 myoblasts (Wu et al.,

2000). It will be important to establish how SHPS-1 becomes tyrosyl-phosphorylated during C2C12 myogenesis; this information may also yield insight into how the p38 MAPK pathway is activated.

In summary, we have identified that during C2C12 myogenesis, SHP-2 forms a complex with SHPS-1 and Gab-1. We have found that SHP-2/SHPS-1 interactions are induced as a specific consequence of myogenic progression in C2C12 myoblasts. Further investigation is now underway to determine the functional role of SHP-2, in the context of its interactions with both Gab-1 and SHPS-1, in myoblast growth and differentiation.

We thank Drs Agnes Vignery, Paul Lombroso, William Sessa and members of the Bennett laboratory for their critical reading of this manuscript. We also thank Drs Agnes Vignery, Benjamin G. Neel, Gen-Sheng Feng, Stephen Tapscoff and Peter Houghton for their kind gifts of reagents. This work was supported by a Hellman Family Foundation Award and NIH grant R01-AR65402 to A.M.B. and NIH training grants to M.I.K. (T32-GM07324) and X.L. (T32-NS07136).

REFERENCES

- Abosch, A. and Lagenaur, C. (1993). Sensitivity of neurite outgrowth to microfilament disruption varies with adhesion molecule substrate. *J. Neurobiol.* **24**, 344-355.
- Adachi, M., Sekiya, M., Miyachi, T., Matsuno, K., Hinoda, Y., Imai, K. and Yachi, A. (1992). Molecular cloning of a novel protein-tyrosine phosphatase SH-PTP3 with sequence similarity to the *src*-homology region 2. *FEBS Lett.* **314**, 335-339.
- Ahmad, S., Banville, D., Zhao, Z., Fischer, E. H. and Shen, S.-H. (1993). A widely expressed human protein tyrosine phosphatase contains src homology 2 domains. *Proc. Natl. Acad. Sci. USA* **90**, 2197-2201.
- Bennett, A. M. and Tonks, N. K. (1997). Regulation of distinct stages of skeletal muscle differentiation by mitogen-activated protein kinases. *Science* **278**, 1288-1291.
- Comu, S., Weng, W., Olinsky, S., Ishwad, P., Mi, Z., Hempel, J., Watkins, S., Lagenaur, C. F. and Narayanan, V. (1997). The murine P84 neural adhesion molecule is SHPS-1, a member of the phosphatase-binding protein family. *J. Neurosci.* **17**, 8702-8710.
- Coolican, S. A., Samuel, D. S., Ewton, D. Z., McWade, F. J. and Florini, J. R. (1997). The mitogenic and myogenic actions of insulin-like growth factors utilize distinct signaling pathways. *J. Biol. Chem.* **272**, 6653-6662.
- Cuenda, A. and Cohen, P. (1999). Stress-activated Protein Kinase-2/p38 and a Rapamycin-sensitive Pathway Are Required for C2C12 Myogenesis. *J. Biol. Chem.* **274**, 4341-4346.
- Dias, P., Dilling, M. and Houghton, P. (1994). The molecular basis of skeletal muscle differentiation. *Semin. Diagn. Pathol.* **11**, 3-14.
- Feng, G.-S., Hui, C.-C. and Pawson, T. (1993). SH2-containing phosphotyrosine phosphatase as a target of protein-tyrosine kinases. *Science* **259**, 1607-1611.
- Florini, J. R., Ewton, D. Z. and Coolican, S. A. (1996). Growth hormone and the insulin-like growth factor system in myogenesis. *Endocr. Rev.* **17**, 481-517.
- Florini, J. R., Ewton, D. Z., Magri, K. A. and Mangiacapra, F. J. (1993). IGFs and muscle differentiation. *Adv. Exp. Med. Biol.* **343**, 319-326.
- Florini, J. R., Magri, K. A., Ewton, D. Z., James, P. L., Grindstaff, K. and Rotwein, P. S. (1991). 'Spontaneous' differentiation of skeletal myoblasts is dependent upon autocrine secretion of insulin-like growth factor-II. *J. Biol. Chem.* **266**, 15917-15923.
- Freeman, R. M., Plutzky, J. and Neel, B. G. (1992). Identification of a human src homology 2-containing protein tyrosine-phosphatase: A putative homolog of *Drosophila* corkscrew. *Proc. Natl. Acad. Sci. USA* **89**, 11239-11243.
- Fujio, Y., Guo, K., Mano, T., Mitsuchi, Y., Testa, J. R. and Walsh, K. (1999). Cell cycle withdrawal promotes myogenic induction of Akt, a positive modulator of myocyte survival. *Mol. Cell. Biol.* **19**, 5073-5082.
- Fujioka, Y., Matozaki, T., Noguchi, T., Iwamatsu, A., Yamao, T., Takahashi, N., Tsuda, M., Takada, T. and Kasuga, M. (1996). A novel

- membrane glycoprotein, SHPS-1, that binds the SH2-domain-containing protein tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. *Mol. Cell. Biol.* **16**, 6887-6899.
- Gredinger, E., Gerber, A. N., Tamir, Y., Tapscott, S. J. and Bengal, E.** (1998). Mitogen-activated protein kinase pathway is involved in the differentiation of muscle cells. *J. Biol. Chem.* **273**, 10436-10444.
- Gu, H., Pratt, J. C., Burakoff, S. J. and Neel, B. G.** (1998). Cloning of p97/Gab2, the major SHP2-binding protein in hematopoietic cells, reveals a novel pathway for cytokine-induced gene activation. *Mol. Cell* **2**, 729-740.
- Gutch, M. J., Flint, A. J., Keller, J., Tonks, N. K. and Hengartner, M. O.** (1998). The *Caenorhabditis elegans* SH2 domain-containing protein tyrosine phosphatase PTP-2 participates in signal transduction during oogenesis and vulval development. *Genes Dev.* **12**, 571-585.
- Han, J., Jiang, Y., Li, Z., Kravchenko, V. and Ulevitch, R.** (1997). Activation of the transcription factor MEF2C by the MAP kinase p38 in inflammation. *Nature* **386**, 296-299.
- Han, X., Sterling, H., Chen, Y., Saginario, C., Brown, E. J., Frazier, W. A., Lindberg, F. P. and Vignery, A.** (2000). CD47, a ligand for the Macrophage Fusion Receptor, prevents macrophage multinucleation. *J. Biol. Chem.* **275**, 37984-37992.
- Herbst, R., Carroll, P. M., Allard, J. D., Schilling, J., Raabe, T. and Simon, M. A.** (1996). Daughter of sevenless is a substrate of the phosphotyrosine phosphatase corkscrew and functions during sevenless signaling. *Cell* **85**, 899-909.
- Holado-Madruga, M., Moscatello, D. K., Emlet, D. R., Dieterich, R. and Wong, A. J.** (1997). Grb2-associated binder-1 mediates phosphatidylinositol 3-kinase activation and the promotion of cell survival by nerve growth factor. *Proc. Natl. Acad. Sci. USA* **94**, 12419-12424.
- Hollenberg, S. M., Cheng, P. F. and Weintraub, H.** (1993). Use of a conditional MyoD transcription factor in studies of MyoD trans-activation and muscle determination. *Proc. Natl. Acad. Sci. USA* **90**, 8028-8032.
- Inagaki, K., Yamao, T., Noguchi, T., Matozaki, T., Fukunaga, K., Takada, T., Hosooka, T., Akira, S. and Kasuga, M.** (2000). SHPS-1 regulates integrin-mediated cytoskeletal reorganization and cell motility. *EMBO J.* **19**, 6721-6731.
- Jackson, D. E., Ward, C. M., Wang, R. and Newman, P. J.** (1997). The protein-tyrosine phosphatase SHP-2 binds platelet/endothelial cell adhesion molecule-1 (PECAM-1) and forms a distinct signaling complex during platelet aggregation. Evidence for a mechanistic link between PECAM-1 and integrin-mediated cellular signaling. *J. Biol. Chem.* **272**, 6986-6993.
- Jiang, B. H., Aoki, M., Zheng, J. Z., Li, J. and Vogt, P. K.** (1999). Myogenic signaling of phosphatidylinositol 3-kinase requires the serine-threonine kinase Akt/protein kinase B. *Proc. Natl. Acad. Sci. USA* **96**, 2077-2081.
- Kaliman, P., Canicio, J., Shepherd, P. R., Beeton, C. A., Testar, X., Palacin, M. and Zorzano, A.** (1998). Insulin-like growth factors require phosphatidylinositol 3-kinase to signal myogenesis: dominant negative p85 expression blocks differentiation of L6E9 muscle cells. *Mol. Endocrinol.* **12**, 66-77.
- Kaliman, P., Vinals, F., Testar, X., Palacin, M. and Zorzano, A.** (1996). Phosphatidylinositol 3-kinase inhibitors block differentiation of skeletal muscle cells. *J. Biol. Chem.* **271**, 19146-19151.
- Kato, Y., Kravchenko, V. V., Tapping, R. I., Han, J., Ulevitch, R. J. and Lee, J.-D.** (1997). BMK1/ERK5 regulates serum-induced early gene expression through transcription factor MEF2C. *EMBO J.* **16**, 7054-7066.
- Kharitonov, A., Chen, Z., Sures, I., Wang, H., Schilling, J. and Ullrich, A.** (1997). A family of proteins that inhibit signalling through tyrosine kinase receptors. *Nature* **386**, 181-186.
- Kou, K. and Rotwein, P.** (1993). Transcriptional activation of the insulin-like growth factor-II gene during myoblast differentiation. *Mol. Endocrinol.* **7**, 291-302.
- Kouhara, H., Hadari, Y. R., Spivak-Kroizman, T., Schilling, J., Bar-Sagi, D., Lax, I. and Schlessinger, J.** (1997). A lipid-anchored Grb2-binding protein that links FGF-receptor activation to the Ras/MAPK signaling pathway. *Cell* **89**, 693-702.
- Kuhné, M. R., Pawson, T., Lienhard, G. V. and Feng, G.-S.** (1993). The insulin receptor substrate 1 associates with the SH2-containing phosphotyrosine phosphatase Syp. *J. Biol. Chem.* **268**, 11479-11481.
- Lassar, A. B., Skapek, S. X. and Bennett, N.** (1994). Regulatory mechanisms that coordinate skeletal muscle differentiation and cell cycle withdrawal. *Curr. Opin. Cell Biol.* **6**, 788-794.
- Lechleider, R. J., Freeman, R. M. and Neel, B. G.** (1993). Tyrosyl-phosphorylation and growth factor receptor association of the human corkscrew homolog, SH-PTP2. *J. Biol. Chem.* **268**, 13434-23438.
- Lechner, C., Zahalka, M. A., Giot, J.-F., Moller, N. P. H. and Ullrich, A.** (1996). ERK6, a mitogen activated protein kinase involved in C2C12 myoblast differentiation. *Proc. Natl. Acad. Sci. USA* **93**, 4355-4359.
- Lehr, S., Kotzka, J., Herkner, A., Klein, E., Siethoff, C., Knebel, B., Noelle, V., Bruning, J. C., Klein, H. W., Meyer, H. E. et al.** (1999). Identification of tyrosine phosphorylation sites in human Gab-1 protein by EGF receptor kinase in vitro. *Biochemistry* **38**, 151-159.
- Lienard, H., Bruhns, P., Malbec, O., Fridman, W. H. and Daeron, M.** (1999). Signal regulatory proteins negatively regulate immunoreceptor-dependent cell activation. *J. Biol. Chem.* **274**, 32493-32499.
- Maroun, C. R., Hogado-Madruga, M., Royal, I., Naujokas, M. A., Fournier, T. M., Wong, A. J. and Park, M.** (1999). The Gab1 PH domain is required for localization of Gab1 at sites of cell-cell contact and epithelial morphogenesis downstream from the Met receptor tyrosine kinase. *Mol. Cell. Biol.* **19**, 1784-1799.
- Mei, L., Kachinsky, A. M., Seiden, J. E., Kuncl, R. W., Miller, J. B. and Haganir, R. L.** (1996). Differential Expression of PTP1D, a Protein Tyrosine Phosphatase with Two SH2 Domains, in Slow and Fast Skeletal Muscle Fibers. *Exp. Cell Res.* **224**, 379-390.
- Neel, B. G.** (1993). Structure and function of SH2-domain containing tyrosine phosphatases. *Semin. Cell Biol.* **4**, 419-432.
- Neel, B. G. and Tonks, N. K.** (1997). Protein tyrosine phosphatases in signal transduction. *Curr. Opin. Cell Biol.* **9**, 193-204.
- Nishida, K., Yoshida, Y., Itoh, M., Fukada, T., Ohtani, T., Shirogane, T., Atsumi, T., Takahashi-Tezuka, M., Ishihara, K., Hibbi, M. et al.** (1999). Gab-family adapter proteins act downstream of cytokine and growth factor receptors and T- and B-cell antigen receptors. *Blood* **93**, 1809-1816.
- O'Reilly, A., Pluskey, S., Shoelson, S. E. and Neel, B. G.** (2000). Activated mutants of SHP-2 preferentially induce elongation of xenopus animal caps. *Mol. Cell. Biol.* **20**, 299-311.
- Ochi, F., Matozaki, T., Noguchi, T., Fujioka, Y., Yamao, T., Takada, T., Tsuda, M., Takeda, H., Fukunaga, K., Okabayashi, Y. et al.** (1997). Epidermal growth factor stimulates the tyrosine phosphorylation of SHPS-1 and association of SHPS-1 with SHP-2, a SH2 domain-containing protein tyrosine phosphatase. *Biochem. Biophys. Res. Commun.* **239**, 483-487.
- Oh, E. S., Gu, H., Saxton, T. M., Timms, J. F., Hausdorff, S., Frevert, E. U., Kahn, B. B., Pawson, T., Neel, B. G. and Thomas, S. M.** (1999). Regulation of early events in integrin signaling by protein tyrosine phosphatase SHP-2. *Mol. Cell. Biol.* **19**, 3205-3215.
- Olson, E. and Klein, W.** (1994). bHLH factors in muscle development: dead lines and commitments, what to leave in and what to leave out. *Genes Dev.* **8**, 1-8.
- Perkins, L. A., Larsen, I. and Perrimon, N.** (1992). corkscrew encodes a putative protein tyrosine phosphatase that functions to transduce the terminal signal from the receptor tyrosine kinase torso. *Cell* **70**, 225-236.
- Pinsat, C., Garcia, A., Rouse, S., Dubois, C. and Montarras, D.** (1997). Wortmannin inhibits IGF-dependent differentiation in the mouse myogenic cell line C2. *CR Acad. Sci. III* **320**, 367-374.
- Raabe, T., Riesgo-Escovar, J., Liu, X., Bausenwein, B. S., Deak, P., Maroy, P. and Hafen, E.** (1996). DOS, a novel pleckstrin homology domain-containing protein required for signal transduction between sevenless and Ras1 in *Drosophila*. *Cell* **85**, 911-920.
- Rommel, C., Clarke, B. A., Zimmermann, S., Nunez, L., Rossman, R., Reid, K., Moelling, K., Yancopoulos, G. D. and Glass, D. J.** (1999). Differentiation stage-specific inhibition of the raf-MEK-ERK pathway by Akt [In Process Citation]. *Science* **286**, 1738-1741.
- Rosen, K. M., Wentworth, B. M., Rosenthal, N. and Villa-Komaroff, L.** (1993). Specific, temporally regulated expression of the insulin-like growth factor II gene during muscle cell differentiation. *Endocrinology* **133**, 474-481.
- Sagawa, K., Kimura, T., Swieter, M. and Siraganian, R. P.** (1997). The protein-tyrosine phosphatase SHP-2 associates with tyrosine-phosphorylated adhesion molecule PECAM-1. *J. Biol. Chem.* **272**, 31086-31091.
- Saginario, C., Qian, H. Y. and Vignery, A.** (1995). Identification of an inducible surface molecule specific to fusing macrophages. *Proc. Natl. Acad. Sci. USA* **92**, 12210-12214.
- Saginario, C., Sterling, H., Beckers, C., Kobayashi, R., Solimena, M., Ullu, E. and Vignery, A.** (1998). MFR, a putative receptor mediating the fusion of macrophages. *Mol. Cell. Biol.* **18**, 6213-6223.
- Sano, S., Ohnishi, H., Omori, A., Hasegawa, J. and Kubota, M.** (1997). BIT, an immune antigen receptor-like molecule in the brain. *FEBS Lett.* **411**, 327-334.
- Sastry, S. K., Lakonishok, M., Thomas, D. A., Muschler, J. and Horwitz, A. F.** (1996). Integrin alpha subunit ratios, cytoplasmic domains, and growth

- factor synergy regulate muscle proliferation and differentiation. *J. Cell Biol.* **133**, 169-184.
- Saxton, T. M., Henkemeyer, M., Gasca, S., Shen, R., Rossi, D. J., Shalaby, F., Feng, G. S. and Pawson, T.** (1997). Abnormal mesoderm patterning in mouse embryos mutant for the SH2 tyrosine phosphatase Shp-2. *EMBO J.* **16**, 2352-64.
- Stewart, C. E. and Rotwein, P.** (1996). Insulin-like growth factor-II is an autocrine survival factor for differentiating myoblasts. *J. Biol. Chem.* **271**, 11330-11338.
- Takeda, H., Matozaki, T., Fujioka, Y., Takada, T., Noguchi, T., Yamao, T., Tsuda, M., Ochi, F., Fukunaga, K., Narumiya, S. et al.** (1998). Lysophosphatidic acid-induced association of SHP-2 with SHPS-1: roles of RHO, FAK, and a SRC family kinase. *Oncogene* **16**, 3019-3027.
- Tang, T. L., Freeman, R. M., O'Reilly, A. M., Neel, B. G. and Sokol, S. Y.** (1995). The SH2-containing protein tyrosine phosphatase SH-PTP2 is required upstream of MAP kinase for early xenopus development. *Cell* **80**, 473-483.
- Timms, J. F., Carlberg, K., Gu, H., Chen, H., Kamatkar, S., Nadler, M. J. S., Rohrschneider, L. R. and Neel, B. G.** (1998). Identification of major binding proteins and substrates for the SH2-containing protein tyrosine phosphatase SHP-1 in macrophages. *Mol. Cell. Biol.* **18**, 3838-3850.
- Tonks, N. K. and Neel, B. G.** (1996). From form to function: Signaling by protein tyrosine phosphatases. *Cell* **87**, 365-368.
- Tsuda, M., Matozaki, T., Fukunaga, K., Fujioka, Y., Imamoto, A., Noguchi, T., Takada, T., Yamao, T., Takeda, H., Ochi, F. et al.** (1998). Integrin-mediated tyrosine phosphorylation of SHPS-1 and its association with SHP-2. Roles of Fak and Src family kinases. *J. Biol. Chem.* **273**, 13223-13229.
- Vogel, W., Lammers, R., Huang, J. and Ullrich, A.** (1993). Activation of a phosphotyrosine phosphatases by tyrosine phosphorylation. *Science* **259**, 1611-1614.
- Weidner, K. M., Di Cesare, S., Sachs, M., Brinkmann, V., Behrens, J. and Birchmeier, W.** (1996). Interaction between Gab1 and the c-Met receptor tyrosine kinase is responsible for epithelial morphogenesis. *Nature* **384**, 173-176.
- Weintraub, H.** (1993). The MyoD family and myogenesis: Redundancy, networks, and thresholds. *Cell* **75**, 1241-1244.
- Weyman, C. M. and Wolfman, A.** (1998). Mitogen-activated protein kinase (MEK) activity is required for inhibition of skeletal muscle differentiation by insulin-like growth factor 1 or fibroblast growth factor 2. *Endocrinology* **139**, 1794-1800.
- Wu, Z., Woodring, P. J., Bhakta, K. S., Tamura, K., Wen, F., Feramisco, J. R., Karin, M., Wang, J. Y. and Puri, P. L.** (2000). p38 and extracellular signal-regulated kinases regulate the myogenic program at multiple steps. *Mol. Cell. Biol.* **20**, 3951-3964.
- Xu, H., Lee, K. W. and Goldfarb, M.** (1998). Novel recognition motif on fibroblast growth factor receptor mediates direct association and activation of SNT adapter proteins. *J. Biol. Chem.* **273**, 17987-17990.
- Yu, D. H., Qu, C. K., Henegariu, O., Lu, X. and Feng, G. S.** (1998). Protein-tyrosine phosphatase Shp-2 regulates cell spreading, migration, and focal adhesion. *J. Biol. Chem.* **273**, 21125-21131.
- Zetser, A., Gredinger, E. and Bengal, E.** (1999). p38 Mitogen-activated Protein Kinase Pathway Promotes Skeletal Muscle Differentiation. Participation of the MEF2C transcription factor. *J. Biol. Chem.* **274**, 5193-5200.
- Zhao, C., Yu, D. H., Shen, R. and Feng, G. S.** (1999a). Gab2, a New Pleckstrin Homology Domain-containing Adapter Protein, Acts to Uncouple Signaling from ERK Kinase to Elk-1. *J. Biol. Chem.* **274**, 19649-19654.
- Zhao, M., New, L., Kravchenko, V. V., Kato, Y., Gram, H., di Padova, F., Olson, E. N., Ulevitch, R. J. and Han, J.** (1999b). Regulation of the MEF2 family of transcription factors by p38. *Mol. Cell. Biol.* **19**, 21-30.