



CYP24A1 variant modifies the association between use of oestrogen plus progestogen therapy and colorectal cancer risk

Citation

Garcia-Albeniz, X., A. Rudolph, C. Hutter, E. White, Y. Lin, S. A. Rosse, J. C. Figueiredo, et al. 2016. "CYP24A1 variant modifies the association between use of oestrogen plus progestogen therapy and colorectal cancer risk." *British Journal of Cancer* 114 (2): 221-229. doi:10.1038/bjc.2015.443. <http://dx.doi.org/10.1038/bjc.2015.443>.

Published Version

doi:10.1038/bjc.2015.443

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:30371175>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Keywords: GWAS; colorectal cancer; menopausal hormone therapy; polymorphisms; gene–environment interaction; cytochrome P450

CYP24A1 variant modifies the association between use of oestrogen plus progestogen therapy and colorectal cancer risk

Xabier Garcia-Albeniz^{1,2,36}, Anja Rudolph^{3,36}, Carolyn Hutter^{4,36}, Emily White^{5,6}, Yi Lin^{5,6}, Stephanie A Rosse^{5,6}, Jane C Figueiredo⁷, Tabitha A Harrison^{5,6}, Shuo Jiao^{5,6}, Hermann Brenner^{8,9,10}, Graham Casey⁷, Thomas J Hudson¹¹, Mark Thornquist^{5,6}, Loic Le Marchand¹², John Potter^{5,6}, Martha L Slattery¹³, Brent Zanke¹⁴, John A Baron¹⁵, Bette J Caan¹⁶, Stephen J Chanock¹⁷, Sonja I Berndt¹⁷, Deanna Stelling^{5,6}, Charles S Fuchs^{18,19,20}, Michael Hoffmeister⁸, Katja Butterbach⁸, Mengmeng Du^{5,6}, W James Gauderman²¹, Marc J Gunter²², Mathieu Lemire¹¹, Shuji Ogino^{18,19,20}, Jennifer Lin², Richard B Hayes²³, Robert W Haile²⁴, Robert E Schoen²⁵, Greg S Warnick^{5,6}, Mark A Jenkins²⁶, Stephen N Thibodeau²⁷, Fredrick R Schumacher⁷, Noralane M Lindor²⁸, Laurence N Kolonel¹², John L Hopper²⁶, Jian Gong^{5,6}, Daniela Seminara⁴, Bethann M Pflugeisen²⁹, Cornelia M Ulrich^{5,9,30}, Conghui Qu²⁹, David Duggan³¹, Michelle Cotterchio³², Peter T Campbell³³, Christopher S Carlson^{5,6}, Polly A Newcomb^{5,6}, Edward Giovannucci^{1,2}, Li Hsu^{5,6}, Andrew T Chan^{*2,34,37}, Ulrike Peters^{*5,6,37} and Jenny Chang-Claude^{*3,35,37}

Background: Menopausal hormone therapy (MHT) use has been consistently associated with a decreased risk of colorectal cancer (CRC) in women. Our aim was to use a genome-wide gene–environment interaction analysis to identify genetic modifiers of CRC risk associated with use of MHT.

Methods: We included 10835 postmenopausal women (5419 cases and 5416 controls) from 10 studies. We evaluated use of any MHT, oestrogen-only (E-only) and combined oestrogen–progestogen (E + P) hormone preparations. To test for multiplicative interactions, we applied the empirical Bayes (EB) test as well as the Wald test in conventional case–control logistic regression as primary tests. The Cocktail test was used as secondary test.

Results: The EB test identified a significant interaction between rs964293 at 20q13.2/CYP24A1 and E + P (interaction OR (95% CIs) = 0.61 (0.52–0.72), $P = 4.8 \times 10^{-9}$). The secondary analysis also identified this interaction (Cocktail test OR = 0.64 (0.52–0.78), $P = 1.2 \times 10^{-5}$ (alpha threshold = 3.1×10^{-4})). The ORs for association between E + P and CRC risk by rs964293 genotype were as follows: C/C, 0.96 (0.61–1.50); A/C, 0.61 (0.39–0.95) and A/A, 0.40 (0.22–0.73), respectively.

Conclusions: Our results indicate that rs964293 modifies the association between E + P and CRC risk. The variant is located near CYP24A1, which encodes an enzyme involved in vitamin D metabolism. This novel finding offers additional insight into downstream pathways of CRC etiopathogenesis.

*Correspondence: Professor AT Chan; E-mail: achan@partners.org or Professor U Peters; E-mail: upeters@fhcrc.org or Professor J Chang-Claude; E-mail: j.chang-claude@dkfz.de

³⁶Shared first authorship.

³⁷Shared last authorship.

Received 19 August 2015; revised 26 November 2015; accepted 30 November 2015; published online 14 January 2016

© 2016 Cancer Research UK. All rights reserved 0007–0920/16

The use of menopausal hormone therapy (MHT) has been consistently associated with a reduced risk of developing colorectal cancer (CRC; Calle *et al*, 1995; Fernandez *et al*, 1998; Grodstein *et al*, 1998, 1999; Anderson *et al*, 2004; Campbell *et al*, 2007; Johnson *et al*, 2009; Rennert *et al*, 2009; Green *et al*, 2012; Lin *et al*, 2012). According to a recently published meta-analysis, the relative risk (RR) of CRC was 0.74 (95% confidence interval (CI) 0.68–0.81) for ever use of oestrogen plus progestogen (E + P) therapy and 0.79 (95% CI 0.69–0.91) for ever use of oestrogen-only (E-only; Lin *et al*, 2012). Compared to placebo, menopausal women randomised to combined E + P hormone therapy in the Women's Health Initiative (WHI) also had a lower risk of CRC (Chlebowski *et al*, 2004), although their cancers tended to be of higher stage with poorer prognosis (Simon *et al*, 2012). However, randomisation to conjugated E-only in the WHI trial was not associated with risk of CRC (Anderson *et al*, 2004).

The underlying mechanisms of how MHT use influences colon carcinogenesis are largely unknown. Insight into potential biological pathways could be gained by investigating genetic modifiers of CRC risk associated with use of MHT. Furthermore, certain loci associated with susceptibility for CRC may only be evident in presence or absence of a specific environmental factor such as use of MHT. Thus, studies that specifically examine the association of MHT use with CRC risk in the context of varying genetic backgrounds are needed. Previous studies that have investigated gene–environment (G × E) interactions between single nucleotide polymorphisms (SNPs), MHT use and CRC risk have been largely based on candidate genes, which encompass only limited genetic variance (Lin *et al*, 2010, 2011; Rudolph *et al*, 2011; Slattery *et al*, 2011). A genome-wide scan to examine G × E interactions is a crucial next step to comprehensively examine additional variants and identify novel interactions with MHT. We therefore carried out a genome-wide association analysis to assess G × E interactions with use of MHT, using recently developed statistical methods and data from several studies comprising in total 5419 CRC cases and 5416 controls.

MATERIALS AND METHODS

Study participants. Our overall genome-wide association study (GWAS) design has been described previously (Hutter *et al*, 2012; Peters *et al*, 2013). In brief, this analysis is based on 10 studies (a case–control study from the Colon Cancer Family Registry (CCFR), and nine studies from the Genetics and Epidemiology of Colorectal Cancer Consortium (GECCO)). Study-specific details are described in Supplementary Methods. All cases were defined as colorectal adenocarcinoma, and confirmed by medical records, pathology reports or death certificate. All participants provided written informed consent, and studies were approved by their respective institutional review boards.

Harmonisation of environmental data. Information on basic demographics and environmental risk factors was collected by using in-person interviews and/or structured questionnaires, as detailed previously (The Women's Health Initiative Study Group, 1998; Colditz and Hankinson, 2005; Newcomb *et al*, 2007; Hoffmeister *et al*, 2009). Postmenopausal status was defined by using either (i) study-derived menopausal status, if available; (ii) self-reported menopausal status, if study derived was not available; or (iii) age ≥ 55 , if study derived and self-report were not available. MHT use was considered either as any MHT use, E-only use or E + P use at reference time. Non-users (of any MHT) at reference time were used as reference group. The reference time for nested case–control studies was time of enrolment into the cohort (VITAL, WHI and PLCO) or blood draw (NHS). For case–control studies, the reference time was at diagnosis and 2 years prior to

diagnosis for CCFR and DAL5. The harmonisation procedure is described in more detail in the Supplementary Methods.

Genotyping, quality assurance/quality control and imputation. All analyses were based on genotyped data generated from genome-wide association scans and imputation to HapMap II. Genotyped SNPs were excluded if they were triallelic, not assigned an rs number, or were reported or observed as not performing consistently across platforms. Furthermore, genotyped SNPs were excluded based on call rate ($< 98\%$), lack of Hardy-Weinberg equilibrium in controls (HWE, $P < 1 \times 10^{-4}$) and minor allele frequency (MAF $< 5\%$). Further details on DNA extraction, genotyping and quality assurance/quality control for each of the involved studies can be found in the Supplementary Methods and in Peters *et al* (2012).

All autosomal SNPs of all studies were imputed to the CEU population in HapMap II release 24, with the exception of OFCCR, which was imputed to HapMap II release 22. CCFR set 1 was imputed using IMPUTE (Marchini *et al*, 2007), OFCCR was imputed using BEAGLE (Browning and Browning, 2007), and all other studies were imputed using MACH (Li *et al*, 2010). Imputed data were merged with genotype data such that genotype data were used if a SNP had both types of data, unless there was a difference in terms of reference allele frequency (> 0.1) or position (> 100 base pairs), in which case imputed data were used. SNPs were restricted based on per study MAF $\geq 5\%$ of samples and per study imputation accuracy ($r^2 > 0.3$). After imputation and QC analyses, a total of about 2.7 M SNPs were used in the analysis. The inflation factor λ (a measurement of the overdispersion of the test statistics from the marginal association tests) obtained from the fixed-effect meta-analysis of GWAS scans was 1.029, 1.027 and 1.027 for the samples used for any MHT use, E + P and E-only analyses, respectively, indicating little evidence of population substructure (Quantile–quantile (Q–Q) plots in Supplementary Figure 1).

Statistical Methods. Models were adjusted for age at reference time, centre and the first three principal components from EIGENSTRAT (Price *et al*, 2006) to account for population substructure. Each directly genotyped SNP was coded as 0, 1 or 2 copies of the variant allele. For imputed SNPs, we used the expected number of copies of the variant allele (the 'dosage'), which has been shown to give unbiased test statistics (Jiao *et al*, 2011). SNPs are treated as continuous variables (i.e., assuming log-additive effects). Each study was analysed separately using logistic regression, and study-specific results were combined using fixed-effect meta-analysis to obtain summary odds ratios (ORs) and 95% CIs across studies. We calculated the heterogeneity P -values by Woolf's test (Woolf, 1955). Q–Q plots were assessed to determine whether the distribution of the P -values was consistent with the null distribution (except for the extreme tail). To test for multiplicative interactions between SNPs and environmental risk factors, we primarily used the empirical Bayes (EB) test (Mukherjee and Chatterjee, 2008; Mukherjee *et al*, 2008, 2010) given that this test can be more powerful than the conventional case–control logistic regression analysis while maintaining the desired type I error. It calculates the interaction log OR that corresponds to a weighted average of the case-only and case–control estimators. Thus, the method makes use of the greater precision of the case-only estimator by simultaneously reducing the chance of generating biased estimates due to violations of the assumption of gene–environment independence in controls (Mukherjee and Chatterjee, 2008). We modelled the SNP by environment (G × E) interaction by the product of the SNP and the dichotomised environmental variable. A two-sided P -value $< 5 \times 10^{-8}$ was considered genome-wide significant, yielding a genome-wide significance level of 0.05 (Risch and Merikangas, 1996; International HapMap Consortium, 2005; Wellcome Trust Case Control Consortium, 2007; Dudbridge and Gusnanto, 2008; Hoggart *et al*, 2008; Pe'er *et al*, 2008).

We secondarily used the multiplicative Cocktail method (Hsu *et al*, 2012), to evaluate how robust the findings were under the use of an alternative test to evaluate a multiplicative interaction (Supplementary Methods).

To estimate the effects of the environment variable stratified by genotype, we fit the following model: $\text{logit}(d) = b_0 + b_1e + c_1p_1 + c_2p_2 + \beta_1p_1e + \beta_2p_2e + \text{covariates}$, where p_1 and p_2 are the imputation posterior probabilities for genotype A/B, B/B. Then the stratified effects of environment variable can be estimated as \hat{b}_1 , $\hat{b}_1\hat{\beta}_1$ and $\hat{b}_1\hat{\beta}_2$ for genotypes A/A, A/B and B/B, respectively. The s.e. are estimated by using the standard formula for a linear combination of two parameters based on the covariance matrix of these parameters.

We estimated the CRC incidence rates associated with E + P use among individuals with each genotype of SNPs, which would provide more direct interpretation for $G \times E$ interaction effects on public health. We based the estimation of the incidence rate on the Surveillance, Epidemiology, and End Results (SEER) age-adjusted CRC incidence rate (denoted by 'I') 1992–2010 among the White population, which is 59.5 per 100 000 women per year. By using I, we estimated the reference incidence rate of CRC (denoted by ' I_{ref} ') using the population attributable risk (PAR), which is estimated by one minus the average of the inverse of estimated risk score $\exp(-X\beta)$ in cases (Bruzzi *et al*, 1985). Specifically, the formula for computing the PAR estimator is:

$$\text{PAR} = 1 - \frac{\sum_j Y_j \exp(X_j\beta)}{\sum_j Y_j},$$

where $Y_j = 1$ for case and 0 for control; $X_j = \text{covariates}$; $\beta =$ estimated regression coefficients in the logistic regression analysis. We can then estimate the reference incidence rate of CRC for $X = 0$ by $I_{\text{ref}} = (1 - \text{PAR}) \times I$ (Gail *et al*, 1989). On the basis of this reference incidence rate of CRC (i.e., I_{ref}), we further calculated the CRC incidence rate for each subgroup defined by genotypes of the SNP according to E + P use or non-use by multiplying the I_{ref} with each corresponding OR estimate. We calculated the 95% CIs using a resampling technique with 1000 weighted bootstraps. Since SEER incidence rates are based on a large number of individuals, the uncertainty of 'I' is negligible compared to the uncertainty from the PAR estimate, and hence was not considered in the calculation of 95% CIs.

Methods used for functional follow-up on promising loci are described in the Supplementary Methods.

RESULTS

Our study population comprised 10 835 menopausal women: 5419 cases and 5416 controls. For all 10 835 women information on use of any MHT was available and information on the type of MHT preparation was available for 9004 participants. At reference time, 3384 women used any MHT (31.2%), 1283 (11.8%) used E + P and 1606 (14.8%) used E-only (Supplementary Table 1). MHT use was inversely associated with the risk of CRC. Compared to non-users of any MHT at reference time, the OR for CRC was 0.70 (95% CI 0.62–0.79, $P = 1.9 \times 10^{-9}$, P for heterogeneity (phet) = 0.14) for women using any MHT preparation, 0.76 (95% CI 0.64–0.90, $P = 0.0015$, phet = 0.049) for women who used E + P and 0.71 (95% CI 0.61–0.84, $P = 7.1 \times 10^{-5}$, phet = 0.017) for women who used E-only.

The EB test identified a significant interaction of the variant rs964293 with E + P, with OR = 0.61 (95% CI 0.52–0.72, $P = 4.8 \times 10^{-9}$). This interaction showed borderline evidence of heterogeneity across studies (phet = 0.044; Figure 1A). The same variant (rs964293) showed an interaction with E + P use on CRC

risk in our secondary analysis, using the Cocktail test (OR = 0.64, 95% CI 0.52–0.78, $P = 1.2 \times 10^{-5}$, alpha threshold for significance in the group where the variant is assigned according to its rank = 3.1×10^{-4}). Table 1 summarises the results of the interaction analyses of rs964293 with MHT use. When testing for interaction between rs964293 and use of any MHT or E-only, no significant results were observed (Table 1).

Table 2 presents the association of the different strata of MHT by rs964293 with CRC risk. The OR of CRC for women taking E + P compared with women not using MHT is 0.96 (95% CI 0.61–1.50, $P = 0.84$), 0.61 (95% CI 0.39–0.95, $P = 0.03$) and 0.40 (95% CI 0.22–0.73, $P = 0.0026$) for women with C/C, A/C and A/A genotype of rs964293, respectively (Table 2). The per study ORs of E + P on CRC stratified by rs964293 genotype are shown in Figure 2A–C. No significant heterogeneity between study wise estimates was observed (phet = 0.3, 0.66 and 0.23, respectively).

The variant rs964293 is located in an intergenic region 28 kb upstream of *CYP24A1* on chromosome 20q13.2. Supplementary Figure 2 shows the interaction P -value (EB test) of rs964293 as well as SNPs surrounding rs964293 in region 20q13.2. The variant rs964293 was directly genotyped in five of the study sets included in the analysis and imputation accuracy was high in the remaining six study sets (r^2 ranging from 0.82 to 0.99). The MAF ranged from 0.34 to 0.38 in the 11 study sets.

The SNP rs964293 is in moderate LD ($r^2 = 0.61$ in 1000 Genomes pilot CEU) with a strong functional candidate, rs6023015. Our *in silico* functional analysis demonstrates that this SNP is located in a region with strong DNase hypersensitivity and histone methylation patterns consistent with enhancer activity. Supplementary Figure 3 shows how these two patterns of DNase hypersensitivity and histone methylation are stronger in some cancerous cell lines (including the CRC cell lines CACO2 and HCT-116) relative to non-cancerous cell lines. ChIP-seq experiments indicate that several transcription factors bind to this region (Supplementary Figure 4). In data from 169 tissue samples of the transverse colon available through the GTEx portal (Arldie *et al*, 2015), both variants, rs964293 and rs6023015, are expression quantitative trait loci (eQTL) for *CYP24A1* expression ($P = 0.037$ and 0.018, respectively). Expression of *CYP24A1* varied slightly stronger by genotypes of another SNP rs2256649 in LD with rs964293 ($r^2 = 0.60$ in 1000 Genomes pilot CEU), with $P = 0.0069$. Compared to homozygous carriers of the wild-type allele, expression of *CYP24A1* was higher in homozygous carriers of the minor allele for all three SNPs. Supplementary Figure 5 displays the distribution of the rank normalised gene expression of *CYP24A1* by genotypes of rs964293, rs6023015 and rs2256649.

We estimated absolute risks for CRC according to the use of E + P and genotypes of rs964293 and rs6023015 (Table 3). Compared to not-using E + P and carrying the wild-type CC genotype of rs964293, using E + P was associated with 30.6 fewer cases of CRC within a year among carriers of the AA genotype. Likewise, carrying the CC genotype of rs6023015 and using E + P was associated with 32.4 fewer cases of CRC within a year when compared to carrying the GG genotype of rs6023015 and not-using E + P.

Aside from results reported above, we did not observe any additional genome-wide significant $G \times E$ interactions with use of E + P and none for any MHT and E-only use.

DISCUSSION

In this study, including over 10 000 women, we evaluated genome-wide $G \times E$ interactions with the use of any MHT preparation as well as separately with use of E + P and of E-only. Employing the

powerful yet robust EB method, we found evidence of an interaction between the variant rs964293 at 20q13.2/*CYP24A1* and use of E + P.

Although most prior studies of G × E interactions in CRC have utilised a candidate gene approach (Slattery *et al*, 2010; Zhong *et al*, 2013), Figueiredo *et al* (2011), examined G × E interactions using a genome-wide scan and 14 environmental variables, including MHT use, and did not observe any genome-wide significant associations within the CCFR and OFCCR study. However, that study had limited power to detect any significant G × MHT interactions since it included only 572 highly selected CRC cases of menopausal women (age < 50 years or with a family history) and did not assess the possible differential interaction with use of MHT according to type of preparation.

Our finding of a genome-wide significant interaction between rs964293 and use of E + P has convincing biologic plausibility. This SNP is located in an intergenic region 28 kb upstream of *CYP24A1*. As it is not in strong LD with any coding variants, we hypothesised that the underlying causal variant(s) exerts its effect through a regulatory mechanism. We found that rs964293 is in moderate LD with rs6023015 ($r^2 > 0.61$ in CEU), which lies in a putative enhancer region for *CYP24A1*. The rs6023015 SNP was imputed in all 11 studies with high accuracy (mean $r^2 = 0.95$, range 0.87–1.00), and interaction and association of E + P across strata defined by number of minor alleles of rs6023015 paralleled that of rs964293 (Supplementary Table 4 and Figure 2D–F). Also the estimated ORs for interaction of rs6023015 with use of E + P were comparable to those found for rs964293 (Figure 1C and D), but *P*-values observed with the EB test were less extreme ($P = 2.8 \times 10^{-6}$). Both SNPs are eQTL for *CYP24A1* expression in 169 normal tissue samples of the transverse colon. *CYP24A1* expression was increased in homozygous carriers of the minor allele compared to homozygous carriers of the wild-type allele (Supplementary Figure 5). Details of the *in silico* functional analyses of rs6023015 are provided in the Supplementary Methods.

CYP24A1 codes for a protein that belongs to the cytochrome P450 family. These mitochondrial proteins are monooxygenases that catalyse several reactions involved in drug metabolism and

synthesis of cholesterol, steroids, sex hormones and other lipids. Specifically, *CYP24A1* plays a key role in the metabolism of the steroid hormone vitamin D by degrading its active form. *CYP24A1* is highly expressed in malignant colon tumours as compared with healthy colonic epithelium at both the mRNA (Bareis *et al*, 2001) and protein level (Matusiak and Benya, 2007), and some variants in *CYP24A1* have been associated with CRC risk in candidate gene studies (Dong *et al*, 2009). Vitamin D refers to a group of lipid soluble cholesterol-based compounds that, in contrast with other vitamins, can be synthesised endogenously. The active form of vitamin D exerts several functions relevant to regulation of tumour pathogenesis and progression, such as the activation of apoptotic pathways, anti-proliferative effects and angiogenesis inhibition (Deeb *et al*, 2007). Substantial experimental and epidemiological data support an inverse association with risk of CRC (Chan and Giovannucci, 2010). Moreover, there is evidence linking the vitamin D pathway and sex hormones in CRC aetiology. A secondary analysis of data from the WHI suggested that the association between low-dose vitamin D plus calcium supplementation and CRC risk is modified by MHT (E-only and E + P; Ding *et al*, 2008). Women randomised to receive calcium and 400 international units (IU) vitamin D3 supplementation in the placebo arms of the factorial oestrogen therapy trials were at suggestively decreased risk of developing CRC (HR = 0.71, 95% CI 0.46–1.09) compared to women randomised to receive calcium, vitamin D3 and concurrent MHT (HR = 1.30 (95% CI 0.83–2.03), *P* for interaction = 0.05; Ding and Giovannucci, 2009). Furthermore, the difference in the association of calcium and vitamin D supplementation with CRC between users and non-users of MHT appeared to be more evident for E + P than for E-only.

The results observed here suggest that *CYP24A1* may be regulated by sex hormones exposure and that the modifying effect observed with rs964293 may be caused by a disruption of the increased *CYP24A1* expression induced by sex hormones exposure. Alternatively, given that we found an interaction only with E + P exposure and not with E-only intake, *CYP24A1* may be a metabolising enzyme for progestogens but not oestrogen. Taken together, this evidence provides a strong rationale for additional




Figure 1. Forest plot for meta-analysis of the interaction between SNP and oestrogen plus progestogen use, using the empirical Bayes (A and C) and case-control logistic regression method (B and D) for rs964293 (A and B) and rs6023015 (C and D). DAL5 and PLCO studies are not plotted because they do not have information on the type of hormone compound.

Table 1. Result of the interaction tests for any MHT use, E + P use and E-only use and rs964293

Test	Any MHT preparation				E + P				E-only			
	OR (95% CI)	P	P ^a	phet	OR (95% CI)	P	P ^a	phet	OR (95% CI)	P	P ^a	phet
Empirical Bayes	0.91 (0.80–1.02)	0.11	5 × 10 ⁻⁸	0.11	0.61 (0.52–0.72)	4.8 × 10 ⁻⁹	5 × 10 ⁻⁸	0.043	1.01 (0.85–1.20)	0.90	5 × 10 ⁻⁸	0.27
Case-control	1.02 (0.89–1.17)	0.81	5 × 10 ⁻⁸	0.021	0.64 (0.52–0.78)	1.2 × 10 ⁻⁵	5 × 10 ⁻⁸	0.028	1.08 (0.90–1.30)	0.38	5 × 10 ⁻⁸	0.068
Cocktail ^b	1.02 (0.89–1.17)	0.81	0.005 (1)	0.021	0.64 (0.52–0.78)	1.2 × 10 ⁻⁵	3.1 × 10 ⁻⁴ (3)	0.028	1.08 (0.90–1.30)	0.38	Not in top 9 groups	0.068

Abbreviations: CI = confidence interval; E-only = oestrogen-only; E + P = oestrogen plus progestogen; MHT = menopausal hormone therapy; OR = odds ratio.
^aP^a: alpha threshold for significance, in brackets: group in weighted testing.
^brs964293 was selected based on correlation screen, the case-control test was used in the testing step.

Table 2. Associations with colorectal cancer risk stratified by E + P use and genotypes of rs964293

Strata of E + P use	rs964293 genotype										per allele OR ^a (95% CI) for rs964293	P
	CC			AC			AA			P		
	N Ca/Co ^b	OR ^a (95% CI)	P	N Ca/Co ^b	OR ^a (95% CI)	P	N Ca/Co ^b	OR ^a (95% CI)	P			
No	1219/1309	1 (Ref.)		1504/1455	1.04 (0.74–1.47)	0.82	477/472	1.30 (0.77–2.18)	0.33	1.15 (0.89–1.50)	0.28	
Yes	269/295	0.97 (0.62–1.52)	0.90	241/343	0.64 (0.36–1.12)	0.12	44/92	0.49 (0.22–1.06)	0.071	0.52 (0.26–1.01)	0.052	
OR ^a (95% CI) for E + P use		0.96 (0.61–1.50)	0.84		0.61 (0.39–0.95)	0.03		0.40 (0.22–0.73)	0.0026			

Abbreviations: CI = confidence interval; E + P = oestrogen plus progestogen; OR = odds ratio.
^aORs are adjusted for age, centre and the three principal components from EIGENSTRAT.
^bNumbers are expected frequencies based on imputed data.




Figure 2. Forest plot for meta-analysis of the marginal association of oestrogen plus progestogen with colorectal cancer risk in strata defined by zero, one or two minor alleles of rs964293 (A, B and C, respectively) and rs6023015 (D, E and F, respectively).

functional studies to determine the role of rs964293/rs6023015 in CYP24A1, particularly in relation to exposure to exogenous E + P in colorectal carcinogenesis.

Our study has several strengths. First, our large sample size facilitated the detection of genome-wide G × E interactions even accounting for the stringent threshold for significance necessary for control of type I error and the generally small magnitude of effect modification that can be reasonably expected. We followed a

‘whole set’ approach rather than a ‘discovery-validation’ strategy to maximise efficiency (Skol *et al*, 2006). Second, as previously described (Hutter *et al*, 2012), we have carefully harmonised data on a range of environmental variables, including MHT use across 10 studies. Third, two different methods (EB and Cocktail) to evaluate G × E interaction identified the same variant as having a significant G × E interaction, providing greater confidence that this association is not a false positive finding (although the EB test can

Table 3. Estimated incidence rates (95% CI) per 100 000 individuals stratified by genotypes of rs964293 and rs6023015 and use of E + P

rs964293 genotype			
Strata of E + P use	CC	AC	AA
No	59.81 (46.44–73.18)	62.27 (50.22–74.33)	77.58 (38.27–116.88)
Yes	58.06 (33.39–82.73)	38.09 (21.55–54.63)	29.18 (6.90–51.47)
rs6023015 genotype			
Strata of E + P use	GG	CG	CC
No	59.64 (44.80–74.48)	63.6 (50.46–76.74)	69.17 (41.50–96.84)
Yes	52.43 (28.98–75.89)	45.6 (25.97–65.22)	27.26 (9.58–44.93)

Abbreviations: CI = confidence interval; E + P = oestrogen plus progestogen. Use SEER 13 Regs Research Data, Nov 2014 Sub (1992–2012; Katrina/Rita Population Adjustment). The SEER age-adjusted colorectal cancer incidence rate based on 1992–2010 cases among White population is 59.5 per 100,000 women per year. Rates are age-adjusted to the 2000 US Standard Population.

be part of the Cocktail test itself, this was not the case for rs964293). The magnitude of the interaction between rs964293 and the use of E + P yielded by the traditional case–control logistic regression analysis was similar to the one found by the EB test (OR = 0.64 and 0.61, respectively; Figure 1 and Table 1), though it did not reach the threshold of genome-wide significance ($P = 1.2 \times 10^{-5}$). We observed some degree of heterogeneity among studies for the interaction of rs964293 with E + P ($\text{phet} = 0.044$). Figure 1 shows that the OR for interaction is ≤ 1 for all the studies but the VITAL, which constitutes about 2.5% of the total sample size. We do not consider this heterogeneity as a strong limitation to the results of the study.

The use of E + P for the purpose of chemoprevention is not routinely recommended for the general population due to concerns about the potential adverse consequences of long-term exposure. However, our results suggest the possibility that the benefit of E + P may be enhanced in women carrying genetic variants at rs964293 (Table 3). In conjunction with other strategies for risk-stratification and after its evaluation across other relevant clinical outcomes, this finding could be exploited to more specifically identify individuals for whom the benefits of chemoprevention may outweigh potential risks (Collins, 2015).

In summary, we have identified a *CYP24A1*-related variant as effect modifier of CRC risk associated with use of E + P using a genome-wide approach. This finding offers important insight into the role of E + P and its downstream pathways, including its potential interaction with vitamin D in the etiopathogenesis of CRC and supports the need for further studies to confirm the involvement of *CYP24A1* in modulating CRC risk.

ACKNOWLEDGEMENTS

DACHS: we thank all participants and cooperating clinicians, and Ute Handte-Daub, Renate Hettler-Jensen, Utz Benschaid, Muhabet Celik and Ursula Eilber for excellent technical assistance. GECCO: we would like to thank all those at the GECCO Coordinating Center for helping bring together the data and people that made this project possible. NHS: we would like to acknowledge Patrice Soule and Hardeep Ranu of the Dana Farber Harvard Cancer Center High-Throughput Polymorphism Core who assisted in the genotyping for NHS under the supervision of Dr Immaculata Devivo and Dr David Hunter, Qin (Carolyn) Guo and Lixue Zhu who assisted in programming for NHS. We would like to thank the participants and staff of the Nurses' Health Study for their valuable contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND,

OH, OK, OR, PA, RI, SC, TN, TX, VA, WA and WY. PLCO: we thank Drs Christine Berg and Philip Prorok, Division of Cancer Prevention, National Cancer Institute; the Screening Center investigators and staff or the Prostate, Lung, Colorectal, and Ovarian (PLCO) Cancer Screening Trial; Mr Tom Riley and staff, Information Management Services, Inc.; Ms Barbara O'Brien and staff, Westat, Inc.; and Drs Bill Kopp, Wen Shao and staff, SAIC-Frederick. Most importantly, we acknowledge the study participants for their contributions to making this study possible. PMH: we would like to thank the study participants and staff of the Hormones and Colon Cancer study. WHI: we thank the WHI investigators and staff for their dedication, and the study participants for making the programme possible. A full listing of WHI investigators can be found at: <https://cleo.whi.org/researchers/Documents%20Write%20a%20Paper/WHI%20Investigator%20Short%20List.pdf>. Funding details are as follows: GECCO: National Cancer Institute, National Institutes of Health, U.S. Department of Health and Human Services (U01 CA137088; R01 CA059045). Mengmeng Du is supported by grants R25 CA094880 and P30 CA008748 from NCI. CCFR: National Institutes of Health (RFA # CA-95-011) and through cooperative agreements with members of the Colon Cancer Family Registry and P.I.s. This genome-wide scan was supported by the National Cancer Institute, National Institutes of Health by U01 CA122839. The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centres in the CFRs, nor does mention of trade names, commercial products, or organisations imply endorsement by the US Government or the CFR. The following Colon CFR centres contributed data to this manuscript and were supported by National Institutes of Health: Australasian Colorectal Cancer Family Registry (U01 CA097735), Ontario Registry for Studies of Familial Colorectal Cancer (U01 CA074783) and Seattle Colorectal Cancer Family Registry (U01 CA074794). DACHS: German Research Council (Deutsche Forschungsgemeinschaft, BR 1704/6-1, BR 1704/6-3, BR 1704/6-4 and CH 117/1-1), and the German Federal Ministry of Education and Research (01KH0404 and 01ER0814). DALs: National Institutes of Health (R01 CA48998 to MLS); NHS is supported by the National Institutes of Health (R01 CA137178, P01 CA 087969 and P50 CA 127003). MEC: National Institutes of Health (R37 CA54281, P01 CA033619 and R01 CA63464). OFCCR: National Institutes of Health, through funding allocated to the Ontario Registry for Studies of Familial Colorectal Cancer (U01 CA074783); see CCFR section above. As subset of ARCTIC, OFCCR is supported by a GL2 grant from the Ontario Research Fund, the Canadian Institutes of Health Research, and the Cancer Risk Evaluation (CaRE) Program grant from the Canadian Cancer Society Research Institute. TJH and BZ are recipients of Senior Investigator Awards from the Ontario Institute for Cancer Research, through generous support from the Ontario Ministry

of Research and Innovation. PLCO: Intramural Research Program of the Division of Cancer Epidemiology and Genetics and supported by contracts from the Division of Cancer Prevention, National Cancer Institute, NIH, DHHS. In addition, a subset of control samples were genotyped as part of the Cancer Genetic Markers of Susceptibility (CGEMS) Prostate Cancer GWAS, Colon CGEMS pancreatic cancer scan (PanScan) and the Lung Cancer and Smoking study. The prostate and PanScan study data sets were accessed with appropriate approval through the dbGaP online resource (<http://cgems.cancer.gov/data/>) accession numbers phs000207v.1p1 and phs000206.v3.p2, respectively, and the lung data sets were accessed from the dbGaP website (<http://www.ncbi.nlm.nih.gov/gap>) through accession number phs000093 v2.p2. Funding for the Lung Cancer and Smoking study was provided by National Institutes of Health (NIH), Genes, Environment and Health Initiative (GEI) Z01 CP 010200, NIH U01 HG004446 and NIH GEI U01 HG 004438. For the lung study, the GENEVA Coordinating Center provided assistance with genotype cleaning and general study coordination, and the Johns Hopkins University Center for Inherited Disease Research conducted genotyping. PMH: National Institutes of Health (R01 CA076366 to PAN). VITAL: National Institutes of Health (K05 CA154337). WHI: The WHI programme is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts HHSN268201100046C, HHSN268201100001C, HHSN268201100002C, HHSN268201100003C, HHSN268201100004C and HHSN271201100004C. XG-A is a recipient of an ASISA Fellowship and SEOM (Sociedad Española de Oncología Médica) grant. ATC is a Damon Runyon Clinical Investigator and is also supported by NIDDK K24DK098311. WJG is supported by grant #HL115606. SO is supported by grant R35 CA197735.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

REFERENCES

- Anderson GL, Limacher M, Assaf AR, Bassford T, Beresford SA, Black H, Bonds D, Brunner R, Brzyski R, Caan B, Chlebowski R, Curb D, Gass M, Hays J, Heiss G, Hendrix S, Howard BV, Hsia J, Hubbell A, Jackson R, Johnson KC, Judd H, Kotchen JM, Kuller L, LaCroix AZ, Lane D, Langer RD, Lasser N, Lewis CE, Manson J, Margolis K, Ockene J, O'Sullivan MJ, Phillips L, Prentice RL, Ritenbaugh C, Robbins J, Rossouw JE, Sarto G, Stefanick ML, Van Horn L, Wactawski-Wende J, Wallace R, Wassertheil-Smoller S (2004) Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* **291**: 1701–1712.
- Ardlie KG, Deluca DS, Segre AV, Sullivan TJ, Young TR, Gelfand ET, Trowbridge CA, Maller JB, Tukiainen T, Lek M, Ward LD, Kheradpour P, Iriarte B, Meng Y, Palmer CD, Esko T, Winckler W, Hirschhorn JN, Kellis M, MacArthur DG, Getz G, Shabalina AA, Li G, Zhou Y-H, Nobel AB, Rusyn I, Wright FA, Lappalainen T, Ferreira PG, Ongen H, Rivas MA, Battle A, Mostafavi S, Monlong J, Sammeth M, Mele M, Reverter F, Goldmann JM, Koller D, Guigo R, McCarthy MI, Dermitzakis ET, Gamazon ER, Im HK, Konkashbaev A, Nicolae DL, Cox NJ, Flutre T, Wen X, Stephens M, Pritchard JK, Tu Z, Zhang B, Huang T, Long Q, Lin L, Yang J, Zhu J, Liu J, Brown A, Mesticelli B, Tidwell D, Lo E, Salvatore M, Shad S, Thomas JA, Lonsdale JT, Moser MT, Gillard BM, Karasik E, Ramsey K, Choi C, Foster BA, Syron J, Fleming J, Magazine H, Hasz R, Walters GD, Bridge JP, Miklos M, Sullivan S, Barker LK, Traino HM, Mosavel M, Siminoff LA, Valley DR, Rohrer DC, Jewell SD, Branton PA, Sobin LH, Barcus M, Qi L, McLean J, Hariharan P, Um KS, Wu S, Tabor D, Shive C, Smith AM, Buia SA, Undale AH, Robinson KL, Roche N, Valentino KM, Britton A, Burges R, Bradbury D, Hambright KW, Seleski J, Korzeniewski GE, Erickson K, Marcus Y, Tejada J, Taherian M, Lu C, Basile M, Mash DC, Volpi S, Struewing JP, Temple GF, Boyer J, Colantuoni D, Little R, Koester S, Carithers LJ, Moore HM, Guan P, Compton C, Sawyer SJ, Demchok JP, Vaught JB, Rabiner CA, Lockhart NC (2015) The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**: 648–660.
- Bareis P, Bises G, Bischof MG, Cross HS, Peterlik M (2001) 25-Hydroxy-vitamin D metabolism in human colon cancer cells during tumor progression. *Biochem Biophys Res Commun* **285**: 1012–1017.
- Browning SR, Browning BL (2007) Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am J Hum Genet* **81**: 1084–1097.
- Bruzzi P, Green SB, Byar DP, Brinton LA, Schairer C (1985) Estimating the population attributable risk for multiple risk factors using case-control data. *Am J Epidemiol* **122**: 904–914.
- Calle EE, Miracle-McMahill HL, Thun MJ, Heath Jr CW (1995) Estrogen replacement therapy and risk of fatal colon cancer in a prospective cohort of postmenopausal women. *J Natl Cancer Inst* **87**: 517–523.
- Campbell PT, Newcomb P, Gallinger S, Cotterchio M, McLaughlin JR (2007) Exogenous hormones and colorectal cancer risk in Canada: associations stratified by clinically defined familial risk of cancer. *Cancer Causes Control* **18**: 723–733.
- Chan AT, Giovannucci EL (2010) Primary prevention of colorectal cancer. *Gastroenterology* **138**: 2029–2043.e10.
- Chlebowski RT, Wactawski-Wende J, Ritenbaugh C, Hubbell FA, Ascensao J, Rodabough RJ, Rosenberg Ca, Taylor VM, Harris R, Chen C, Adams-Campbell LL, White E (2004) Estrogen plus progestin and colorectal cancer in postmenopausal women. *N Engl J Med* **350**: 991–1004.
- Colditz GA, Hankinson SE (2005) The Nurses' Health Study: lifestyle and health among women. *Nat Rev Cancer* **5**: 388–396.
- Collins FS (2015) Precision medicine: who benefits from aspirin to prevent colorectal cancer? Available at: <http://directorsblog.nih.gov/2015/03/24/precision-medicine-who-benefits-from-aspirin-to-prevent-colorectal-cancer/> (accessed on 07 August 2015).
- Deeb KK, Trump DL, Johnson CS (2007) Vitamin D signalling pathways in cancer: potential for anticancer therapeutics. *Nat Rev Cancer* **7**: 684–700.
- Ding EL, Giovannucci EL (2009) Reply to comment on: interaction of hormone replacement therapy with calcium and Vitamin D supplementation on colorectal cancer risk. *Int J Cancer* **124**: 1737–1738.
- Ding EL, Mehta S, Fawzi WW, Giovannucci EL (2008) Interaction of estrogen therapy with calcium and vitamin D supplementation on colorectal cancer risk: reanalysis of Women's Health Initiative randomized trial. *Int J Cancer* **122**: 1690–1694.
- Dong LM, Ulrich CM, Hsu L, Duggan DJ, Benitez DS, White E, Slattery ML, Farin FM, Makar KW, Carlson CS, Caan BJ, Potter JD, Peters U (2009) Vitamin D related genes, CYP24A1 and CYP27B1, and colon cancer risk. *Cancer Epidemiol Biomarkers Prev* **18**: 2540–2548.
- Dudbridge F, Gusnanto A (2008) Estimation of significance thresholds for genome-wide association scans. *Genet Epidemiol* **32**: 227–234.
- Fernandez E, La Vecchia C, Braga C, Talamini R, Negri E, Parazzini F, Franceschi S (1998) Hormone replacement therapy and risk of colon and rectal cancer. *Cancer Epidemiol Biomarkers Prev* **7**: 329–333.
- Figueiredo JC, Lewinger JP, Song C, Campbell PT, Conti DV, Edlund CK, Duggan DJ, Rangrej J, Lemire M, Hudson T, Zanke B, Cotterchio M, Gallinger S, Jenkins M, Hopper J, Haile R, Newcomb P, Potter J, Baron JA, Le Marchand L, Casey G (2011) Genotype-environment interactions in microsatellite stable/microsatellite instability-low colorectal cancer: results from a genome-wide association study. *Cancer Epidemiol Biomarkers Prev* **20**: 758–766.
- Gail MH, Brinton LA, Byar DP, Corle DK, Green SB, Schairer C, Mulvihill JJ (1989) Projecting individualized probabilities of developing breast cancer for white females who are being examined annually. *J Natl Cancer Inst* **81**: 1879–1886.
- Green J, Czanner G, Reeves G, Watson J, Wise L, Roddam A, Beral V (2012) Menopausal hormone therapy and risk of gastrointestinal cancer: Nested case-control study within a prospective cohort, and meta-analysis. *Int J Cancer* **130**: 2387–2396.
- Grodstein F, Martinez ME, Platz EA, Giovannucci E, Colditz GA, Kautzky M, Fuchs C, Stampfer MJ (1998) Postmenopausal hormone use and risk for colorectal cancer and adenoma. *Ann Intern Med* **128**: 705–712.

- Grodstein F, Newcomb PA, Stampfer MJ (1999) Postmenopausal hormone therapy and the risk of colorectal cancer: a review and meta-analysis. *Am J Med* **106**: 574–582.
- Hoffmeister M, Raum E, Krtischil A, Chang-Claude J, Brenner H (2009) No evidence for variation in colorectal cancer risk associated with different types of postmenopausal hormone therapy. *Clin Pharmacol Ther* **86**: 416–424.
- Hoggart CJ, Clark TG, De Iorio M, Whittaker JC, Balding DJ (2008) Genome-wide significance for dense SNP and resequencing data. *Genet Epidemiol* **32**: 179–185.
- Hsu L, Jiao S, Dai JY, Hutter C, Peters U, Kooperberg C (2012) Powerful cocktail methods for detecting genome-wide gene-environment interaction. *Genet Epidemiol* **36**: 183–194.
- Hutter CM, Chang-Claude J, Slattery ML, Pflugeisen BM, Lin Y, Duggan D, Nan H, Lemire M, Rangrej J, Figueiredo JC, Jiao S, Harrison TA, Liu Y, Chen LS, Stelling DL, Warnick GS, Hoffmeister M, Küry S, Fuchs CS, Giovannucci E, Hazra A, Kraft P, Hunter DJ, Gallinger S, Zanke BW, Brenner H, Frank B, Ma J, Ulrich CM, White E, Newcomb PA, Kooperberg C, LaCroix AZ, Prentice RL, Jackson RD, Schoen RE, Chanock SJ, Berndt SI, Hayes RB, Caan BJ, Potter JD, Hsu L, Béziau S, Chan AT, Hudson TJ, Peters U (2012) Characterization of gene-environment interactions for colorectal cancer susceptibility loci. *Cancer Res* **72**: 2036–2044.
- International HapMap Consortium (2005) A haplotype map of the human genome. *Nature* **437**: 1299–1320.
- Jiao S, Hsu L, Hutter CM, Peters U (2011) The use of imputed values in the meta-analysis of genome-wide association studies. *Genet Epidemiol* **35**: 597–605.
- Johnson JR, Lacey JV, Lazovich D, Geller Ma, Schairer C, Schatzkin A, Flood A (2009) Menopausal hormone therapy and risk of colorectal cancer. *Cancer Epidemiol Biomarkers Prev* **18**: 196–203.
- Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR (2010) MaCH: Using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genet Epidemiol* **34**: 816–834.
- Lin J, Zee RYL, Liu KY, Zhang SM, Lee IM, Manson JE, Giovannucci E, Buring JE, Cook NR (2010) Genetic variation in sex-steroid receptors and synthesizing enzymes and colorectal cancer risk in women. *Cancer Causes Control* **21**: 897–908.
- Lin JH, Manson JE, Kraft P, Cochrane BB, Gunter MJ, Chlebowski RT, Zhang SM (2011) Estrogen and progesterone-related gene variants and colorectal cancer risk in women. *BMC Med Genet* **12**: 78.
- Lin KJ, Cheung WY, Lai JYC, Giovannucci EL (2012) The effect of estrogen vs. combined estrogen-progesterone therapy on the risk of colorectal cancer. *Int J Cancer* **130**: 419–430.
- Marchini J, Howie B, Myers S, McVean G, Donnelly P (2007) A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet* **39**: 906–913.
- Matusiak D, Benya RV (2007) CYP27A1 and CYP24 expression as a function of malignant transformation in the colon. *J Histochem Cytochem* **55**: 1257–1264.
- Mukherjee B, Ahn J, Gruber SB, Ghosh M, Chatterjee N (2010) Case-control studies of gene-environment interaction: Bayesian design and analysis. *Biometrics* **66**: 934–948.
- Mukherjee B, Ahn J, Gruber SB, Rennert G, Moreno V, Chatterjee N (2008) Tests for gene-environment interaction from case-control data: a novel study of type I error, power and designs. *Genet Epidemiol* **32**: 615–626.
- Mukherjee B, Chatterjee N (2008) Exploiting gene-environment independence for analysis of case-control studies: an empirical Bayes-type shrinkage estimator to trade-off between bias and efficiency. *Biometrics* **64**: 685–694.
- Newcomb PA, Baron J, Cotterchio M, Gallinger S, Grove J, Haile R, Hall D, Hopper JL, Jass J, Le Marchand L, Limburg P, Lindor N, Potter JD, Templeton AS, Thibodeau S, Seminara D (2007) Colon cancer family registry: an international resource for studies of the genetic epidemiology of colon cancer. *Cancer Epidemiol Biomarkers Prev* **16**: 2331–2343.
- Pe'er I, Yelensky R, Altshuler D, Daly MJ (2008) Estimation of the multiple testing burden for genomewide association studies of nearly all common variants. *Genet Epidemiol* **32**: 381–385.
- Peters U, Hutter CM, Hsu L, Schumacher FR, Conti DV, Carlson CS, Edlund CK, Haile RW, Gallinger S, Zanke BW, Lemire M, Rangrej J, Vijayaraghavan R, Chan AT, Hazra A, Hunter DJ, Ma J, Fuchs CS, Giovannucci EL, Kraft P, Liu Y, Chen L, Jiao S, Makar KW, Taverna D, Gruber SB, Rennert G, Moreno V, Ulrich CM, Woods MO, Green RC, Parfrey PS, Prentice RL, Kooperberg C, Jackson RD, Lacroix AZ, Caan BJ, Hayes RB, Berndt SI, Chanock SJ, Schoen RE, Chang-Claude J, Hoffmeister M, Brenner H, Frank B, Béziau S, Küry S, Slattery ML, Hopper JL, Jenkins Ma, Le Marchand L, Lindor NM, Newcomb Pa, Seminara D, Hudson TJ, Duggan DJ, Potter JD, Casey G (2012) Meta-analysis of new genome-wide association studies of colorectal cancer risk. *Hum Genet* **131**: 217–234.
- Peters U, Jiao S, Schumacher FR, Hutter CM, Aragaki AK, Baron Ja, Berndt SI, Béziau S, Brenner H, Butterbach K, Caan BJ, Campbell PT, Carlson CS, Casey G, Chan AT, Chang-Claude J, Chanock SJ, Chen LS, Coetzee Ga, Coetzee SG, Conti DV, Curtis KR, Duggan D, Edwards T, Fuchs CS, Gallinger S, Giovannucci EL, Gogarten SM, Gruber SB, Haile RW, Harrison Ta, Hayes RB, Henderson BE, Hoffmeister M, Hopper JL, Hudson TJ, Hunter DJ, Jackson RD, Jee SH, Jenkins Ma, Jia WH, Kolonel LN, Kooperberg C, Küry S, Lacroix AZ, Laurie CC, Laurie Ca, Le Marchand L, Lemire M, Levine D, Lindor NM, Liu Y, Ma J, Makar KW, Matsuo K, Newcomb Pa, Potter JD, Prentice RL, Qu C, Rohan T, Rosse Sa, Schoen RE, Seminara D, Shrubsole M, Shu XO, Slattery ML, Taverna D, Thibodeau SN, Ulrich CM, White E, Xiang Y, Zanke BW, Zeng YX, Zhang B, Zheng W, Hsu L (2013) Identification of genetic susceptibility loci for colorectal tumors in a genome-wide meta-analysis. *Gastroenterology* **144**: 799–807.
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick Na, Reich D (2006) Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet* **38**: 904–909.
- Rennert G, Rennert HS, Pinchev M, Lavie O, Gruber SB (2009) Use of hormone replacement therapy and the risk of colorectal cancer. *J Clin Oncol* **27**: 4542–4547.
- Risch N, Merikangas K (1996) The future of genetic studies of complex human diseases. *Science* **273**: 1516–1517.
- Rudolph A, Sainz J, Hein R, Hoffmeister M, Frank B, Försti A, Brenner H, Hemminki K, Chang-Claude J (2011) Modification of menopausal hormone therapy-associated colorectal cancer risk by polymorphisms in sex steroid signaling, metabolism and transport related genes. *Endocr Relat Cancer* **18**: 371–384.
- Simon MS, Chlebowski RT, Wactawski-Wende J, Johnson KC, Muskovitz A, Kato I, Young A, Hubbell FA, Prentice RL (2012) Estrogen plus progestin and colorectal cancer incidence and mortality. *J Clin Oncol* **30**: 3983–3990.
- Skol AD, Scott LJ, Abecasis GR, Boehnke M (2006) Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet* **38**: 209–213.
- Slattery ML, Herrick J, Curtin K, Samowitz W, Wolff RK, Caan BJ, Duggan D, Potter JD, Peters U (2010) Increased risk of colon cancer associated with a genetic polymorphism of SMAD7. *Cancer Res* **70**: 1479–1485.
- Slattery ML, Lundgreen A, Herrick JS, Kadlubar S, Caan BJ, Potter JD, Wolff RK (2011) Variation in the CYP19A1 gene and risk of colon and rectal cancer. *Cancer Causes Control* **22**: 955–963.
- The Women's Health Initiative Study Group (1998) Design of the Women's Health Initiative clinical trial and observational study. The Women's Health Initiative Study Group. *Control Clin Trials* **19**: 61–109.
- Wellcome Trust Case Control Consortium (2007) Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* **447**: 661–678.
- Woolf B (1955) On estimating the relation between blood group and disease. *Ann Hum Genet* **19**: 251–253.
- Zhong R, Liu L, Zou L, Sheng W, Zhu B, Xiang H, Chen W, Chen J, Rui R, Zheng X, Yin J, Duan S, Yang B, Sun J, Lou J, Liu L, Xie D, Xu Y, Nie S, Miao X (2013) Genetic variations in the TGFβ signaling pathway, smoking and risk of colorectal cancer in a Chinese population. *Carcinogenesis* **34**: 936–942.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 4.0 Unported License.

¹Department of Epidemiology, Harvard T.H., Chan School of Public Health, Boston, MA 02115, USA; ²Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA 02115, USA; ³Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany; ⁴Division of Cancer Control and Population Sciences, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892, USA; ⁵Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109-1024, USA; ⁶Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY 10017, USA; ⁷Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA; ⁸Division of Clinical Epidemiology and Aging Research, German Cancer Research Center (DKFZ), 69120 Heidelberg, Germany; ⁹Division of Preventive Oncology, German Cancer Research Center (DKFZ) and National Center for Tumor Diseases (NCT), Heidelberg, Germany; ¹⁰German Cancer Consortium (DKTK), 69120 Heidelberg, Germany; ¹¹Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada; ¹²Epidemiology Program, University of Hawaii Cancer Center, Honolulu, HI 96813, USA; ¹³Department of Internal Medicine, University of Utah Health Sciences Center, Salt Lake City, UT 84132, USA; ¹⁴Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, ON K1Y 4E9, Canada; ¹⁵Division of Gastroenterology and Hepatology, University of North Carolina School of Medicine, Chapel Hill, NC 27599-7080, USA; ¹⁶Division of Research, Kaiser Permanente Medical Care Program, Oakland, CA 94612, USA; ¹⁷Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892-4608, USA; ¹⁸Department of Medical Oncology, Dana Farber Cancer Institute, Boston, MA 02215, USA; ¹⁹Department of Pathology, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02215, USA; ²⁰Department of Epidemiology, Harvard T.H., Chan School of Public Health, Boston, MA 02215, USA; ²¹Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA; ²²Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London W2 1PG, UK; ²³Division of Epidemiology, Department of Population Health, New York University School of Medicine, New York, NY 10016, USA; ²⁴Department of Medicine, Stanford University, Stanford, CA 94304, USA; ²⁵Department of Medicine and Epidemiology, University of Pittsburgh Medical Center, Pittsburgh, PA 15213-2582, USA; ²⁶Melbourne School of Population Health, The University of Melbourne, Melbourne, VIC 3010, Australia; ²⁷Departments of Laboratory Medicine, Pathology and Laboratory Genetics, Mayo Clinic, Scottsdale, AZ 85259, USA; ²⁸Department of Health Sciences Research, Mayo Clinic, Scottsdale, AZ 85259, USA; ²⁹Department of Cancer Prevention, Fred Hutchinson Cancer Research Center, Seattle, WA 98109-1024, USA; ³⁰Department of Population Sciences, Huntsman Cancer Institute, University of Utah, Salt Lake City, UT 84112, USA; ³¹Translational Genomics Research Institute (Tgen), Phoenix, AZ 85004, USA; ³²Prevention and Cancer Control, Cancer Care Ontario, Toronto, ON M5G 2L7, Canada; ³³Epidemiology Research Program, American Cancer Society, Atlanta, GA 30303, USA; ³⁴Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA and ³⁵University Cancer Center Hamburg (UCCH), University Medical Center Hamburg-Eppendorf, 20246 Hamburg, Germany

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)