



What Do We Really Know about the Role of Microorganisms in Iron Sulfide Mineral Formation?

Citation

Picard, Aude, Amy Gartman, and Peter R. Girguis. 2016. What Do We Really Know About the Role of Microorganisms in Iron Sulfide Mineral Formation? *Front. Earth Sci.* 4. doi:10.3389/feart.2016.00068.

Published Version

doi:10.3389/feart.2016.00068

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:30653602>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



What Do We Really Know about the Role of Microorganisms in Iron Sulfide Mineral Formation?

Aude Picard^{1*}, Amy Gartman^{2*} and Peter R. Girguis¹

¹ Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA, USA, ² U. S. Geological Survey, Santa Cruz, CA, USA

OPEN ACCESS

Edited by:

Dina M. Bower,
NASA Goddard SFC/University of
Maryland, USA

Reviewed by:

Jakob Zopfi,
University of Basel, Switzerland
William Patrick Gilhooly III,
Indiana University–Purdue University
Indianapolis, USA

*Correspondence:

Aude Picard
apicard@fas.harvard.edu;
Amy Gartman
agartman@usgs.gov

Specialty section:

This article was submitted to
Biogeoscience,
a section of the journal
Frontiers in Earth Science

Received: 09 November 2015

Accepted: 25 May 2016

Published: 15 June 2016

Citation:

Picard A, Gartman A and Girguis PR
(2016) What Do We Really Know
about the Role of Microorganisms in
Iron Sulfide Mineral Formation?
Front. Earth Sci. 4:68.
doi: 10.3389/feart.2016.00068

Iron sulfide mineralization in low-temperature systems is a result of biotic and abiotic processes, though the delineation between these two modes of formation is not always straightforward. Here we review the role of microorganisms in the precipitation of extracellular iron sulfide minerals. We summarize the evidence that links sulfur-metabolizing microorganisms and sulfide minerals in nature and we present a critical overview of laboratory-based studies of the nucleation and growth of iron sulfide minerals in microbial cultures. We discuss whether biologically derived minerals are distinguishable from abiotic minerals, possessing attributes that are uniquely diagnostic of biomineralization. These inquiries have revealed the need for additional thorough, mechanistic and high-resolution studies to understand microbially mediated formation of a variety of sulfide minerals across a range of natural environments.

Keywords: iron sulfide mineral, biomineralization, pyrite, mackinawite, greigite, microbial sulfate reduction, microbial sulfur disproportionation, sulfate-reducing prokaryotes

CO-OCCURENCE OF MICROORGANISMS AND SULFIDE MINERALS IN NATURE

Throughout Earth's history the burial of solid phases of Fe and S has controlled the redox state of Earth's surface environments (Berner, 1984). While iron is one of the most abundant elements on Earth, sulfur represents <1% of the Earth by mass (Allègre et al., 1995), although its importance to life and earth systems is greater than its abundance would suggest. Though the sulfur cycle was the first elemental cycle to be studied (Beijerinck, 1895), research on sulfur biogeochemistry is far from complete, and novel aspects of sulfur's transformations on Earth are still being discovered (Canfield et al., 2010). In Earth's biosphere, sulfur may be gaseous (e.g., sulfur dioxide), dissolved (e.g., sulfide, polysulfides, thiosulfate, sulfite, or sulfate) or solid (e.g., metal sulfides, elemental sulfur). Much of the interest in sulfur is due to its redox versatility—from sulfide (−2) to sulfate (+6), with numerous redox transformations possible in between (Zopfi et al., 2004). Microorganisms can take advantage of this diversity of oxidation states for energy conservation, which can be achieved by: (1) coupling the oxidation of organic compounds or dihydrogen to the reduction of oxidized organic and inorganic sulfur compounds (e.g., dimethyl sulfoxide, sulfate, elemental sulfur, and thiosulfate) (Widdel and Bak, 1992; Rabus et al., 2013); (2) disproportionating elemental sulfur, thiosulfate and sulfite (Bak and Pfennig, 1987; Thamdrup et al., 1993); and (3) oxidizing organosulfur compounds, hydrogen sulfide, sulfur, sulfite, and thiosulfate chemosynthetically with oxygen and nitrate during respiration, or by anoxygenic photosynthesis (Jørgensen and Nelson, 2004). Sulfate is the dominant sulfur species at 28 mM in

the modern oxic oceans, while reduced sulfur species, including hydrogen sulfide and organosulfur compounds, are often abundant where oxygen is low or absent. In these low-oxygen environments, sulfur and iron can be immobilized in the form of iron sulfide minerals, primarily the iron(II) monosulfide mackinawite (tetragonal FeS), the iron(II,III) sulfide greigite (Fe₃S₄), and the iron (II) disulfide pyrite (FeS₂) (Schoonen, 2004; Rickard, 2012a,c).

Ninety-seven percent of the sulfide produced on Earth is attributable to the activity of sulfate-reducing prokaryotes (SRP) in low-temperature environments (Trudinger et al., 1985; Rickard, 2012b), while the remaining three percent are produced at volcanoes and deep-sea hydrothermal vents (Elderfield and Schultz, 1996; Andres and Kasgnoc, 1998). SRP are present in an enormous diversity of environments including freshwater (Ramamoorthy et al., 2009; Sass et al., 2009), hypersaline (Foti et al., 2007), hydrothermal sediment (Jørgensen et al., 1992), polar (Ravenschlag et al., 2000; Karr et al., 2006), and oceanic crust (Robador et al., 2015) habitats. Although black sulfide precipitates in anoxic sediments are used as an indicator for the presence of SRP, their abundance isn't necessarily linked to large metal sulfide deposits. There are few environmental studies closely linking microbial sulfate-reducing activity and specific sulfide minerals, even though it has been 100 years since the first suggestion that SRP might be responsible for metal sulfide ore formation (Siebenthal, 1915). In the Cu-Zn Kidd Creek Mine, the presence of SRP corresponded to the enrichment of pyrite in the sediments (Fortin and Beveridge, 1997), while in the Zn-Pb Piquette mine, only sphalerite (ZnS) precipitated in a SRP-rich biofilm to the exclusion of other metal sulfides (Labrenz et al., 2000; Druschel et al., 2002; Labrenz and Banfield, 2004). In the Evander Au mine, ZnS rich biofilms were observed in association with pyrite framboids, and the minerals occurred inside organic matrices such as exopolysaccharides (Maclean et al., 2007, 2008). In sediments contaminated by mine tailing drainage, microbial cell surfaces were associated with FeS and NiS (Ferris et al., 1987). In hydrothermal vents, which result in the formation of submarine ores, both metabolically active SRP and dissimilatory sulfite reductase (*dsr*) genes have been found in the outer walls of chimneys and in seafloor massive sulfide deposits, where seawater sulfate is entrained (Nakagawa et al., 2004; Kormas et al., 2006; Frank et al., 2013; Kato et al., 2015). Nonetheless, the role of microorganisms in sulfide mineral formation at vents is considered quantitatively unimportant, due to the abundant abiotic sulfide provided by the vents.

In low-temperature sedimentary marine environments, where microbial sulfate reduction is the most important pathway for organic matter remineralization (Jørgensen, 1982), the biogeochemical cycles of iron and sulfur are intimately linked to carbon cycling (Bernier, 1982). The interactions between extracellular sulfide mineral precipitates and microorganisms are thus of great interest to understand the pathways of organic matter preservation in modern and ancient systems. For example, studies of marine sediments in Aarhus Bay in Denmark found that the highest sulfide production rates are attributed to members of the *Desulfobacteraceae* and that Fe₃S₄ is most abundant in the sediment just below these communities (Leloup

et al., 2009; Holmkvist et al., 2011). Although the focus of this mini review is on extracellular iron sulfide minerals, it should be noted that some magnetotactic bacteria (MTB) form intracellular iron sulfides ranging in size 35–120 nm, that are contained in bilayer membranes within an organelle called the magnetosome (Lefevre and Bazylinski, 2013). These sulfide-mineral-forming MTB are typically found below the oxic-anoxic interface both in marine and freshwater environments, where sulfide is found in the millimolar range (Farina et al., 1990; Bazylinski et al., 1995; Simmons et al., 2004; Lefevre et al., 2011). The intracellular mineral primarily formed is Fe₃S₄ (Posfai et al., 2013b), but tetragonal and cubic FeS have also been described as intermediate phases in the formation of greigite (Posfai et al., 1998).

Despite these discoveries in intracellular sulfide biomineralization, the role of microorganisms in the nucleation and growth of extracellular sulfide minerals is still ripe for exploration and explanation. In the next section, we examine the efforts made to understand the potential microbial mechanisms and pathways of sulfide mineral formation through laboratory experiments.

EXAMINING THE ROLE OF MICROORGANISMS IN EXTRACELLULAR IRON SULFIDE MINERAL FORMATION VIA LABORATORY STUDIES

SRP are the major producers of free sulfide in low-temperature environments, and thus have been the subject of most laboratory studies considering extracellular iron sulfide mineralization. Early studies demonstrated that Fe(II) salts react with sulfide produced in cultures of SRP, forming black precipitates, without further mineralogical characterization (Miller, 1950; Baas Becking and Moore, 1961; Freke and Tate, 1961; Temple and Le Roux, 1964). Later studies used energy-dispersive spectroscopy (EDS) in association with scanning electron microscopy (SEM) to identify the elemental ratios of iron sulfide precipitates (**Table 1**) (Fortin et al., 1994; Herbert et al., 1998; Benning et al., 1999; Donald and Southam, 1999; Li et al., 2004, 2006; Williams et al., 2005; Kim et al., 2015). However, determining mineralogy using EDS only can be contentious, as hydrogen sulfide can potentially adsorb onto iron sulfide aggregates, and thus alter the measured Fe:S ratio of the precipitates. X-ray diffraction (XRD), selected area electron diffraction (SAED), or X-ray absorption spectroscopy (XAS), revealed primarily the formation of mackinawite and greigite in microbial cultures to which Fe has been provided as soluble Fe(II) (**Table 1**) (Rickard, 1969; Herbert et al., 1998; Benning et al., 1999; Watson et al., 2000; Neal et al., 2001; Li et al., 2004; Williams et al., 2005; Gramp et al., 2009, 2010; Zhou et al., 2014). While greigite is typically found as rod-shaped and 100- to 300-nm platelet structures (Herbert et al., 1998; Gramp et al., 2010), mackinawite does not have a specific morphology, and both minerals appear (via XRD and SEM) to be disordered and poorly crystalline and tend to assemble in μm-sized aggregates (Fortin et al., 1994; Herbert et al., 1998; Benning et al., 1999; Watson et al., 2000; Kim et al., 2015). Additionally, using X-ray photoelectron

spectroscopy (XPS), pyrrhotite (Fe_{1-x}S , $0 < x < 0.2$) has been detected in association with mackinawite (Herbert et al., 1998) or at the surface of iron oxides, when these were used in culture medium as a source of iron (Neal et al., 2001). Notably, the formation of pyrite in microbial cultures has been reported only in two instances (Rickard, 1969; Donald and Southam, 1999).

Given the limited variety of isolates investigated in these studies (Table 1), it is unclear if there is a relationship between the type of SRP and the iron sulfide minerals that form in microbial cultures. The mineralogy of iron sulfide minerals seems to differ when Fe is provided as soluble Fe(II) vs. when it is provided as Fe(III) minerals (Rickard, 1969; Neal et al., 2001). Also, in microbial cultures, the crystallinity of iron sulfide minerals, in particular mackinawite and greigite, increases with incubation time (Rickard, 1969; Gramp et al., 2010; Zhou et al., 2014), and with increasing incubation temperature (Gramp et al., 2010). The size of mackinawite crystals decreases when initial pH deviates from the optimal pH for growth of the strain, and the largest crystals form when the sulfide production rates are highest (Zhou et al., 2014). However, to establish whether microorganisms have a more specific role in sulfide mineral nucleation and growth, it is essential to perform control experiments by reacting free sulfide with Fe in the culture medium in the absence of microorganisms. Among the few studies that did such controls, sulfide was not added at concentrations or rates similar to those occurring in microbial cultures (Rickard, 1969; Donald and Southam, 1999; Neal et al., 2001; Li et al., 2004, 2006; Gramp et al., 2010). Based on comparative XRD data for biotic and abiotic control experiments, Rickard (1969) concluded that no differences can be observed between synthetic and biotic mackinawite formed extracellularly in cultures of SRP. However, when pyrite and marcasite formed in cultures, he noted the association of an oil-like material, assumed to derive from the decomposition of cells, with the minerals. Gramp et al. (2010) found no difference in the mineralogy of abiotic vs. biotic metal sulfides, but reported differences in the morphology of the iron sulfide precipitates, suggesting a role of extracellular microbial substances. Finally, Donald and Southam (1999) proposed the involvement of organic sulfur, released during cell autolysis, in the formation of iron sulfide minerals.

Microbial cell walls are highly reactive and have a net negative charge at circumneutral pH, due to deprotonation of organic ligands contained in various polymers, such as peptidoglycan, lipopolysaccharides, teichoic acid, or murein (Beveridge, 1989). They can provide templates for metal binding and/or microenvironments for supersaturation of elements, leading to the nucleation and growth of minerals (Beveridge, 1989). Very few sulfide biomineralization studies have investigated cell-mineral associations (Table 1). Cells appear encrusted in sulfide minerals only in a few cases (Figure 1) (Fortin et al., 1994; Herbert et al., 1998; Donald and Southam, 1999; Watson et al., 2000; Williams et al., 2005). If cells interact with Fe while being metabolically inactive, then soluble Fe(II) could bind to reactive groups at the surface of cells (Figure 1A). Sulfide produced afterwards will then be able to react with iron, forming a crust of extracellular iron sulfides (Figure 1A). The apparent differences observed between Gram + and Gram -

bacteria (Figure 1A) could be explained by variations among cell envelope structure and composition. Alternatively, if Fe is at distance from metabolically active cells, or is provided when cells have already produced sulfide, then iron sulfide minerals could precipitate away from the cells (Figure 1B). The underlying causes of the variability of cell encrustation generally remain to be addressed.

UNRESOLVED QUESTIONS AND FUTURE DIRECTIONS

If the role of microbes in metal sulfide formation is to be assessed on a broader scale, and with greater depth, the field must use microbial isolates and enrichments whose phylogeny and physiology are well characterized. Future studies should use defined rather than organic-rich and strongly buffered cultivation media, in order to mimic more closely conditions of natural environments. Additionally, while SRP essentially occupy the entire range of temperatures at which life exists, sulfide biomineralization studies have been limited to ambient pressure and temperature conditions. For example, while most sulfide mineral formation in hydrothermal environments is likely abiotically derived, the extent to which thermophilic SRP are involved remains unknown. Studies of other mineral systems suggest additional possibilities for SRP mediated mineral formation (Burns et al., 2000; Braissant et al., 2007; Gallagher et al., 2012; Zammit et al., 2015). For example, the localized microenvironments which result from SRP metabolism cause increases in pH and production of exopolymeric substances (EPS), which have been demonstrated to affect the morphology and mineralogy of carbonate minerals (Braissant et al., 2007). EPS may locally bind metal ions to a number of functional groups such as carboxylic acids, sulfinic or sulfonic acids, or amino acids, and the presence of metal ions in solution may also affect EPS production, resulting in a feedback between soluble metals and EPS binding (Braissant et al., 2007). In addition, SRP are not the only microorganisms that may contribute to metal sulfide mineralization. Sulfur-disproportionating bacteria that can produce sulfide and sulfate from thiosulfate, sulfite (Bak and Pfennig, 1987), or elemental sulfur (Thamdrup et al., 1993), as well as sulfide oxidizing bacteria that can excrete polysulfides (Griesbeck et al., 2000; Prange et al., 2004), both produce reduced sulfur products that may contribute to sulfide mineralization. Iron sulfides, notably pyrite, can indeed form as a result of S(0)-disproportionation in microbial cultures and in marine sediments (Canfield et al., 1998; Finster et al., 1998; Zopf et al., 2008). Microbially produced organosulfides and even complexation of metals by microbially-produced organic matter may also be relevant, including at conditions where high temperatures preclude direct microbial involvement.

The correct application of methods can aid in resolving the spatial distribution of microorganisms and iron sulfide minerals. The use of bulk methods, like XRD, for mineralogical characterization will remove ambiguities inherent in elemental ratio analyses such as EDS. Besides XRD, Raman spectroscopy is a powerful method for the identification of sulfide minerals

TABLE 1 | Laboratory studies investigating the formation of extracellular Fe sulfide minerals in microbial cultures of sulfate-reducing bacteria.

Inoculum (origin)	Type of incubation Culture medium Carbon source Temperature	Metal provided Concentration Fe:S ratio	Analytical methods (mineralogy and morphology)	Composition, mineralogy and properties of precipitates Morphology of minerals	Interactions with cells	References
<i>Desulfovibrio desulfuricans</i> strain Canet 41	Batch Mineral medium Lactate 30°C	Hexagonal green rust 2 182 mM S in excess	XRD	At pH 6-8: initial precipitate = M At pH 6: 3 m. = G+M, 6 m. = G+M At pH 7: 3 m. = G+M, 6 m. = G At pH 8: 3 m. = M, 6 m. = M, 9 m. = pyrrhotite	ND	Rickard, 1969
<i>Desulfovibrio desulfuricans</i> strain Canet 41	Batch Mineral medium Lactate 30°C	Goethite NI S in excess	XRD	At pH 6-8: initial precipitate = M At pH 6: 3 m. = marcasite + pyrite, 6 m. = marcasite + pyrite At pH 7: 3 m. = marcasite + pyrite + M, 6 m. = marcasite + pyrite At pH 8: 3 m. = M, 6 m. = pyrite + marcasite Minerals are all fine-grained Pyrite is floating in "oily" phase	ND	Rickard, 1969
<i>Desulfohalobium</i> sp. strain DF-1 (metal refining plant)	Batch Freshwater medium Tryptone RT	Fe(II) sulfate 0.1–0.5 g/l S in excess	TEM-EDS	FeS aggregates with no specific morphology	Extensive mineralization of the surface of the cells	Fortin et al., 1994
Enrichment (aquifer)	Batch Postgate medium C Lactate + acetate RT	Fe(II) sulfate 18 μM + 1 Fe(0) nail S in excess	SEM-EDS XRD XPS	Disordered M (XRD) Potentially G, oxidized M and/or pyrrhotite (XPS) Platy particles (100–300 nm) aggregating in rosettes (1–2 μm)	No precipitates at the cell surface visible in SEM images	Herbert et al., 1998
Enrichment (sediment)	Batch Defined medium Lactate 25°C	Fe(II) sulfate 0.5 g/l (1.62 mM) S in excess	SEM-EDS TEM	Cubic and hexagonal crystals of FeS ₂ assigned as pyrite/marcasite (based on EDS data)	Precipitates at the cell surface	Donald and Southam, 1999
<i>Desulfovibrio desulfuricans</i> ATCC 29578	Batch Seawater medium Lactate 34°C	Fe(II) <1 mM S in excess	SEM-EDS XRD	Aggregates of M and G (15 w)	No precipitates at the cell surface visible in SEM images	Benning et al., 1999

(Continued)

TABLE 1 | Continued


Inoculum (origin)	Type of incubation Culture medium Carbon source Temperature	Metal provided Concentration Fe:S ratio	Analytical methods (mineralogy and morphology)	Composition, mineralogy and properties of precipitates Morphology of minerals	Interactions with cells	References
Enrichment ("natural source")	Chemostat Dilution rate 0.1 h-1 Postgate medium C Lactate 32°C	Fe "sufficient to maximize the production of iron sulfide"	TEM-SAED XANES EXAFS	Very small and irregular particles or clusters of particles of amorphous M and G (strongly magnetic)	100-nm thick layer of fine-grained precipitates at the cell surface	Watson et al., 2000
<i>Desulfovibrio desulfuricans</i> strains G-20 and Essex6	Batch Postgate medium C Lactate RT	Hematite NI S in excess	TEM Fluorescence microscopy XRD XPS	FeS precipitates at the surface of hematite assigned as pyrrhotite	ND Cells attached to hematite	Neal et al., 2001
<i>Desulfovibrio</i> sp. strain G-11	Batch PHELPS medium Lactate 30°C	Nontronite NAU-2 30 mM Fe (mineral) S in excess	TEM-EDS XRD	Textural and structural alterations of nontronite Aggregates of FeS (mineralogy not characterized), calcium carbonates	No precipitates at the cell surface visible in SEM images	Li et al., 2004
<i>Desulfovibrio vulgaris</i>	Sand-packed column Anoxic feed solution Lactate RT	Fe(II) sulfate 360 µM S in excess	TEM-EDS TEM-SAED Geophysical methods	M, 3-nm average crystal size forming 10–20 µm aggregates/clusters—phase shift in resistivity and acoustic wave deviation at the site of biomineralization	Sulfide-encrusted cells coating sand grains and cementing the pores of the column	Williams et al., 2005
<i>Desulfovibrio desulfuricans</i> strain G-20	Batch Caccavo's medium Lactate 30°C	Ferrihydrite, goethite, hematite, magnetite	TEM-EDS	FeS with no specific morphology	Mineral precipitates surrounding cells, EPS and Fe oxides	Li et al., 2006
Enrichments (acid mine drainage and compost)	Batch Baar's medium Lactate 22, 45, 60°C	Fe(II) chloride 8.95 mM Fe probably in excess (Sulfate provided as sulfate minerals)	XRD	At 60°C: (2 w. with schwertmannite, jarosite and gypsum): G	ND	Gramp et al., 2009

(Continued)

TABLE 1 | Continued

Inoculum (origin)	Type of incubation Culture medium Carbon source Temperature	Metal provided Concentration Fe:S ratio	Analytical methods (mineralogy and morphology)	Composition, mineralogy and properties of precipitates Morphology of minerals	Interactions with cells	References
Enrichments (acid mine drainage and compost)	Batch Baat's medium Lactate 22, 45 and 60°C	Fe(II) chloride Variable to adjust to the Fe:S ratios 1:0.5, 1:0.65, 1:1.3, 1:3.25, 1:10, 1:20 and 1:50	XRD SEM	M at 22°C M and G at 45 and 60°C Disordered morphology for both M and G Fine particles of M without specific pattern, rod-shaped aggregates and encrusted platelet structures of G	ND	Gramp et al., 2010
<i>Desulfovibrio vulgaris</i>	Batch ATCC 1249 medium Lactate 30°C	Fe(II) ammonium sulfate 3.5 mM S in excess	XRD	Crystalline M, G when sulfide is in excess	ND	Zhou et al., 2014
Biofilm containing <i>Desulfovibrio</i> and <i>Desulfuromonas</i> (acid mine drainage)	Batch Biofilm+fluids Glucose RT	Akaganeite and schwertmannite (contained in biofilm) NI	TEM-EDS	40–60 nm rod-shaped FeS particles	ND	Kim et al., 2015

Technical abbreviations: TEM, transmission electron microscopy; SEM, scanning electron microscopy; SAED, selective area electron diffraction; EDS, electron dispersive spectroscopy; XRD, X-ray diffraction; XPS, X-ray photon spectroscopy; XANES, X-ray absorption near-edge structure; EXAFS, extended X-ray absorption fine structure. Other abbreviations: M, mackinawite; G, greigite; RT, Room temperature; NI, not indicated; ND, not determined; w., weeks; m., months.



(White, 2009), as well as for the distinction between different various compositions of mackinawite and greigite (Bourdoiseau et al., 2008, 2011). Synchrotron X-ray methods, such as X-ray microscopy coupled to XAS may provide details about the microbe-mineral interface at the microscale that are overlooked in bulk studies (Templeton and Knowles, 2009). In the specific case of iron sulfide minerals, soft X-ray scanning transmission X-ray microscopy (STXM) can provide the distribution of C, Fe, and S (Behrens et al., 2012; Cosmidis and Benzerara, 2014). Using soft or hard X-rays, X-ray absorption near-edge structure (XANES), also called near-edge x-ray absorption fine structure (NEXAFS), spectroscopy can resolve the characterization of major biomolecules, such as lipids, polysaccharides and proteins, and the oxidation state of Fe and S and the associated mineralogy at the sub-micron scale, respectively (Templeton and Knowles, 2009). As newly precipitated iron sulfide particles are generally in the nm-size range, transmission electron microscopy (TEM) imaging and associated techniques, such as electron energy loss spectroscopy (EELS) and SAED, are essential to characterize the morphology and crystal structure of sulfide mineral crystals at the sub-micron scale (Posfai et al., 2013a). While bulk sulfur isotopic analyses provide the best line of evidence regarding the provenance of sulfide in sulfide minerals,

e.g., chemical sulfide vs. sulfide from sulfate reduction or from sulfur disproportionation (Chambers and Trudinger, 1979; Zopfi et al., 2008), considerable variations in isotopic compositions between—and even within—individual pyrite grains can occur (Kohn et al., 1998). In future studies, more spatially precise work using SIMS and nanoSIMS may provide critical information at the sub-micron scale to understand small spatial scale isotope fractionation and its possible causes (Fike et al., 2008; Orphan and House, 2009; Wacey et al., 2010). In highly resolved elemental analyses, co-localization of elements (especially C, N, Fe, S) must be considered. For example, nano-scale chemical mapping of organic carbon and nitrogen has detected relict pyrite framboids likely formed in microbial biofilms in larger pyrite grains (Wacey et al., 2015). In culture experiments, abiotic controls where pH, Eh, and reactant supply—including sulfide oxidation intermediates—are accurately controlled or quantified are critical to constraining unambiguous biotic influence on mineral rates of formation or morphology. Regarding rates of mineral formation, one study supports the conclusion that SRP may increase them (Mohagheghi et al., 1985), and that pyrite formation might be accelerated by $\text{S}(0)$ -disproportionating microorganisms (Canfield et al., 1998). Nevertheless, more studies with a greater variety of microorganisms and

conditions should be performed to confirm or constrain this result.

If the goal is to arrive at a better understanding of the effects SRP and other microbes have on the formation of extracellular metal sulfides in natural environments, the “holy grail” could be described as studies that link microbial phylogeny and physiology with mineralogy, mineral morphology, rates of mineral formation, and isotope and element chemistry. Our understanding of the evolution of the Earth’s surface environment is mainly based on analyses of iron sulfides preserved in ancient or modern sediments. Additionally the presence of pyritized organisms in the sedimentary record of Archaean to Jurassic age implies a potential role for Fe sulfides in the preservation of micro- and macrofossils (Sagemann et al., 1999; Grimes et al., 2001; Schieber, 2002; Cosmidis et al., 2013; Vietti et al., 2015). Deciphering the role of microorganisms in the formation of sulfide minerals is therefore crucial to better constrain the evolution of the biogeochemical cycles of Fe, S, and C.

CONCLUDING REMARKS

Sulfide minerals are of great importance for diverse fields of research, e.g., paleontology and reconstruction of past environments, material sciences, and the potential use of sulfide minerals for industrial applications, and economic geology. Two mechanisms of extracellular, biologically induced mineralization have been established: (1) SRP, and potentially other microorganisms, produce free sulfide, which reacts abiotically with dissolved metals to form sulfide minerals, and

(2) the association of iron with their cell walls and EPS, which then react with sulfide on these surfaces to form metal sulfide minerals. These data, however, do not preclude the possibility that microbes play a more direct role in extracellular iron sulfide formation, perhaps in a manner similar to the intracellular metal sulfide formation found in MTB. The limits of analytical tools and resolution faced by previous investigators may have hindered their ability to better characterize the mineral phases observed during laboratory experiments, but recent advances may provide a timely opportunity to revisit this long-standing question. We suggest that in order to fully consider the microbial role in the formation of metal sulfides, a structured approach that queries both microbial processes and minerals in culture, looks for evidence in natural environments, and attempts to link microbial processes and metal sulfides across spatial scales should be followed. Such an approach is likely to be both challenging and enlightening.

AUTHOR CONTRIBUTIONS

All authors listed, have made substantial, direct and intellectual contribution to the work, and approved it for publication.

ACKNOWLEDGMENTS

This material is based upon work supported by the National Science Foundation under Grant No. 1344241. We thank Alyssa J. Findlay and Elizabeth D. Swanner for providing comments and suggestions regarding the manuscript.

REFERENCES

- Allègre, C. J., Poirier, J. P., Humler, E., and Hofmann, A. W. (1995). The chemical composition of the Earth. *Earth Planet. Sci. Lett.* 134, 515–526.
- Andres, R., and Kasgnoc, A. (1998). A time-averaged inventory of subaerial volcanic sulfur emissions. *J. Geophys. Res.* 103, 25251–25261.
- Baas Becking, L. G. M., and Moore, D. (1961). Biogenic sulfides. *Econ. Geol.* 56, 259–272.
- Bak, F., and Pfennig, N. (1987). Chemolithotrophic growth of *Desulfovibrio sulfodismutans* sp. nov. by disproportionation of inorganic sulfur compounds. *Arch. Microb.* 147, 184–189.
- Bazylinski, D. A., Frankel, R. B., Heywood, B. R., Mann, S., King, J. W., Donaghay, P. L., et al. (1995). Controlled biomineralization of magnetite (Fe₃O₄) and greigite (Fe₃S₄) in a magnetotactic bacterium. *Appl. Environ. Microbiol.* 61, 3232–3239.
- Behrens, S., Kappler, A., and Obst, M. (2012). Linking environmental processes to the *in situ* functioning of microorganisms by high-resolution secondary ion mass spectrometry (NanoSIMS) and scanning transmission X-ray microscopy (STXM). *Environ. Microbiol.* 14, 2851–2869. doi: 10.1111/j.1462-2920.2012.02724.x
- Beijerinck, W. M. (1895). Ueber *Spirillum desulfuricans* als Ursache von Sulfatreduktion. *Zentralbl. Bakteriol.* 1, 1–9, 49–59, 104–114.
- Benning, L. G., Wilkin, R. T., and Konhauser, K. O. (1999). “Iron monosulfide stability: experiments with sulphate-reducing bacteria,” in *International Symposium on the Geochemistry of Earth’s surface*, ed H. Armannsson (Reykjavik: Balkema), 429–432.
- Berner, R. A. (1982). Burial of organic carbon and pyrite sulfur in the modern ocean: its geochemical and environmental significance. *Am. J. Sci.* 282, 451–473. doi: 10.2475/ajs.282.4.451
- Berner, R. A. (1984). Sedimentary pyrite formation: An update. *Geochim. Cosmochim. Acta* 48, 605–615.
- Beveridge, T. J. (1989). Role of cellular design in metal accumulation and mineralization. *Annu. Rev. Microbiol.* 43, 147–171.
- Bourdoiseau, J. A., Jeannin, M., Remazeilles, C., Sabot, R., and Refait, P. (2011). The transformation of mackinawite into greigite studied by Raman spectroscopy. *J. Raman Spectrosc.* 42, 496–504. doi: 10.1002/jrs.2729
- Bourdoiseau, J. A., Jeannin, M., Sabot, R., Remazeilles, C., and Refait, P. (2008). Characterisation of mackinawite by Raman spectroscopy: effects of crystallisation, drying and oxidation. *Corros. Sci.* 50, 3247–3255. doi: 10.1016/j.corsci.2008.08.041
- Braissant, O., Decho, A. W., Dupraz, C., Glunk, C., Przekop, K. M., and Visscher, P. T. (2007). Exopolymeric substances of sulfate-reducing bacteria: interactions with calcium at alkaline pH and implication for formation of carbonate minerals. *Geobiology* 5, 401–411. doi: 10.1111/j.1472-4669.2007.00117.x
- Burns, S. J., McKenzie, J. A., and Vasconcelos, C. (2000). Dolomite formation and biogeochemical cycles in the phanerozoic. *Sedimentology* 47, 49–61. doi: 10.1046/j.1365-3091.2000.00004.x
- Canfield, D. E., Stewart, F. J., Thamdrup, B., De Brabandere, L., Dalsgaard, T., Delong, E. F., et al. (2010). A cryptic sulfur cycle in oxygen-minimum-zone waters off the Chilean coast. *Science* 330, 1375–1378. doi: 10.1126/science.1196889
- Canfield, D. E., Thamdrup, B., and Fleischer, S. (1998). Isotope fractionation and sulfur metabolism by pure and enrichment cultures of elemental sulfur-disproportionating bacteria. *Limnol. Oceanogr.* 43, 253–264.
- Chambers, L., and Trudinger, P. (1979). Microbiological fractionation of stable sulfur isotopes: a review and critique. *Geomicrobiol. J.* 1, 249–293. doi: 10.1080/01490457909377735

- Cosmidis, J., and Benzerara, K. (2014). "Soft X-ray scanning transmission spectromicroscopy," in *Biomining Sourcebook: Characterization of Biominerals and Biomimetic Materials*, eds E. DiMasi and L. B. Gower (London: CRC Press), 115–133.
- Cosmidis, J., Benzerara, K., Gheerbrant, E., Esteve, I., Bouya, B., and Amaghaz, M. (2013). Nanometer-scale characterization of exceptionally preserved bacterial fossils in Paleocene phosphorites from Ouled Abdoun (Morocco). *Geobiology* 11, 139–153. doi: 10.1111/gbi.12022
- Donald, R., and Southam, G. (1999). Low temperature anaerobic bacterial diagenesis of ferrous monosulfide to pyrite. *Geochim. Cosmochim. Acta* 63, 2019–2023.
- Druschel, G. K., Labrenz, M., Thomsen-Ebert, T., Fowle, D. A., and Banfield, J. F. (2002). Geochemical modeling of ZnS in biofilms: an example of ore depositional processes. *Econ. Geol.* 97, 1319–1329. doi: 10.2113/gsecongeo.97.6.1319
- Elderfield, H., and Schultz, A. (1996). Mid-ocean ridge hydrothermal fluxes and the chemical composition of the ocean. *Annu. Rev. Earth Planet. Sci.* 24, 191–224.
- Farina, M., Esquivel, D. M. S., and De Barros, H. G. P. L. (1990). Magnetic iron-sulphur crystals from a magnetotactic microorganism. *Nature* 343, 256–258. doi: 10.1038/343256a0
- Ferris, F. G., Fyfe, W. S., and Beveridge, T. J. (1987). Bacteria as nucleation sites for authigenic minerals in a metal-contaminated lake sediment. *Chem. Geol.* 63, 225–232.
- Fike, D. A., Gammon, C. L., Ziebis, W., and Orphan, V. J. (2008). Micron-scale mapping of sulfur cycling across the oxycline of a cyanobacterial mat: a paired nanoSIMS and CARD-FISH approach. *ISME J.* 2, 749–759. doi: 10.1038/ismej.2008.39
- Finster, K., Liesack, W., and Thamdrup, B. (1998). Elemental sulfur and thiosulfate disproportionation by *Desulfocapsa sulfoexigens* sp nov, a new anaerobic bacterium isolated from marine surface sediment. *Appl. Environ. Microbiol.* 64, 119–125.
- Fortin, D., and Beveridge, T. J. (1997). Microbial sulfate reduction within sulfidic mine tailings: formation of diagenetic Fe sulfides. *Geomicrobiol. J.* 14, 1–21. doi: 10.1080/01490459709378030
- Fortin, D., Southam, G., and Beveridge, T. J. (1994). Nickel sulfide, iron-nickel sulfide and iron sulfide precipitation by a newly isolated *Desulfotomaculum* species and its relation to nickel resistance. *FEMS Microbiol. Ecol.* 14, 121–132.
- Foti, M., Sorokin, D. Y., Lomans, B., Mussman, M., Zacharova, E. E., Pimenov, N. V., et al. (2007). Diversity, activity, and abundance of sulfate-reducing bacteria in saline and hypersaline soda lakes. *Appl. Environ. Microbiol.* 73, 2093–2100. doi: 10.1128/AEM.02622-06
- Frank, K. L., Rogers, D. R., Olins, H. C., Vidoudez, C., and Girguis, P. R. (2013). Characterizing the distribution and rates of microbial sulfate reduction at Middle Valley hydrothermal vents. *ISME J.* 7, 1391–1401. doi: 10.1038/ismej.2013.17
- Freke, A. M., and Tate, D. (1961). The formation of magnetic iron sulfide by bacterial reduction of iron solutions. *J. Biochem. Microbiol.* 3, 29–39.
- Gallagher, K. L., Kading, T. J., Braissant, O., Dupraz, C., and Visscher, P. T. (2012). Inside the alkalinity engine: the role of electron donors in the organomineralization potential of sulfate-reducing bacteria. *Geobiology* 10, 518–530. doi: 10.1111/j.1472-4669.2012.00342.x
- Gramp, J. P., Bigham, J. M., Jones, F. S., and Tuovinen, O. H. (2010). Formation of Fe-sulfides in cultures of sulfate-reducing bacteria. *J. Hazard. Mater.* 175, 1062–1067. doi: 10.1016/j.jhazmat.2009.10.119
- Gramp, J. P., Wang, H. M., Bigham, J. M., Jones, F. S., and Tuovinen, O. H. (2009). Biogenic synthesis and reduction of Fe(III)-hydroxysulfates. *Geomicrobiol. J.* 26, 275–280. doi: 10.1080/01490450902892597
- Griesbeck, C., Hauska, G., and Schütz, M. (2000). "Biological sulfide oxidation: sulfide-quinone reductase (SQR), the primary reaction," in *Recent Research Developments in Microbiology*, ed S. G. Pandalai (Trivandrum: Research Signpost), 179–203.
- Grimes, S. T., Brock, F., Rickard, D., Davies, K. L., Edwards, D., Briggs, D. E., et al. (2001). Understanding fossilization: experimental pyritization of plants. *Geology* 29, 123–126. doi: 10.1130/0091-7613(2001)029<0123:UFEPPOP>2.0.CO;2
- Herbert, R. B. Jr., Benner, S. G., Pratt, A. R., and Blowes, D. W. (1998). Surface chemistry and morphology of poorly crystalline iron sulfides precipitated in media containing sulfate-reducing bacteria. *Chem. Geol.* 144, 87–97.
- Holmkvist, L., Ferdelman, T. G., and Jørgensen, B. B. (2011). A cryptic sulfur cycle driven by iron in the methane zone of marine sediment (Aarhus Bay, Denmark). *Geochim. Cosmochim. Acta* 75, 3581–3599. doi: 10.1016/j.gca.2011.03.033
- Jørgensen, B. B. (1982). Mineralization of organic matter in the sea bed—the role of sulphate reduction. *Nature* 296, 643–645. doi: 10.1038/296643a0
- Jørgensen, B. B., Isaksen, M. F., and Jannasch, H. W. (1992). Bacterial sulfate reduction above 100°C in deep-sea hydrothermal vent sediments. *Science* 258, 1756–1757. doi: 10.1126/science.258.5089.1756
- Jørgensen, B. B., and Nelson, D. C. (2004). Sulfide oxidation in marine sediments: geochemistry meets microbiology. *Geol. Soc. Am. Spec. Pap.* 379, 63–81. doi: 10.1130/0-8137-2379-5.63
- Karr, E. A., Ng, J. M., Belchik, S. M., Sattley, W. M., Madigan, M. T., and Achenbach, L. A. (2006). Biodiversity of methanogenic and other archaea in the permanently frozen lake Fryxell, Antarctica. *Appl. Environ. Microbiol.* 72, 1663–1666. doi: 10.1128/AEM.72.2.1663-1666.2006
- Kato, S., Ikehata, K., Shibuya, T., Urabe, T., Ohkuma, M., and Yamagishi, A. (2015). Potential for biogeochemical cycling of sulfur, iron and carbon within massive sulfide deposits below the seafloor. *Environ. Microbiol.* 17, 1817–1835. doi: 10.1111/1462-2920.12648
- Kim, Y., Lee, Y., and Roh, Y. (2015). Microbial synthesis of iron sulfide (FeS) and iron carbonate (FeCO₃) nanoparticles. *J. Nanosci. Nanotechnol.* 15, 5794–5797. doi: 10.1166/jnn.2015.10457
- Kohn, M. J., Riciputi, L. R., Stakes, D., and Orange, D. L. (1998). Sulfur isotope variability in biogenic pyrite: reflections of heterogeneous bacterial colonization? *Am. Mineral.* 83, 1454–1468. doi: 10.2138/am-1998-11-1234
- Kormas, K. A., Tivey, M. K., Von Damm, K., and Teske, A. (2006). Bacterial and archaeal phylotypes associated with distinct mineralogical layers of a white smoker spire from a deep-sea hydrothermal vent site (9 degrees N, East Pacific Rise). *Environ. Microbiol.* 8, 909–920. doi: 10.1111/j.1462-2920.2005.00978.x
- Labrenz, M., and Banfield, J. F. (2004). Sulfate-reducing bacteria-dominated biofilms that precipitate ZnS in a subsurface circumneutral-pH mine drainage system. *Microb. Ecol.* 47, 205–217. doi: 10.1007/s00248-003-1025-8
- Labrenz, M., Druschel, G. K., Thomsen-Ebert, T., Gilbert, B., Welch, S. A., Kemner, K. M., et al. (2000). Formation of sphalerite (ZnS) deposits in natural biofilms of sulfate-reducing bacteria. *Science* 290, 1744–1747. doi: 10.1126/science.290.5497.1744
- Lefevre, C. T., and Bazylynski, D. A. (2013). Ecology, diversity, and evolution of magnetotactic bacteria. *Microbiol. Mol. Biol. R.* 77, 497–526. doi: 10.1128/MMBR.00021-13
- Lefevre, C. T., Menguy, N., Abreu, F., Lins, U., Posfai, M., Prozorov, T., et al. (2011). A cultured greigite-producing magnetotactic bacterium in a novel group of sulfate-reducing bacteria. *Science* 334, 1720–1723. doi: 10.1126/science.1212596
- Leloup, J., Fossing, H., Kohls, K., Holmkvist, L., Borowski, C., and Jørgensen, B. B. (2009). Sulfate-reducing bacteria in marine sediment (Aarhus Bay, Denmark): abundance and diversity related to geochemical zonation. *Environ. Microbiol.* 11, 1278–1291. doi: 10.1111/j.1462-2920.2008.01855.x
- Li, Y. L., Vali, H., Sears, S. K., Yang, J., Deng, B., and Zhang, C. L. (2004). Iron reduction and alteration of nontronite NAu-2 by a sulfate-reducing bacterium. *Geochim. Cosmochim. Acta* 68, 3251–3260. doi: 10.1016/j.gca.2004.03.004
- Li, Y. L., Vali, H., Yang, J., Phelps, T. J., and Zhang, C. L. (2006). Reduction of iron oxides enhanced by a sulfate-reducing bacterium and biogenic H₂S. *Geomicrobiol. J.* 23, 103–117. doi: 10.1080/01490450500533965
- Maclean, L. C. W., Pray, T. J., Onstott, T. C., Brodie, E. L., Hazen, T. C., and Southam, G. (2007). Mineralogical, chemical and biological characterization of an anaerobic biofilm collected from a borehole in a deep gold mine in South Africa. *Geomicrobiol. J.* 24, 491–504. doi: 10.1080/01490450701572416
- Maclean, L. C. W., Tylliszczak, T., Gilbert, P. U. P. A., Zhou, D., Pray, T. J., Onstott, T. C., et al. (2008). A high-resolution chemical and structural study of framboidal pyrite formed within a low-temperature bacterial biofilm. *Geobiology* 6, 471–480. doi: 10.1111/j.1472-4669.2008.00174.x
- Miller, L. P. (1950). Tolerance of sulfate-reducing bacteria to hydrogen sulfide. *Contrib. Boyce Thompson Inst.* 16, 73–83.
- Mohagheghi, A., Updegraff, D. M., and Goldhaber, M. B. (1985). The role of sulfate-reducing bacteria in the deposition of sedimentary uranium ores. *Geomicrobiol. J.* 4, 153–173.

- Nakagawa, T., Ishibashi, J.-I., Maruyama, A., Yamanaka, T., Morimoto, Y., Kimura, H., et al. (2004). Analysis of dissimilatory sulfite reductase and 16S rRNA gene fragments from deep-sea hydrothermal sites of the Suiyo seamount, Izu-Bonin-arc, Western Pacific. *Appl. Environ. Microbiol.* 70, 393–403. doi: 10.1128/AEM.70.1.393-403.2004
- Neal, A. L., Techkarnjanaruk, S., Dohnalkova, A., McCready, D., Peyton, B. M., and Geesey, G. G. (2001). Iron sulfides and sulfur species produced at hematite surfaces in the presence of sulfate-reducing bacteria. *Geochim. Cosmochim. Acta* 65, 223–235. doi: 10.1016/S0016-7037(00)00537-8
- Orphan, V., and House, C. (2009). Geobiological investigations using secondary ion mass spectrometry: microanalysis of extant and paleo-microbial processes. *Geobiology* 7, 360–372. doi: 10.1111/j.1472-4669.2009.00201.x
- Posfai, M., Buseck, P. R., Bazylinski, D. A., and Frankel, R. B. (1998). Reaction sequence of iron sulfide minerals in bacteria and their use as biomarkers. *Science* 280, 880–883. doi: 10.1126/science.280.5365.880
- Posfai, M., Kasama, T., and Dunin-Borkowski, R. E. (2013a). “Biomaterials at the nanoscale: transmission electron microscopy methods for studying the special properties of biomaterials,” in *Minerals at the Nanoscale*, eds F. Nieto, K. T. V. Livi, and R. Oberti (London: Mineralogical Society of Great Britain and Ireland), 377–435.
- Posfai, M., Lefevre, C. T., Trubitsyn, D., Bazylinski, D. A., and Frankel, R. B. (2013b). Phylogenetic significance of composition and crystal morphology of magnetosome minerals. *Front. Microbiol.* 4:344. doi: 10.3389/fmicb.2013.00344
- Prange, A., Engelhardt, H., Truper, H. G., and Dahl, C. (2004). The role of the sulfur globule proteins of *Allochromatium vinosum*: mutagenesis of the sulfur globule protein genes and expression studies by real-time RT-PCR. *Arch. Microbiol.* 182, 165–174. doi: 10.1007/s00203-004-0683-3
- Rabus, R., Hansen, T., and Widdel, F. (2013). “Dissimilatory sulfate- and sulfur-reducing prokaryotes,” in *The Prokaryotes*, eds E. Rosenberg, E. DeLong, S. Lory, E. Stackebrandt, and F. Thompson (Heidelberg: Springer), 309–404.
- Ramamoorthy, S., Piotrowski, J. S., Langner, H. W., Holben, W. E., Morra, M. J., and Rosenzweig, R. F. (2009). Ecology of sulfate-reducing bacteria in an iron-dominated, mining-impacted freshwater sediment. *J. Environ. Qual.* 38, 675–684. doi: 10.2134/jeq2007.0577
- Ravenschlag, K., Sahn, K., Knoblauch, C., Jørgensen, B. B., and Amann, R. (2000). Community structure, cellular rRNA content, and activity of sulfate-reducing bacteria in marine Arctic sediments. *Appl. Environ. Microbiol.* 66, 3592–3602. doi: 10.1128/AEM.66.8.3592-3602.2000
- Rickard, D. T. (1969). The microbiological formation of iron sulphides. *Stockholm Contrib. Geol.* 20, 49–66.
- Rickard, D. T. (2012a). “Metastable sedimentary iron sulfides,” in *Developments in Sedimentology*, ed D. T. Rickard (Oxford: Elsevier), 195–231.
- Rickard, D. T. (2012b). “Microbial sulfate reduction in sediments,” in *Developments in Sedimentology*, ed D. T. Rickard (Oxford: Elsevier), 319–351.
- Rickard, D. T. (2012c). “Sedimentary pyrite,” in *Developments in Sedimentology*, ed D. T. Rickard (Oxford: Elsevier), 233–285.
- Robador, A., Jungbluth, S. P., Larowe, D., Bowers, R., Rappe, M., Amend, J., et al. (2015). Activity and phylogenetic diversity of sulfate-reducing microorganisms in low-temperature subsurface fluids within the upper oceanic crust. *Front. Microbiol.* 5:748. doi: 10.3389/fmicb.2014.00748
- Sagemann, J., Bale, S. J., Briggs, D. E., and Parkes, R. J. (1999). Controls on the formation of authigenic minerals in association with decaying organic matter: an experimental approach. *Geochim. Cosmochim. Acta* 63, 1083–1095. doi: 10.1016/S0016-7037(99)00087-3
- Sass, H., Ramamoorthy, S., Yarwood, C., Langner, H., Schumann, P., Kroppenstedt, R. M., et al. (2009). *Desulfovibrio idahonensis* sp. nov., sulfate-reducing bacteria isolated from a metal(loid)-contaminated freshwater sediment. *Int. J. Syst. Evol. Microbiol.* 59, 2208–2214. doi: 10.1099/ijs.0.016709-0
- Schieber, J. (2002). Sedimentary pyrite: a window into the microbial past. *Geology* 30, 531–534. doi: 10.1130/0091-7613(2002)030<0531:SPAUIT>2.0.CO;2
- Schoonen, M. A. (2004). Mechanisms of sedimentary pyrite formation. *Geol. Soc. Am. Spec. Pap.* 379, 117–134. doi: 10.1130/0-8137-2379-5.117
- Siebert, C. E. (1915). *Origin of the Zinc and Lead Deposits of the Joplin Region, Missouri, Kansas, and Oklahoma*. Washington, DC: Govt. Print. Off.
- Simmons, S. L., Sievert, S. M., Frankel, R. B., Bazylinski, D. A., and Edwards, K. J. (2004). Spatiotemporal distribution of marine magnetotactic bacteria in a seasonally stratified coastal salt pond. *Appl. Environ. Microbiol.* 70, 6230–6239. doi: 10.1128/AEM.70.10.6230-6239.2004
- Temple, K. L., and Le Roux, N. W. (1964). Syngensis of sulfide ores: desorption of adsorbed metal ions and their precipitation as sulfides. *Econ. Geol.* 59, 647–655. doi: 10.2113/gsecongeo.59.4.647
- Templeton, A., and Knowles, E. (2009). Microbial transformations of minerals and metals: recent advances in geomicrobiology derived from synchrotron-based X-ray spectroscopy and X-ray microscopy. *Annu. Rev. Earth Planet. Sci.* 37, 367–391. doi: 10.1146/annurev.earth.36.031207.124346
- Thamdrup, B., Finster, K., Hansen, J. W., and Bak, F. (1993). Bacterial disproportionation of elemental sulfur coupled to chemical reduction of iron or manganese. *Appl. Environ. Microbiol.* 59, 101–108.
- Trudinger, P. A., Chambers, L. A., and Smith, J. W. (1985). Low-temperature sulphate reduction: biological versus abiological. *Can. J. Earth Sci.* 22, 1910–1918.
- Vietti, L. A., Bailey, J. V., Fox, D., and Rogers, R. R. (2015). Rapid formation of framboidal sulfides on bone surfaces from a simulated marine carcass fall. *Palaios* 30, 327–334. doi: 10.2110/palo.2014.027
- Wacey, D., Kilburn, M. R., Saunders, M., Cliff, J. B., Kong, C., Liu, A. G., et al. (2015). Uncovering framboidal pyrite biogenicity using nano-scale CN_{org} mapping. *Geology* 43, 27–30. doi: 10.1130/G36048.1
- Wacey, D., McLoughlin, N., Whitehouse, M. J., and Kilburn, M. R. (2010). Two coexisting sulfur metabolisms in a ca. 3400 Ma sandstone. *Geology* 38, 1115–1118. doi: 10.1130/G31329.1
- Watson, J. H. P., Cressey, B. A., Roberts, A. P., Ellwood, D. C., Charnock, J. M., and Soper, A. K. (2000). Structural and magnetic studies on heavy-metal-adsorbing iron sulphide nanoparticles produced by sulphate-reducing bacteria. *J. Magn. Mater.* 214, 13–30. doi: 10.1016/S0304-8853(00)00025-1
- White, S. N. (2009). Laser Raman spectroscopy as a technique for identification of seafloor hydrothermal and cold seep minerals. *Chem. Geol.* 259, 240–252. doi: 10.1016/j.chemgeo.2008.11.008
- Widdel, F., and Bak, F. (1992). “Gram-negative mesophilic sulfate-reducing bacteria,” in *The Prokaryotes*, eds A. Balows, H. Trüper, M. Dworkin, W. Harder, and K. H. Schleifer (New York, NY: Springer), 3352–3378.
- Williams, K. H., Ntarlagiannis, D., Slater, L. D., Dohnalkova, A., Hubbard, S. S., and Banfield, J. F. (2005). Geophysical imaging of stimulated microbial biomineralization. *Environ. Sci. Technol.* 39, 7592–7600. doi: 10.1021/es0504035
- Zammit, C. M., Shuster, J. P., Gagen, E. J., and Southam, G. (2015). The geomicrobiology of supergene metal deposits. *Elements* 11, 337–342. doi: 10.2113/gselements.11.5.337
- Zhou, C., Vannela, R., Hayes, K. F., and Rittmann, B. E. (2014). Effect of growth conditions on microbial activity and iron-sulfide production by *Desulfovibrio vulgaris*. *J. Hazard. Mater.* 272, 28–35. doi: 10.1016/j.jhazmat.2014.02.046
- Zopf, J., Böttcher, M. E., and Jørgensen, B. B. (2008). Biogeochemistry of sulfur and iron in *Thioploca*-colonized surface sediments in the upwelling area off central Chile. *Geochim. Cosmochim. Acta* 72, 827–843. doi: 10.1016/j.gca.2007.11.031
- Zopf, J., Ferdelman, T., and Fossing, H. (2004). Distribution and fate of sulfur intermediates - sulfite, tetrathionate, thiosulfate, and elemental sulfur - in marine sediments. *Geol. Soc. Am. Spec. Pap.* 379, 97–116. doi: 10.1130/0-8137-2379-5.97

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Picard, Gartman and Girguis. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.