



Novel genetic loci associated with hippocampal volume

Citation

Hibar, D. P., H. H. H. Adams, N. Jahanshad, G. Chauhan, J. L. Stein, E. Hofer, M. E. Renteria, et al. 2017. "Novel genetic loci associated with hippocampal volume." Nature Communications 8 (1): 13624. doi:10.1038/ncomms13624. http://dx.doi.org/10.1038/ncomms13624.

Published Version

doi:10.1038/ncomms13624

Permanent link

http://nrs.harvard.edu/urn-3:HUL.InstRepos:31731854

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA

Share Your Story

The Harvard community has made this article openly available. Please share how this access benefits you. <u>Submit a story</u>.

<u>Accessibility</u>



ARTICLE

Received 23 Aug 2016 | Accepted 18 Oct 2016 | Published 18 Jan 2017

DOI: 10.1038/ncomms13624

OPEN

Novel genetic loci associated with hippocampal volume

Derrek P. Hibar, Hieab H.H. Adams, Neda Jahanshad, Ganesh Chauhan, Jason L. Stein, Edith Hofer, Miguel E. Renteria, Joshua C. Bis *et al.*[#]

The hippocampal formation is a brain structure integrally involved in episodic memory, spatial navigation, cognition and stress responsiveness. Structural abnormalities in hippocampal volume and shape are found in several common neuropsychiatric disorders. To identify the genetic underpinnings of hippocampal structure here we perform a genome-wide association study (GWAS) of 33,536 individuals and discover six independent loci significantly associated with hippocampal volume, four of them novel. Of the novel loci, three lie within genes (*ASTN2, DPP4* and *MAST4*) and one is found 200 kb upstream of *SHH*. A hippocampal subfield analysis shows that a locus within the *MSRB3* gene shows evidence of a localized effect along the dentate gyrus, subiculum, CA1 and fissure. Further, we show that genetic variants associated with decreased hippocampal volume are also associated with increased risk for Alzheimer's disease ($r_g = -0.155$). Our findings suggest novel biological pathways through which human genetic variation influences hippocampal volume and risk for neuropsychiatric illness.

Correspondence and requests for materials should be addressed to P.M.T. (email: pthomp@usc.edu) or to M.A.I. (email:m.a.ikram@erasmusmc.nl). #A full list of authors and their affiliations appears at the end of the paper.

rain structural abnormalities in the hippocampal formation are found in many complex neurological and psychiatric disorders including temporal lobe epilepsy¹, vascular dementia², Alzheimer's disease³, major depression⁴, bipolar disorder⁵, schizophrenia⁶ and post-traumatic stress disorder⁷, among others. The diverse functions of the hippocampus, including episodic memory⁸, spatial navigation⁹, cognition¹⁰ and stress responsiveness¹¹ are commonly impaired in a broad range of diseases and disorders of the brain that are associated with insults to the hippocampal structure. Further, the cytoarchitectural subdivisions (or 'subfields') of the hippocampus are associated with distinct functions. For example, the dentate gyrus (DG) and sectors 3 and 4 of the cornu ammonis (CA) are involved in declarative memory acquisition¹², the subiculum and CA1 play a role in disambiguation during working memory processes¹³, and the CA2 is implicated in animal models of episodic time encoding¹⁴ and social memory¹⁵. The anterior hippocampus, which includes the fimbria, CA subregions and hippocampal -amygdaloid transition area (HATA), may be involved in the mediation of cognitive processes including imagination, recall and visual perception¹⁶ and anxiety-related behaviours¹⁷.

Environmental factors, such as stress, affect the hippocampus¹⁸, but genetic differences across individuals account for most of the population variation in its size; the heritability of hippocampal volume is high at around 70% (refs 19–21). High heritability and a crucial role in healthy and diseased brain function make the hippocampus an ideal target for genetic analysis. We formed a large global partnership to empower the quest for mechanistic insights into neuropsychiatric disorders associated with hippocampal abnormalities and to chart, in depth, the genetic underpinnings of the hippocampal structure.

Here we perform a GWAS meta-analysis of mean bilateral hippocampal volume in 33,536 individuals scanned at 65 sites around the world as a joint effort between the Enhancing Neuroimaging Genetics through Meta-analysis (ENIGMA) and the Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) consortia. Our primary goal is to find common genetic determinants of hippocampal volume with previously unobtainable power. We make considerable efforts to coordinate data analysis across all sites from both consortia to maximize the comparability of both genetic and imaging data. Standardized protocols for image analysis and genetic imputation are freely available online (see URLs). In the most powerful imaging study of the hippocampus to date, we shed light on the common genetic determinants of hippocampal structure and allow for a deepened understanding of the biological workings of the brain's memory centre. We confirm previously identified loci influencing hippocampal volume, identify four novel loci and determine genome-wide overlap with Alzheimer's disease.

Results

Novel genome-wide markers associated with hippocampal volume. Our combined meta-analysis (n = 26,814 individuals of European ancestry) revealed six independent, genome-wide significant loci associated with hippocampal volume (Fig. 1; Table 1). Four are novel: with index SNPs rs11979341 (7q36.3; $P = 1.42 \times 10^{-11}$), rs7020341 (9q33.1; $P = 3.04 \times 10^{-11}$), rs2268894 (2q24.2; $P = 5.89 \times 10^{-11}$), and rs2289881 (5q12.3; $P = 2.73 \times 10^{-8}$). The other two loci have been previously characterized in detail: with index SNPs rs77956314 (12q24.22, $P = 2.06 \times 10^{-25}$), in linkage disequilibrium (LD) $(r^2 = 0.901)$ in European samples from the 1000 Genomes Project, Phase 1v3) with our previously identified variant at this locus (rs7294919) and rs61921502 (12q14.3, $P = 1.94 \times 10^{-19}$), in LD ($r^2 = 0.459$) with previous top locus rs17178006 (refs 22-24; Fig. 2a-f). In addition to these SNPs, we identified nine independent loci with a statistically suggestive influence on hippocampal volume $(P < 1 \times 10^{-6};$ Supplementary Data 4). All pathway results and gene-based P values are summarized in Supplementary Data 6 and 7.

Variance explained in hippocampal volume by common variants. Common variants genotyped from across the whole-genome explained as much as 18.76% (s.e. 1.56%) of the observed variance in human hippocampal volume, based on LDSCORE regression²⁵ (Supplementary Fig. 3). Common genetic variants account for around a quarter of the overall heritability, estimated in twin studies to be around 70% (refs 19–21). Further partitioning the genome into functional categories using LDSCORE²⁶ revealed significant over-representation of regions evolutionarily conserved in mammals (P = 0.0026): 2.6% of the variants accounted for 43.3% of the 18.76% variance explained (Fig. 3).




Figure 1 | **Common genetic variants associated with hippocampal volume (**N = 26,814 of European ancestry). A Manhattan plot displays the association P value for each single-nucleotide polymorphism (SNP) in the genome (displayed as $-\log_{10}$ of the P-value). Genome-wide significance is shown for the $P = 5 \times 10^{-8}$ threshold (solid line) and also for the suggestive significance threshold of $P = 1 \times 10^{-6}$ (dotted line). The most significant SNP within an associated locus is labeled. For the significant loci and age-dependent loci (Chromosome 19) we labeled the nearest gene, which is not necessarily the gene of action.




Figure 2 | Functional annotation within genome-wide significant loci. For each panel (**a**-**f**), zoomed-in Manhattan plots (\pm 400 kb from top SNP) are shown with gene models below (GENCODE version 19). Plots below are zoomed to highlight the genomic region that likely harbors the causal variant(s) ($r^2 > 0.8$ from the top SNP). Genomic annotations from the Roadmap Epigenomics Consortium⁵³ are displayed to indicate potential functionality (see Methods for detailed track information). Each plot was made using the LocusTrack software⁵⁵ (see URLs).

Effects of top variants on hippocampal subfield volume. To test for differential effects on individual subfields of the hippocampal formation, we examined the six significant variants influencing whole hippocampal volume in a large cohort (n = 5,368). We found that the top SNP from our primary analysis, rs77956314, has a broad, nonspecific effect on hippocampal subfield volumes with the greatest effect in the right hippocampal tail $(P = 1.27 \times 10^{-8})$. rs61921502 showed strong lateral effects across right hippocampal subfields with the largest effect in the right hippocampal fissure $(P = 6.45 \times 10^{-9})$. rs7020341 showed greatest effects bilaterally in the subiculum (left: $P = 1.59 \times 10^{-8}$; right: P = 1.42 $\times 10^{-8}$). rs2268894 show left-lateralized effects across hippocampal subfields with the strongest effect in the left hippocampal tail $(P = 1.76 \times 10^{-5})$. The remaining two variants (rs11979341 and rs2289881) did not show significant evidence of association across any of the hippocampal subfields. The full set of results from the hippocampal subfield analysis is tabulated in Supplementary Data 8.

Genetic overlap with hippocampal volume. We used LDSCORE²⁷ regression to quantify the degree of common genetic overlap between variants influencing the hippocampus and those influencing Alzheimer's disease. We found significant evidence of a moderate, negative relationship whereby variants associated with a decrease in hippocampal volume are associated with an increased risk for Alzheimer's disease ($r_g = -0.155$ (s.e. 0.0529), P = 0.0034; see Methods).

Discussion

We identified six genome-wide significant, independent loci associated with hippocampal volume in 26,814 subjects of European ancestry. Of the six loci, four were novel: rs11979341 (7q36.3; $P = 1.42 \times 10^{-11}$), rs7020341 (9q33.1; $P = 3.04 \times 10^{-11}$), rs2268894 (2q24.2; $P = 5.89 \times 10^{-11}$) and rs2289881 (5q12.3; $P = 2.73 \times 10^{-8}$). We previously discovered two of the novel loci, rs7020341 and rs2268894 (ref. 24), but in this higher-powered analysis they now surpassed the genome-wide significance. In addition to the four novel loci, we replicated two loci associated with hippocampal volume: rs7492919 and rs17178006 (refs 23,24). Hibar *et al.*²² previously reported additional support for the rs17178006 association with hippocampal volume.

Each novel locus identified has unique functions and has previously been linked to diseases of the brain. Variant rs7020341 lies within an intron of the *astrotactin 2* (*ASTN2*) gene (Fig. 2d) which encodes for a protein involved in glial-mediated neuronal migration in the developing brain²⁸. Rare deletions overlapping this locus near the 3' end of *ASTN2* have been observed in patients with autism spectrum disorder and attention-deficit/ hyperactivity disorder²⁹. Common variants near this site are associated with autism spectrum disorders²⁹ and migraine³⁰. Variant rs2268894 is located in an intron of *DPP4* (Fig. 2e) that encodes dipeptidyl peptidase IV; an enzyme regulating response to the ingestion of food³¹, and an established target of a treatment for type 2 diabetes mellitus (vildagliptin)³². In addition, rs2268894 is in strong LD ($r^2 = 0.83$) with a genome-wide significant locus associated with a decreased risk for schizophrenia




Figure 3 | Analysis of variance explained, functional annotation, and pathway analysis. LDSCORE regression analysis for different functional annotation²⁶ categories (described further in Finucane *et al.*²⁶). Plotted values are the proportion of h_g^2 explained divided by the proportion of SNPs in a given functional category. Values are significantly over- or under-represented if they differ significantly from 1. Values are plotted with a standard error calculated with a jackknife in LDSCORE. Evolutionarily conserved regions across mammals significantly contributed to the heritability of hippocampal volume (indicated by **).

RSID	Chr	Pos	Nearest gene	Allele1	Allele2	Freq	Z-score	N	P value
rs77956314	12	117,323,367	4 kb 5' to HRK	Т	С	0.9160	- 10.418	26,814	2.06×10^{-25}
rs61921502	12	65,832,468	Intron of MSRB3	Т	G	0.8466	9.017	26,814	1.94×10^{-19}
rs11979341	7	155,797,978	200 kb 5′ to SHH	С	G	0.6837	- 6.755	24,484	1.42×10^{-11}
rs7020341	9	119,247,974	Intron of ASTN2	С	G	0.3590	6.645	26,700	3.04×10^{-11}
rs2268894	2	162,856,148	Intron of DPP4	Т	С	0.5412	- 6.546	26,814	5.89×10^{-11}
rs2289881	5	66,084,260	Intron of MAST4	Т	G	0.3544	- 5.558	26,814	2.73×10^{-8}

 $(rs2909457)^{33}$; however, the allele that increases risk for schizophrenia also increases hippocampal volume even though patients with schizophrenia show decreased hippocampal volume relative to controls⁶. Variant rs11979341 lies in an intergenic region (Fig. 2c) around 200 kb upstream of the *sonic hedgehog* (*SHH*) gene, crucial for neural tube formation³⁴. Adult brain expression data provide some evidence that rs11979341-C increases the expression of *SHH* in adult human hippocampus³⁵ (*P*=0.0089). Finally, variant rs2289881 lies within an intron of the microtubule-associated serine/threonine kinase family member 4 (*MAST4*) gene (Fig. 2f). The protein product of *MAST4* modulates the microtubule scaffolding; the gene has

been linked to susceptibility for atherosclerosis in HIV-infected men³⁶, and atypical frontotemporal dementia³⁷.

Effect sizes from the full sample were almost identical to those obtained from a subset meta-analysis (Pearson's $r^2 > 0.99$; n = 22,761) that removed all patients diagnosed with a neuropsychiatric disorder. Observed effects are therefore not likely to be driven by inclusion of patients with brain disorders. All significant loci are tabulated in Table 1. We found little evidence that these effects could be generalized to populations of African, Japanese, and Mexican-American ancestry, which could be due to the limited power from smaller non-European sample sizes available (n = 6,722; Supplementary Data 5).

We estimated that 18.76% (s.e. 1.56%) of the variance in hippocampal volume could be explained by genotyped common genetic variation. This effect was only tested within populations of European ancestry and does not necessarily reflect the level of explained variance in other populations worldwide. This is a substantial fraction of the overall genetic component of variance determined by twin heritability studies, and the heritability of hippocampal volume is relatively high at around 70% (refs 19-21). With the same LDSCORE method, we estimated the amount of variance explained by common gene variants belonging to known functional cell categories²⁶. We discovered enrichment of genomic regions conserved across mammals, which may have a strong evolutionary role in the hippocampal formation, a structure much more extensively developed in mammals than in other vertebrates³⁸. Given that hippocampal atrophy is a hallmark of Alzheimer's disease pathology³⁹, we were motivated to examine common genetic overlap between hippocampal volume and Alzheimer's disease risk. We found a significant negative relationship ($r_{\rm g} = -0.155$ (s.e. 0.0529), P = 0.0034), through which loci associated with decreased hippocampal volume also increase risk for AD. This confirms a shared etiological component between AD and hippocampal volume whereby genetic variants influencing hippocampal volume also modify the risk for developing AD.

As the hippocampal formation is a complex structure comprised of diverse functional units, we sought to examine the genetic variants identified in our analysis for focal effects on hippocampal subfield volumes. When assessing 13 subfields of the hippocampus (26 total, left and right) we found that two of the top variants from our analysis (rs77956314 and rs7020341) had largely non-specific effects: most of the subfield volumes showed significant evidence of association (Supplementary Data 8). The variant rs61921502 showed a lateralized effect across the body of the right hippocampal formation, which includes the DG, subiculum, CA1 and fissure. Volume losses are frequently observed across the hippocampal body in AD⁴⁰, major depression⁴¹, bipolar disorder⁴² and temporal lobe epilepsy⁴³. Prior pathway analyses have implicated the rs61921502 with MSR3B, a gene related to oxidative stress²⁴. Genetic variation at MSR3B may influence neurogenesis specifically within the dentate regions of the hippocampal body, where cell proliferation is known to continue into adulthood in healthy humans⁴⁴. However, further functional validation is required to test this hypothesis. Finally, the variant rs2268894 was associated with volume differences in the left hippocampal tail, a subfield that has previously shown shape abnormalities⁴⁵ and volume differences⁴⁶ in schizophrenia.

Here we identified four novel loci associated with hippocampal volume and examined each variant for localized effects in hippocampal subfields. When partitioning the full genome-wide association results into functionally annotated categories, we discovered that SNPs in evolutionarily conserved regions were significantly over-represented in their contribution to hippocampal volume. Further, we found significant evidence of shared genetic overlap between hippocampal volume and Alzheimer's disease. This large international effort shows that by mapping out the genetic influences on brain structure, we may begin to derive mechanistic hypotheses for brain regions causally implicated in the risk for neuropsychiatric disorders.

Methods

Subjects and sites. High-resolution MRI brain scans and genome-wide genotyping data were available for 33,536 individuals from 65 sites in two large consortia: the ENIGMA Consortium and the CHARGE Consortium. Full details and demographics for each participating cohort are given in Supplementary Data 1. All participants (or their legal representatives) provided written informed consent.

The institutional review board of the University of Southern California and the local ethics board of Erasmus MC University Medical Center approved this study.

Imaging analysis and quality control. Hippocampal volumes were estimated using the automated and previously validated segmentation algorithms, FSL FIRST⁴⁷ from the FMRIB Software Library (FSL) and FreeSurfer⁴⁸. Hippocampal segmentations were visually examined at each site, and poorly segmented scans were excluded. Sites also generated histogram plots to identify any volume outliers. Individuals with a volume more than three standard deviations away from the mean were visually inspected to verify proper segmentation. Statistical outliers were included in analysis if they were properly segmented; otherwise, they were removed. Average bilateral hippocampal volume was highly correlated across automated procedures used to measure it (Pearson's r = 0.74)²². A measure of head size-intracranial volume (ICV)-was used as a covariate in these analyses to adjust for volumetric differences due to differences in head size alone. Most sites measured ICV for each participant using the inverse of the determinant of the transformation matrix required to register the subject's MRI scan to a common template and then multiplied by the template volume (1,948,105 mm³). Full details of image acquisition and processing performed at each site are given in Supplementary Data 2.

Genetic imputation and quality control. Genetic data were obtained at each site using commercially available genotyping platforms. Before imputation, genetic homogeneity was assessed in each sample using multi-dimensional scaling (MDS). Ancestry outliers were excluded by visual inspection of the first two components. The primary analysis and all data presented in this main text were derived from subjects with European ancestry. Replication attempts in subjects of additional ancestries are presented in Supplementary Data 5. Data were further cleaned and filtered to remove single-nucleotide polymorphisms (SNPs) with low minor allele frequency (MAF < 0.01), deviations from Hardy-Weinberg Equilibrium (HWE; $P < 1 \times 10^{-6}$), and poor genotyping call rate (<95%). Cleaned and filtered data sets were imputed to the 1000 Genomes Project reference panel (phase 1, version 3) using freely available and validated imputation software (MaCH/minimac, IMPUTE2, BEAGLE, GenABLE). After imputation, genetic data were further quality checked to remove poorly imputed SNPs (estimated $R^2 < 0.5$) or low MAF <0.5%). Details on filtering criteria, quality control, and imputation at each site may be found in Supplementary Data 3.

Genome-wide association analysis and statistical models. GWAS were performed at each site, as follows. Mean bilateral hippocampal volume ((left + right)/2) was the trait of interest, and the additive dosage value of a SNP was the predictor of interest, while controlling for 4 MDS components, age, age², sex, intracranial volume and diagnosis (when applicable). For studies with data collected from multiple centres or scanners, additional covariates were also included in the model to adjust for any scanning site effects. Sites with family data (NTR-Adults, BrainSCALE, QTIM, SYS, GOBS, ASPSFam, ERF, GeneSTAR, NeuroIMAGE, OATS, RSIx) used mixed-effects models to account for familial relationships, in addition to covariates stated previously. The primary analyses for this paper focused on the full set of individuals, including datasets with patients, to maximize power. We re-analysed the data excluding patients to verify that detected effects were not due to disease alone. The regression coefficients for SNPs with $P < 1 \times 10^{-5}$ in the model including all patients were almost perfectly correlated with the regression coefficients from the model including only healthy individuals (Pearson's r = 0.996). Full details for the software used at each site are given in Supplementary Data 3.

The GWAS of mean hippocampal volume was performed at each site, and the resulting summary statistics uploaded to a centralized site for meta-analysis. Before meta-analysis, GWAS results from each site were checked for genomic inflation and errors using Quantile–Quantile (QQ) plots (Supplementary Figs 1 and 2). GWAS results from each site were combined using a fixed-effects sample size-weighted meta-analysis framework as implemented in METAL⁴⁹. Data were meta-analysed first in the ENIGMA and CHARGE Consortia separately and then combined into a final meta-analysed result file. After the final meta-analysis, SNPs were excluded if the SNP was available for fewer than 5,000 individuals.

Variance explained and genetic overlap in hippocampal volume. The common genetic overlap, total variance explained by the GWAS, and the partitioned heritability analyses were estimated using LDSCORE^{25,26}. Following from the polygenic model, an association test statistic at a given locus includes signal from all linked loci. Given a heritable polygenic trait, a SNP in high LD with, or tagging, a large number of SNPs is on average likely to show stronger association than a SNP that is not. The magnitude of information conveyed by each variant (a function of the number of SNPs tagged taking into account the strength of the tagging) is summarized as an LD score. By regressing the LD scores on the test statistics, we estimated the proportion of variance in the trait explained by the variants included in the analysis. As an extension, two LD score models for two separate traits can be used to estimate the covariance (and correlation) structure to yield an estimate of the common genetic overlap (r_e) between any two trait pairs.

ARTICLE

Here we estimated the common genetic overlap between hippocampal volume and Alzheimer's disease⁵⁰. Standard errors were estimated using a block jackknife.

Genomic partitioning into functional categories. As well as estimating the total variance explained, the genomic heritability (h_{g}^{2}) can be partitioned into specific subsets of variants. The functional annotation partitioning used the pre-prepared LDSCORE and annotation (.annot) files available online (see URLs) following the method of Finucane *et al.*²⁶. These analyses use the following 24 functional classes not specifically unique to any cell type: coding, UTR, promoter, intron, histone marks H3K4me1, H3K4me3, H3K9ac5 and two versions of H3K27ac, open chromatin DNase I hypersensitivity Site (DHS) regions, combined chromHMM/ Segway predictions, regions conserved in mammals, super-enhancers and active enhancers from the FANTOM5 panel of samples (Finucane et al., page 4)² Annotated coordinates are determined by a combination of all cell types from ENCODE. As in Finucane et al.²⁶, to avoid bias, we included the 500 bp windows surrounding the variants included in the functional classes. The chromosomepartitioned analyses were conducted using LDSCOREs calculated for each chromosome. Following the method of Bulik-Sullivan et al.25, these analyses focus on the variants within HapMap3 as these SNPs are typically well imputed across cohorts. Enrichment of a given partition is calculated as the proportion of h^2 explained by that partition divided by the proportion of variants in the GWAS that fall into that partition. All LDSCORE analyses used non-genomic controlled metaanalyses.

Gene annotation and pathway analysis. Gene annotation, gene-based test statistics, and pathway analysis were performed using the KGG2.5 software package⁵¹ (Supplementary Data 6 and 7). LD was calculated based on RSID numbers using the 1000 Genomes Project European samples as a reference (see URLs). For annotation, SNPs were considered 'within' a gene, if they fell within 5 kb of the 3'/5' UTR based on human genome (hg19) coordinates. Gene-based tests were performed using the GATES test⁵¹ without weighting *P* values by predicted functional relevance. Pathway analysis was performed using the HYST test of association⁵². For all gene-based tests and pathway analyses, results were considered significant if they exceeded a Bonferroni correction threshold accounting for the number of pathways in the REACTOME database tested such that $P_{thresh} = 0.05/(671 \text{ pathways}) = 7.45 \times 10^{-5}$.

Annotation of SNPs with epigenetic factors. In Fig. 2, all tracks were taken from the UCSC Genome Browser Human hg19 assembly. SNPs (top 5%) shows the top 5% associated SNPs within the locus and are coloured by their correlation to the top SNP. Genes shows the gene models from GENCODE version 19. Hippocampus gives the predicted chromatin states based on computational integration of ChIP-seq data for 18 chromatin marks in human hippocampal tissue derived from the Roadmap Epigenomics Consortium⁵³. The 18 chromatin states from the hippocampus track are as follows: TssA (Active TSS), TssFlnk (Flanking Active TSS), TssFlnkU (Flanking TSS Upstream), TssFlnkD (Flanking TSS Downstream), Tx (Strong transcription), TxWk (Weak transcription), EnhG1 (Genic Enhancers 1), EnhG2 (Genic Enhancers 2), EnhA1 (Active Enhancers 1), EnhA2 (Active Enhancers 2), EnhWk (Weak Enhancers), ZNF/Rpts (ZNF genes & repeats), Het (Heterochromatin), TssBiv (Bivalent/Poised TSS), EnhBiv (Bivalent Enhancer), ReprPC (Repressed PolyComb), ReprPCWk (Weak Repressed PolyComb), Quies (Quiescent/Low). Additional information about the 18 state chromatin model is detailed elsewhere⁵³. Conservation is the basewise conservation score over 100 vertebrates estimated by PhyloP from the UCSC Genome Browser Human hg19 assembly.

Analysis of hippocampal subfields. We segmented the hippocampal formation into 13 subfield regions: CA1, CA3, CA4, fimbria, Granule Layer + Molecular Layer + Dentate Gyrus Boundary (GC_ML_DG), hippocampal-amygdaloid transition area (HATA), hippocampal tail, hippocampal fissure, molecular layer (HP), parasubiculum, presubiculum and subiculum using a freely available, validated algorithm distributed with the FreeSurfer image analysis package⁵⁴. We measured the hippocampal subfield volumes within the Rotterdam (n = 4,491) and HUNT (n = 877) cohorts. Volumes from the 26 subfield regions (13 in each hemisphere) were the phenotypes of interest and individually assessed for significance with the top variants from our primary analysis while correcting for the following nuisance variables: 4 MDS components, age, age², sex, intracranial volume. Association statistics from each of the tests in the Rotterdam and HUNT cohorts were meta-analysed using a fixed-effects inverse variance-weighted model yielding the final results. We declare an individual test significant if the P value is less than a Bonferroni-corrected P value threshold accounting for the total number of tests: $P_{\text{thresh}} = 0.05/(26 \text{ subfields} \times 6$ $SNPs) = 3.21 \times 10^{-4}$.

Data availability. The genome-wide summary statistics that support the findings of this study are available upon request from the corresponding authors MAI and PMT (see URLs). The data are not publicly available due to them containing information that could compromise research participant privacy/consent.

URLs

https://github.com/bulik/ldsc http://enigma.usc.edu/protocols/genetics-protocols/ http://gump.qimr.edu.au/general/gabrieC/LocusTrack/ http://enigma.ini.usc.edu/download-enigma-gwas-results/ http://www.internationalgenome.org/data

References

- 1. Van Paesschen, W. *et al.* Quantitative hippocampal MRI and intractable temporal lobe epilepsy. *Neurology* **45**, 2233–2240 (1995).
- Kim, G. H. et al. Hippocampal volume and shape in pure subcortical vascular dementia. *Neurobiol. Aging* 36, 485–491 (2015).
- 3. Thompson, P. M. et al. Mapping hippocampal and ventricular change in Alzheimer disease. NeuroImage 22, 1754–1766 (2004).
- Schmaal, L. *et al.* Subcortical brain alterations in major depressive disorder: findings from the ENIGMA major depressive disorder working group. *Mol Psychiatry* 21, 806–812 (2015).
- Hibar, D. P. et al. Subcortical volumetric abnormalities in bipolar disorder. Molecular Psychiatry 21, 1710–1716 (2016).
- van Erp, T. G. *et al.* Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Mol Psychiatry* 21, 547–553 (2015).
- Smith, M. E. Bilateral hippocampal volume reduction in adults with posttraumatic stress disorder: a meta-analysis of structural MRI studies. *Hippocampus* 15, 798–807 (2005).
- Blis, T. V. & Collingridge, G. L. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 361, 31–39 (1993).
- 9. Maguire, E. A. *et al.* Knowing where and getting there: a human navigation network. *Science* **280**, 921–924 (1998).
- Vinogradova, O. S. Hippocampus as comparator: role of the two input and two output systems of the hippocampus in selection and registration of information. *Hippocampus* 11, 578–598 (2001).
- McEwen, B. S., Eiland, L., Hunter, R. G. & Miller, M. M. Stress and anxiety: structural plasticity and epigenetic regulation as a consequence of stress. *Neuropharmacology* 62, 3–12 (2012).
- Coras, R. *et al.* Differential influence of hippocampal subfields to memory formation: insights from patients with temporal lobe epilepsy. *Brain* 137, 1945–1957 (2014).
- Newmark, R. E., Schon, K., Ross, R. S. & Stern, C. E. Contributions of the hippocampal subfields and entorhinal cortex to disambiguation during working memory. *Hippocampus* 23, 467–475 (2013).
- 14. Navratilova, Z. & Battaglia, F. P. CA2: it's about time-and episodes. *Neuron* 85, 8–10 (2015).
- Hitti, F. L. & Siegelbaum, S. A. The hippocampal CA2 region is essential for social memory. *Nature* 508, 88–92 (2014).
- McCormick, C., St-Laurent, M., Ty, A., Valiante, T. A. & McAndrews, M. P. Functional and effective hippocampal-neocortical connectivity during construction and elaboration of autobiographical memory retrieval. *Cereb. Cortex.* 25, 1297–1305 (2015).
- Strange, B. A., Witter, M. P., Lein, E. S. & Moser, E. I. Functional organization of the hippocampal longitudinal axis. *Nat. Rev. Neurosci.* 15, 655–669 (2014).
- Lupien, S. J., McEwen, B. S., Gunnar, M. R. & Heim, C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat. Rev. Neurosci.* 10, 434–445 (2009).
- Renteria, M. E. *et al.* Genetic architecture of subcortical brain regions: common and region-specific genetic contributions. *Genes Brain Behav.* 13, 821–830 (2014).
- Kremen, W. S. *et al.* Genetic and environmental influences on the size of specific brain regions in midlife: the VETSA MRI study. *NeuroImage* 49, 1213–1223 (2010).
- den Braber, A. *et al.* Heritability of subcortical brain measures: a perspective for future genome-wide association studies. *NeuroImage* 83, 98–102 (2013).
- Hibar, D. P. et al. Common genetic variants influence human subcortical brain structures. Nature 520, 224–229 (2015).
- 23. Stein, J. L. et al. Identification of common variants associated with human hippocampal and intracranial volumes. Nat. Genet. 44, 552–561 (2012).
- 24. Bis, J. C. et al. Common variants at 12q14 and 12q24 are associated with hippocampal volume. Nat. Genet. 44, 545 (2012).
- Bulik-Sullivan, B. K. et al. LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nat. Genet.* 47, 291–295 (2015).
- Finucane, H. K. et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat. Genet. 47, 1228–1235 (2015).
- 27. Bulik-Sullivan, B. et al. An atlas of genetic correlations across human diseases and traits. Nat. Genet. 47, 1236–1241 (2015).
- Wilson, P. M., Fryer, R. H., Fang, Y. & Astn2, Hatten M. E. a novel member of the astrotactin gene family, regulates the trafficking of ASTN1 during glialguided neuronal migration. *J. Neurosci.* **30**, 8529–8540 (2010).

- Lionel A. C., et al. Disruption of the ASTN2/TRIM32 locus at 9q33.1 is a risk factor in males for autism spectrum disorders, ADHD and other neurodevelopmental phenotypes. Hum. Mol. Genet. 23, 2752–2768 (2014).
- D. Freilinger, T. et al. Genome-wide association analysis identifies susceptibility loci for migraine without aura. Nat. Genet. 44, 777–U205 (2012).
- Pratley, R. E. & Salsali, A. Inhibition of DPP-4: a new therapeutic approach for the treatment of type 2 diabetes. *Curr. Med. Res. Opin.* 23, 919–931 (2007).
- 32. Pratley, R. E., Jauffret-Kamel, S., Galbreath, E. & Holmes, D. Twelve-week monotherapy with the DPP-4 inhibitor vildagliptin improves glycemic control in subjects with type 2 diabetes. *Horm. Metab. Res.* 38, 423–428 (2006).
- Schizophrenia Working Group of the Psychiatric Genomics C. Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511, 421–427 (2014).
- Dessaud, E., McMahon, A. P. & Briscoe, J. Pattern formation in the vertebrate neural tube: a sonic hedgehog morphogen-regulated transcriptional network. *Development* 135, 2489–2503 (2008).
- 35. Ramasamy, A. *et al.* Genetic variability in the regulation of gene expression in ten regions of the human brain. *Nat. Neurosci.* **17**, 1418–1428 (2014).
- Shrestha, S. et al. A genome-wide association study of carotid atherosclerosis in HIV-infected men. AIDS 24, 583–592 (2010).
- Martins-de-Souza, D. *et al.* Proteomic analysis identifies dysfunction in cellular transport, energy, and protein metabolism in different brain regions of atypical frontotemporal lobar degeneration. *J. Proteome. Res.* 11, 2533–2543 (2012).
- Garcia-Verdugo, J. M. *et al.* The proliferative ventricular zone in adult vertebrates: a comparative study using reptiles, birds, and mammals. *Brain Res. Bull.* 57, 765–775 (2002).
- Apostolova, L. G. *et al.* Conversion of mild cognitive impairment to Alzheimer disease predicted by hippocampal atrophy maps. *Arch. Neurol.* 63, 693–699 (2006).
- Mueller, S. G. & Weiner, M. W. Selective effect of age, Apo e4, and Alzheimer's disease on hippocampal subfields. *Hippocampus* 19, 558–564 (2009).
- Huang, Y. *et al.* Structural changes in hippocampal subfields in major depressive disorder: a high-field magnetic resonance imaging study. *Biol. Psychiatry* 74, 62–68 (2013).
- 42. Haukvik, U. K. et al. In vivo hippocampal subfield volumes in schizophrenia and bipolar disorder. Biol. Psychiatry 77, 581–588 (2015).
- Schoene-Bake, J. C. *et al. In vivo* mapping of hippocampal subfields in mesial temporal lobe epilepsy: relation to histopathology. *Hum. Brain Mapp.* 35, 4718–4728 (2014).
- Erickson, K. I. et al. Exercise training increases size of hippocampus and improves memory. Proc. Natl Acad. Sci. USA 108, 3017–3022 (2011).
- Styner, M., Lieberman, J. A., Pantazis, D. & Gerig, G. Boundary and medial shape analysis of the hippocampus in schizophrenia. *Med. Image Anal.* 8, 197–203 (2004).
- Maller, J. J. et al. Hippocampal volumetrics in treatment-resistant depression and schizophrenia: the devil's in de-tail. *Hippocampus* 22, 9–16 (2012).
- Patenaude, B., Smith, S. M., Kennedy, D. N. & Jenkinson, M. A Bayesian model of shape and appearance for subcortical brain segmentation. *NeuroImage* 56, 907–922 (2011).
- Fischl, B. et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. Neuron 33, 341–355 (2002).
- WillerLi, C. J. & Abecasis, Y. GR. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26, 2190–2191 (2010).

- Lambert, J. C. et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. Nat. Genet. 45, 1452–1458 (2013).
- Li, M. X., Gui, H. S., Kwan, J. S. & Sham, P. C. GATES: a rapid and powerful gene-based association test using extended Simes procedure. *Am. J. Hum. Genet.* 88, 283–293 (2011).
- Li, M. X., Kwan, J. S. & Sham, P. C. HYST: a hybrid set-based test for genomewide association studies, with application to protein-protein interaction-based association analysis. *Am. J. Hum. Genet.* **91**, 478–488 (2012).
- 53. Roadmap Epigenomics, C. et al. Integrative analysis of 111 reference human epigenomes. Nature 518, 317-330 (2015).
- Iglesias, J. E. et al. A computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI: application to adaptive segmentation of *in vivo* MRI. *NeuroImage* 115, 117–137 (2015).
- Cuellar-Partida, G., Renteria, M. E. & MacGregor, S. LocusTrack: integrated visualization of GWAS results and genomic annotation. *Source Code Biol. Med.* 10, 1 (2015).

Acknowledgements

See Supplementary Note 2 for information on funding sources. Data used in preparing this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, many investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report (see Supplementary Note 1). A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/ how_to_apply/ADNI_Acknowledgement_List.pdf

Author contributions

See Supplementary Note 3 for author contribution statements.

Additional information

Supplementary Information accompanies this paper at http://www.nature.com/ naturecommunications

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at http://npg.nature.com/ reprintsandpermissions/

How to cite this article: Hibar, D. P. et al. Novel genetic loci associated with hippocampal volume. Nat. Commun. 8, 13624 doi: 10.1038/ncomms13624 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2017

Derrek P. Hibar^{1,*}, Hieab H.H. Adams^{2,3,*}, Neda Jahanshad^{1,*}, Ganesh Chauhan^{4,*}, Jason L. Stein^{1,5,*}, Edith Hofer^{6,7,*}, Miguel E. Renteria^{8,*}, Joshua C. Bis^{9,*}, Alejandro Arias-Vasquez^{10,11,12,13}, M. Kamran Ikram^{2,14,15,16,17}, Sylvane Desrivières¹⁸, Meike W. Vernooij^{2,3}, Lucija Abramovic¹⁹, Saud Alhusaini^{20,21}, Najaf Amin², Micael Andersson²², Konstantinos Arfanakis^{23,24,25}, Benjamin S. Aribisala^{26,27,28}, Nicola J. Armstrong^{29,30}, Lavinia Athanasiu^{31,32}, Tomas Axelsson³³, Ashley H. Beecham^{34,35}, Alexa Beiser^{36,37,38}, Manon Bernard³⁹, Susan H. Blanton^{34,35}, Marc M. Bohlken¹⁹, Marco P. Boks¹⁹, Janita Bralten^{10,13}, Adam M. Brickman⁴⁰, Owen Carmichael⁴¹, M. Mallar Chakravarty^{42,43}, Qiang Chen⁴⁴, Christopher R.K. Ching^{1,45}, Vincent Chouraki^{36,38,46}, Gabriel Cuellar-Partida⁸, Fabrice Crivello⁴⁷, Anouk Den Braber⁴⁸, Nhat Trung Doan³¹, Stefan Ehrlich^{49,50,51}, Sudheer Giddaluru^{52,53}, Aaron L. Goldman⁴⁴, Rebecca F. Gottesman⁵⁴, Oliver Grimm⁵⁵, Michael E. Griswold⁵⁶, Tulio Guadalupe^{57,58}, Boris A. Gutman¹, Johanna Hass⁵⁹, Unn K. Haukvik^{31,60}, David Hoehn⁶¹, Avram J. Holmes^{50,62}, Martine Hoogman^{10,13}, Deborah Janowitz⁶³, Tianye Jia¹⁸, Kjetil N. Jørgensen^{31,60}, Nazanin Karbalai⁶¹, Dalia Kasperaviciute^{64,65}, Sungeun Kim^{66,67,68}, Marieke Klein^{10,13}, Bernd Kraemer⁶⁹, Phil H. Lee^{50,70,71,72,73}, David C.M. Liewald⁷⁴, Lorna M. Lopez⁷⁴, Michelle Luciano⁷⁴, Christine Macare¹⁸, Andre F. Marquand^{13,75}, Mar Matarin^{64,76}, Karen A. Mather²⁹, Manuel Mattheisen^{77,78,79}, David R. McKay^{80,81}, Yuri Milaneschi⁸², Susana Muñoz Maniega^{26,28,74}, Kwangsik Nho^{66,67,68}, Allison C. Nugent⁸³, Paul Nyquist⁸⁴, Loes M. Olde Loohuis⁸⁵, Jaap Oosterlaan⁸⁶, Martina Papmeyer^{87,88}, Lukas Pirpamer⁶, Benno Pütz⁶¹, Adaikalavan Ramasamy^{76,89,90}, Jennifer S. Richards^{12,13,91}, Shannon L. Risacher^{66,68}, Roberto Roiz-Santiañez^{92,93}, Nanda Rommelse^{11,13,91}, Stefan Ropele⁶, Emma J. Rose⁹⁴, Natalie A. Royle^{26,28,74,95}, Tatjana Rundek^{96,97}, Philipp G. Sämann⁶¹, Arvin Saremi¹, Claudia L. Satizabal^{36,38}, Lianne Schmaal^{98,99,100}, Andrew J. Schork^{101,102}, Li Shen^{66,67,68}, Jean Shin³⁹, Elena Shumskaya^{10,13,75}, Albert V. Smith^{103,104}, Emma Sprooten^{80,81,105}, Lachlan T. Strike^{8,106}, Alexander Teumer¹⁰⁷, Diana Tordesillas-Gutierrez^{93,108}, Roberto Toro¹⁰⁹, Daniah Trabzuni^{76,110}, Stella Trompet¹¹¹, Dhananjay Vaidya¹¹², Jeroen Van der Grond¹¹³, Sven J. Van der Lee², Dennis Van der Meer¹¹⁴, Marjolein M.J. Van Donkelaar^{10,13}, Kristel R. Van Eijk¹¹⁵, Theo G.M. Van Erp¹¹⁶, Daan Van Rooij^{12,13,114}, Esther Walton^{49,117}, Lars T. Westlye^{32,117}, Christopher D. Whelan^{1,21}, Beverly G. Windham¹¹⁸, Anderson M. Winkler^{80,119}, Katharina Wittfeld^{63,120}, Girma Woldehawariat⁸³, Christiane Wolf¹²¹, Thomas Wolfers^{10,13}, Lisa R. Yanek¹¹², Jingyun Yang^{24,122}, Alex Zijdenbos¹²³, Marcel P. Zwiers^{13,75}, Ingrid Agartz^{31,60,124}, Laura Almasy^{125,126,127}, David Ames^{128,129}, Philippe Amouyel⁴⁶, Ole A. Andreassen^{31,32}, Sampath Arepalli¹³⁰, Amelia A. Assareh²⁹, Sandra Barral⁴⁰, Mark E. Bastin^{26,28,74,95}, Diane M. Becker¹¹², James T. Becker¹³¹, David A. Bennett^{24,122}, John Blangero¹²⁵, Hans van Bokhoven^{10,13}, Dorret I. Boomsma⁴⁸, Henry Brodaty^{29,132}, Rachel M. Brouwer¹⁹, Han G. Brunner^{10,13,133}, Randy L. Buckner^{50,134}, Jan K. Buitelaar^{12,13,91}, Kazima B. Bulayeva¹³⁵, Wiepke Cahn¹⁹, Vince D. Calhoun^{136,137}, Dara M. Cannon^{83,138}, Gianpiero L. Cavalleri²¹, Ching-Yu Cheng^{14,15,139}, Sven Cichon^{140,141,142}, Mark R. Cookson¹³⁰, Aiden Corvin⁹⁴, Benedicto Crespo-Facorro^{92,93}, Joanne E. Curran¹²⁵, Michael Czisch⁶¹, Anders M. Dale^{143,144}, Gareth E. Davies¹⁴⁵, Anton J.M. De Craen¹⁴⁶, Eco J.C. De Geus⁴⁸, Philip L. De Jager^{71,147,148,149,150}, Greig I. De Zubicaray¹⁵¹, Ian J. Deary⁷⁴, Stéphanie Debette^{4,36,152}, Charles DeCarli¹⁵³, Norman Delanty^{21,154}, Chantal Depondt¹⁵⁵, Anita DeStefano^{37,38}, Allissa Dillman¹³⁰, Srdjan Djurovic^{52,156}, Gary Donohoe^{157,158}, Wayne C. Drevets^{83,159}, Ravi Duggirala¹²⁵, Thomas D. Dyer¹²⁵, Christian Enzinger⁶, Susanne Erk¹⁶⁰, Thomas Espeseth^{32,117}, Iryna O. Fedko⁴⁸, Guillén Fernández^{12,13}, Luigi Ferrucci¹⁶¹, Simon E. Fisher^{13,57}, Debra A. Fleischman^{24,162}, Ian Ford¹⁶³, Myriam Fornage¹⁶⁴, Tatiana M. Foroud^{68,165}, Peter T. Fox¹⁶⁶, Clyde Francks^{13,57}, Masaki Fukunaga¹⁶⁷, J. Raphael Gibbs^{76,130}, David C. Glahn^{80,81}, Randy L. Gollub^{50,51,71}, Harald H.H. Göring¹²⁵, Robert C. Green^{71,168}, Oliver Gruber⁶⁹, Vilmundur Gudnason^{103,104}, Sebastian Guelfi⁷⁶, Asta K. Håberg^{169,170}, Narelle K. Hansell^{8,106}, John Hardy⁷⁶, Catharina A. Hartman¹¹⁴, Ryota Hashimoto^{171,172}, Katrin Hegenscheid¹⁷³, Andreas Heinz¹⁶⁰, Stephanie Le Hellard^{52,53}, Dena G. Hernandez^{76,130,174}, Dirk J. Heslenfeld¹⁷⁵, Beng-Choon Ho¹⁷⁶, Pieter J. Hoekstra¹¹⁴, Wolfgang Hoffmann^{107,120},

Albert Hofman¹⁷⁸, Florian Holsboer^{61,177}, Georg Homuth¹⁷⁸, Norbert Hosten¹⁷³, Jouke-Jan Hottenga⁴⁸, Matthew Huentelman¹⁷⁹, Hilleke E. Hulshoff Pol¹⁹, Masashi Ikeda¹⁸⁰, Clifford R. Jack Jr¹⁸¹, Mark Jenkinson¹¹⁹, Robert Johnson¹⁸², Erik G. Jönsson^{31,124}, J. Wouter Jukema¹¹¹, René S. Kahn¹⁹, Ryota Kanai^{183,184,185}, Iwona Kloszewska¹⁸⁶, David S. Knopman¹⁸⁷, Peter Kochunov¹⁸⁸, John B. Kwok^{189,190}, Stephen M. Lawrie⁸⁷, Hervé Lemaître¹⁹¹, Xinmin Liu^{83,192}, Dan L. Longo¹⁹³, Oscar L. Lopez¹⁹⁴, Simon Lovestone^{195,196}, Oliver Martinez¹⁵³, Jean-Luc Martinot¹⁹¹, Venkata S. Mattay^{44,54,197}, Colm McDonald¹³⁸, Andrew M. McIntosh^{74,87}, Francis J. McMahon⁸³, Katie L. McMahon¹⁹⁸, Patrizia Mecocci¹⁹⁹, Ingrid Melle^{31,32}, Andreas Meyer-Lindenberg⁵⁵, Sebastian Mohnke¹⁶⁰, Grant W. Montgomery⁸, Derek W. Morris¹⁵⁷, Thomas H. Mosley¹¹⁸, Thomas W. Mühleisen^{141,142}, Bertram Müller-Myhsok^{61,200,201}, Michael A. Nalls¹³⁰, Matthias Nauck^{202,203}, Thomas E. Nichols^{119,204}, Wiro J. Niessen^{3,205,206}, Markus M. Nöthen^{141,207}, Lars Nyberg²², Kazutaka Ohi¹⁷¹, Rene L. Olvera¹⁶⁶, Roel A. Ophoff^{19,85}, Massimo Pandolfo¹⁵⁵, Tomas Paus^{208,209,210}, Zdenka Pausova^{39,211}, Brenda W.J. H. Penninx¹⁰⁰, G. Bruce Pike^{212,213}, Steven G. Potkin¹¹⁶, Bruce M. Psaty²¹⁴, Simone Reppermund^{29,215}, Marcella Rietschel⁵⁵, Joshua L. Roffman⁵⁰, Nina Romanczuk-Seiferth¹⁶⁰, Jerome I. Rotter²¹⁶, Mina Ryten^{76,89}, Ralph L. Sacco^{35,96,97,217}, Perminder S. Sachdev^{29,218}, Andrew J. Saykin^{66,68,165}, Reinhold Schmidt⁶, Helena Schmidt²¹⁹, Peter R. Schofield^{189,190}, Sigurdur Sigursson¹⁰³, Andrew Simmons^{220,221,222}, Andrew Singleton¹³⁰, Sanjay M. Sisodiya⁶⁴, Colin Smith²²³, Jordan W. Smoller^{50,70,71,72}, Hilkka Soininen^{224,225}, Vidar M. Steen^{52,53}, David J. Stott²²⁶, Jessika E. Sussmann⁸⁷, Anbupalam Thalamuthu²⁹, Arthur W. Toga²²⁷, Bryan J. Traynor¹³⁰, Juan Troncoso²²⁸, Magda Tsolaki²²⁹, Christophe Tzourio^{4,230}, Andre G. Uitterlinden^{2,231}, Maria C. Valdés Hernández^{26,28,74,95}, Marcel Van der Brug²³², Aad van der Lugt³, Nic J.A. van der Wee²³³, Neeltje E.M. Van Haren¹⁹, Dennis van 't Ent⁴⁸, Marie-Jose Van Tol²³⁴, Badri N. Vardarajan⁴⁰, Bruno Vellas²³⁵, Dick J. Veltman¹⁰⁰, Henry Völzke¹⁰⁷, Henrik Walter¹⁶⁰, Joanna M. Wardlaw^{26,28,74,95}, Thomas H. Wassink²³⁶, Michael E. Weale⁸⁹, Daniel R. Weinberger^{44,237}, Michael W. Weiner²³⁸, Wei Wen^{29,218}, Eric Westman²³⁹, Tonya White^{3,240}, Tien Y. Wong^{14,15,139}, Clinton B. Wright^{96,97,217}, Ronald H. Zielke¹⁸², Alan B. Zonderman²⁴¹, Nicholas G. Martin⁸, Cornelia M. Van Duijn², Margaret J. Wright^{106,198}, W.T. Longstreth²⁴², Gunter Schumann^{18,**}, Hans J. Grabe^{63,**}, Barbara Franke^{10,11,13,**}, Lenore J. Launer^{243,**}, Sarah E. Medland^{8,**}, Sudha Seshadri^{36,38,**}, Paul M. Thompson^{1,**} & M. Arfan Ikram^{2,3,244,**}

¹Imaging Genetics Center, USC Mark and Mary Stevens Neuroimaging & Informatics Institute, Keck School of Medicine of University of Southern California, Los Angeles, California 90292, USA. ² Department of Epidemiology, Erasmus University Medical Center, 3015 CE Rotterdam, The Netherlands. ³ Department of Radiology and Nuclear Medicine, Erasmus MC, 3015 CE Rotterdam, The Netherlands. ⁴ INSERM Unit U1219, University of Bordeaux, 33076 Bordeaux, France. ⁵ Department of Genetics & UNC Neuroscience Center, University of North Carolina (UNC), Chapel Hill, North Carolina, 27599, USA. ⁶ Department of Neurology, Clinical Division of Neurogeriatrics, Medical University Graz, Auenbruggerplatz 22, 8036 Graz, Austria.⁷ Institute of Medical Informatics, Statistics and Documentation, Medical University Graz, Auenbruggerplatz 22, 8036 Graz, Austria. ⁸ QIMR Berghofer Medical Research Institute, Brisbane, Queensland 4006, Australia. ⁹Cardiovascular Health Research Unit, Department of Medicine, University of Washington, 1730 Minor Avenue/Suite 1360. Seattle, Washington 98101, USA. ¹⁰ Department of Human Genetics, Radboud University Medical Center, 6525 GA Niimegen, The Netherlands. ¹¹ Department of Psychiatry, Radboud University Medical Center, 6525 GA Nijmegen, The Netherlands. ¹² Department of Cognitive Neuroscience, Radboud University Medical Center, 6525 GA Nijmegen, The Netherlands. ¹³ Donders Institute for Brain, Cognition and Behaviour, Radboud University, 6525 GA Nijmegen, The Netherlands. ¹⁴ Academic Medicine Research Institute, Duke-NUS Graduate Medical School, Singapore, 169857, Singapore. ¹⁵ Singapore Eye Research Institute, Singapore National Eye Centre, Singapore, 168751, Singapore. ¹⁶ Memory Aging & Cognition Centre (MACC), National University Health System, Singapore, 119228, Singapore. ¹⁷ Department of Pharmacology, National University of Singapore, Singapore, 119077, Singapore. ¹⁸ MRC-SGDP Centre, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE5 8AF, UK.¹⁹ Brain Center Rudolf Magnus, Department of Psychiatry, UMC Utrecht, 3584 CX Utrecht, The Netherlands. ²⁰ Department of Neurology and Neurosurgery, Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada H3A 2B4. ²¹ The Royal College of Surgeons in Ireland, 123 St Stephen's Green, Dublin 2, Ireland. ²² Department of Integrative Medical Biology and Umeå Center for Functional Brain Imaging, Umeå University, 901 87 Umeå, Sweden. 23 Department of Biomedical Engineering, Illinois Institute of Technology, Chicago, Illinois 60616, USA. ²⁴ Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, Illinois 60612, USA. ²⁵ Department of Diagnostic Radiology and Nuclear Medicine, Rush University Medical Center, Chicago, Illinois 60616, USA. ²⁶ Brain Research Imaging Centre, University of Edinburgh, Edinburgh EH4 2XU, UK. ²⁷ Department of Computer Science, Lagos State University, Lagos, P.M.B. 01 LASU, Nigeria. ²⁸ Scottish Imaging Network, A Platform for Scientific Excellence (SINAPSE) Collaboration, Department of Neuroimaging Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK. ²⁹ Centre for Healthy Brain Ageing, School of Psychiatry, University of New South Wales, Sydney, New South Wales 2052, Australia. ³⁰ Mathematics and Statistics, Murdoch University, Perth, Western Australia, 6150, Australia. ³¹ NORMENT—KG Jebsen Centre, Institute of Clinical Medicine, University of Oslo, 0315 Oslo, Norway. 32 NORMENT—KG Jebsen Centre, Division of Mental Health and Addiction, Oslo University Hospital, 0424 Oslo, Norway. ³³ Department of Medical Sciences, Molecular Medicine and Science for Life Laboratory, Uppsala University, Box 1432, SE-751 44 Uppsala, Sweden. ³⁴ Dr John T. Macdonald Foundation Department of Human Genetics, University of Miami, Miller School of Medicine, Miami, Florida, 33136, USA.

³⁵ John P. Hussman Institute for Human Genomics, University of Miami, Miller School of Medicine, Miami, Florida, 33136, USA. ³⁶ Department of Neurology, Boston University School of Medicine, Boston, Massachusetts,02118, USA. 37 Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts 02118 USA. ³⁸ Framingham Heart Study, 17 Mount Wayte Avenue, Framingham, Massachusetts 01703 USA. ³⁹ Hospital for Sick Children, University of Toronto, Toronto, Ontario, Canada M5G 1X8, ⁴⁰ Taub Institute for Research on Alzheimer's Disease and the Aging Brain: G.H. Sergievsky Center: Department of Neurology, Columbia University Medical Center, 639 West 1168th Street, New York, New York 10032, USA. ⁴¹ Pennington Biomedical Research Center, Baton Rouge, Louisiana 70808, USA. ⁴² Cerebral Imaging Centre, Douglas Mental Health University Institute, Montreal, Quebec, Canada H4H 1R3. ⁴³ Department of Psychiatry and Biomedical Engineering, McGill University, Montreal, Quebec, Canada H3A 2B4. ⁴⁴ Lieber Institute for Brain Development, Baltimore, Maryland 21205, USA. ⁴⁵ Interdepartmental Neuroscience Graduate Program, UCLA School of Medicine, Los Angeles, California 90095, USA. 46 Lille University, Inserm, CHU Lille, Institut Pasteur de Lille, U1167-RID-AGE-Risk factors and molecular determinants of aging-related diseases, F-59000 Lille, France. ⁴⁷ IMN UMR5293, GIN, CNRS, CEA, University of Bordeaux, 146 rue Léo Saignat, 33076 Bordeaux, France. ⁴⁸ Biological Psychology, Amsterdam Neuroscience, Vrije Universiteit & Vrije Universiteit Medical Center, 1081 BT Amsterdam, The Netherlands. ⁴⁹ Division of Psychological and Social Medicine and Developmental Neurosciences, Faculty of Medicine, TU Dresden, 01307 Dresden, Germany. ⁵⁰ Department of Psychiatry, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. ⁵¹ Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown, Massachusetts 02129, USA. 52 NORMENT—KG Jebsen Centre for Psychosis Research, Department of Clinical Science, University of Bergen, 5021 Bergen, Norway. ⁵³ Dr Einar Martens Research Group for Biological Psychiatry, Center for Medical Genetics and Molecular Medicine, Haukeland University Hospital, 5021 Bergen, Norway.⁵⁴ Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland 21287, USA. ⁵⁵Central Institute of Mental Health, Medical Faculty Mannheim, University Heidelberg, 68159 Mannheim, Germany. ⁵⁶Department of Data Science, University of Mississippi Medical Center, Jackson, Mississippi, 39216, USA. ⁵⁷ Language and Genetics Department, Max Planck Institute for Psycholinguistics, 6525 XD Nijmegen, The Netherlands. ⁵⁸ International Max Planck Research School for Language Sciences, 6525 XD Nijmegen, The Netherlands. ⁵⁹ Department of Child and Adolescent Psychiatry, Faculty of Medicine of the TU Dresden, 01307 Dresden, Germany. ⁶⁰ Department of Research and Development, Diakonhjemmet Hospital, 0319 Oslo, Norway. ⁶¹ Max Planck Institute of Psychiatry, 80804 Munich, Germany. ⁶² Department of Psychology, Yale University, New Haven, Connecticut 06520, USA. ⁶³ Department of Psychiatry, University Medicine Greifswald, 17489 Greifswald, Germany. ⁶⁴ UCL Institute of Neurology, London, United Kingdom and Epilepsy Society, Bucks, SL9 ORJ, UK. ⁶⁵ Department of Medicine, Imperial College London, London SW7 2AZ, UK. ⁶⁶ Center for Neuroimaging, Radiology and Imaging Sciences, Indiana University School of Medicine, Indianapolis, Indiana 46202, USA. ⁶⁷ Center for Computational Biology and Bioinformatics, Indiana University School of Medicine, Indianapolis, Indiana 46202, USA. 68 Indiana Alzheimer Disease Center, Indiana University School of Medicine, Indianapolis, Indiana 46202, USA. 69 Section for Experimental Psychopathology and Neuroimaging, Department of General Psychiatry, Heidelberg University, Heidelberg, 69120, Germany. ⁷⁰ Psychiatric and Neurodevelopmental Genetics Unit, Center for Human Genetic Research, Massachusetts General Hospital, Boston, Massachusetts 02114, USA. ⁷¹ Harvard Medical School, Boston, Massachusetts 02115, USA. ⁷² Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Boston, Massachusetts 02141, USA. ⁷³ Lurie Center for Autism, Massachusetts General Hospital, Harvard Medical School, Lexington, Massachusetts, 02421, USA. ⁷⁴ Centre for Cognitive Ageing and Cognitive Epidemiology, Psychology, University of Edinburgh, Edinburgh EH8 9JZ, UK. ⁷⁵ Donders Centre for Cognitive Neuroimaging, Radboud University, Nijmegen, 6525 EN, The Netherlands. ⁷⁶ Reta Lila Weston Institute and Department of Molecular Neuroscience, UCL Institute of Neurology, London WC1N 3BG, UK. ⁷⁷ Department of Biomedicine, Aarhus University, DK-8000 Aarhus, Denmark. ⁷⁸ The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, DK-8000 Aarhus and Copenhagen, Denmark. ⁷⁹ Center for integrated Sequencing, iSEQ, Aarhus University, DK-8000 Aarhus, Denmark. ⁸⁰ Department of Psychiatry, Yale University, New Haven, Connecticut 06511, USA. ⁸¹ Olin Neuropsychiatric Research Center, Hartford, Connecticut 06114, USA. ⁸² Department of Psychiatry, EMGO Institute for Health and Care Research and Neuroscience Campus Amsterdam, VU University Medical Center/GGZ inGeest, 1081 HL Amsterdam, The Netherlands. ⁸³ Human Genetics Branch, National Institute of Mental Health Intramural Research Program, 35 Convent Drive, Rm 1A202, Bethesda, Marvland 20892-3719, USA. ⁸⁴ Department of Neurology, Department of Anesthesia/Critical Care Medicine, Department of Neurosurgery, Johns Hopkins, USA600 N. Wolfe St, Baltimore, Maryland 21287, USA.⁸⁵ Center for Neurobehavioral Genetics, University of California, Los Angeles, California 90095, USA. ⁸⁶ Department of Clinical Neuropsychology, VU University Amsterdam, Amsterdam, 1081 HV, The Netherlands. ⁸⁷ Division of Psychiatry, Royal Edinburgh Hospital, University of Edinburgh, Edinburgh EH10 5HF, UK. ⁸⁸ Division of Systems Neuroscience of Psychopathology, Translational Research Center, University Hospital of Psychiatry, University of Bern, Bern, 3060, Switzerland. ⁸⁹ Department of Medical and Molecular Genetics, King's College London, London SE1 9RT, UK. ⁹⁰ The Jenner Institute Laboratories, University of Oxford, Oxford OX3 7DQ, UK. ⁹¹ Karakter Child and Adolescent Psychiatry University Center, Nijmegen, 6525 GC, The Netherlands. ⁹² Department of Medicine and Psychiatry, University Hospital Marqués de Valdecilla, School of Medicine, University of Cantabria-IDIVAL, 39008 Santander, Spain. 93 CIBERSAM (Centro Investigación Biomédica en Red Salud Mental), Santander, 39011, Spain. ⁹⁴ Psychosis Research Group, Department of Psychiatry & Trinity Translational Medicine Institute, Trinity College, Dublin, Dublin 2, Ireland. ⁹⁵ Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK. 96 Department of Neurology, University of Miami, Miller School of Medicine, Miami, Florida, 33136, USA. 97 Department of Epidemiology and Public Health Sciences, University of Miami, Miller School of Medicine, Miami, Florida, 33136, USA. ⁹⁸ Orygen, The National Centre of Excellence in Youth Mental Health, Melbourne, Victoria, 3502, Australia. ⁹⁹ Centre for Youth Mental Health, The University of Melbourne, Melbourne, Victoria, 3502, Australia.¹⁰⁰ Department of Psychiatry, Neuroscience Campus Amsterdam, VU University Medical Center, 1007 MB Amsterdam, The Netherlands.¹⁰¹ Multimodal Imaging Laboratory, Department of Neurosciences, University of California, San Diego, California 92093, USA. ¹⁰² Department of Cognitive Sciences, University of California, San Diego, California 92161, USA. ¹⁰³ Icelandic Heart Association, Kopavogur, 201, Iceland. ¹⁰⁴ Faculty of Medicine, University of Iceland, Reykjavik, 101, Iceland. ¹⁰⁵ Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, New York, 10029, USA. ¹⁰⁶ Queensland Brain Institute, University of Queensland, Brisbane, Queensland 4072, Australia. ¹⁰⁷ Institute for Community Medicine, University Medicine Greifswald, 17489 Greifswald, Germany. ¹⁰⁸ Neuroimaging Unit, Technological Facilities. Valdecilla Biomedical Research Institute IDIVAL, Santander, Cantabria, 39011, Spain. ¹⁰⁹ Institut Pasteur, 75015 Paris, France. ¹¹⁰ Department of Genetics, King Faisal Specialist Hospital and Research Centre, Riyadh 11211, Saudi Arabia.¹¹¹ Department of Cardiology, Leiden University Medical Center, Leiden, 2300RC, The Netherlands.¹¹² GeneSTAR Research Center, Department of Medicine, Johns Hopkins University School of Medicine, 1830 E Monument St Suite 8028, Baltimore, Maryland 21287, USA. ¹¹³ Department of Radiology, Leiden University Medical Center, Leiden, 2300RC, The Netherlands. ¹¹⁴ Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, 9700RB, The Netherlands. ¹¹⁵ Brain Center Rudolf Magnus, Human Neurogenetics Unit, UMC Utrecht, 3584 CG Utrecht, The Netherlands. ¹¹⁶ Department of Psychiatry and Human Behavior, University of California-Irvine, Irvine, California 92617, USA. ¹¹⁷ Department of Psychology, Georgia State University, Atlanta, Georgia 30302, USA. ¹¹⁸ NORMENT—KG Jebsen Centre, Department of Psychology, University of Oslo, 0317 Oslo, Norway. ¹¹⁹ Department of Medicine, University of Mississippi Medical Center, Jackson, Mississippi, 39216, USA. ¹²⁰ FMRIB Centre, University of Oxford, Oxford OX3 9DU, UK. ¹²¹German Center for Neurodegenerative Diseases (DZNE) Rostock/Greifswald, 17487 Greifswald, Germany. ¹²² University of Wuerzburg, Department of Psychiatry, Psychosomatics and Psychotherapy, Wuerzburg, 97080, Germany. ¹²³ Department of Neurological Sciences, Rush University Medical Center, Chicago, Illinois 60612, USA. ¹²⁴ Biospective Inc, Montreal, Quebec, Canada, 6100 Avenue Royalmount, Montréal, Québec, Canada H4P 2R2.¹²⁵ Department of Clinical Neuroscience, Centre for Psychiatric Research, Karolinska Institutet, SE-171 77 Stockholm, Sweden. ¹²⁶ South Texas Diabetes and Obesity Institute, University of Texas Rio Grande Valley School of Medicine, Brownsville/Edinburg/San Antonio, Texas, 78250, USA. 127 Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA.

¹²⁸ Department of Biomedical and Health Informatics, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania 29104, USA. ¹²⁹ National Ageing Research Institute, Royal Melbourne Hospital, Melbourne, Victoria 3052, Australia. ¹³⁰ Academic Unit for Psychiatry of Old Age, University of Melbourne, Melbourne, Victoria 3101, Australia. ¹³¹Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, Maryland 20892, USA. ¹³² Departments of Psychiatry, Neurology, and Psychology, University of Pittsburgh, 3501 Forbes Ave., Suite 830, Pittsburgh, Pennsylvania 15213, USA. ¹³³ Dementia Collaborative Research Centre—Assessment and Better Care, University of New South Wales, Sydney, New South Wales 2052, Australia. 134 Department of Clinical Genetics and GROW School for Oncology and Developmental Biology, Maastricht University Medical Center, 6200 MD Maastricht, The Netherlands. ¹³⁵ Department of Psychology, Center for Brain Science, Harvard University, Cambridge, Massachusetts 02138, USA. ¹³⁶ Department of Evolution and Genetics, Dagestan State University, Makhachkala 367000, Dagestan, Russia. ¹³⁷ The Mind Research Network & LBERI, Albuquerque, New Mexico 87106, USA. ¹³⁸ Department of ECE, University of New Mexico, Albuquerque, New Mexico 87131, USA. ¹³⁹ Centre for Neuroimaging & Cognitive Genomics (NICOG), Clinical Neuroimaging Laboratory, NCBES Galway Neuroscience Centre, College of Medicine Nursing and Health Sciences, National University of Ireland Galway, H91 TK33 Galway, Ireland. ¹⁴⁰ Department of Ophthalmology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, 119077, Singapore. ¹⁴¹ Division of Medical Genetics, Department of Biomedicine, University of Basel, 4031 Basel, Switzerland. ¹⁴² Institute of Human Genetics, University of Bonn, 53127 Bonn, Germany. ¹⁴³ Institute of Neuroscience and Medicine (INM-1), Research Centre Jülich, 52425 Jülich, Germany.¹⁴⁴ Center for Multimodal Imaging and Genetics, University of California, San Diego, California 92093, USA.¹⁴⁵ Departments of Neurosciences, Radiology, Psychiatry, and Cognitive Science, University of California, San Diego, California 92093, USA. 146 Avera Institute for Human Genetics, Sioux Falls, South Dakota 57108, USA. ¹⁴⁷ Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, 2300RC, The Netherlands. ¹⁴⁸ Program in Translational NeuroPsychiatric Genomics, Departments of Neurology & Psychiatry, Brigham and Women's Hospital, Boston, Massachusetts, 02115, USA. ¹⁴⁹ Harvard Medical School, Boston, Massachusetts, 02115, USA. ¹⁵⁰ Program in Medical and Population Genetics, Broad Institute, Cambridge, Massachusetts, 02142, USA. 151 Broad Institute, Cambridge, Massachusetts, 02142, USA. 152 Faculty of Health and Institute of Health and Biomedical Innovation, Queensland University of Technology (QUT), Brisbane, Queensland 4059, Australia. ¹⁵³ Department of Neurology, Bordeaux University Hospital, Bordeaux, 33076, France. 154 Imaging of Dementia and Aging (IDeA) Laboratory, Department of Neurology and Center for Neuroscience, University of California at Davis, 4860 Y Street, Suite 3700, Sacramento, California 95817, USA. 155 Neurology Division, Beaumont Hospital, Dublin 9, Ireland, ¹⁵⁶ Department of Neurology, Hopital Erasme, Universite Libre de Bruxelles, 1070 Brussels, Belgium. ¹⁵⁷ Department of Medical Genetics, Oslo University Hospital, 0420 Oslo, Norway.¹⁵⁸ Cognitive Genetics and Cognitive Therapy Group, Neuroimaging, Cognition & Genomics Centre (NICOG) & NCBES Galway Neuroscience Centre, School of Psychology and Discipline of Biochemistry, National University of Ireland Galway, H91 TK33, Galway, Ireland. ¹⁵⁹ Neuropsychiatric Genetics Research Group, Department of Psychiatry and Trinity College Institute of Psychiatry, Trinity College Dublin, Dublin 8, Ireland. ¹⁶⁰ Janssen Research & Development, LLC, Titusville, New Jersey 08560, USA. ¹⁶¹ Charité - Universitätsmedizin Berlin, Campus Charité Mitte, Department of Psychiatry and Psychotherapy, Charitéplatz 1, 10117 Berlin, Germany.¹⁶² Intramural Research Program of the National Institute on Aging, Baltimore, Maryland, 21224, USA. ¹⁶³ Department of Neurological Sciences & Department of Behavioral Sciences, Rush University Medical Center, Chicago, Illinois 60616, USA. ¹⁶⁴ Robertson Centre for Biostatistics, University of Glasgow, Glasgow, G41 4DQ, UK. ¹⁶⁵ Institute of Molecular Medicine and Human Genetics Center, University of Texas Health Science Center at Houston, Houston, Texas, 77030, USA. ¹⁶⁶ Medical and Molecular Genetics, Indiana University School of Medicine, Indianapolis, Indiana 46202, USA. ¹⁶⁷ University of Texas Health Science Center, San Antonio, Texas 78229, USA. ¹⁶⁸ Division of Cerebral Integration, National Institute for Physiological Sciences, Aichi, 444-8585, Japan.¹⁶⁹ Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Boston, Massachusetts 02115, USA. ¹⁷⁰ Department of Neuroscience, Faculty of Medicine, Norwegian University of Science and Technology (NTNU), Trondheim, 7491, Norway. ¹⁷¹ Department of Radiology, St. Olav's Hospital, Trondheim University Hospital, Trondheim, 7030, Norway. ¹⁷² Department of Psychiatry, Osaka University Graduate School of Medicine, Osaka 565-0871, Japan. ¹⁷³ Molecular Research Center for Children's Mental Development, United Graduate School of Child Development, Osaka University, Osaka, 565-0871, Japan. ¹⁷⁴ Institute of Diagnostic Radiology and Neuroradiology, University Medicine Greifswald, 17489 Greifswald, Germany. ¹⁷⁵ German Center for Neurodegenerative Diseases (DZNE), Tübingen, 72076, Germany. ¹⁷⁶ Department of Psychology, VU University Amsterdam, 1081 BT Amsterdam, The Netherlands. ¹⁷⁷ Department of Psychiatry, University of Iowa, lowa City, Iowa 52242, USA. ¹⁷⁸ Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts 02115 USA. ¹⁷⁹ HMNC Brain Health, Munich, 80539, Germany. ¹⁸⁰ Interfaculty Institute for Genetics and Functional Genomics, University Medicine Greifswald, 17489 Greifswald, Germany. ¹⁸¹ Translational Genomics Research Institute, Neurogenomics Division, 445N Fifth Street, Phoenix, Arizona 85004, USA. ¹⁸² Department of Psychiatry, Fujita Health University School of Medicine, Toyoake 470-1192, Japan.¹⁸³ Department of Radiology, Mayo Clinic, Rochester, Minnesota 55905, USA. ¹⁸⁴ NICHD Brain and Tissue Bank for Developmental Disorders, University of Maryland Medical School, Baltimore, Maryland 21201, USA. ¹⁸⁵ School of Psychology, University of Sussex, Brighton BN1 9QH, UK. ¹⁸⁶ Institute of Cognitive Neuroscience, University College London, London WC1N 3AR, UK. ¹⁸⁷ Department of Neuroinformatics, Araya Brain Imaging, Tokyo, 102-0093, Japan. ¹⁸⁸ Medical University of Lodz, 90-419 Lodz, Poland. ¹⁸⁹ Department of Neurology, Mayo Clinic, Rochester, Minnesota, 55905, USA. ¹⁹⁰ Maryland Psychiatric Research Center, Department of Psychiatry, University of Maryland School of Medicine, Baltimore, Maryland, 21228, USA. 191 Neuroscience Research Australia, Sydney, New South Wales 2031, Australia. 192 School of Medical Sciences, UNSW, Sydney, New South Wales 2052, Australia. ¹⁹³ INSERM UMR 1000 "Neuroimaging and Psychiatry", Service Hospitalier Frédéric Joliot; University Paris-Sud, Université Paris-Saclay, University Paris Descartes, Maison de Solenn, Paris, 91400, France. ¹⁹⁴Columbia University Medical Center, New York, New York 10032, USA. ¹⁹⁵ Laboratory of Genetics, National Institute on Aging, National Institutes of Health, Baltimore, Maryland 21224, USA. ¹⁹⁶ Departments of Neurology and Psychiatry, University of Pittsburgh, 3501 Forbes Ave., Suite 830, Pittsburgh Pennsylvania 15213, USA. ¹⁹⁷ Department of Psychiatry, University of Oxford, Oxford OX3 7JX, UK. ¹⁹⁸ Department of Radiology, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA. ¹⁹⁹ Centre for Advanced Imaging, University of Queensland, Brisbane, Queensland 4072, Australia. ²⁰⁰ Section of Gerontology and Geriatrics, Department of Medicine, University of Perugia, 06132 Perugia, Italy. ²⁰¹ Munich Cluster for Systems Neurology (SyNergy), 81377 Munich, Germany. ²⁰² University of Liverpool, Institute of Translational Medicine, Liverpool L69 3BX, UK. ²⁰³ Institute of Clinical Chemistry and Laboratory Medicine, University Medicine Greifswald, 17489 Greifswald, Germany. ²⁰⁴ German Center for Cardiovascular Research (DZHK e.V.), partner site Greifswald, Greifswald, 17475, Germany. 205 Department of Statistics & WMG, University of Warwick, Coventry CV4 7AL, UK. 206 Department of Medical Informatics Erasmus MC, 3015 CE Rotterdam, The Netherlands. ²⁰⁷ Faculty of Applied Sciences, Delft University of Technology, Delft, 2628 CD, The Netherlands. ²⁰⁸ Department of Genomics, Life & Brain Center, University of Bonn, 53127 Bonn, Germany. ²⁰⁹ Rotman Research Institute, University of Toronto, Toronto, Ontario, Canada M6A 2E1. ²¹⁰ Departments of Psychology and Psychiatry, University of Toronto, Toronto, Ontario, Canada M5T 1R8. ²¹¹ Child Mind Institute, New York, New York, 10022, USA. ²¹² Departments of Physiology and Nutritional Sciences, University of Toronto, Toronto, Ontario, Canada M5S 3E2. ²¹³ Department of Radiology, University of Calgary, Calgary, Alberta, Canada T2N 4N1.²¹⁴ Department of Clinical Neuroscience, University of Calgary, Calgary, Alberta, Canada T2N 4N1. ²¹⁵ Departments of Epidemiology, Medicine and Health Services, University of Washington, Seattle, WA, USA Group Health Research Institute, Group Health, 1730 Minor Avenue/Suite 1360, Seattle, Washington 98101, USA. ²¹⁶ Department of Developmental Disability Neuropsychiatry, School of Psychiatry, University of New South Wales, Sydney, New South Wales 2052, Australia. ²¹⁷ Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute and Pediatrics at Harbor-UCLA Medical Center, Torrance, California 90502, USA. ²¹⁸ Evelyn F. McKnight Brain Institute, University of Miami, Miller School of Medicine, Miami, Florida, 33136, USA. ²¹⁹ Neuropsychiatric Institute, Prince of Wales Hospital, Randwick, New South Wales 2031, Australia. ²²⁰ Institute of Molecular Biology and Biochemistry, Medical University Graz, Harrachgasse 21/III, 8010 Graz, Austria.

²²¹Department of Neuroimaging, Institute of Psychiatry, King's College London, London SE5 8AF, UK. ²²² Biomedical Research Centre for Mental Health, King's College London, London SE5 8AF, UK. 223 Biomedical Research Unit for Dementia, King's College London, London SE5 8AF, UK. 224 MRC Edinburgh Brain Bank, University of Edinburgh, Academic Department of Neuropathology, Centre for Clinical Brain Sciences, Edinburgh, EH16 4SB UK. 225 Institute of Clinical Medicine, Neurology, University of Eastern Finland, FI-70211 Kuopio, Finland. ²²⁶ Neurocentre Neurology, Kuopio University Hospital, FI-70211 Kuopio, Finland. ²²⁷ Institute of Cardiovascular and Medical Sciences, Faculty of Medicine, University of Glasgow, Glasgow, G4 OSF, UK. ²²⁸ Laboratory of Neuro Imaging, Institute for Neuroimaging and Informatics, Keck School of Medicine of the University of Southern California, Los Angeles, California 90033, USA. ²²⁹ Department of Pathology, Johns Hopkins University, Baltimore, Maryland 21205, USA. ²³⁰ 3rd Department of Neurology, "G. Papanicolaou", Hospital, Aristotle University of Thessaloniki, Thessaloniki, 57010, Greece. 231 Univ. Bordeaux, Inserm, Bordeaux Population Health Research Center, UMR1219, Bordeaux, F-33000, France. ²³² Department of Internal Medicine, Erasmus MC, 3015 CE Rotterdam, The Netherlands. ²³³ Genentech Inc., South San Francisco, California 94080, USA. ²³⁴ Department of Psychiatry and Leiden Institute for Brain and Cognition, Leiden University Medical Center, 2333 ZA Leiden, The Netherlands. ²³⁵ University of Groningen, University Medical Center Groningen, Department of Neuroscience, 9713 AW Groningen, the Netherlands. ²³⁶ Department of Internal Medicine and Geriatric Medicine, INSERM U1027, University of Toulouse, Toulouse, 31024, France. ²³⁷ Department of Psychiatry, Carver College of Medicine, University of Iowa, Iowa City, Iowa 52242, USA. ²³⁸ Departments of Psychiatry, Neurology, Neuroscience and the Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA. ²³⁹ Center for Imaging of Neurodegenerative Disease, San Francisco VA Medical Center, University of California, San Francisco, California 94121, USA. ²⁴⁰ Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, SE-141 57 Huddinge, Sweden.²⁴¹ Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC-Sophia Children's Hospital, 3015 CE Rotterdam, The Netherlands. ²⁴² Laboratory of Epidemiology & Population Sciences, National Institute on Aging, National Institutes of Health, Bethesda, Maryland 20892, USA. ²⁴³Departments of Neurology and Epidemiology, University of Washington, 325 Ninth Avenue, Seattle, Washington 98104-2420, USA. 244 Intramural Research Program, NIA, NIH, 7201 Wisconsin Ave, Suite 3C-309, Bethesda, Maryland 20892, USA. ²⁴⁵ Department of Neurology, Erasmus MC, Rotterdam 3015 CE, The Netherlands. * These authors contributed equally to this work... ** These authors jointly supervised the study