



# Foliations of Hilbert Modular Surfaces

## Citation

McMullen, Curtis T. 2007. Foliations of Hilbert modular surfaces. American Journal of Mathematics 129(1): 183-215.

## Published Version

<http://www.math.jhu.edu/~ajm/>

## Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:3446012>

## Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

## Share Your Story

The Harvard community has made this article openly available.  
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

# Foliations of Hilbert modular surfaces

Curtis T. McMullen\*

21 February, 2005

## Abstract

The Hilbert modular surface  $X_D$  is the moduli space of Abelian varieties  $A$  with real multiplication by a quadratic order of discriminant  $D > 1$ . The locus where  $A$  is a product of elliptic curves determines a finite union of algebraic curves  $X_D(1) \subset X_D$ .

In this paper we show the lamination  $X_D(1)$  extends to an essentially unique foliation  $\mathcal{F}_D$  of  $X_D$  by complex geodesics. The geometry of  $\mathcal{F}_D$  is related to Teichmüller theory, holomorphic motions, polygonal billiards and Lattès rational maps. We show every leaf of  $\mathcal{F}_D$  is either closed or dense, and compute its holonomy. We also introduce refinements  $T_N(\nu)$  of the classical modular curves on  $X_D$ , leading to an explicit description of  $X_D(1)$ .

## Contents

1	Introduction . . . . .	1
2	Quaternion algebras . . . . .	5
3	Modular curves and surfaces . . . . .	12
4	Laminations . . . . .	17
5	Foliations of Teichmüller space . . . . .	20
6	Genus two . . . . .	23
7	Holomorphic motions . . . . .	25
8	Quasiconformal dynamics . . . . .	27
9	Further results . . . . .	30

---

\*Research supported in part by the NSF and the Guggenheim Foundation.

# 1 Introduction

Let  $D > 1$  be an integer congruent to 0 or 1 mod 4, and let  $\mathcal{O}_D$  be the real quadratic order of discriminant  $D$ . The *Hilbert modular surface*

$$X_D = (\mathbb{H} \times \mathbb{H}) / \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$$

is the moduli space for principally polarized Abelian varieties

$$A_\tau = \mathbb{C}^2 / (\mathcal{O}_D \oplus \mathcal{O}_D^\vee \tau)$$

with real multiplication by  $\mathcal{O}_D$ .

Let  $X_D(1) \subset X_D$  denote the locus where  $A_\tau$  is isomorphic to a polarized product of elliptic curves  $E_1 \times E_2$ . The set  $X_D(1)$  is a finite union of disjoint, irreducible algebraic curves (§4), forming a *lamination* of  $X_D$ . Note that  $X_D(1)$  is preserved by the twofold symmetry  $\iota(\tau_1, \tau_2) = (\tau_2, \tau_1)$  of  $X_D$ .

In this paper we will show:

**Theorem 1.1** *Up to the action of  $\iota$ , the lamination  $X_D(1)$  extends to a unique foliation  $\mathcal{F}_D$  of  $X_D$  by complex geodesics.*

(Here a Riemann surface in  $X_D$  is a *complex geodesic* if it is isometrically immersed for the Kobayashi metric.)

**Holomorphic graphs.** The preimage  $\tilde{X}_D(1)$  of  $X_D(1)$  in the universal cover of  $X_D$  gives a lamination of  $\mathbb{H} \times \mathbb{H}$  by the graphs of countably many Möbius transformations. To foliate  $X_D$  itself, in §6 we will show:

**Theorem 1.2** *For any  $(\tau_1, \tau_2) \notin \tilde{X}_D(1)$ , there is a unique holomorphic function*

$$f : \mathbb{H} \rightarrow \mathbb{H}$$

*such that  $f(\tau_1) = \tau_2$  and the graph of  $f$  is disjoint from  $\tilde{X}_D(1)$ .*

The graphs of such functions descend to  $X_D$ , and form the leaves of the foliation  $\mathcal{F}_D$  (§7). The case  $D = 4$  is illustrated in Figure 1.

**Modular curves.** To describe the lamination  $X_D(1)$  explicitly, recall that the Hilbert modular surface  $X_D$  is populated by infinitely many *modular curves*  $F_N$  [Hir], [vG]. The endomorphism ring of a generic Abelian variety in  $F_N$  is a quaternionic order  $R$  of discriminant  $N^2$ .

In general  $F_N$  can be reducible, and  $R$  is not determined up to isomorphism by  $N$ . In §3 we introduce a refinement  $F_N(\nu)$  of the traditional modular curves, such that the isomorphism class of  $R$  is constant along




Figure 1. Foliation of the Hilbert modular surface  $X_D$ ,  $D = 4$ .

$F_N(\nu)$  and  $F_N = \bigcup F_N(\nu)$ . The additional finite invariant  $\nu$  ranges in the ring  $\mathcal{O}_D/(\sqrt{D})$  and its norm satisfies  $N(\nu) = -N \pmod{D}$ . The curves  $T_N = \bigcup F_{N/\ell^2}$  can be refined similarly, and we obtain:

**Theorem 1.3** *The locus  $X_D(1) \subset X_D$  is given by*

$$X_D(1) = \bigcup T_N((e + \sqrt{D})/2),$$

where the union is over all integral solutions to  $e^2 + 4N = D$ ,  $N > 0$ .

**Remark.** Although  $X_D(1) = \bigcup T_{(D-e^2)/4}$  when  $D$  is prime, in general (e.g. for  $D = 12, 16, 20, 21, \dots$ ) the locus  $X_D(1)$  cannot be expressed as a union of the traditional modular curves  $T_N$  (§3).

Here is a corresponding description of the lamination  $\tilde{X}_D(1)$ . Given  $N > 0$  such that  $D = e^2 + 4N$ , let

$$\Lambda_D^N = \left\{ U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} : \begin{array}{l} a, b \in \mathbb{Z}, \mu \in \mathcal{O}_D, \det(U) = N \\ \text{and } \mu \equiv \pm(e + \sqrt{D})/2 \text{ in } \mathcal{O}_D/(\sqrt{D}) \end{array} \right\}.$$

Let  $\Lambda_D$  be the union of all such  $\Lambda_D^N$ . Choosing a real place  $\iota_1 : \mathcal{O}_D \rightarrow \mathbb{R}$ , we can regard  $\Lambda_D$  as a set of matrices in  $\mathrm{GL}_2^+(\mathbb{R})$ , acting by Möbius transformations on  $\mathbb{H}$ .

**Theorem 1.4** *The lamination  $\tilde{X}_D(1)$  of  $\mathbb{H} \times \mathbb{H}$  is the union of the loci  $\tau_2 = U(\tau_1)$  over all  $U \in \Lambda_D$ .*

We also obtain a description of the locus  $X_D(E) \subset X_D$  where  $A_\tau$  admits an action of both  $\mathcal{O}_D$  and  $\mathcal{O}_E$  (§3).

**Quasiconformal dynamics.** Although its leaves are Riemann surfaces,  $\mathcal{F}_D$  is not a holomorphic foliation. Its transverse dynamics is given instead by quasiconformal maps, which can be described as follows.

Let  $q = q(z) dz^2$  be a meromorphic quadratic differential on  $\mathbb{H}$ . We say a homeomorphism  $f : \mathbb{H} \rightarrow \mathbb{H}$  is a *Teichmüller mapping* relative to  $q$  if it satisfies  $\bar{\partial}f/\partial f = \alpha q/|q|$  for some complex number  $|\alpha| < 1$ ; equivalently, if  $f$  has the form of an orientation-preserving real-linear mapping

$$f(x + iy) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = D_q(f) \begin{pmatrix} x \\ y \end{pmatrix}$$

in local charts where  $q = dz^2 = (dx + i dy)^2$ .

Fix a transversal  $\mathbb{H}_s = \{s\} \times \mathbb{H}$  to  $\tilde{\mathcal{F}}_D$ . Any  $g \in \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  acts on  $\mathbb{H} \times \mathbb{H}$ , permuting the leaves of  $\tilde{\mathcal{F}}_D$ . The permutation of leaves is recorded by the *holonomy map*

$$\phi_g : \mathbb{H}_s \rightarrow \mathbb{H}_s,$$

characterized by the property that  $g(s, z)$  and  $(s, \phi_g(z))$  lie on the same leaf of  $\tilde{\mathcal{F}}_D$ .

In §8 we will show:

**Theorem 1.5** *The holonomy acts by Teichmüller mappings relative to a fixed meromorphic quadratic differential  $q$  on  $\mathbb{H}_s$ . For  $s = i$  and  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , we have*

$$D_q(\phi_g) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{PSL}_2(\mathbb{R}).$$

On the other hand, for  $z \in \partial\mathbb{H}_s$  we have

$$\phi_g(z) = (a'z - b')/(-c'z + d');$$

in particular, the holonomy acts by Möbius transformations on  $\partial\mathbb{H}_s$ .

Here  $(x + y\sqrt{D})' = (x - y\sqrt{D})$ . Note that both Galois conjugate actions of  $g$  on  $\mathbb{R}^2$  appear, as different aspects of the holonomy map  $\phi_g$ .

**Quantum Teichmüller curves.** For comparison, consider an isometrically immersed *Teichmüller curve*

$$f : V \rightarrow \mathcal{M}_g,$$

generated by a holomorphic quadratic differential  $(Y, q)$  of genus  $g$ . For simplicity assume  $\text{Aut}(Y)$  is trivial. Then the pullback of the universal curve  $X = f^*(\mathcal{M}_{g,1})$  gives an algebraic surface

$$p : X \rightarrow V$$

with  $p^{-1}(v) = Y$  for a suitable basepoint  $v \in V$ . The surface  $X$  carries a canonical foliation  $\mathcal{F}$ , transverse to the fibers of  $p$ , whose leaves map to Teichmüller geodesics in  $\mathcal{M}_{g,1}$ . The holonomy of  $\mathcal{F}$  determines a map

$$\pi_1(V, v) \rightarrow \text{Aff}^+(Y, q)$$

giving an action of the fundamental group by Teichmüller mappings; and its linear part yields the isomorphism

$$\pi_1(V, v) \cong \text{PSL}(Y, q) \subset \text{PSL}_2(\mathbb{R}),$$

where  $\text{PSL}(Y, q)$  is the stabilizer of  $(Y, q)$  in the bundle of quadratic differentials  $Q\mathcal{M}_g \rightarrow \mathcal{M}_g$ . (See e.g. [V1], [Mc4, §2].)

The foliated Hilbert modular surface  $(X_D, \mathcal{F}_D)$  presents a similar structure, with the fibration  $p : X \rightarrow V$  replaced by the holomorphic foliation  $\mathcal{A}_D$  coming from the level sets of  $\tau_1$  on  $\tilde{X}_D = \mathbb{H} \times \mathbb{H}$ . This suggests that one should regard  $(X_D, \mathcal{A}_D, \mathcal{F}_D)$  as a *quantum* Teichmüller curve, in the same sense that a 3-manifold with a measured foliation can be regarded as a quantum Teichmüller geodesic [Mc3].

**Question.** Does every fibered surface  $p : X \rightarrow C$  admit a foliation  $\mathcal{F}$  by Riemann surfaces transverse to the fibers of  $p$ ?

**Complements.** We conclude in §9 by presenting the following related results.

1. Every leaf of  $\mathcal{F}_D$  is either closed or dense.
2. When  $D \neq d^2$ , there are infinitely many eigenforms for real multiplication by  $\mathcal{O}_D$  that are isoperiodic but not isomorphic.
3. The Möbius transformations  $\Lambda_D$  give a maximal top-speed holomorphic motion of a discrete subset of  $\mathbb{H}$ .
4. The foliation  $\mathcal{F}_4$  also arises as the motion of the Julia set in a Lattès family of iterated rational maps.

The link with complex dynamics was used to produce Figure 1.

**Notes and references.** The foliation  $\mathcal{F}_D$  is constructed using the connection between polygonal billiards and Hilbert modular surfaces presented in [Mc4]. For more on the interplay of dynamics, holomorphic motions and quasiconformal mappings, see e.g. [MSS], [BR], [Sl], [Mc2], [Sul], [McS], [EKK] and [Dou]. A survey of the theory of *holomorphic* foliations of surfaces appears in [Br1]; see also [Br2] for the Hilbert modular case.

I would like to thank G. van der Geer, B. Gross and the referees for useful comments and suggestions.

## 2 Quaternion algebras

In this section we consider a real quadratic order  $\mathcal{O}_D$  acting on a symplectic lattice  $L$ , and classify the quaternionic orders  $R \subset \text{End}(L)$  extending  $\mathcal{O}_D$ .

**Quadratic orders.** Given an integer  $D > 0$ ,  $D \equiv 0$  or  $1 \pmod{4}$ , the *real quadratic order* of discriminant  $D$  is given by

$$\mathcal{O}_D = \mathbb{Z}[T]/(T^2 + bT + c), \quad \text{where } D = b^2 - 4c.$$

Let  $K_D = \mathcal{O}_D \otimes \mathbb{Q}$ . Provided  $D$  is not a square,  $K_D$  is a real quadratic field. Fixing an embedding  $\iota_1 : K_D \rightarrow \mathbb{R}$ , we obtain a unique basis

$$K_D = \mathbb{Q} \cdot 1 \oplus \mathbb{Q} \cdot \sqrt{D}$$

such that  $\iota_1(\sqrt{D}) > 0$ . The conjugate real embedding  $\iota_2 : K_D \rightarrow \mathbb{R}$  is given by  $\iota_2(x) = \iota_1(x')$ , where  $(a + b\sqrt{D})' = (a - b\sqrt{D})$ .

**Square discriminants.** The case  $D = d^2$  can be treated similarly, so long as we regard  $x = \sqrt{d^2}$  as an element of  $K_D$  satisfying  $x^2 = d^2$  but  $x \notin \mathbb{Q}$ . In this case the algebra  $K_D \cong \mathbb{Q} \oplus \mathbb{Q}$  is not a field, so we must take care to distinguish between elements of the algebra such as

$$x = d - \sqrt{d^2} \in K_D,$$

and the corresponding real numbers

$$\iota_1(x) = d - d = 0, \quad \text{and} \quad \iota_2(x) = d + d = 2d.$$

**Trace, norm and different.** For simplicity of notation, we fix  $D$  and denote  $\mathcal{O}_D$  and  $K_D$  by  $K$  and  $\mathcal{O}$ .

The trace and norm on  $K$  are the rational numbers  $\text{Tr}(x) = x + x'$  and  $\text{N}(x) = xx'$ . The *inverse different* is the fractional ideal

$$\mathcal{O}^\vee = \{x \in K : \text{Tr}(xy) \in \mathbb{Z} \forall y \in \mathcal{O}\}.$$

It is easy to see that  $\mathcal{O}^\vee = D^{-1/2} \mathcal{O}$ , and thus the *different*  $\mathcal{D} = (\mathcal{O}^\vee)^{-1} \subset \mathcal{O}$  is the principal ideal  $(\sqrt{D})$ . The trace and norm descend to give maps

$$\text{Tr}, \text{N} : \mathcal{O} / \mathcal{D} \rightarrow \mathbb{Z} / D,$$

satisfying

$$\text{Tr}(x)^2 = 4 \text{N}(x) \bmod D. \quad (2.1)$$

When  $D$  is odd,  $\text{Tr} : \mathcal{O} / \mathcal{D} \rightarrow \mathbb{Z} / D$  is an isomorphism, and thus (2.1) determines the norm on  $\mathcal{O} / \mathcal{D}$ . On the other hand, when  $D = 4E$  is even, we have an isomorphism

$$\mathcal{O} / \mathcal{D} \cong \mathbb{Z} / 2E \oplus \mathbb{Z} / 2$$

given by  $a + b\sqrt{E} \mapsto (a, b)$ , and the trace and norm on  $\mathcal{O} / \mathcal{D}$  are given by

$$\text{Tr}(a, b) = 2a \bmod D, \quad \text{N}(a, b) = a^2 - Eb^2 \bmod D.$$

**Symplectic lattices.** Now let  $L \cong (\mathbb{Z}^{2g}, \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix})$  be a unimodular symplectic lattice of genus  $g$ . (This lattice is isomorphic to the first homology group  $H_1(\Sigma_g, \mathbb{Z})$  of an oriented surface of genus  $g$  with the symplectic form given by the intersection pairing.)

Let  $\text{End}(L) \cong \text{M}_{2g}(\mathbb{Z})$  denote the endomorphism ring of  $L$  as a  $\mathbb{Z}$ -module. The *Rosati involution*  $T \mapsto T^*$  on  $\text{End}(L)$  is defined by the condition  $\langle Tx, y \rangle = \langle x, T^*y \rangle$ ; it satisfies  $(ST)^* = T^*S^*$ , and we say  $T$  is *self-adjoint* if  $T = T^*$ .

Specializing to the case  $g = 2$ , let  $L$  denote the lattice

$$L = \mathcal{O} \oplus \mathcal{O}^\vee$$

with the unimodular symplectic form

$$\langle x, y \rangle = \text{Tr}(x \wedge y) = \text{Tr}_{\mathbb{Q}}^K(x_1y_2 - x_2y_1).$$

A standard symplectic basis for  $L$  (satisfying  $\langle a_i \cdot b_j \rangle = \delta_{ij}$ ) is given by

$$(a_1, a_2, b_1, b_2) = ((1, 0), (\gamma, 0), (0, -\gamma'/\sqrt{D}), (0, 1/\sqrt{D})), \quad (2.2)$$



where  $\gamma = (D + \sqrt{D})/2$ .

The lattice  $L$  comes equipped with a proper, self-adjoint action of  $\mathcal{O}$ , given by

$$k \cdot (x_1, x_2) = (kx_1, kx_2). \quad (2.3)$$

Conversely, any proper, self-adjoint action of  $\mathcal{O}$  on a symplectic lattice of genus two is isomorphic to this model (see e.g. [Ru], [Mc7, Thm 4.1]). (Here an action of  $R$  on  $L$  is *proper* if it is indivisible: if whenever  $T \in \text{End}(L)$  and  $mT \in R$  for some integer  $m \neq 0$ , then  $T \in R$ .)

**Matrices.** The natural embedding of  $L = \mathcal{O} \oplus \mathcal{O}^\vee$  into  $K \oplus K$  determines an embedding of matrices

$$M_2(K) \rightarrow \text{End}(L \otimes \mathbb{Q}),$$

and hence a diagonal inclusion

$$K \rightarrow \text{End}(L \otimes \mathbb{Q})$$

extending the natural action (2.3) of  $\mathcal{O}$  on  $L$ . Every  $T \in \text{End}(L \otimes \mathbb{Q})$  can be uniquely expressed in the form

$$T(x) = Ax + Bx', \quad A, B \in M_2(K),$$

where  $(x_1, x_2)' = (x'_1, x'_2)$ ; and we have

$$T^*(x) = A^\dagger x + (B^\dagger)'x', \quad (2.4)$$

where  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}^\dagger = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ .

The automorphisms of  $L$  as a symplectic  $\mathcal{O}$ -module are given, as a subgroup of  $M_2(K)$ , by

$$\text{SL}(\mathcal{O} \oplus \mathcal{O}^\vee) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \begin{pmatrix} \mathcal{O} & \mathcal{D} \\ \mathcal{O}^\vee & \mathcal{O} \end{pmatrix} : ad - bc = 1 \right\}.$$

Compare [vG, p.12].

**Integrality.** An endomorphism  $T \in \text{End}(L \otimes \mathbb{Q})$  is *integral* if it satisfies  $T(L) \subset L$ .

**Lemma 2.1** *The endomorphism  $\phi(x) = ax + bx'$  of  $K$  satisfies  $\phi(\mathcal{O}) \subset \mathcal{O}$  iff  $a, b \in \mathcal{O}^\vee$  and  $a + b \in \mathcal{O}$ .*

**Proof.** Since  $x - x' \in \sqrt{D}\mathbb{Z}$  for all  $x \in \mathcal{O}$ , the conditions on  $a, b$  imply  $\phi(x) = a(x - x') + (a + b)x' \in \mathcal{O}$  for all  $x \in \mathcal{O}$ . Conversely, if  $\phi$  is integral, then  $\phi(1) = a + b \in \mathcal{O}$ , and thus  $a(x - x') \in \mathcal{O}$  for all  $x \in \mathcal{O}$ , which implies  $a \in D^{-1/2}\mathcal{O} = \mathcal{O}^\vee$ . ■

**Corollary 2.2** *The endomorphism  $T(x) = kx + \begin{pmatrix} a & bD \\ c & d \end{pmatrix} x'$  is integral iff we have*

$$a, b, c, d, k \in \mathcal{O}^\vee \quad \text{and} \quad k + a, k - d \in \mathcal{O}.$$

**Proof.** This follows from the preceding Lemma, using the fact that  $kx + dx'$  maps  $\mathcal{O}^\vee$  to  $\mathcal{O}^\vee$  iff  $kx - dx'$  maps  $\mathcal{O}$  to  $\mathcal{O}$ . ■

**Quaternion algebras.** A rational *quaternion algebra* is a central simple algebra of dimension 4 over  $\mathbb{Q}$ . Every such algebra has the form

$$Q \cong \mathbb{Q}[i, j]/(i^2 = a, j^2 = b, ij = -ji) = \left( \frac{a, b}{\mathbb{Q}} \right)$$

for suitable  $a, b \in \mathbb{Q}^*$ . Any  $q \in Q$  satisfies a quadratic equation

$$q^2 - \text{Tr}(q)q + N(q) = 0,$$

where  $\text{Tr}, N : Q \rightarrow \mathbb{Q}$  are the *reduced trace and norm*.

An *order*  $R \subset Q$  is a subring such that, as an additive group, we have  $R \cong \mathbb{Z}^4$  and  $\mathbb{Q} \cdot R = Q$ . Its *discriminant* is the square integer

$$N^2 = |\det(\text{Tr}(q_i q_j))| > 0,$$

where  $(q_i)_1^4$  is an integral basis for  $R$ . The discriminants of a pair of orders  $R_1 \subset R_2$  are related by  $N_1/N_2 = |R_2/R_1|^2$ .

**Generators.** We say  $V \in \text{End}(L)$  is a *quaternionic generator* if:

1.  $V^* = -V$ ,
2.  $V^2 = -N \in \mathbb{Z}$ ,  $N \neq 0$ ,
3.  $Vk = k'V$  for all  $k \in K$ , and
4.  $k + D^{-1/2}V \in \text{End}(L)$  for some  $k \in K$ .

These conditions imply that  $Q = K \oplus KV$  is a quaternion algebra isomorphic to  $\left( \frac{D, -N}{\mathbb{Q}} \right)$ . Conversely, we have:

**Theorem 2.3** *Any Rosati-invariant quaternion algebra  $Q$  with*

$$K \subset Q \subset \text{End}(L \otimes \mathbb{Q})$$

*contains a unique pair of primitive quaternionic generators  $\pm V$ .*

(A generator is *primitive* unless  $(1/m)V, m > 1$  is also a generator.)

**Proof.** By a standard application of the Skolem-Noether theorem, we can write  $Q = K \oplus KW$  with  $0 \neq W^2 \in \mathbb{Q}$  and  $Wk = k'W$  for all  $k \in K$ . Then  $KW$  coincides with the subalgebra of  $Q$  anticommuting with the self-adjoint element  $\sqrt{D}$ , so it is Rosati-invariant. The eigenspaces of  $*|KW$  are exchanged by multiplication by  $\sqrt{D}$ , so up to a rational multiple there is a unique nonzero  $V \in KW$  with  $V^* = -V$ . A suitable integral multiple of  $V$  is then a generator, and a rational multiple is primitive. ■

**Corollary 2.4** *Quaternionic extensions  $K \subset Q \subset \text{End}(L)$  correspond bijectively to pairs of primitive generators  $\pm V \in \text{End}(L)$ .*

**Generator matrices.** We say  $U \in M_2(K)$  is a *quaternionic generator matrix* if it has the form

$$U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} \quad (2.5)$$

with  $a, b \in \mathbb{Z}$ ,  $\mu \in \mathcal{O}$  and  $N = \det(U) \neq 0$ .

**Theorem 2.5** *The endomorphism  $V(x) = Ux'$  is a quaternionic generator iff  $U$  is a quaternionic generator matrix.*

**Proof.** By (2.4) the condition  $V = -V^*$  is equivalent to  $U^\dagger = -U'$ , and thus  $U$  can be written in the form (2.5) with  $a, b \in \mathbb{Q}$  and  $\mu \in K$ . Assuming  $U^\dagger = -U'$ , we have

$$N = \det(U) = UU^\dagger = -UU' = -V^2,$$

so  $V^2 \neq 0 \iff \det(U) \neq 0$ . The condition that  $D^{-1/2}(k+V)$  is integral for some  $k$  implies, by Corollary 2.2, that the coefficients of  $U$  satisfy  $a, b \in \mathbb{Z}$  and  $\mu \in \mathcal{O}$ ; and given such coefficients for  $U$ , the endomorphism  $D^{-1/2}(k+V)$  is integral when  $k = -\mu$ . ■

**The invariant  $\nu(U)$ .** Given generator matrix  $U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix}$ , let  $\nu(U)$  denote the image of  $\mu$  in the finite ring  $\mathcal{O}/D$ . It is easy to check that

$$\nu(U) = \pm \nu(g'Ug^{-1})$$

for all  $g \in \text{SL}(\mathcal{O} \oplus \mathcal{O}^\vee)$ , and that its norm satisfies

$$N(\nu(U)) \equiv -N \pmod{D}. \quad (2.6)$$

**Quaternionic orders.** Let  $V(x) = Ux'$ , and let

$$R_U = (K \oplus KV) \cap \text{End}(L).$$

Then  $R_U$  is a Rosati-invariant order in the quaternion algebra generated by  $V$ . Clearly  $\mathcal{O} \subset R_U$ , so we can also regard  $(R_U, *)$  as an involutive algebra over  $\mathcal{O}$ . We will show that  $N = \det(U)$  and  $\nu(U)$  determine  $(R_U, *)$  up to isomorphism.

**Models.** We begin by constructing a model algebra  $(R_N(\nu), *)$  over  $\mathcal{O}_D$  for every  $\nu \in \mathcal{O}/\mathcal{D}$  with  $N(\nu) = -N \not\equiv 0 \pmod{D}$ .

Let  $Q_N = K \oplus KV$  be the abstract quaternion algebra with the relations  $V^2 = -N$  and  $Vk = k'V$ . Define an involution on  $Q_N$  by  $(k_1 + k_2V)^* = (k_1 - k_2'V)$ , and let  $R_N(\nu)$  be the order in  $Q_N$  defined by

$$R_N(\nu) = \{\alpha + \beta V : \alpha, \beta \in \mathcal{O}^\vee, \alpha + \beta\nu \in \mathcal{O}\}. \quad (2.7)$$

Note that  $\mathcal{O}^\vee \cdot \mathcal{D} \subset \mathcal{O}$ , so the definition of  $R_N(\nu)$  depends only on the class of  $\nu$  in  $\mathcal{O}/\mathcal{D}$ . To check that  $R_N(\nu)$  is an order, note that

$$(\alpha + \beta V)(\gamma + \delta V) = (\kappa + \lambda V) = (\alpha\gamma - N\beta\delta') + (\alpha\delta + \beta\gamma')V;$$

since  $-N \equiv N(\nu) = \nu\nu' \pmod{D}$ , we have

$$\begin{aligned} \kappa + \nu\lambda &\equiv (\alpha\gamma + \nu\nu'\beta\delta') + \nu(\alpha\delta + \beta\gamma') \\ &= (\alpha + \beta\nu)(\gamma' + \delta'\nu') + \alpha(\gamma - \gamma' + \nu\delta - \nu'\delta') \\ &\equiv 0 + 0 \pmod{\mathcal{O}}, \end{aligned}$$

and thus  $R_U$  is closed under multiplication.

**Theorem 2.6** *The quaternionic order  $R_N(\nu)$  has discriminant  $N^2$ .*

**Proof.** Note that the inclusions

$$\mathcal{O} \oplus \mathcal{O}V \subset R_N(\nu) \subset \mathcal{O}^\vee \oplus \mathcal{O}^\vee V$$

each have index  $D$ . The quaternionic order  $\mathcal{O} \oplus \mathcal{O}V$  has discriminant  $D^2N^2$ , since  $V^2 = -N$  and  $\text{Tr}|\mathcal{O}V = 0$ , and thus  $R_N(\nu)$  has discriminant  $N^2$ . ■

**Theorem 2.7** *We have  $(R_N(\nu), *) \cong (R_M(\mu), *)$  iff  $N = M$  and  $\nu = \pm\mu$ .*

**Proof.** The element  $V \in R_N(\nu)$  is, up to sign, the order's unique primitive generator, in the sense that  $V^* = -V$ ,  $Vk = k'V$  for all  $k \in \mathcal{O}_D$ ,  $V^2 \neq 0$ ,  $k + D^{-1/2}V \in R_N(\nu)$  for some  $k \in K$ , and  $V$  is not a proper multiple of another element in  $R_N(\nu)$  with the same properties. Thus the structure of  $(R_N(\nu), *)$  as an  $\mathcal{O}_D$ -algebra determines  $V \in R_N(\nu)$  up to sign, and  $V$  determines  $N = -V^2$  and the constant  $\nu \in \mathcal{O}/\mathcal{D}$  in the relation  $\alpha + \beta\nu \in \mathcal{O}$  defining  $R_N(\nu) \subset K \oplus KV$ . ■

**Theorem 2.8** *If  $U$  is a primitive generator matrix, then we have*

$$(R_U, *) \cong (R_N(\nu), *)$$

where  $N = \det(U)$  and  $\nu = \nu(U)$ .

**Proof.** Setting  $V(x) = Ux'$ , we need only verify that  $(K \oplus KV) \cap \text{End}(L)$  coincides with the order  $R_N(\nu)$  defined by (2.7). To see this, let

$$T(x) = \alpha x + \beta V(x) = \alpha x + \beta \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} x'$$

in  $K \oplus KV$ . By Corollary 2.2,  $T$  is integral iff

- (i)  $a\beta, b\beta, \mu\beta, \mu'\beta \in \mathcal{O}^\vee$ ,
- (ii)  $\alpha \in \mathcal{O}^\vee$ ,
- (iii)  $\alpha + \beta\mu \in \mathcal{O}$  and
- (iv)  $\alpha + \beta\mu' \in \mathcal{O}$ .

Using (iii), condition (iv) can be replaced by

$$(iv') \quad \beta(\mu - \mu')/\sqrt{D} \in \mathcal{O}^\vee.$$

Since  $U$  is primitive, the ideal  $(a, b, \mu, (\mu - \mu')/\sqrt{D})$  is equal to  $\mathcal{O}$ . Thus (i) and (iv') together are equivalent to the condition  $\beta \in \mathcal{O}^\vee$ , and we are left with the definition of  $R_N(\nu)$ . ■

**Remark.** In general, the invariants  $\det(U)$  and  $\nu(U)$  do not determine the embedding  $R_U \subset \text{End}(L)$  up to conjugacy. For example, when  $D$  is odd, the generator matrices  $U_1 = \begin{pmatrix} 0 & D^2 \\ -D & 0 \end{pmatrix}$  and  $U_2 = \begin{pmatrix} 0 & D^3 \\ -1 & 0 \end{pmatrix}$  have the same invariants, but the corresponding endomorphisms are not conjugate in  $\text{End}(L)$  because

$$L/V_1(L) \cong (\mathbb{Z}/D \times \mathbb{Z}/D^2)^2$$

while

$$L/V_2(L) \cong \mathbb{Z}/D \times \mathbb{Z}/D^2 \times \mathbb{Z}/D^3.$$

**Extra quadratic orders.** Finally we determine when the algebra  $R_N(\nu)$  contains a second, independent quadratic order  $\mathcal{O}_E$ .

**Theorem 2.9** *The algebra  $(R_N(\nu), *)$  contains a self-adjoint element  $T \notin \mathcal{O}_D$  generating a copy of  $\mathcal{O}_E$  iff there exist  $e, \ell \in \mathbb{Z}$  such that*

$$ED = e^2 + 4N\ell^2, \quad \ell \neq 0$$

and  $(e + E\sqrt{D})/2 + \ell\nu = 0 \pmod{\mathcal{D}}$ .

**Proof.** Given  $e, \ell$  as above, let

$$T = \alpha + \beta V = D^{-1/2} \left( \frac{e + E\sqrt{D}}{2} + \ell V \right).$$

Then we have  $T = T^*$ ,  $T \in R_N(\nu)$  and  $T^2 - eT + (E - E^2)/4 = 0$ ; therefore  $\mathbb{Z}[T] \cong \mathcal{O}_E$ . A straightforward computation shows that, conversely, any independent copy of  $\mathcal{O}_E$  in  $R_N(\nu)$  arises as above.  $\blacksquare$

For additional background on quaternion algebras, see e.g. [Vi], [MR] and [Mn].

### 3 Modular curves and surfaces

In this section we describe modular curves on Hilbert modular surfaces from the perspective of the Abelian varieties they determine.

**Abelian varieties.** A *principally polarized Abelian variety* is a complex torus  $A \cong \mathbb{C}^g/L$  equipped with a unimodular symplectic form  $\langle x, y \rangle$  on  $L \cong \mathbb{Z}^{2g}$ , whose extension to  $L \otimes \mathbb{R} \cong \mathbb{C}^g$  satisfies

$$\langle x, y \rangle = \langle ix, iy \rangle \quad \text{and} \quad \langle x, ix \rangle \geq 0.$$

The ring  $\text{End}(A) = \text{End}(L) \cap \text{End}(\mathbb{C}^g)$  is Rosati invariant, and coincides with the endomorphism ring of  $A$  as a complex Lie group. We have  $\text{Tr}(TT^*) \geq 0$  for all  $T \in \text{End}(A)$ .

Every Abelian variety can be presented in the form

$$A = \mathbb{C}^g / (\mathbb{Z}^g \oplus \Pi \mathbb{Z}^g),$$

where  $\Pi$  is an element of the Siegel upper halfplane

$$\mathfrak{H}_g = \{\Pi \in M_g(\mathbb{C}) : \Pi^t = \Pi \text{ and } \text{Im}(\Pi) \text{ is positive-definite}\}.$$

The symplectic form on  $L = \mathbb{Z}^g \oplus \Pi \mathbb{Z}^g$  is given by  $\begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$ . Any two such presentations of  $A$  differ by an automorphism of  $L$ , so the moduli space of abelian varieties of genus  $g$  is given by the quotient space

$$\mathcal{A}_g = \mathfrak{H}_g / \text{Sp}_{2g}(\mathbb{Z}).$$

**Real multiplication.** As in §2, let  $D > 0$  be the discriminant of a real quadratic order  $\mathcal{O}_D$ , and let  $K = \mathcal{O} \otimes \mathbb{Q}$ . Fix a real place  $\iota_1 : K \rightarrow \mathbb{R}$ , and set  $\iota_2(k) = \iota_1(k')$ .

We will regard  $K$  as a subfield of the reals, using the fixed embedding  $\iota_1 : K \subset \mathbb{R}$ . The case  $D = d^2$  is treated with the understanding that the real numbers  $(k, k')$  implicitly denote  $(\iota_1(k), \iota_2(k))$ ,  $k \in K$ .

An Abelian variety  $A \in \mathcal{A}_2$  admits *real multiplication* by  $\mathcal{O}_D$  if there is a self-adjoint endomorphism  $T \in \text{End}(A)$  generating a proper action of  $\mathbb{Z}[T] \cong \mathcal{O}_D$  on  $A$ . Any such variety can be presented in the form

$$A_\tau = \mathbb{C}^2 / (\mathcal{O}_D \oplus \mathcal{O}_D^\vee \tau) = \mathbb{C}^2 / \phi_\tau(L), \quad (3.1)$$

where  $\tau = (\tau_1, \tau_2) \in \mathbb{H} \times \mathbb{H}$  and where  $L = \mathcal{O} \oplus \mathcal{O}^\vee$  is embedded in  $\mathbb{C}^2$  by the map

$$\phi_\tau(x_1, x_2) = (x_1 + x_2 \tau_1, x_1' + x_2' \tau_2).$$

As in §2, the symplectic form on  $L$  is given by  $\langle x, y \rangle = \text{Tr}_{\mathbb{Q}}^K(x \wedge y)$ , and the action of  $\mathcal{O}_D$  on  $\mathbb{C}^2 \supset L$  is given simply by  $k \cdot (z_1, z_2) = (kz_1, k'z_2)$ .

**Eigenforms.** The Abelian variety  $A_\tau$  comes equipped with a distinguished pair of normalized *eigenforms*  $\eta_1, \eta_2 \in \Omega(A_\tau)$ . Using the isomorphism  $H_1(A_\tau, \mathbb{Z}) \cong L$ , these forms are characterized by the property that

$$\phi_\tau(C) = \left( \int_C \eta_1, \int_C \eta_2 \right). \quad (3.2)$$

**Modular surfaces.** If we change the identification  $L \cong H_1(A_\tau, \mathbb{Z})$  by an automorphism  $g$  of  $L$ , we obtain an isomorphic Abelian variety  $A_{g \cdot \tau}$ . Thus the moduli space of Abelian varieties with real multiplication by  $\mathcal{O}_D$  is given by the Hilbert modular surface

$$X_D = (\mathbb{H} \times \mathbb{H}) / \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee).$$

The point  $g(\tau)$  is characterized by the property that

$$\phi_{g \cdot \tau} = \chi(g, \tau) \phi_\tau \circ g^{-1}$$

for some matrix  $\chi(g, \tau) \in \mathrm{GL}_2(\mathbb{C})$ ; explicitly, we have

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (\tau_1, \tau_2) = \left( \frac{a\tau_1 - b}{-c\tau_1 + d}, \frac{a'\tau_2 - b'}{-c'\tau_2 + d'} \right) \quad (3.3)$$

and

$$\chi(g, \tau) = \begin{pmatrix} (d - c\tau_1)^{-1} & 0 \\ 0 & (d' - c'\tau_2)^{-1} \end{pmatrix}. \quad (3.4)$$

A point  $[\tau] \in X_D$  gives an Abelian variety  $[A_\tau] \in \mathcal{A}_2$  with a *chosen* embedding  $\mathcal{O}_D \rightarrow \mathrm{End}(A_\tau)$ . Similarly, a point  $\tau \in \tilde{X}_D = \mathbb{H} \times \mathbb{H}$  gives an Abelian variety with a distinguished isomorphism or *marking*,  $L \cong H_1(A_\tau, \mathbb{Z})$ , sending  $\mathcal{O}_D$  into  $\mathrm{End}(A_\tau)$ .

**Modular embedding.** The *modular embedding*

$$p_D : X_D \rightarrow \mathcal{A}_2$$

is given by  $[\tau] \mapsto [A_\tau]$ . To write  $p_D$  explicitly, note that the embedding  $\phi_\tau : L \rightarrow \mathbb{C}^2$  can be expressed with respect to the basis  $(a_1, a_2, b_1, b_2)$  for  $L$  given in (2.2) by the matrix

$$\phi_\tau = \begin{pmatrix} 1 & \gamma & -\tau_1\gamma'/\sqrt{D} & \tau_1/\sqrt{D} \\ 1 & \gamma' & \tau_2\gamma/\sqrt{D} & -\tau_2/\sqrt{D} \end{pmatrix} = (A, B).$$

Consequently we have  $A_\tau \cong \mathbb{C}^2 / (\mathbb{Z}^2 \oplus \Pi\mathbb{Z}^2)$ , where

$$\Pi = \widetilde{p}_D(\tau) = A^{-1}B = \frac{1}{D} \begin{pmatrix} \tau_1(\gamma')^2 + \tau_2\gamma^2 & -\tau_1\gamma' - \tau_2\gamma \\ -\tau_1\gamma' - \tau_2\gamma & \tau_1 + \tau_2 \end{pmatrix}.$$

The map  $X_D \rightarrow p_D(X_D)$  has degree two.



**Modular curves.** Given a matrix  $U(x) = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in M_2(K) \cap \text{End}(L)$  such that  $U' = -U^*$ , let  $V(x) = Ux'$  and define

$$\mathbb{H}_U = \{\tau \in \mathbb{H} \times \mathbb{H} : V \in \text{End}(A_\tau)\}.$$

It is straightforward to check that

$$\mathbb{H}_U = \left\{ (\tau_1, \tau_2) : \tau_2 = \frac{d\tau_1 + b}{c\tau_1 + a} \right\}; \quad (3.5)$$

indeed, when  $\tau_1$  and  $\tau_2$  are related as above, the map  $\phi_\tau : L \rightarrow \mathbb{C}^2$  satisfies

$$\phi_\tau(V(x)) = \begin{pmatrix} 0 & a + c\tau_1 \\ a' + c'\tau_2 & 0 \end{pmatrix} \phi_\tau(x),$$

exhibiting the complex-linearity of  $V$ . Note that  $\mathbb{H}_U = \emptyset$  if  $\det(U) < 0$ .

We now restrict attention to the case where  $U$  is a generator matrix. Then by the results of §2, we have:

**Theorem 3.1** *The ring  $\text{End}(A_\tau)$  contains a quaternionic order extending  $\mathcal{O}_D$  if and only if  $\tau \in \mathbb{H}_U$  for some generator matrix  $U$ .*

Let  $F_U \subset X_D$  denote the projection of  $\mathbb{H}_U$  to the quotient  $(\mathbb{H} \times \mathbb{H}) / \text{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ . Following [Hir, §5.3], we define the *modular curve*  $F_N$  by

$$F_N = \bigcup \{F_U : U \text{ is a primitive generator matrix with } \det(U) = N\}.$$

It can be shown that  $F_N$  is an algebraic curve on  $X_D$ .

To describe this curve more precisely, let

$$F_N(\nu) = \{F_U : U \text{ is primitive, } \det(U) = N \text{ and } \nu(U) = \pm\nu\},$$

where  $\nu \in \mathcal{O}_D / \mathcal{D}_D$ . Note that we have

$$F_N(\nu) \neq \emptyset \iff N(\nu) = -N \pmod{D}$$

by equation (2.6),  $F_N(\nu) = F_N(-\nu)$ , and  $F_N = \bigcup F_N(\nu)$ .

The results of §2 give the structure of the quaternion ring generated by  $V(x) = Ux'$ .

**Theorem 3.2** *The curve  $F_N(\nu) \subset X_D$  coincides with the locus of Abelian varieties such that*

$$\mathcal{O}_D \subset R \subset \text{End}(A_\tau),$$

*for some properly embedded quaternionic order  $(R, *)$  isomorphic to  $(R_N(\nu), *)$ .*

**Corollary 3.3** *The curve  $F_N$  is the locus where  $\mathcal{O}_D \subset \text{End}(A_\tau)$  extends to a properly embedded, Rosati-invariant quaternionic order of discriminant  $N^2$ .*

**Two quadratic orders.** We can now describe the locus  $X_D(E)$  of Abelian varieties with an independent, self-adjoint action of  $\mathcal{O}_E$ . (We do not require the action of  $\mathcal{O}_E$  to be proper.)

To state this description, it is useful to define:

$$T_N = \bigcup \{F_U : \det(U) = N\} = \bigcup F_{N/\ell^2},$$

and

$$T_N(\nu) = \bigcup \{F_U : \det(U) = N, \nu(U) = \pm\nu\}.$$

Then Theorem 2.9 implies:

**Theorem 3.4** *The locus  $X_D(E)$  is given by*

$$X_D(E) = \bigcup T_N((e + E\sqrt{D})/2),$$

where the union is over all  $N > 0$  and  $e \in \mathbb{Z}$  such that  $ED = e^2 + 4N$ .

**Corollary 3.5** *We have  $X_D(1) = \bigcup \{T_N((e + \sqrt{D})/2) : e^2 + 4N = D\}$ .*

**Refined modular curves.** To conclude we show that in general the expression  $F_N = \bigcup F_N(\nu)$  gives a proper refinement of  $F_N$ . First note:

**Theorem 3.6** *We have  $F_N(\nu) = F_N$  iff  $\pm\nu$  are the only solutions to*

$$N(\xi) = -N \pmod{D}, \quad \xi \in \mathcal{O}_D/\mathcal{D}_D.$$

**Corollary 3.7** *If  $D = p$  is prime, then  $F_N = F_N(\nu)$  whenever  $F_N(\nu) \neq \emptyset$ .*

**Proof.** In this case, according to (2.1), the norm map

$$N : \mathcal{O}_D/\mathcal{D}_D \stackrel{\text{Tr}}{\cong} \mathbb{Z}/p \rightarrow \mathbb{Z}/p$$

is given by  $N(\xi) = \xi^2/4$ . Since  $F_N(\nu) \neq \emptyset$ , we have  $N(\nu) = -N$ ; and since  $\mathbb{Z}/p$  is a field,  $\pm\nu$  are the only solutions to this equation.  $\blacksquare$

**Corollary 3.8** *When  $D$  is prime, we have  $X_D(E) = \bigcup T_{(ED-e^2)/4}$ .*

Now consider the case  $D = 21$ , the first odd discriminant which is not a prime. Then the norm map is still given by  $N(\xi) = \xi^2/4$  on  $\mathcal{O}_D/\mathcal{D}_D \cong \mathbb{Z}/D$ , but now  $\mathbb{Z}/D$  is not a field. For example, the equation  $\xi^2 = 1 \pmod{D}$  has four solutions, namely  $\xi = 1, 8, 13$  or  $20$ . These give four solutions to the equation  $N(\xi) = -5$ , and hence contribute two distinct terms to the expression

$$F_5 = \bigcup F_5(\nu) = F_5((1 + \sqrt{21})/2) \cup F_5((8 + \sqrt{21})/2).$$

Only one of these terms appears in the expression for  $X_D(1)$ . In fact, since  $21 = 1^2 + 4 \cdot 5 = 3^2 + 4 \cdot 3$ , by Corollary 3.5 we have

$$\begin{aligned} X_{21}(1) &= F_3 \cup F_5((1 + \sqrt{21})/2) \\ &\neq F_3 \cup F_5. \end{aligned}$$

(The full curve  $F_3$  appears because the only solutions to  $N(\xi) = \xi^2/4 = -3 \pmod{21}$  are  $\xi = \pm 3$ .)

Using Theorem 3.6, it is similarly straightforward to check other small discriminants; for example:

**Theorem 3.9** *For  $D \leq 30$  we have  $X_D(1) = \bigcup_{e^2+4N=D} T_N$  when  $D = 4, 5, 8, 9, 13, 17, 25$  and  $29$ , but not when  $D = 12, 16, 20, 21, 24$  or  $28$ .*

**Notes.** For more background on modular curves and surfaces, see [Hir], [HZ2], [HZ1], [BL], [Mc7, §4] and [vG]. Our  $U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix}$  corresponds to the skew-Hermitian matrix  $B = \sqrt{D} \begin{pmatrix} a & \mu \\ \mu' & bD \end{pmatrix}$  in [vG, Ch. V]. Note that (3.3) agrees with the standard action  $(a\tau + b)/(c\tau + d)$  up to the automorphism  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a & -b \\ -c & d \end{pmatrix}$  of  $\mathrm{SL}_2(K)$ . We remark that  $X_D$  can also be presented as the quotient  $(\mathbb{H} \times -\mathbb{H})/\mathrm{SL}_2(\mathcal{O}_D)$ , using the fact that  $\sqrt{D}' = -\sqrt{D}$ ; on the other hand, the surfaces  $(\mathbb{H} \times \mathbb{H})/\mathrm{SL}_2(\mathcal{O}_D)$  and  $X_D$  are generally not isomorphic (see e.g. [HH].)

It is known that the intersection numbers  $\langle T_N, T_M \rangle$  form the coefficients of a modular form [HZ1], [vG, Ch. VI]. The results of [GKZ] suggest that the intersection numbers of the refined modular curves  $T_N(\nu)$  may similarly yield a Jacobi form.

## 4 Laminations

In this section we show algebraically that  $\tilde{X}_D(1)$  gives a lamination of  $\mathbb{H} \times \mathbb{H}$  by countably many disjoint hyperbolic planes. We also describe these

laminations explicitly for small values of  $D$ . Another proof of laminarity appears in §7.

**Jacobian varieties.** Let  $\Omega(X)$  denote the space of holomorphic 1-forms on a compact Riemann surface  $X$ . The *Jacobian* of  $X$  is the Abelian variety  $\text{Jac}(X) = \Omega(X)^*/H_1(X, \mathbb{Z})$ , polarized by the intersection pairing on 1-cycles.

In the case of genus two, any principally polarized Abelian variety  $A$  is either a Jacobian or a product of polarized elliptic curves. The latter case occurs iff  $A$  admits real multiplication by  $\mathcal{O}_1$ , generated by projection to one of the factors of  $A \cong B_1 \times B_2$ . In particular, we have:

**Theorem 4.1** *For any  $D \geq 4$ , the locus of Jacobian varieties in  $X_D$  is given by  $X_D - X_D(1)$ .*

**Laminations.** To describe  $X_D(1)$  in more detail, given  $N > 0$  such that  $D = e^2 + 4N$  let

$$\Lambda_D^N = \{U \in M_2(K) : U \text{ is a generator matrix, } \det(U) = N \text{ and } \nu(U) \equiv \pm(e + \sqrt{D})/2 \pmod{\mathcal{D}_D}\},$$

and let  $\Lambda_D$  be the union of all such  $\Lambda_D^N$ . Note that if  $U$  is in  $\Lambda_D$ , then  $-U, U'$  and  $U^*$  are also in  $\Lambda_D$ .

By Corollary 3.5, the preimage of  $X_D(1)$  in  $\tilde{X}_D = \mathbb{H} \times \mathbb{H}$  is given by:

$$\tilde{X}_D(1) = \bigcup \{\mathbb{H}_U : U \in \Lambda_D\}.$$

Note that each  $\mathbb{H}_U$  is the graph of a Möbius transformation.

**Theorem 4.2** *The locus  $\tilde{X}_D(1)$  gives a lamination of  $\mathbb{H} \times \mathbb{H}$  by countably many hyperbolic planes.*

(This means any two planes in  $\tilde{X}_D(1)$  are either identical or disjoint.)

For the proof, it suffices to show that the difference  $g \circ h^{-1}$  of two Möbius transformations in  $\Lambda_D$  is never elliptic. Since  $\Lambda_D$  is invariant under  $U \mapsto U^* = (\det U)U^{-1}$ , this in turn follows from:

**Theorem 4.3** *For any  $U_1, U_2 \in \Lambda_D$ , we have  $\text{Tr}(U_1 U_2)^2 \geq 4 \det(U_1 U_2)$ .*

**Proof.** By the definition of  $\Lambda_D$ , we can write  $D = e_i^2 + 4 \det(U_i) = e_i^2 + 4N_i$ , where  $e_i \geq 0$ . We can also assume that

$$U_i = \begin{pmatrix} \mu_i & b_i D \\ -a_i & -\mu'_i \end{pmatrix}$$

satisfies

$$\mu_i \equiv (x_i + y_i\sqrt{D})/2 \equiv (e_i + \sqrt{D})/2 \pmod{\mathcal{D}_D}$$

(replacing  $U_i$  with  $-U_i$  if necessary). It follows that  $y_i$  is odd and  $x_i \equiv e_i \pmod{D}$ , which implies

$$\mathrm{Tr}(U_1U_2) \equiv \mathrm{Tr}(\mu_1\mu_2) = (x_1x_2 + Dy_1y_2)/2 \equiv (e_1e_2 - D)/2 \pmod{D}. \quad (4.1)$$

(The factor of  $1/2$  presents no difficulties, because  $x_i$  is even when  $D$  is even.)

Now suppose

$$\mathrm{Tr}(U_1U_2)^2 < 4 \det(U_1U_2) = 4N_1N_2. \quad (4.2)$$

Then we have  $|\mathrm{Tr}(U_1U_2)| < 2\sqrt{N_1N_2} \leq D/2$ , and thus (4.1) implies

$$\mathrm{Tr}(U_1U_2) = (e_1e_2 - D)/2.$$

But this implies

$$\begin{aligned} 4 \mathrm{Tr}(U_1U_2)^2 &= (D - e_1e_2)^2 \\ &\geq (D - e_1^2)(D - e_2^2) = (4N_1)(4N_2) = 16 \det(U_1U_2), \end{aligned}$$

contradicting (4.2). ■

**Small discriminants.** To conclude we record a few cases where  $\Lambda_D$  admits a particularly economical description.

For concreteness, we will present  $\Lambda_D$  as a set matrices in  $\mathrm{GL}_2^+(\mathbb{R})$  using the chosen real place  $\iota_1 : K \rightarrow \mathbb{R}$ . This works even when  $D = d^2$ , since both  $\mu$  and  $\mu'$  appear on the diagonal of  $U \in \Lambda_D$  (no information is lost). Under the standard action  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot z = (az + b)/(cz + d)$  of  $\mathrm{GL}_2^+(\mathbb{R})$  on  $\mathbb{H}$ , we can then write

$$\tilde{X}_D(1) = \bigcup_{\Lambda_D} \{(\tau_1, \tau_2) : \tau_2 = U(\tau_1)\}.$$

This holds despite the twist in the definition (3.5) of  $\mathbb{H}_U$ , because  $\Lambda_D$  is invariant under  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} d & b \\ c & a \end{pmatrix}$ .

**Theorem 4.4** *For  $D = 4, 5, 8, 9$  and  $13$  respectively, we have:*

$$\begin{aligned} \Lambda_4 &= \{U \in \mathrm{M}_2(\mathbb{Z}) : \det(U) = 1 \text{ and } U \equiv \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} \pmod{4}\}, \\ \Lambda_5 &= \{U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} : \det(U) = 1\}, \\ \Lambda_8 &= \Lambda_8^1 \cup \Lambda_8^2 = \left\{U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} : \det(U) = 1 \text{ or } 2\right\}, \\ \Lambda_9 &= \{U \in \mathrm{M}_2(\mathbb{Z}) : \det(U) = 2 \text{ and } U \equiv \begin{pmatrix} * & 0 \\ * & * \end{pmatrix} \pmod{9}\}, \quad \text{and} \\ \Lambda_{13} &= \Lambda_{13}^1 \cup \Lambda_{13}^3 = \left\{U = \begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix} : \det(U) = 1 \text{ or } 3\right\}, \end{aligned}$$

where it is understood that  $a, b \in \mathbb{Z}$  and  $\mu \in \mathcal{O}_D$ .

**Proof.** Recall from Theorem 3.9 that  $X_D(1) = \bigcup_{e^2+4N=D} T_N$  when  $D = 4, 5, 8, 9$  and  $13$ . When this equality holds, we can ignore the condition on  $\nu(U)$  in the definition of  $\Lambda_D$ . The cases  $D = 5, 8$  and  $13$  then follow directly from the definition of  $\Lambda_D^N$ . For  $D = 9$ , we note that any integral matrix satisfying  $\det \begin{pmatrix} x & 9b \\ -a & y \end{pmatrix} = 2$  also satisfies  $x + y = 0 \pmod{3}$ , and thus it can be written in the form  $\begin{pmatrix} \mu & bD \\ -a & -\mu' \end{pmatrix}$  with

$$\mu = \frac{(x - y) + (x + y)\sqrt{9/3}}{2}.$$

Similar considerations apply when  $D = 4$ . ■

## 5 Foliations of Teichmüller space

In this section we introduce a family of foliations  $\mathcal{F}_i$  of Teichmüller space, related to normalized Abelian differentials and their periods  $\tau_{ij} = \int_{b_i} \omega_j$ . We then show:

**Theorem 5.1** *There is a unique holomorphic section of the period map*

$$\tau_{ii} : \mathcal{T}_g \rightarrow \mathbb{H}$$

*through any  $Y \in \mathcal{T}_g$ . Its image is the leaf of  $\mathcal{F}_i$  containing  $Y$ .*

The case  $g = 2$  will furnish the desired foliations of Hilbert modular surfaces.

**Abelian differentials.** Let  $Z_g$  be a smooth oriented surface of genus  $g$ . Let  $\mathcal{T}_g$  be the Teichmüller space of Riemann surfaces  $Y$ , each equipped with an isotopy class of homeomorphism or *marking*  $Z_g \rightarrow Y$ . The marking determines a natural identification between  $H_1(Z_g)$  and  $H_1(Y)$  used frequently below.

Let  $\Omega\mathcal{T}_g \rightarrow \mathcal{T}_g$  denote the bundle of nonzero Abelian differentials  $(Y, \omega)$ ,  $\omega \in \Omega(Y)$ . For each such form we have a *period map*

$$I(\omega) : H_1(Z_g, \mathbb{Z}) \rightarrow \mathbb{C}$$

given by  $I(\omega) : C \rightarrow \int_C \omega$ . There is a natural action of  $\mathrm{GL}_2^+(\mathbb{R})$  on  $\Omega\mathcal{T}_g$ , satisfying

$$I(A \cdot \omega) = A \circ I(\omega) \tag{5.1}$$

under the identification  $\mathbb{C} = \mathbb{R}^2$  given by  $x + iy = (x, y)$ .

Each orbit  $\mathrm{GL}_2^+(\mathbb{R}) \cdot (Y, \omega)$  projects to a *complex geodesic*

$$f : \mathbb{H} \rightarrow \mathcal{T}_g,$$

which can be normalized so that  $f(i) = Y$  and

$$\nu = \left. \frac{df}{dt} \right|_{t=i} = \frac{i \bar{\omega}}{2\omega}.$$

The subspace of  $H^1(Z_g, \mathbb{R})$  spanned by  $(\mathrm{Re} \omega, \mathrm{Im} \omega)$  is constant along each orbit (cf. [Mc7, §3]).

**Symplectic framings.** Now let  $(a_1, \dots, a_g, b_1, \dots, b_g)$  be a real symplectic basis for  $H_1(Z_g, \mathbb{R})$  (with  $\langle a_i, b_i \rangle = -\langle b_i, a_i \rangle = 1$  and all other products zero). Then for each  $Y \in \mathcal{T}_g$ , there exists a unique basis  $(\omega_1, \dots, \omega_g)$  of  $\Omega(Y)$  such that  $\int_{a_i} \omega_j = \delta_{ij}$ . The *period matrix*

$$\tau_{ij}(Y) = \int_{b_i} \omega_j$$

then determines an embedding

$$\tau : \mathcal{T}_g \rightarrow \mathfrak{H}_g.$$

This agrees with the usual Torelli embedding, up to composition with an element of  $\mathrm{Sp}_{2g}(\mathbb{R})$ . Note that  $\mathrm{Im}(\tau_{ii}(Y)) > 0$  since  $\mathrm{Im} \tau$  is positive definite.

The normalized 1-forms  $(\omega_i)$  give a splitting

$$\Omega(Y) = \bigoplus_1^g \mathbb{C} \omega_i = \bigoplus_1^g F_i(Y),$$

and corresponding subbundles  $F_i \mathcal{T}_g \subset \Omega \mathcal{T}_g$ .

**Complex subspaces.** Let  $(a_i^*, b_i^*)$  denote the dual basis for  $H^1(Z_g, \mathbb{R})$ , and let  $S_i$  be the span of  $(a_i^*, b_i^*)$ . It is easy to check that the following conditions are equivalent:

1.  $S_i$  is a complex subspace of  $H^1(Y, \mathbb{R}) \cong \Omega(Y)$ .
2.  $S_i$  is spanned by  $(\mathrm{Re} \omega_i, \mathrm{Im} \omega_i)$ .
3. The period matrix  $\tau(Y)$  satisfies  $\tau_{ij} = 0$  for all  $j \neq i$ .

Let  $\mathcal{T}_g(S_i) \subset \mathcal{T}_g$  denote the locus where these conditions hold. Note that condition (3) defines a totally geodesic subset

$$H_i \cong \mathbb{H} \times \mathfrak{H}_{g-1} \subset \mathfrak{H}_g$$

such that  $\mathcal{T}_g(S_i) = \tau^{-1}(H_i)$ .

**Foliations.** Next we show that the complex geodesics generated by the forms  $(Y, \omega_i)$  give a foliation of Teichmüller space.

**Theorem 5.2** *The sub-bundle  $F_i\mathcal{T}_g \subset \Omega\mathcal{T}_g$  is invariant under the action of  $\mathrm{GL}_2^+(\mathbb{R})$ , as is its restriction to  $\mathcal{T}_g(S_i)$ .*

**Proof.** The invariance of  $F_i\mathcal{T}_g$  is immediate from (5.1). To handle the restriction to  $\mathcal{T}_g(S_i)$ , recall that the span  $W$  of  $(\mathrm{Re} \omega_i, \mathrm{Im} \omega_i)$  is constant along orbits; thus the condition  $W = S_i$  characterizing  $\mathcal{T}_g(S_i)$  is preserved by the action of  $\mathrm{GL}_2^+(\mathbb{R})$ . ■

**Corollary 5.3** *The foliation of  $F_i\mathcal{T}_g$  by  $\mathrm{GL}_2^+(\mathbb{R})$  orbits projects to a foliation  $\mathcal{F}_i$  of  $\mathcal{T}_g$  by complex geodesics.*

**Corollary 5.4** *The locus  $\mathcal{T}_g(S_i)$  is also foliated by  $\mathcal{F}_i$ : any leaf meeting  $\mathcal{T}_g(S_i)$  is entirely contained therein.*

**Proof of Theorem 5.1.** The proof uses Ahlfors' variational formula [Ah] and follows the same lines as the proof of [Mc4, Thm. 4.2]; it is based on the fact that the leaves of  $\mathcal{F}_i$  are the geodesics along which the periods of  $\omega_i$  change most rapidly.

Let  $s : \mathbb{H} \rightarrow \mathcal{T}_g$  be a holomorphic section of  $\tau_{ii}$ . Let  $v \in T\mathbb{H}$  be a unit tangent vector with respect to the hyperbolic metric  $\rho = |dz|/(2\mathrm{Im} z)$  of constant curvature  $-4$ , mapping to  $Ds(v) \in T_Y\mathcal{T}_g$ . By the equality of the Teichmüller and Kobayashi metrics [Gd, Ch. 7],  $Ds(v)$  is represented by a Beltrami differential  $\nu = \nu(z)d\bar{z}/dz$  on  $Y$  with  $\|\nu\|_\infty \leq 1$ . But  $s$  is a section, so the composition

$$\tau_{ii} \circ s : \mathbb{H} \rightarrow \mathbb{H}$$

is the identity; thus the norm of its derivative, given by Ahlfors' formula as

$$\|D(\tau_{ii} \circ s)(\nu)\| = \left| \int_Y \omega_i^2 \nu \right| / \int_Y |\omega_i|^2,$$

is one. It follows that  $\nu = \bar{\omega}_i/\omega_i$  up to a complex scalar of modulus one, and thus  $Ds(v)$  is tangent to the complex geodesic generated by  $(Y, \omega_i)$ . Equivalently,  $s(\mathbb{H})$  is everywhere tangent to the foliation  $\mathcal{F}_i$ ; therefore its image is the unique leaf through  $Y$ . ■



## 6 Genus two

We can now obtain results on Hilbert modular surfaces by specializing to the case of genus two. In this section we will show:

**Theorem 6.1** *There is a unique holomorphic section of  $\tau_1$  passing through any given point of  $\mathbb{H} \times \mathbb{H} - \tilde{X}_D(1)$ .*

Here  $\tau_1 : \mathbb{H} \times \mathbb{H} \rightarrow \mathbb{H}$  is simply projection onto the first factor. This result is a restatement of Theorem 1.2; as in §1, we assume  $D \geq 4$ .

**Framings for real multiplication.** Let  $g = 2$ , and choose a symplectic isomorphism

$$L = H_1(Z_g, \mathbb{Z}) \cong \mathcal{O}_D \oplus \mathcal{O}_D^\vee.$$

We then have an action of  $\mathcal{O}_D$  on  $H_1(Z_g, \mathbb{Z})$ , and the elements  $\{a, b\} = \{(1, 0), (0, 1)\}$  in  $L$  give a distinguished basis for

$$H_1(Z_g, \mathbb{Q}) = L \otimes \mathbb{Q} \cong K^2$$

as a vector space over  $K = \mathcal{O}_D \otimes \mathbb{Q}$ . Using the two Galois conjugate embeddings  $K \rightarrow \mathbb{R}$ , we obtain an orthogonal splitting

$$H_1(Z_g, \mathbb{R}) = L \otimes \mathbb{R} = V_1 \oplus V_2$$

such that  $k \cdot (C_1, C_2) = (kC_1, k'C_2)$ . The projections  $(a_i, b_i)$  of  $a, b \in L$  to each summand yield bases for  $V_i$ , which taken together give a standard symplectic basis for  $H_1(Z_g, \mathbb{R})$ . (Note that  $(a_i, b_i)$  is generally *not* an integral symplectic basis; indeed, when  $K$  is a field, the elements  $(a_i, b_i)$  do not even lie in  $H_1(Z_g, \mathbb{Q})$ .)

Let  $S_i^D \subset H^1(Z_g, \mathbb{R})$  be the span of the dual basis  $a_i^*, b_i^*$ .

**Theorem 6.2** *The ring  $\mathcal{O}_D \subset \text{End}(L)$  acts by real multiplication on  $\text{Jac}(Y)$  if and only if  $Y \in \mathcal{T}_g(S_1^D)$ .*

**Proof.** Since  $g = 2$  we have  $S_2^D = (S_1^D)^\perp$ , and thus  $\mathcal{T}_g(S_1^D) = \mathcal{T}_g(S_2^D)$ . But  $\text{Jac}(Y)$  has real multiplication iff  $S_1^D$  and  $S_2^D$  are complex subspaces of  $H^1(Y, \mathbb{R}) \cong \Omega(Y)$  so the result follows. (Cf. [Mc4, Lemma 7.4].) ■

**Sections.** Let  $E_D = X_D - X_D(1)$  denote the space of Jacobians in  $X_D$ , and  $\tilde{E}_D = \mathbb{H} \times \mathbb{H} - \tilde{X}_D(1)$  its preimage in the universal cover. (The notation comes from [Mc7, §4], where we consider the space of eigenforms  $\Omega E_D$  as a closed,  $\mathrm{GL}_2^+(\mathbb{R})$ -invariant subset of  $\Omega \mathcal{M}_g$ .)

By the preceding result, the Jacobian of any  $Y \in \mathcal{T}_g(S_1^D)$  is an Abelian variety with real multiplication. Moreover, the marking of  $Y$  determines a marking

$$L \cong H_1(Y, \mathbb{Z}) \cong H_1(\mathrm{Jac}(Y), \mathbb{Z})$$

of its Jacobian, and thus a map

$$\mathrm{Jac} : \mathcal{T}_g(S_1^D) \rightarrow \tilde{E}_D = \tilde{X}_D - \tilde{X}_D(1).$$

The basis  $(a_i, b_i)$  yields a pair of normalized forms  $\omega_1, \omega_2 \in \Omega(Y)$ . Similarly, we have a pair of normalized eigenforms  $\eta_1, \eta_2 \in \Omega(A_\tau)$  for each  $\tau \in \tilde{X}_D$ , characterized by (3.2). Under the identification  $\Omega(Y) = \Omega(\mathrm{Jac}(Y))$ , we find:

**Theorem 6.3** *The forms  $\omega_i$  and  $\eta_i$  are equal for any  $Y \in \mathcal{T}_g(S_1^D)$ . Thus  $\mathrm{Jac}(Y) = A_{(\tau_1, \tau_2)}$ , where*

$$\begin{pmatrix} \tau_1 & 0 \\ 0 & \tau_2 \end{pmatrix} = \tau_{ij}(Y) = \left( \int_{b_i} \omega_j \right). \quad (6.1)$$

**Proof.** The period map  $\phi_\tau : L \rightarrow \mathbb{C}^2$  for  $A_\tau = \mathrm{Jac}(Y)$  is given by

$$\phi_\tau(C) = \left( \int_C \eta_1, \int_C \eta_2 \right) = (x_1 + x_2\tau_1, x'_1 + x'_2\tau_2),$$

where  $C = (x_1, x_2) \in \mathcal{O}_D \oplus \mathcal{O}_D^\vee$ ; in particular, we have

$$\phi_\tau(a) = \phi_\tau(1, 0) = (1, 1).$$

Since  $\phi_\tau$  diagonalizes the action of  $K$ , we also have

$$\phi_\tau(C) = \left( \int_{C_1} \eta_1, \int_{C_2} \eta_2 \right)$$

for any  $C = C_1 + C_2 \in L \otimes \mathbb{R} = V_1 \oplus V_2$ . Setting  $C = a$ , this implies  $\phi_\tau(a_1) = (1, 0)$  and  $\phi_\tau(a_2) = (0, 1)$ ; thus  $\int_{a_i} \eta_j = \delta_{ij}$ , and therefore  $\eta_i = \omega_i$  for  $i = 1, 2$ . Similarly, we have

$$\phi_\tau(b) = (\tau_1, \tau_2) = (\tau_{11}, \tau_{22}),$$

which implies  $Y$  and  $A_\tau$  are related by (6.1). ■

**Corollary 6.4** *We have a commutative diagram*

$$\begin{array}{ccc}
 \mathcal{T}_g(S_1^D) & \xrightarrow{\text{Jac}} & \tilde{E}_D \\
 & \searrow \tau_{11} & \downarrow \tau_1 \\
 & & \mathbb{H}.
 \end{array}$$

**Proof of Theorem 6.1.** Using the Torelli theorem, it follows easily that  $\text{Jac} : \mathcal{T}_g(S_1^D) \rightarrow \tilde{E}_D$  is a holomorphic covering map. Since  $\mathbb{H}$  is simply-connected, any section  $s$  of  $\tau_1$  lifts to a section  $\text{Jac}^{-1} \circ s$  of  $\tau_{11}$ . Thus Theorem 5.1 immediately implies Theorem 6.1. ■

## 7 Holomorphic motions

In this section we use the theory of holomorphic motions to define and characterize the foliation  $\mathcal{F}_D$ .

**Holomorphic motions.** Given a set  $E \subset \hat{\mathbb{C}}$  and a basepoint  $s \in \mathbb{H}$ , a *holomorphic motion* of  $E$  over  $(\mathbb{H}, s)$  is a family of injective maps

$$F_t : E \rightarrow \hat{\mathbb{C}}, \quad t \in \mathbb{H},$$

such that  $F_s(z) = z$  and  $F_t(z)$  is a holomorphic function of  $t$ .

A holomorphic motion of  $E$  has a unique extension to a holomorphic motion of its closure  $\bar{E}$ ; and each map  $F_t : E \rightarrow \hat{\mathbb{C}}$  extends to a quasiconformal homeomorphism of the sphere. In particular,  $F_t|_{\text{int}(E)}$  is quasiconformal (see e.g. [Dou]).

These properties imply:

**Theorem 7.1** *Let  $P$  be a partition of  $\mathbb{H} \times \mathbb{H}$  into disjoint graphs of holomorphic functions. Then:*

1.  $P$  is the set of leaves of a transversally quasiconformal foliation  $\mathcal{F}$  of  $\mathbb{H} \times \mathbb{H}$ ; and
2. If we adjoin the graphs of the constant functions  $f : \mathbb{H} \rightarrow \partial\mathbb{H}$  to  $P$ , we obtain a continuous foliation of  $\mathbb{H} \times \bar{\mathbb{H}}$ .

**The foliation  $\mathcal{F}_D$ .** Recall that every component of  $\tilde{X}_D(1) \subset \mathbb{H} \times \mathbb{H}$  is the graph of a Möbius transformation. By Theorem 6.1, there is a unique partition of  $\mathbb{H} \times \mathbb{H} - \tilde{X}_D(1)$  into the graphs of holomorphic maps as well.

Taken together, these graphs form the leaves of a foliation  $\tilde{\mathcal{F}}_D$  of  $\mathbb{H} \times \mathbb{H}$  by the preceding result. Since  $\tilde{X}_D(1)$  is invariant under  $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$ , the foliation  $\tilde{\mathcal{F}}_D$  descends to a foliation  $\mathcal{F}_D$  of  $X_D$ .

To characterize  $\mathcal{F}_D$ , recall that the surface  $X_D$  admits a holomorphic involution  $\iota(\tau_1, \tau_2) = (\tau_2, \tau_1)$  which preserves  $X_D(1)$ .

**Theorem 7.2** *The only leaves shared by  $\mathcal{F}_D$  and  $\iota(\mathcal{F}_D)$  are the curves in  $X_D(1)$ .*

**Proof.** Let  $f : \mathbb{H} \rightarrow \mathbb{H}$  be a holomorphic function whose graph  $F$  is both a leaf of  $\tilde{\mathcal{F}}_D$  and  $\iota(\tilde{\mathcal{F}}_D)$ . Then  $\iota(F)$  is also a graph, so  $f$  is an isometry. But if  $F \cap \tilde{X}_D(1) = \emptyset$ , then  $F$  lifts to a leaf of the foliation  $\mathcal{F}_1$  of Teichmüller space, and hence  $f$  is a contraction by [Mc4, Thm. 4.2]. ■

**Corollary 7.3** *The only leaves of  $\tilde{\mathcal{F}}_D$  that are graphs of Möbius transformations are those belonging to  $\tilde{X}_D(1)$ .*

**Complex geodesics.** Let us say  $\mathcal{F}$  is a foliation by *complex geodesics* if each leaf is a hyperbolic Riemann surface, isometrically immersed for the Kobayashi metric. We can then characterize  $\mathcal{F}_D$  as follows.

**Theorem 7.4** *Up to the action of  $\iota$ ,  $\mathcal{F}_D$  is the unique extension of the lamination  $X_D(1)$  to a foliation of  $X_D$  by complex geodesics.*

**Proof.** Let  $\mathcal{F}$  be a foliation by complex geodesics extending  $X_D(1)$ . Then every leaf of its lift  $\tilde{\mathcal{F}}$  to  $\tilde{X}_D$  is a Kobayashi geodesic for  $\mathbb{H} \times \mathbb{H}$ . But a complex geodesic in  $\mathbb{H} \times \mathbb{H}$  is either the graph of a holomorphic function or its inverse, so every leaf belongs to either  $\tilde{\mathcal{F}}_D$  or  $\iota(\tilde{\mathcal{F}}_D)$ . Consequently every leaf of  $\mathcal{F}$  is a leaf of  $\mathcal{F}_D$  or  $\iota(\mathcal{F}_D)$ . Since these foliations have no leaves in common on the open set  $U = X_D - X_D(1)$ ,  $\mathcal{F}$  coincides with one or the other. ■

**Stable curves.** The Abelian varieties  $E \times F$  in  $X_D(1)$  are the Jacobians of certain *stable curves* with real multiplication, namely the nodal curves  $Y = E \vee F$  obtained by gluing  $E$  to  $F$  at a single point. If we adjoin these stable curves to  $\mathcal{M}_2$ , we obtain a partial compactification  $\mathcal{M}_2^*$  which maps isomorphically to  $\mathcal{A}_2$ . The locus  $X_D(1)$  can then be regarded as the projection to  $X_D$  of a finite set of  $\mathrm{GL}_2^+(\mathbb{R})$  orbits in  $\Omega\mathcal{M}_2^*$ , giving another proof that it is a lamination.

## 8 Quasiconformal dynamics

In this section we use the relative period map  $\rho = \int_{y_1}^{y_2} \eta_1$  to define a meromorphic quadratic differential  $q = (d\rho)^2$  transverse to  $\mathcal{F}_D$ . We then show the transverse dynamics of  $\mathcal{F}_D$  is given by Teichmüller mappings relative to  $q$ .

**Absolute periods.** The level sets of  $\tau_1$  form the leaves of a holomorphic foliation  $\tilde{\mathcal{A}}_D$  on  $\mathbb{H} \times \mathbb{H}$  which covers foliation  $\mathcal{A}_D$  of  $X_D$ . By (3.2), every  $\tau = (\tau_1, \tau_2)$  determines a pair of eigenforms  $\eta_1, \eta_2 \in \Omega(A_\tau)$  such that the *absolute periods*

$$\int_C \eta_1, \quad C \in H_1(A_\tau, \mathbb{Z})$$

are constant along the leaves of  $\tilde{\mathcal{A}}_D$ . Since every leaf of  $\tilde{\mathcal{F}}_D$  is the graph of a function  $f : \mathbb{H} \rightarrow \mathbb{H}$ , we have:

**Theorem 8.1** *The foliation  $\mathcal{A}_D$  is transverse to  $\mathcal{F}_D$ .*

**The Weierstrass curve.** Recall that  $E_D \subset X_D$  denotes the locus of Jacobians with real multiplication by  $\mathcal{O}_D$ . For  $[A_\tau] = \text{Jac}(Y) \in E_D$  we can regard the eigenforms  $\eta_1, \eta_2$  as holomorphic 1-forms in  $\Omega(Y) \cong \Omega(A_\tau)$ .

Let  $W_D \subset E_D$  denote the locus where  $\eta_1$  has a double zero on  $Y$ . By [Mc5] we have:

**Theorem 8.2** *The locus  $W_D$  is an algebraic curve with one or two irreducible components, each of which is a leaf of  $\mathcal{F}_D$ .*

We refer to  $W_D$  as the *Weierstrass curve*, since  $\eta_1$  vanishes at a Weierstrass point of  $Y$ .

**Relative periods.** Let  $E_D(1, 1) = X_D - (W_D \cup X_D(1))$  denote the Zariski open set where  $\eta_1$  has a pair of simple zeros, and let  $\tilde{E}_D(1, 1)$  be its preimage in the universal cover  $\tilde{X}_D$ . Let

$$\mathbb{H}_s = \{s\} \times \mathbb{H} \subset \mathbb{H} \times \mathbb{H},$$

and let  $\mathbb{H}_s^* = \mathbb{H}_s \cap \tilde{E}_D(1, 1)$ .

For each  $\tau \in \mathbb{H}_s^*$ , let  $y_1, y_2$  denote the zeros of the associated form  $\eta_1 \in \Omega(Y)$ . We can then define the (multivalued) *relative period map*  $\rho_s : \mathbb{H}_s^* \rightarrow \mathbb{C}$  by

$$\rho_s(\tau) = \int_{y_1}^{y_2} \eta_1.$$

To make  $\rho_s(\tau)$  single-valued, we must (locally) choose (i) an ordering of the zeros  $y_1$  and  $y_2$ , and (ii) a path on  $Y$  connecting them.

**Quadratic differentials.** Let  $z$  be a local coordinate on  $\mathbb{H}_s$ , and recall that the absolute periods of  $\eta_1$  are constant along  $\mathbb{H}_s$ . Thus if we change the choice of path from  $y_1$  to  $y_2$ , the derivative  $d\rho/dz$  remains the same; and if we interchange  $y_1$  and  $y_2$ , it changes only by sign. Thus the *quadratic differential*

$$q = (d\rho/dz)^2 dz^2$$

is globally well-defined on  $\mathbb{H}_s^*$ .

**Theorem 8.3** *The form  $q$  extends to a meromorphic quadratic differential on  $\mathbb{H}_s$ , with simple zeros where  $\mathbb{H}_s$  meets  $\widetilde{W}_D$ , and simple poles where it meets  $\widetilde{X}_D(1)$ .*

**Proof.** It is a general result that the period map provides holomorphic local coordinates on any stratum of  $\Omega\mathcal{M}_g$  (see [V2], [MS, Lemma 1.1], [KZ]). Thus  $\rho_s|_{\mathbb{H}_s^*}$  is holomorphic with  $d\rho_s \neq 0$ , and hence  $q|_{\mathbb{H}_s^*}$  is a nowhere vanishing holomorphic quadratic differential.

To see  $q$  acquires a simple zero when  $\eta_1$  acquires a double zero, note that the relative period map

$$\rho(t) = \int_{-\sqrt{t}}^{\sqrt{t}} (z^2 - t) dz = (-4/3)t^{3/2}$$

of the local model  $\eta_t = (z^2 - t) dz$  satisfies  $(d\rho/dt)^2 = 4t$ . Similarly, a point of  $\mathbb{H}_s \cap \widetilde{X}_D(1)$  is locally modeled by the family of connected sums

$$(Y_t, \eta_t) = (E_1, \omega_1) \#_I (E_2, \omega_2),$$

with  $I = [0, \rho(t)] = [0, \pm\sqrt{t}]$ . Since  $(d\rho/dt)^2 = 1/(4t)$ , at these points  $q$  has simple poles. ■

See [Mc7, §6] for more on connected sums.

**Teichmüller maps.** Now let  $f : \mathbb{H}_s \rightarrow \mathbb{H}_t$  be a quasiconformal map. We say  $f$  is a *Teichmüller map*, relative to a holomorphic quadratic differential  $q$ , if its complex dilatation satisfies

$$\mu(f) = \left( \frac{\partial f / \partial \bar{z}}{\partial f / \partial z} \right) \frac{d\bar{z}}{dz} = \alpha \frac{\bar{q}}{|q|}$$

for some  $\alpha \in \mathbb{C}^*$ . This is equivalent to the condition that  $w = f(z)$  is real-linear in local coordinates where  $q = dz^2$  and  $dw^2$  respectively. In such charts we can write

$$w = w_0 + D_q(f) \cdot z,$$

with  $D_q(f) \in \mathrm{SL}_2(\mathbb{R})$ . We refer to  $D_q(f)$  as the *linear part* of  $f$ ; it is only well-defined up to sign, since  $z \mapsto -z$  preserves  $dz^2$ .

**Theorem 8.4** *Given  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  and  $s \in \mathbb{H}$ , let  $\mathbb{H}_t = g(\mathbb{H}_s)$ . Then the linear part of  $g : \mathbb{H}_s \rightarrow \mathbb{H}_t$  is given by  $D_q(g) \cdot z = (d - cs)^{-1}z$ .*

**Proof.** Since the Riemann surfaces  $Y$  at corresponding points of  $\mathbb{H}_s$  and  $\mathbb{H}_t$  differ only by marking, the relative period maps  $\rho_s$  and  $\rho_t$  differ only by the normalization of  $\eta_1$ . This discrepancy is accounted for by equation (3.4), which gives  $\rho_t/\rho_s = \chi(g, s) = (d - cs)^{-1}$ . Since the coordinates  $\rho_s$  and  $\rho_t$  linearize  $q$ , the map  $D_q(g)$  is given by multiplication by  $(d - cs)^{-1}$ . ■

Now let  $C_{st} : \mathbb{H}_s \rightarrow \mathbb{H}_t$  be the unique map such that  $z$  and  $C_{st}(z)$  lie on the same leaf of  $\tilde{\mathcal{F}}_D$ .

**Theorem 8.5** *The linear part of  $C_{st}$  is given by  $D_q(C_{st}) = A_t A_s^{-1}$ , where  $A_u = \begin{pmatrix} 1 & \mathrm{Re}(u) \\ 0 & \mathrm{Im}(u) \end{pmatrix} \in \mathrm{PSL}_2(\mathbb{R})$ .*

**Proof.** By the definition of  $\mathcal{F}_D$ , the forms  $\eta_1$  at corresponding points of  $\mathbb{H}_s^*$  and  $\mathbb{H}_t^*$  are related by some element  $B \in \mathrm{GL}_2^+(\mathbb{R})$  acting on  $\Omega\mathcal{T}_g$ . Thus  $\rho_t = B \circ \rho_s$  and therefore  $D_q(C_{st}) = B$ . Since the action of  $B$  on the absolute periods of  $\eta_1$  satisfies

$$B(\mathcal{O}_D \oplus \mathcal{O}_D^\vee s) = \mathcal{O}_D \oplus \mathcal{O}_D^\vee t$$

(in the sense of equation (3.1)), we have  $B(1) = 1$  and  $B(s) = t$ , and thus  $B = A_t A_s^{-1}$  as above. ■

**Dynamics.** Every leaf of  $\tilde{\mathcal{F}}_D$  meets the transversal  $\mathbb{H}_s$  in a single point. Thus the action of  $g \in \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  on the space of leaves determines a *holonomy map*

$$\phi_g : \mathbb{H}_s \rightarrow \mathbb{H}_s,$$

characterized by the property that  $(s, \phi_g(z))$  lies on the same leaf as  $g(s, z)$ .

**Theorem 8.6** *The group  $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  acts on  $\mathbb{H}_s$  by Teichmüller mappings, satisfying  $D_q(\phi_g) = g$  in the case  $s = i$ .*

(As usual we regard  $g$  as a real matrix using  $\iota_1 : K \rightarrow \mathbb{R}$ .)

**Proof.** Let  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , and  $t = (as - b)/(-cs + d)$ ; then  $\mathbb{H}_t = g(\mathbb{H}_s)$ .

Since  $\phi_g(z)$  is obtained from  $g(s, z)$  by combing it along the leaves of  $\tilde{\mathcal{F}}_D$  back into  $\mathbb{H}_s$ , we have  $\phi_g(s, z) = C_{ts}(g(s, z))$ . Thus the chain rule implies

$$D_q(\phi_g) \cdot z = B \cdot z = A_s \circ A_t^{-1}(z/(-cs + d)).$$

Now assume  $s = i$ . Then we have  $B(ai - b) = A_t^{-1}(t) = i$  and  $B(-ci + d) = A_t^{-1}(1) = 1$ ; therefore  $B^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$  and thus  $B = \begin{pmatrix} a & b \\ c & d \end{pmatrix} = g$ .  $\blacksquare$

**Corollary 8.7** *The foliation  $\mathcal{F}_D$  carries a natural transverse invariant measure.*

**Proof.** Since  $\det D_q(\phi_g) = 1$  for all  $g$ , the form  $|q|$  gives a holonomy-invariant measure on the transversal  $\mathbb{H}_s$ .  $\blacksquare$

Finally we show that, although  $\phi_g|_{\mathbb{H}_s}$  is quasiconformal, its continuous extension to  $\partial\mathbb{H}_s$  is a Möbius transformation.

**Theorem 8.8** *For any  $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  and  $z \in \partial\mathbb{H}_s$ , we have*

$$\phi_g(z) = (a'z - b')/(-c'z + d').$$

**Proof.** By Theorem 7.1, the combing maps  $C_{st}$  extend to the identity on  $\partial\mathbb{H}_s$ . Thus  $(t, \phi_g(z)) = g(s, z)$ , and the result follows from equation (3.3).  $\blacksquare$

Note: if we use the transversal  $\mathbb{H}_t$  instead of  $\mathbb{H}_s$ , the holonomy simply changes by conjugation by  $C_{st}$ .

## 9 Further results

In this section we summarize related results on the density of leaves, isoperiodic forms, holomorphic motions and iterated rational maps.

**I. Density of leaves.** By [Mc7], the closure of the complex geodesic  $f : \mathbb{H} \rightarrow \mathcal{M}_2$  generated by a holomorphic 1-form is either an algebraic curve, a Hilbert modular surface or the whole moduli space. Since the leaves of  $\mathcal{F}_D$  are examples of such complex geodesics, we obtain:



**Theorem 9.1** *Every leaf of  $\mathcal{F}_D$  is either a closed algebraic curve, or a dense subset of  $X_D$ .*

It is easy to see that the union of the closed leaves is dense when  $D = d^2$ . On the other hand, the classification of Teichmüller curves in [Mc5] and [Mc6] implies:

**Theorem 9.2** *If  $D$  is not a square, then  $\mathcal{F}_D$  has only finitely many closed leaves. These consist of the components of  $W_D \cup X_D(1)$  and, when  $D = 5$ , the Teichmüller curve generated by the regular decagon.*

**II. Isoperiodic forms.** Next we discuss interactions between the foliations  $\mathcal{F}_D$  and  $\mathcal{A}_D$ . When  $D = d^2$  is a square, the surface  $X_D$  is finitely covered by a product, and hence every leaf of  $\mathcal{A}_D$  is closed.

**Theorem 9.3** *If  $D$  is not a square, then every leaf  $L$  of  $\mathcal{A}_D$  is dense in  $X_D$ , and  $L \cap F$  is dense in  $F$  for every leaf  $F$  of  $\mathcal{F}_D$ .*

**Proof.** The first result follows from the fact that  $\mathrm{SL}(\mathcal{O}_D \oplus \mathcal{O}_D^\vee)$  is a dense subgroup of  $\mathrm{SL}_2(\mathbb{R})$ , and the second follows from the first by transversality of  $\mathcal{A}_D$  and  $\mathcal{F}_D$ . ■

Let us say a pair of 1-forms  $(Y_i, \omega_i) \in \Omega\mathcal{M}_g$  are *isoperiodic* if there is a symplectic isomorphism

$$\phi : H_1(Y_1, \mathbb{Z}) \rightarrow H_1(Y_2, \mathbb{Z})$$

such that the period maps

$$I(\omega_i) : H_1(Y_i, \mathbb{Z}) \rightarrow \mathbb{C}$$

satisfy  $I(\omega_1) = I(\omega_2) \circ \phi$ . Since the absolute periods of  $\eta_1$  are constant along the leaves of  $\mathcal{A}_D$ , from the preceding result we obtain:

**Corollary 9.4** *The  $\mathrm{SL}_2(\mathbb{R})$ -orbit of any eigenform for real multiplication by  $\mathcal{O}_D$ ,  $D \neq d^2$ , contains infinitely many isoperiodic forms.*

For a concrete example, let  $Q \subset \mathbb{C}$  be a regular octagon containing  $[0, 1]$  as an edge. Identifying opposite sides of  $Q$ , we obtain the *octagonal form*

$$(Y, \omega) = (Q, dz) / \sim$$

of genus two.

Let  $\mathbb{Z}[\zeta] \subset \mathbb{C}$  denote the ring generated by  $\zeta = (1+i)/\sqrt{2} = \exp(2\pi i/8)$ , equipped with the symplectic form

$$\langle z_1, z_2 \rangle = \text{Tr}_{\mathbb{Q}}^{\mathbb{Q}(\zeta)}((\zeta + \zeta^2 + \zeta^3)z_1\bar{z}_2/4).$$

Then it is easy to check that:

1. The octagonal form  $\omega$  has a single zero of order 2, and
2. Its period map  $I(\omega)$  sends  $H_1(Y, \mathbb{Z})$  to  $\mathbb{Z}[\zeta]$  by a symplectic isomorphism.

However, these two properties do *not* determine  $(Y, \omega)$  uniquely. Indeed,  $\omega$  is an eigenform for real multiplication by  $\mathcal{O}_8$ , so the preceding Corollary ensures there are infinitely many isoperiodic forms  $(Y_i, \omega_i)$  in its  $\text{SL}_2(\mathbb{R})$  orbit. In other words we have:

**Corollary 9.5** *There are infinite many fake octagonal forms in  $\Omega\mathcal{M}_2$ .*

Note that the forms  $(Y_i, \omega_i)$  cannot be distinguished by their relative periods either, since they all have double zeros.

A similar statement can be formulated for the pentagonal form on the curve  $y^2 = x^5 - 1$ .

**III. Top-speed motions.** Let  $F_t : E \rightarrow \mathbb{H}$  be a holomorphic motion of  $E \subset \mathbb{H}$  over  $(\mathbb{H}, s)$ . By the Schwarz lemma, we have  $\|dF_t(z)/dt\| \leq 1$  with respect to the hyperbolic metric on  $\mathbb{H}$ . Let us say  $F_t$  is a *top-speed* holomorphic motion if equality holds everywhere; equivalently, if  $t \mapsto F_t(z)$  is an isometry of  $\mathbb{H}$  for every  $z \in E$ .

A top-speed holomorphic motion is *maximal* if it cannot be extended to a top-speed motion of a larger set  $E' \supset E$ .

**Theorem 9.6** *For any discriminant  $D \geq 4$ , the map*

$$F_t(U(s)) = U(t), \quad U \in \Lambda_D$$

*gives a maximal top-speed holomorphic motion of  $E = \Lambda_D \cdot s$  over  $(\mathbb{H}, s)$ .*

**Proof.** Let  $t \mapsto f(t) = F_t(z)$  be an extension of the motion to a point  $z \notin E$ . Then the graph of  $f$  is a leaf of  $\tilde{\mathcal{F}}_D$ , since it is disjoint from  $\tilde{X}_D(1)$ . But the only leaves that are graphs of Möbius transformations are those in  $\tilde{X}_D(1)$ , by Corollary 7.3. ■

**Corollary 9.7** *The group  $\Gamma(2) = \{A \in \mathrm{SL}_2(\mathbb{Z}) : A \equiv I \pmod{2}\}$  gives a maximal top-speed holomorphic motion of  $E = \Gamma(2) \cdot s$  over  $(\mathbb{H}, s)$ .*

**Proof.** We have  $\Gamma(2) = g\Lambda_4g^{-1}$ , where  $g = \begin{pmatrix} 1/2 & 0 \\ 0 & 2 \end{pmatrix}$  (Theorem 4.4). ■

**IV. Iterated rational maps.** Finally we explain how the foliation  $\mathcal{F}_4$  of  $X_4$  arises in complex dynamics.

First recall that the moduli space of elliptic curves can be described as the quotient orbifold  $\mathcal{M}_1 = \widetilde{\mathcal{M}}_1/S_3$ , where

$$\widetilde{\mathcal{M}}_1 = \mathbb{H}/\Gamma(2) \cong \mathbb{C} - \{0, 1\}.$$

The deck group  $S_3$  also acts diagonally on  $\widetilde{\mathcal{M}}_1 \times \widetilde{\mathcal{M}}_1$ , preserving the diagonal  $\Delta$ .

**Theorem 9.8** *For  $D = 4$ , we have  $(X_D, X_D(1)) \cong (\widetilde{\mathcal{M}}_1 \times \widetilde{\mathcal{M}}_1, \Delta)/S_3$ .*

**Proof.** Since  $\mathcal{O}_4^\vee = (1/2)\mathcal{O}_4$ , the surface  $X_4$  is isomorphic to  $(\mathbb{H} \times \mathbb{H})/\mathrm{SL}_2(\mathcal{O}_4)$ . In these coordinates we have  $\Lambda_4 = \Gamma(2)$ . Since

$$\mathrm{SL}_2(\mathcal{O}_4) \cong \{(A_1, A_2) \in \mathrm{SL}_2(\mathbb{Z}) : A_1 \equiv A_2 \pmod{2}\}$$

contains  $\Gamma(2) \times \Gamma(2)$  as a subgroup of index 6, the result follows. ■

Now consider, for each  $t \in \widetilde{\mathcal{M}}_1$ , the elliptic curve  $E_t$  defined by  $y^2 = x(x-1)(x-t)$ . There is a unique rational map  $f_t : \mathbb{P}^1 \rightarrow \mathbb{P}^1$  such that

$$x(2P) = f_t(x(P))$$

with respect to the usual group law on  $E_t$ . Indeed, using the fact that  $-2P$  lies on the tangent line to  $E_t$  at  $P$ , we find

$$f_t(z) = \frac{(z^2 - t)^2}{4z(z-1)(z-t)}.$$

Note that the *postcritical set*

$$P(f_t) = \bigcup \{f_t^n(z) : n > 0, f_t'(z) = 0\}$$

coincides with the branch locus  $\{0, 1, t, \infty\}$  of the map  $x : E_t \rightarrow \mathbb{P}^1$ .

The rational maps  $f_t(z)$  form a *stable family of Lattès examples*. It is well-known that the Julia set of any Lattès example is the whole Riemann sphere; and that in any stable family, the Julia set varies by a holomorphic motion respecting the dynamics (see e.g. [MSS], [Mc1, Ch. 4], [Mil].)

**Theorem 9.9** *As  $t$  varies in  $\widetilde{\mathcal{M}}_1$ , the holomorphic motion of  $J(f_t)$  sweeps out the lift of the foliation  $\mathcal{F}_4$  to the covering space  $\widetilde{\mathcal{M}}_1 \times \widetilde{\mathcal{M}}_1$  of  $X_4$ .*

**Proof.** Let  $\mathcal{G}$  be the foliation of  $\widetilde{\mathcal{M}}_1 \times \mathbb{P}^1$  swept out by  $J(f_t)$ . Since the holomorphic motion respects the dynamics, it preserves the post-critical set, and thus the leaves of  $\mathcal{G}$  include the loci  $z = 0, 1, \infty$  as well as the diagonal  $t = z$ . In particular,  $\mathcal{G}$  restricts to a foliation of the finite cover  $\widetilde{\mathcal{M}}_1 \times \widetilde{\mathcal{M}}_1 - \Delta$  of  $X_4 - X_4(1)$ . Since each leaf of  $\mathcal{G}$  lifts to the graph of a holomorphic function in the universal cover  $\mathbb{H} \times \mathbb{H}$ , it lies over a leaf of  $\mathcal{F}_D$  by the uniqueness part of Theorem 1.2. ■

**Algebraic curves.** The loci  $f_t^n(z) = \infty$  form a dense set of algebraic leaves of  $\mathcal{G}$  that can easily be computed inductively. The real points of these curves are graphed in Figure 1; thus the figure depicts the lift of  $\mathcal{F}_4$  to the finite cover  $\widetilde{\mathcal{M}}_1 \times \widetilde{\mathcal{M}}_1$  of  $X_4$ .

## References

- [Ah] L. Ahlfors. The complex analytic structure of the space of closed Riemann surfaces. In *Analytic Functions*, pages 45–66. Princeton Univ. Press, 1960.
- [BR] L. Bers and H. L. Royden. Holomorphic families of injections. *Acta Math.* **157**(1986), 259–286.
- [BL] C. Birkenhake and H. Lange. *Complex Abelian Varieties*. Springer-Verlag, 1992.
- [Br1] M. Brunella. Feuilletages holomorphes sur les surfaces complexes compactes. *Ann. Sci. École Norm. Sup.* **30**(1997), 569–594.
- [Br2] M. Brunella. Subharmonic variation of the leafwise Poincaré metric. *Invent. math.* **152**(2003), 119–148.
- [Dou] A. Douady. Prolongement de mouvements holomorphes (d’après Ślodkowski et autres). In *Séminaire Bourbaki, 1993/94*, pages 7–20. Astérisque, vol. 227, 1995.
- [EKK] C. J. Earle, I. Kra, and S. L. Krushkal. Holomorphic motions and Teichmüller spaces. *Trans. Amer. Math. Soc.* **343**(1994), 927–948.

- [Gd] F. Gardiner. *Teichmüller Theory and Quadratic Differentials*. Wiley Interscience, 1987.
- [vG] G. van der Geer. *Hilbert Modular Surfaces*. Springer-Verlag, 1987.
- [GKZ] B. H. Gross, W. Kohlen, and D. Zagier. Heegner points and derivatives of  $L$ -series. *Math. Ann.* **278**(1987), 497–562.
- [HH] W. F. Hammond and F. Hirzebruch.  $L$ -series, modular embeddings and signatures. *Math. Ann.* **204**(1973), 263–270.
- [Hir] F. Hirzebruch. Hilbert modular surfaces. *Enseign. Math.* **19**(1973), 183–281.
- [HZ1] F. Hirzebruch and D. Zagier. Intersection numbers of curves on Hilbert modular surfaces and modular forms of Nebentypus. *Invent. math.* **36**(1976), 57–113.
- [HZ2] F. Hirzebruch and D. Zagier. Classification of Hilbert modular surfaces. In *Complex Analysis and Algebraic Geometry*, pages 43–77. Cambridge University Press and Iwanami Shoten, 1977.
- [KZ] M. Kontsevich and A. Zorich. Connected components of the moduli spaces of Abelian differentials with prescribed singularities. *Invent. math.* **153**(2003), 631–678.
- [MR] C. Maclachlan and A. W. Reid. *The Arithmetic of Hyperbolic 3-Manifolds*. Springer-Verlag, 2003.
- [MSS] R. Mañé, P. Sad, and D. Sullivan. On the dynamics of rational maps. *Ann. Sci. Éc. Norm. Sup.* **16**(1983), 193–217.
- [Mn] W. R. Mann. Gross-Zagier revisited (Appendix). In *Heegner Points and Rankin  $L$ -Series*, volume 49 of *Math. Sci. Res. Inst. Publ.*, pages 139–162. Cambridge Univ. Press, 2004.
- [MS] H. Masur and J. Smillie. Hausdorff dimension of sets of nonergodic measured foliations. *Annals of Math.* **134**(1991), 455–543.
- [Mc1] C. McMullen. *Complex Dynamics and Renormalization*, volume 135 of *Annals of Math. Studies*. Princeton University Press, 1994.
- [Mc2] C. McMullen. The classification of conformal dynamical systems. In *Current Developments in Mathematics, 1995*, pages 323–360. International Press, 1995.

- [Mc3] C. McMullen. Polynomial invariants for fibered 3-manifolds and Teichmüller geodesics for foliations. *Ann. scient. Éc. Norm. Sup.* **33**(2000), 519–560.
- [Mc4] C. McMullen. Billiards and Teichmüller curves on Hilbert modular surfaces. *J. Amer. Math. Soc.* **16**(2003), 857–885.
- [Mc5] C. McMullen. Teichmüller curves in genus two: Discriminant and spin. *Math. Ann.* **333**(2005), 87–130.
- [Mc6] C. McMullen. Teichmüller curves in genus two: Torsion divisors and ratios of sines. *Invent. math.* **165**(2006), 651–672.
- [Mc7] C. McMullen. Dynamics of  $SL_2(\mathbf{R})$  over moduli space in genus two. *Annals of Math.*, *To appear*.
- [McS] C. McMullen and D. Sullivan. Quasiconformal homeomorphisms and dynamics III: The Teichmüller space of a holomorphic dynamical system. *Adv. Math.* **135**(1998), 351–395.
- [Mil] J. Milnor. On Lattès maps. In P. G. Hjorth and C. L. Petersen, editors, *Dynamics on the Riemann sphere*, pages 9–44. European Math. Soc., 2006.
- [Ru] B. Runge. Endomorphism rings of abelian surfaces and projective models of their moduli spaces. *Tôhoku Math. J.* **51**(1999), 283–303.
- [Sl] Z. Ślodkowski. Holomorphic motions and polynomial hulls. *Proc. Amer. Math. Soc.* **111**(1991), 347–355.
- [Sul] D. Sullivan. Quasiconformal homeomorphisms and dynamics II: Structural stability implies hyperbolicity for Kleinian groups. *Acta Math.* **155**(1985), 243–260.
- [V1] W. Veech. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Invent. math.* **97**(1989), 553–583.
- [V2] W. Veech. Moduli spaces of quadratic differentials. *J. Analyse Math.* **55**(1990), 117–171.
- [Vi] M.-F. Vigneras. *Arithmétique des algèbres de quaternions*, volume 800 of *Lectures Notes in Mathematics*. Springer-Verlag, 1980.

MATHEMATICS DEPARTMENT  
HARVARD UNIVERSITY  
1 OXFORD ST  
CAMBRIDGE, MA 02138-2901