



Corticosteroid Use Following the Onset of Invasive Aspergillosis is Associated with Increased Mortality: A Propensity Score-Matched Study

Citation

Abers, Michael, and Jatin Vyas. 2017. "Corticosteroid Use Following the Onset of Invasive Aspergillosis is Associated with Increased Mortality: A Propensity Score-Matched Study." *Open Forum Infectious Diseases* 4 (Suppl 1): S55. doi:10.1093/ofid/ofx162.125. <http://dx.doi.org/10.1093/ofid/ofx162.125>.

Published Version

doi:10.1093/ofid/ofx162.125

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:34493122>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Disclosures. F. M. Marty, Astellas Pharma US: Consultant and Grant Investigator, Consulting fee and Grant recipient; Chimerix: Consultant and Grant Investigator, Consulting fee and Grant recipient; Fate Therapeutics: Scientific Advisor, Consulting fee; Gilead Sciences: Consultant and Grant Investigator, Consulting fee and Grant recipient; LFB: Consultant, Consulting fee; Merck: Consultant, Grant Investigator and Scientific Advisor, Consulting fee and Grant recipient; Roche Molecular Systems: Consultant, Consulting fee; Shire: Consultant and Grant Investigator, Consulting fee and Grant recipient; D. P. Kontoyiannis, Pfizer: Research Contractor, Research support and Speaker honorarium; Astellas: Research Contractor, Research support and Speaker honorarium; Merck: Honorarium, Speaker honorarium; Cidara: Honorarium, Speaker honorarium; Amplyx: Honorarium, Speaker honorarium; F2G: Honorarium, Speaker honorarium

1777. Corticosteroid Use Following the Onset of Invasive Aspergillosis is Associated with Increased Mortality: A Propensity Score-Matched Study
Michael Abers, MD¹; Jatin Vyas, MD, PhD, FIDSA²; ¹Department of Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, Massachusetts; ²Medicine; Infectious Disease, Massachusetts General Hospital, Boston, Massachusetts

Session: 216. The Fungus Among-us – Clinical Advances
Saturday, October 7, 2017: 8:30 AM

Background. The safety of corticosteroid use (CSU) during active infection is controversial. In the invasive aspergillosis (IA) literature, CSU is typically defined using the time period prior to IA onset. Clinicians caring for patients with IA are unable to control prior CSU. The more clinically relevant question is whether CSU after IA onset is harmful.

Methods. Patients hospitalized at our institution from 2004 to 2014 with IA were retrospectively identified. CSU, defined as the average daily prednisone equivalent dose during the 7-day period following IA onset, was calculated for each patient. A CSU cut-off of 7.5 mg was used to assign patients to treatment (>7.5 mg) or control (<7.5 mg, including no CSU) groups. A propensity score (PS) was generated to predict group assignment. Nearest neighbor matching was performed with a caliper width of 0.2. A Cox proportional hazards model was used to assess survival 6 weeks after IA onset.

Results. PS matching generated 61 matched pairs (122 patients). Baseline characteristics did not differ significantly between groups (Table). CSU was associated with increased mortality (PS adjusted hazard ratio [HR] 2.91, 95% CI 1.32–6.40). In the CSU group, a trend towards lower mortality was noted if corticosteroid dose was tapered to 7.5 mg/day (HR 0.68, 95% CI 0.46–1.02).


Conclusion. CSU after IA onset is associated with increased mortality. In IA patients with CSU, efforts to reduce corticosteroid dose may be beneficial.

Table: Propensity matched patients at IA Onset

	CSU (n = 61)	Control (n = 61)	P
Age, years	57.6 (49.2–65.9)	53.2 (42.5–63.2)	.27
Male	59.0% (36/61)	54.1% (33/61)	.72
CSU >7.5mg prior to IA	78.7% (48/61)	70.5% (43/61)	.41
Leukemia	52.5% (32/61)	49.2% (30/61)	.86
Allogeneic bone marrow transplant	26.2% (16/61)	29.5% (18/61)	.84
Graft vs. host disease	3.3% (2/61)	11.5% (7/61)	.16
Neutropenia	48.3% (28/58)	42.9% (24/56)	.58
Solid organ transplant	11.5% (7/61)	6.6% (4/61)	.53
Obstructive lung disease	21.3% (13/61)	24.6% (15/61)	.83
Diabetes mellitus	26.2% (16/61)	29.5% (18/61)	.84
Pulmonary IA	94.8% (55/58)	94.9% (56/59)	.99
Coinfection	23.0% (14/61)	21.3% (13/61)	.99

Data presented as median (interquartile range) or % (n with feature/n with data available)

Figure. Kaplan–Meier curves comparing 6-week survival



Disclosures. All authors: No reported disclosures.

1778. Catheter-free Period Over 2 Days Is Associated with Better Outcome in Catheter-related Bloodstream Infection due to Candida
Takahiro Matsuo, MD¹; Nobuyoshi Mori, MD²; Eri Hoshino, MPA, MPhil²; Aki Sakurai, MD¹; Keiichi Furukawa, MD, FSHEA¹; ¹Infectious Diseases, St. Luke's International Hospital, Tokyo, Japan; ²Center for Clinical Epidemiology, St. Luke's International Hospital, Tokyo, Japan

Session: 216. The Fungus Among-us – Clinical Advances
Saturday, October 7, 2017: 8:30 AM

Background. Regardless of active antifungal drugs, mortality of candidemia remains high. Although it is well-known that central venous catheter (CVC) is one of the most important risk factors of candidemia and should be removed immediately, little is known about optimal timing of CVC replacement after removal. Here, we analyzed contributing risk factors associated with 30-day mortality for catheter-related bloodstream infection (CRBSI) due to candida and optimal timing of CVC replacement.

Methods. We conducted a retrospective cohort study at St. Luke's International Hospital between 2004 and 2015. We compared each clinical component in patients who died within 30 days and were alive at 30 days. Also, catheter-free period (from removal to replacement) was compared between group A and B. Fisher's exact test and Mann–Whitney U test were used in univariate analysis and multivariate linear regression was used for controlling confoundings.

Results. Among 228 patients (pts) with candidemia, 166 patients (73%) were on CVC at diagnosis. Of them, 144 patients (65%) removed CVC after the result of candidemia. Seventy-one patients (31%) replaced CVC. Fifteen patients (6%) died within 30 days (group A) and 56 patients (25%) were alive at 30 days (group B). Median age was 74 in group A and 72 in group B ($P = 0.331$) (Table 1). In univariate analysis, hematological malignancy (OR 6.75, 95% CI 1.01–44.9) and CVC replacement < 2-days after removal (OR 5.63, 95% CI 1.16–27.3) showed statistically significant increase in group A vs group B (Table 2). In multivariate analysis, CVC replacement < 2-days was independently associated with 30-day mortality (Table 3).

Conclusion. This is the first study to demonstrate the optimal timing of CVC replacement in CRBSI due to candida. CVC replacement < 2 days was an independent risk factor for 30-day mortality.

Table 1. Patient demographics

	30-day dead (N=15)	30-day alive (N=56)	Overall	p value
Gender (male), n (%)	9 (60)	34 (60.7)	43 (60.6)	1
Age, median (IQR)	74 (67–84)	72 (63–82)	74 (65–82)	0.331
Malignancy, n (%)	7 (46.7)	20 (35.7)	27 (38)	0.552
Hematological, n (%)	3 (20)	2 (3.6)	5 (7)	0.06
Solid, n (%)	4 (26.7)	18 (32.1)	22 (31)	0.763
DM, n (%)	2 (13.3)	6 (10.7)	8 (11.3)	0.673
COPD, n (%)	0 (0)	3 (5.4)	3 (4.2)	1
Stem cell transplant, n (%)	1 (6.7)	2 (3.6)	3 (4.2)	0.515
Cirrhosis, n (%)	1 (6.7)	3 (5.4)	4 (5.6)	1
Hemodialysis, n (%)	4 (26.7)	9 (16.1)	13 (18.3)	0.452
Eye check, n (%)	14 (93.3)	55 (98.2)	69 (97.2)	0.38
Corticosteroid within 90 days, n (%)	3 (20)	6 (10.7)	9 (12.7)	0.387
Site of infection				
GI, n (%)	1 (6.7)	3 (5.4)	4 (5.6)	1
CLABSI/PLABSI, n (%)	14 (93.3)	52 (92.9)	66 (93)	1
Candida species, n (%)				
C. albicans	6 (40)	34 (60.7)	40 (56.3)	0.241
C. glabrata	4 (26.7)	7 (12.5)	11 (15.5)	0.228
C. parapsilosis	2 (13.3)	6 (10.7)	8 (11.3)	0.673
C. guilliermondii	1 (6.7)	2 (3.6)	3 (4.2)	0.515
C. tropicalis	2 (13.3)	7 (12.5)	9 (12.7)	1
Endophthalmitis, n (%)	4 (26.7)	18 (32.1)	22 (31)	0.763
Chorioretinitis, n (%)	4 (26.7)	16 (28.6)	20 (28.2)	0.74
Vitritis, n (%)	0 (0)	2 (3.6)	2 (2.8)	0.74
ID consultation, n (%)	8 (53.3)	44 (78.6)	52 (73.2)	0.096
Timeframe replacement, median (IQR)	1 (0–1)	1 (0–7.75)	1 (0–7)	1
Replacement < 2 days, n (%)	13 (86.7)	30 (53.6)	43 (60.6)	0.035
Persistent bacteremia, n (%)	2 (13.3)	9 (16.1)	11 (15.5)	1

Table 2. Univariate analysis for 30-day infection related mortality

	OR	p value	95% CI
Gender (male)	0.971	0.96	0.303
Age	0.98	0.394	0.935
Malignancy	1.575	0.44	0.498
Hematological	6.75	0.048	1.014
Solid	0.768	0.684	0.215
DM	1.282	0.776	0.231
Stem cell transplant	1.929	0.602	0.163
Cirrhosis	1.262	0.845	0.122
Hemodialysis	1.899	0.351	0.493
Eye check	0.255	0.344	0.015
Corticosteroid within 90 days	2.083	0.345	0.455
Site of infection			
GI	1.262	0.845	0.122
CLABSI/PLABSI	1.077	0.949	0.111
Candida species			
C. albicans	0.432	0.157	0.135
C. glabrata	2.545	0.188	0.633
C. parapsilosis	1.282	0.776	0.231
C. guilliermondii	1.929	0.602	0.163
C. tropicalis	1.077	0.931	0.199
Endophthalmitis	0.768	0.684	0.215
Chorioretinitis	1.158	0.823	0.32
Vitritis	N/A	N/A	N/A
ID consultation	0.312	0.057	0.094
Timeframe replacement	0.939	0.242	0.846
Replacement < 2 days	5.633	0.032	1.162
Persistent candidemia	0.803	0.795	0.154