



RNA Polymerase II Transcription Attenuation at the Yeast DNA Repair Gene, DEF1, Involves Sen1-Dependent and Polyadenylation Site-Dependent Termination

Citation

Whalen, Courtney, Christine Tuohy, Thomas Tallo, James W. Kaufman, Claire Moore, and Jason N. Kuehner. 2018. "RNA Polymerase II Transcription Attenuation at the Yeast DNA Repair Gene, DEF1, Involves Sen1-Dependent and Polyadenylation Site-Dependent Termination." *G3: Genes|Genomes|Genetics* 8 (6): 2043-2058. doi:10.1534/g3.118.200072. <http://dx.doi.org/10.1534/g3.118.200072>.

Published Version

doi:10.1534/g3.118.200072

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:37298306>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

RNA Polymerase II Transcription Attenuation at the Yeast DNA Repair Gene, *DEF1*, Involves Sen1-Dependent and Polyadenylation Site-Dependent Termination

Courtney Whalen,^{*} Christine Tuohy,[†] Thomas Tallo,[‡] James W. Kaufman,[§] Claire Moore,^{**} and Jason N. Kuehner^{§,1}

^{*}Department of Nutritional Sciences, College of Health and Human Development, The Pennsylvania State University, University Park, PA 16802, [†]Department of Microbiology and Physiological Systems, University of Massachusetts Medical School, Worcester, MA 01605, [‡]Division of Infectious Diseases, Boston Children's Hospital, Harvard Medical School, Boston, MA 02115, [§]Department of Biology, Emmanuel College, Boston, MA 02115, and ^{**}Department of Developmental, Molecular, and Chemical Biology, Tufts University School of Medicine, Boston, MA 02111
ORCID ID: 0000-0001-9432-8506 (J.N.K.)

ABSTRACT Termination of RNA Polymerase II (Pol II) activity serves a vital cellular role by separating ubiquitous transcription units and influencing RNA fate and function. In the yeast *Saccharomyces cerevisiae*, Pol II termination is carried out by cleavage and polyadenylation factor (CPF-CF) and Nrd1-Nab3-Sen1 (NNS) complexes, which operate primarily at mRNA and non-coding RNA genes, respectively. Premature Pol II termination (attenuation) contributes to gene regulation, but there is limited knowledge of its prevalence and biological significance. In particular, it is unclear how much crosstalk occurs between CPF-CF and NNS complexes and how Pol II attenuation is modulated during stress adaptation. In this study, we have identified an attenuator in the *DEF1* DNA repair gene, which includes a portion of the 5'-untranslated region (UTR) and upstream open reading frame (ORF). Using a plasmid-based reporter gene system, we conducted a genetic screen of 14 termination mutants and their ability to confer Pol II read-through defects. The *DEF1* attenuator behaved as a hybrid terminator, relying heavily on CPF-CF and Sen1 but without Nrd1 and Nab3 involvement. Our genetic selection identified 22 *cis*-acting point mutations that clustered into four regions, including a polyadenylation site efficiency element that genetically interacts with its cognate binding-protein Hrp1. Outside of the reporter gene context, a *DEF1* attenuator mutant increased mRNA and protein expression, exacerbating the toxicity of a constitutively active Def1 protein. Overall, our data support a biologically significant role for transcription attenuation in regulating *DEF1* expression, which can be modulated during the DNA damage response.

KEYWORDS

RNA polymerase II termination
attenuation
Def1
cleavage and polyadenylation complex (CPF-CF)
Nrd1/Nab3/Sen1 complex (NNS)

RNA Polymerase II (Pol II) transcribes a wide assortment of transcripts in eukaryotes, including all protein-coding mRNAs and most non-coding RNAs. The high density of Pol II across the genome must be

confined to prevent interference with surrounding protein:DNA transactions (Jensen *et al.* 2013). Genomic partitioning is achieved in part by transcription termination, which releases Pol II and its RNA product from the DNA template. The biological significance of termination is not limited to defining transcriptional boundaries. Due to its connection with 3'-end processing, stability, and export, Pol II termination has the ability to influence RNA fate and function (Kuehner *et al.* 2011; Mischo and Proudfoot 2013; Arndt and Reines 2015; Porrua and Libri 2015). In addition, there is an expanding collection of genes regulated by premature transcription termination (*i.e.*, attenuation), some of which exhibit altered expression during cell metabolism and stress. The significance of Pol II termination has also been revealed in human pathology, including its roles in HIV latency, herpes viral infection, and renal cell carcinoma (Natarajan *et al.* 2013; Rutkowski *et al.* 2015; Grosso *et al.* 2015; Loya and Reines 2016).

Copyright © 2018 Whalen *et al.*

doi: <https://doi.org/10.1534/g3.118.200072>

Manuscript received January 25, 2018; accepted for publication April 16, 2018; published Early Online April 23, 2018.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental material available at Figshare: <https://doi.org/10.25387/g3.6167696>.

¹Corresponding author: Department of Biology, Emmanuel College, 400 The Fenway, Boston, MA 02115, E-mail: kuehnerj@emmanuel.edu

Pol II termination is best understood in the model eukaryote *S. cerevisiae*, which is especially dependent on the process due to its compact genome (Goffeau *et al.* 1996). In yeast, Pol II termination occurs via two major pathways: cleavage and polyadenylation factor (CPF-CF) termination and Nrd1-Nab3-Sen1 (NNS) termination (Kuehner *et al.* 2011; Mischo and Proudfoot 2013; Arndt and Reines 2015; Porrua and Libri 2015). Pol II termination of most mRNAs occurs via CPF-CF termination, during which recognition of a polyadenylation (pA) site contributes to 3'-end processing and Pol II eviction, perhaps via allosteric changes in the elongation complex. Pol II termination of most noncoding RNAs occurs via NNS termination. Noncoding RNAs, including some small nuclear RNAs (snRNAs) and small nucleolar RNAs (snoRNAs), can be processed by endonucleolytic cleavage and exonucleolytic trimming, resulting in stable and abundant products (Bernstein and Toth 2012). Noncoding RNAs can also behave as cryptic unstable transcripts (CUTs) due to NNS-mediated oligoadenylation via the TRAMP complex and exosome-mediated degradation (Schmidt and Butler 2013; Tudek *et al.* 2014). The selection of CPF-CF vs. NNS termination depends on RNA sequence elements and protein factors associated with the Pol II Rpb1 C-terminal domain (CTD), which is modified in accordance with Pol II distance from a transcription start site (TSS) (Rondon *et al.* 2008). There is evidence to support direct roles for a Pol II CTD-binding protein (Pcf11), an RNA exonuclease (Rat1), and an RNA/DNA helicase (Sen1) in the actual Pol II release step, but the termination mechanism remains unresolved (Kuehner *et al.* 2011; Mischo and Proudfoot 2013; Arndt and Reines 2015; Porrua and Libri 2015).

While CPF-CF and NNS termination pathways exhibit distinct behavior, they display substantial overlap among factor requirements, and they may have evolved independently to recognize highly similar sequence elements (Porrua *et al.* 2012). NNS termination typically occurs < 1 kb from a TSS due to Nrd1-mediated recognition of Ser5-phosphorylated CTD, which is predominant when Pol II is proximal to the promoter (Rondon *et al.* 2008). CPF-CF termination typically occurs > 1 kb from a TSS due to Pcf11-mediated recognition of Ser2-phosphorylated CTD, which is predominant when Pol II is distal to the promoter. However, exceptions have been identified that circumvent these trends. NNS terminators can be recognized outside of their normal context and distance threshold, in some cases acting as a fail-safe mechanism to prevent transcription interference (Steinmetz *et al.* 2006a; Gudipati *et al.* 2008; Ghazal *et al.* 2009; Rondón *et al.* 2009; Porrua *et al.* 2012). Likewise, NNS terminators can be dependent on CPF-CF components (Fatica *et al.* 2000; Morlando *et al.* 2002; Dheur *et al.* 2003; Kim *et al.* 2006; Pearson and Moore 2014). The basis of Pol II flexibility in terminator recognition remains unclear, as does the extent of crosstalk and function between CPF-CF and NNS components.

In addition to Pol II termination occurring downstream of genes, premature Pol II termination (attenuation) can regulate mRNA gene expression. Attenuation has long been recognized as a widespread mechanism of bacterial gene regulation (Naville and Gautheret 2010), but the extent of its activity and biological significance has been studied more recently in eukaryotes (Colin *et al.* 2011; Contreras *et al.* 2013). Genome-wide analysis of CUTs and NNS factors suggest that 5–10% of yeast mRNA genes may be regulated by attenuation (Neil *et al.* 2009; Webb *et al.* 2014). The *NRD1* gene was identified as an early target of attenuation in yeast, whereby Nrd1 autoregulates its mRNA expression as part of the NNS termination complex (Arigo *et al.* 2006). Similar autoregulatory schemes appear to operate for RNA processing factor genes *HRP1* and *PCF11* (Kuehner and Brow 2008; Creamer *et al.* 2011). Attenuator recognition and bypass has been linked to changes in

cell metabolism and stress response genes, but in most cases the signaling mechanism is unknown. Examples of mRNA gene attenuation targets include *IMD2* and *URA2* (nucleotide biosynthesis), *FKS2* (cell wall damage), *CLN3* (glucose starvation), *GPH1* (glycogen metabolism), and *GLT1* (nitrogen metabolism) (Jenks *et al.* 2008; Kuehner and Brow 2008; Thiebaut *et al.* 2008; Kwapisz *et al.* 2008; Kim and Levin 2011; Darby *et al.* 2012; Chen *et al.* 2017; Merran and Corden 2017).

In a genome-wide study of yeast polyadenylation site usage, our labs identified *DEF1* as a gene that could be regulated by attenuation (Graber *et al.* 2013). *DEF1* encodes a protein needed for degradation of Pol II stalled at UV-induced lesions, thus providing nucleotide excision repair factors with access to DNA (Woudstra *et al.* 2002). *DEF1* function is regulated at the level of post-translational control, during which UV damage induces Def1 protein processing and nuclear accumulation (Wilson *et al.* 2013; David 2013). In this study, we confirmed that the *DEF1* promoter-proximal pA site is sufficient to behave as a *bona fide* attenuator, supporting an unexpected mechanism for *DEF1* regulation. Using a plasmid-based reporter system, we identified several *cis*- and *trans*-acting requirements, including a putative Hrp1-binding site and factors in both CPF-CF and NNS terminator pathways. Despite its promoter-proximal location, the *DEF1* attenuator behaved most similarly to a hybrid CPF-CF-Sen1 terminator rather than a traditional NNS terminator. To demonstrate biological significance, we have shown that an attenuator mutation results in *DEF1* overexpression, enhancing the toxicity of a constitutively active Def1 protein. To our knowledge, this is the first example of attenuator regulation being linked to a DNA damage response gene.

MATERIALS AND METHODS

Construction of pGAC24-CUP1 and pGAC24-lacZ reporter gene plasmids

PCR using Q5 Hot Start High-Fidelity 2X Master Mix (NEB) was performed to amplify the *DEF1* attenuator region (-187 to +93, relative to the +1 ATG) from yeast strain BY4742 genomic DNA and cloned into the *XhoI* restriction site of the pGAC24-No Term-CUP1 (no terminator) vector (Lesser and Guthrie 1993), which had been treated with Antarctic phosphatase (NEB) to prevent religation of linearized plasmid. The ligation was performed overnight at 16° using T4 DNA ligase (NEB) prior to transformation into 5-alpha Competent *E. coli* cells (NEB) to create pGAC24-*DEF1*-CUP1.

PCR using Q5 Hot Start High-Fidelity 2X Master Mix (NEB) was performed to amplify relevant fragments from plasmids pGAC24-*DEF1*-CUP1 (GAP promoter, *ACT1* intron with *DEF1* attenuator, *ACT1* exons, *AMP^R* gene, *LEU2* gene) and p903 (*lacZ* ORF) (Guarente and Mason 1983). The two PCR products were assembled to create the pGAC24-*DEF1*-*lacZ* vector using a Gibson Assembly Cloning Kit (NEB). The pGAC24-No Term-*lacZ* plasmid (no terminator) was generated by *XhoI* restriction digestion of pGAC24-*DEF1*-*lacZ*, separation of backbone from insert using a ZymoClean Gel DNA recovery kit (Zymo Research), and religation of backbone lacking insert. The *CYC1* terminator (+448 to +528 relative to +1 ATG start codon) and *SNR13* terminator (+125 to +232 relative to +1 TSS) were amplified by PCR and cloned into the *XhoI* site of pGAC24-no Term-*lacZ*. A pGAC24-*DEF1*-*lacZ* (*HIS3*) reporter was generated using Gibson cloning to swap in the *HIS3* selectable marker in place of *LEU2*, and the no Term, *CYC1*, and *SNR13* reporters were generated by PCR cloning and *XhoI* digestion as described above. Transformation of reporter plasmids into yeast strains was performed using a standard lithium acetate method.

Construction of pRS426-DEF1 plasmid and A-1G and C1590A mutants

PCR using Q5 Hot Start High-Fidelity 2X Master Mix (NEB) was performed to amplify the *DEF1* promoter, 5'-UTR, ORF, and 3'-UTR (-428 to +2596 relative to the +1 ATG) from yeast strain BY4742 genomic DNA and cloned into the *NotI* restriction site of the pRS426 (*URA3*) vector. The A-1G and C1590A point mutations were generated using a Quik-Change Lightning Site-Directed Mutagenesis Kit (Agilent).

Construction of pRS314-HRP1 plasmid and the hrp1-5 mutant

PCR using Q5 Hot Start High-Fidelity 2X Master Mix (NEB) was performed to amplify the *HRP1* promoter, 5'-UTR, ORF, and 3'-UTR (-500 to +1848 relative to the +1 ATG) from yeast strain BY4742 genomic DNA and cloned into the *NotI/XhoI* restriction site of the pRS314 (*TRP1*) vector. The L205S point mutation (*hrp1-5*) was generated using a Quik-Change Lightning Site-Directed Mutagenesis Kit (Agilent).

Spot Test Assays

Yeast bearing the reporter plasmids of interest were grown overnight in a shaking incubator (at 25° or 30°) in appropriate selective liquid media. The absorbance at OD₆₀₀ was measured for each strain using a spectrophotometer (Implen Diluphotometer) and cultures were diluted to OD₆₀₀ = 1.0. Each diluted culture (200 μL) was transferred to a 96-well plate, followed by 10-fold serial dilutions (x4) into adjacent wells. All samples were mixed and spotted using a 96-well replica plater (Sigma) on appropriate plate media. The plates were air-dried and incubated for 3-7 days.

Yeast Doubling-Time Growth Assays

The BY4742 *def1Δ* strain bearing the pRS426-*DEF1* plasmids of interest were grown overnight in a shaking 30° incubator in -Ura media. The absorbance at OD₆₀₀ was measured for each strain using a spectrophotometer (Implen Diluphotometer) and cultures were diluted to OD₆₀₀ = 0.15. Each diluted culture was grown for 4 hr at 30° (recovery period) and then split into separate cultures, with half of the culture shifted to 39° and the other half remaining at 30° for 2.5 hr. The absorbance at OD₆₀₀ was measured every half hour, and doubling times were calculated using the following formula: Doubling time = (Time Duration x log(2)) / (log (Final Concentration - log (Initial Concentration))).

Yeast β-galactosidase Microplate Plate Assay

Yeast bearing the reporter plasmids of interest were grown overnight in a shaking incubator (30° for most strains or 25° for *nab3-11* and *hrp1-5*) in appropriate selective liquid media. The cultures were diluted 10-fold and absorbance at OD₆₀₀ was measured for each strain using a plate reader (SpectraMax 190). Cultures were diluted to OD₆₀₀ = 0.15, grown at 30° for 2 hr (recovery period), and shifted to 37° for 2 hr (non-permissive temperature). Each strain (200 μL) was aliquoted into a 96-well plate and media containing no cells served as an instrument blank. Following measurement of absorbance at OD₆₀₀, cell lysis was performed and reporter enzyme activity was measured using the Yeast β-Galactosidase Assay Kit according to manufacturer's instructions (Thermo Scientific). The assay kit working solution was prepared and added separately to the 96-well plate in 100 μL aliquots. The absorbance at OD₄₂₀ was measured every minute for 60 min. to collect a kinetic reaction rate and slope. Slopes were gathered using a kinetic reaction window in a linear range with

strong R² value. Relative β-galactosidase activity was calculated using the equation: ((OD₄₂₀ Slope) ÷ (0.1 mL) x OD₆₀₀) = β-galactosidase activity (Thibodeau *et al.* 2004).

Mutagenesis of the Attenuator

Mutagenesis of the *DEF1* attenuator was carried out by error-prone PCR and *in-vivo* homologous recombination (Muhlrad *et al.* 1992). Primers complementary to pGAC24-*DEF1-CUP1* sequences were used to direct amplification by Taq DNA polymerase (NEB) of a 635 bp fragment containing the *DEF1* 5'-UTR and upstream ORF. PCR amplification was performed for 40 rounds, purified, and repeated for an additional 40 rounds, relying upon the normal error rate of Taq DNA polymerase for the introduction of random mutations. pGAC24-*DEF1-CUP1* was digested with *Bam*HI and *Xho*I, creating a gap of 447 bp flanked by 110 and 78 nt of complementarity to the PCR product, and cotransformed with the PCR product into the 46α yeast strain. After selection for transformants on -Leu plates, colonies were replica plated onto -Leu plates containing 0.4 mM CuSO₄. Plasmids were recovered from copper-resistant colonies and retransformed into yeast to confirm that the copper-resistant phenotype was linked to the plasmid prior to sequencing.

Reverse transcriptase (RT)-PCR

Total RNA was purified from yeast cells by measuring the absorbance at OD₆₀₀ of the desired culture and harvesting 1.5 ODs of cells. The Master Pure Yeast RNA Purification Kit (Epicentre) was used to purify total RNA (including 1 hr DNase I treatment), and RNA quality and yield was determined using a NanoDrop spectrophotometer (Thermo-Fisher). To convert the purified RNA to cDNA, RT-PCR was performed using a OneTaq RT-PCR kit (NEB). Pairwise reactions were set up including 1 μg of total RNA and random primer (mix of hexamer and d(T)₂₃VN primers). A negative control received no reverse transcriptase (-RT) to ensure that the final signal was RNA-dependent and not derived from chromosomal DNA template. Amplification of *CUP1*, *lacZ*, *DEF1*, and 18S RNA was performed using OneTaq DNA polymerase in a 25 μL reaction with 2 μL of diluted cDNA template, as directed by the manufacturer. The amplification cycle number ranged from 12-25 cycles depending on the linearity and intensity of the PCR product signal. The PCR products (10 μL) were loaded into a 2% agarose gel stained with SYBR Safe (Invitrogen). The gel band intensity and ratios of total, read-through, and attenuated transcripts were measured using the Gel Doc EZ System (BioRad) and ImageStudio software (LICOR).

Western blot analysis of Def1 protein levels

The BY4741 *def1Δ* strains containing pRS426-*DEF1* or the *def1* mutants were grown overnight at 30°, diluted back to OD₆₀₀ = 0.4, grown for 4 hr in a 30° shaking incubator (recovery), and shifted to a 39° shaking incubator (non-permissive temperature) for 0, 1, or 2 hr. The cells were harvested (10 OD units), washed in 20% TCA, and stored at -20°. Whole cell protein extracts were prepared using a TCA method (Keogh *et al.* 2006). Briefly, cell pellets were resuspended in 250 μL of 20% TCA and lysed using glass beads and a vortexer in a 4° room (3 × 1 min) with 1 min pauses on ice between runs. The supernatant was transferred to a new tube using a gel-loading tip to avoid the beads. 700 μL of 5% TCA was added to the supernatant (1.25 mL final) and inverted to mix. The sample was microcentrifuged at 13,300 RPM for 10 min at 4°. The supernatant was discarded and the pellet was washed with 750 μL of 100% ice-cold ethanol. The wash buffer was discarded, and the pellet was resuspended in 40 μL of 1 M Tris Cl, pH 8.0. An

additional 80 μ L of 2X SDS reducing sample buffer (66 mM Tris-HCl pH 6.8, 26% (w/v) glycerol, 2.1% SDS, 0.01% bromophenol blue) was added, and the sample was heated to 95° for 5 min, microcentrifuged at top speed for 5 min., and the supernatant was used for Western blot analysis. Western blots were incubated with rabbit polyclonal anti-Def1 antibody (1:5,000; kind gift of Svejstrup lab) and mouse anti- β actin antibody (1:5,000, Abcam) prior to incubation with an anti-rabbit or anti-mouse HRP-conjugated secondary antibody (1:15,000; Jackson ImmunoResearch). Target proteins were detected via chemiluminescence using Clarity ECL Western blotting substrate (BioRad) and a C-Digit blot scanner with Image Studio software (LICOR).

Data Availability

Strains, plasmids, and primer sequences are available upon request. The authors affirm that all data necessary for confirming the conclusions of the article are present within the article, figures, and tables. Supplemental material available at Figshare: <https://doi.org/10.25387/g3.6167696>.

RESULTS

DEF1 promoter-proximal region resembles a transcriptional attenuator

Our labs have previously identified *DEF1* as a gene with both promoter-proximal (pA^1) and promoter-distal (pA^2) polyadenylation sites (Graber *et al.* 2013). This arrangement of signals for *DEF1* 3'-end processing results in multiple mRNA isoforms due to alternative pA site usage. In the absence of stress \sim 70% of *DEF1* transcripts terminate near pA^1 (< 50 nt downstream of the start codon), resulting in a mixture of short RNAs (\sim 150 nt) that are not capable of being translated into Def1 protein. Upon exposure to the DNA damaging agent 4-NQO, pA^1 usage is reduced to \sim 30%, resulting in \sim 70% of polyadenylation occurring at pA^2 , which is \sim 2.5 kb downstream of the TSS (Figure 1A). The switch in pA usage likely contributes to the \sim twofold increase in full-length *DEF1* mRNA produced upon exposure to 4-NQO. Aside from the observed changes in pA /terminator usage, *DEF1* bears additional hallmarks of attenuator regulation. The promoter-proximal *DEF1* transcripts consist of both a polyadenylated fraction and a CUT fraction (Neil *et al.* 2009; Oszolak *et al.* 2010; Graber *et al.* 2013), which is consistent with the CPF-CF and/or NNS termination pathway operating in this region. In addition, the *DEF1* 5'-UTR is highly conserved among yeast species (Figure 1B), suggesting that the RNA takes on an important regulatory function, as observed for *IMD2* and *HRP1* attenuators (Kuehner and Brow 2008).

DEF1 promoter-proximal pA site is sufficient to confer Pol II attenuation in a reporter plasmid

To better understand the signals and factors involved in recognition and read-through of the *DEF1* promoter-proximal pA site, we fused the *DEF1* promoter, 5'-UTR, and pA^1 region (*pDEF1-pA¹*) from -253 to +93 (relative to +1 start codon) to the *CUP1* reporter gene, which confers resistance to copper-containing media in a *cup1 Δ* strain background (Kuehner and Brow 2008). The *pDEF1-pA¹-CUP1* strain was copper sensitive but failed to exhibit copper resistance in any of the termination factor mutants tested (data not shown). The unexpected behavior of *pDEF1-pA¹-CUP1* was likely due to misfolding of the Def1-Cup1 fusion protein upon addition of *DEF1* encoded amino acids (+1 to +93). A fusion of *CUP1* to the *DEF1* promoter and 5'-UTR in the absence of the upstream *DEF1* ORF (-253 to -1) resulted in only modest sensitivity on 0.8 mM copper (data not shown).

To test the activity of the *DEF1* pA^1 site in a context where the ORF region is not translated into protein, we cloned the *DEF1* attenuator

into the intron of the pGAC24-*CUP1* reporter gene plasmid (Figure 2A) (Lesser and Guthrie 1993; Steinmetz and Brow 1996). The *DEF1* attenuator included the *DEF1* 5'-UTR and the pA^1 within the upstream ORF (-187 to +93) but not the consensus TATA box promoter element. In the absence of a transcriptional terminator insert (No Term.), the *cup1 Δ* strain was copper-resistant as expected (Figure 2B). Insertion of the *DEF1* pA^1 into the pGAC24 reporter was sufficient to confer copper sensitivity to 0.3 mM copper (Figure 2B). The *DEF1* pA^1 insert resulted in predominant production (71%) of short RT-PCR products, consistent with transcription attenuation (Figure 2C). Mutations in the NNS protein Sen1 and the CPF-CF proteins Ssu72 and Hrp1 conferred *DEF1-CUP1* resistance to 0.3 mM copper, while a mutation in the NNS Nrd1 protein had no effect (Figure 2B). For comparison, we analyzed transcription termination activity from the mRNA gene *CYC1*, which contains a hybrid CPF-CF-NNS-dependent terminator (+448 to +528 relative to +1 ATG start codon), and the snRNA gene *SNR13*, which contains an NNS-dependent terminator (+125 to +232 relative to +1 TSS). The *CYC1* gene contains a traditional mRNA 3'-end processing site but is somewhat NNS-dependent, at least in part because it is a short gene containing a pA site located < 1 kb from the promoter (Steinmetz *et al.* 2006a). The *sen1*, *ssu72*, and *hrp1* mutants conferred resistance to 0.3 mM copper for *CYC1-CUP1*, and the *sen1*, *nrd1*, and *ssu72* mutants conferred resistance to 0.2 mM copper for *SNR13-CUP1* (Figure 2B). These data for the control terminators were mostly consistent with previous reports (Steinmetz *et al.* 2001; Steinmetz and Brow 2003; Steinmetz *et al.* 2006a; Chen *et al.* 2017).

To extend the versatility of the reporter gene and improve quantitation of read-through defects, we replaced the *CUP1* gene with the *lacZ* gene, which allows β -galactosidase activity to be used as a readout of transcriptional activity. In addition, the *lacZ* reporter in the pGAC24 plasmid is under control of a constitutive promoter instead of a previously described *GAL*-inducible *lacZ* reporter (Hyman *et al.* 1991). The ability to maintain the pGAC24-*DEF1-lacZ* reporter in selective glucose media avoids complications arising from the galactose-sensitivity of some termination mutants (data not shown). In the absence of a terminator (No Term.), the reporter produced \sim 4,500 units of β -gal activity (Figure 2D). The *DEF1-lacZ* reporter produced \sim 800 β -gal units, which is a $>$ fivefold reduction compared to the no terminator control. The *CYC1-lacZ* reporter produced \sim 1500 β -gal units (threefold reduction), and the *SNR13-lacZ* reporter exhibited \sim 40 units of β -gal activity (120-fold reduction). Overall the *DEF1-lacZ* reporter exhibited an intermediate level of termination activity between the stronger *SNR13* terminator and the weaker *CYC1* terminator (Figure 2D).

Mutations in both NNS and CPF-CF pathways confer Pol II read-through of DEF1 attenuator

To investigate the *trans*-acting requirements of *DEF1* attenuation, we measured β -gal reporter activity in response to a variety of Pol II termination mutants. The mutations targeted members of CFI (Pcf11, Rna14, Rna15, Hrp1), CPF (Cft2, Glc7, Ssu72), and NNS (Nrd1, Nab3, Sen1). Our genetic screen also included Ctk1 and Paf1, which promote Pol II modification and recruitment of termination factors (Bowman and Kelly 2014; Van Oss *et al.* 2017). We first classified the effect of mutants on the control terminators to determine their relative impact on NNS vs. CPF-CF pathways in our reporter system. Pol II terminator read-through activity of *SNR13-lacZ*, but not *CYC1-lacZ*, was increased ($>$ twofold) by the *paf1 Δ* , *nrd1-5*, *nab3-11*, *pcf11-2*, *pcf11-9*, and *pcf11-13* mutants, indicating defects in NNS-dependent termination (Figure 3A-F). In contrast, the reporter activity of *CYC1-lacZ*, but not *SNR13-lacZ*, was increased ($>$ twofold)




Figure 1 *DEF1* promoter-proximal region resembles a transcriptional attenuator. (A) Direct RNA sequence tags were mapped to the *DEF1* region of the *S. cerevisiae* genome (Integrative Genomics Viewer) in strains grown +/- the UV-mimetic drug 4-NQO (Graber et al., 2013). In this image, the *DEF1* gene is transcribed to the left (TSS indicated by arrow), with vertical bars indicating the number of pA site reads corresponding to each position. Positions of major TSS and pA sites are numbered relative to the *DEF1* +1 start codon (ATG). Tick marks at the top of the IGV image correspond to 500 bp segments. (B) Alignment of *DEF1* promoter / 5'-UTR in five species of the genus *Saccharomyces*. Sequences were obtained using *Saccharomyces* Genome Database (SGD) Fungal alignment. All non-template strand sequences between -253 and -1 relative to the *DEF1* start codon are shown, with invariant nucleotides indicated with asterisks. The putative TATA box (yellow) and mapped TSS (green) (Zhang et al. 2005; Nagalakshmi et al. 2008) are also indicated.

by the *hrp1-1* and *ctf2-5001* mutants, indicating defects in CPF-CF termination (Figure 3G, H). Interestingly, the reporter activity of *DEF1-lacZ* increased (>twofold) in the *paf1Δ*, *hrp1-1*, and *ctf2-1* mutants (Figure 3A, G, H), indicating a dependence on both NNS- and CPF-CF termination pathways for attenuator recognition. We identified an additional class of mutants (*rna14-5*, *rna15-58*, *glc7-12*, *ssu72-2*, *sen1-1*, *ctk1Δ*) that conferred read-through defects for both *SNR13-lacZ* and *CYC1-lacZ* control reporters (Figure 4A-F). All of these mutants likewise increased *DEF1-lacZ* activity, confirming that the *DEF1* attenuator behaves as a hybrid Sen1- and CPF-CF-dependent terminator.

To confirm that the β -gal activity of the reporter genes was due to increased Pol II read-through of the transcriptional terminator, we analyzed RNA from the *DEF1-lacZ* reporter in *sen1-1* and *rna15-58* strains using RT-PCR to detect attenuated and read-through mRNA (Figure 4G). In wild-type strains, the preponderance of RNA (96%) corresponded to attenuated RNA. The higher efficiency of attenuator activity in *DEF1-lacZ* compared to *DEF1-CUP1* (compare Figure 4G to

Figure 2C) may reflect differences in Pol II elongation through the reporter genes. The *sen1-1* and *rna15-58* mutants increased the level of read-through mRNA ~ 3 fold, which is consistent with the trend observed in the β -gal assay. These data validate the use of β -gal activity from the *DEF1-lacZ* reporter as a proxy for Pol II attenuator recognition and read-through at the RNA level.

To more quantitatively compare the *DEF1* terminator to control terminators, we created a read-through index to rank the level of β -gal activity in mutant vs. wild-type strains. Based on the range of the data, we assigned the mt/WT ratio into 3 categories: little/no effect, ≤ 2 -fold; intermediate effect, 2-10 fold; and strong effect, >10-fold (Figure 5). Ranked in order, the mutants most defective for *DEF1* attenuator recognition were *rna14-5* and *rna15-58*, and those with an intermediate effect were *glc7-12*, *ssu72-2*, *hrp1-1*, *sen1-1*, *ctk1Δ*, *paf1Δ* and *ctf2-5001*. The *DEF1* attenuator behaved more similarly to the *CYC1* terminator than the *SNR13* terminator, with *DEF1* matching *CYC1* in 10/12 cases that allowed for direct comparison within the index. These results indicate




Figure 2 *DEF1* promoter-proximal pA site (pA¹) is sufficient to confer Pol II transcription attenuation in a *CUP1/lacZ* reporter assay. (A) Schematic of *CUP1/lacZ* reporter gene used to measure transcription termination (not to scale). The pGAC24 plasmid contains the actin exon(E1)-intron(exon(E2) fused to a *CUP1* or *lacZ* reporter gene. The promoter-proximal *DEF1* pA site, *CYC1* pA site, or *SNR13* transcription termination site were inserted within the intron. The *DEF1* attenuator includes the 5'-UTR and upstream ORF but not the consensus TATA box promoter element. In the absence of a pA/terminator insert (No Term.), full-length mRNA production confers copper-resistance and high β -galactosidase activity. In the presence of a pA/terminator insert, attenuated non-coding RNA (ncRNA) production confers copper-sensitivity and low β -galactosidase activity. *Trans*-acting mutants that prevent pA/terminator recognition promote copper-resistance and higher β -galactosidase activity. (B) *DEF1* pA¹ site confers copper-sensitivity in a *DEF1-CUP1* reporter, and *sen1*, *ssu72*, and *hrp1* mutants confer copper-resistance. (C) *DEF1* pA¹ site reduces expression of *CUP1* mRNA due to accumulation of attenuated ncRNA. Note that based on the RT-PCR primer locations (F1, R1, and R2 in panel A), the RT-PCR product from spliced mRNA (231 bp) is shorter than the PCR product from attenuated, unspliced transcript (387 bp). The % attenuated vs. full-length was determined by adding up signal intensities for both bands and determining the relative ratio. No Term. = No Terminator. (D) *DEF1* pA¹ site reduces expression of a *lacZ* reporter similarly to known transcription terminators from *CYC1* and *SNR13*. β -galactosidase activity was measured following cell lysis and incubation with ONPG substrate, using absorption at OD₆₀₀ for cell density and OD₄₂₀ for product production. Experiments were performed in biological triplicate, and errors bars show standard deviation.

that the *DEF1* attenuator exhibits hybrid characteristics but is more heavily influenced by the CPF-CF termination pathway.

DEF1 attenuator consists of multiple cis-acting elements spanning the ORF start codon

In order to define the *cis*-acting sequence elements that promote *DEF1* attenuator recognition, we randomly mutagenized the *DEF1-CUP1*

reporter and utilized a genetic selection to identify copper resistant colonies. We identified 22 attenuator point mutations within a 78 bp region (-31 to +47 relative to *DEF1* +1 ATG start codon) (Figure 6A). The most frequently identified mutants were A-1G (n = 8), T+6C (n = 5), A+43G (n = 3), A+40G (n = 2), and T-20C (n = 2). The remaining mutants were each identified once in the genetic selection. To compare the relative level of copper resistance, we conducted growth assays for




Figure 3 Trans-acting mutants *paf1*, *hrp1*, and *cft2* result in Pol II read-through (>twofold) of the *DEF1* attenuator, representing a contribution of both NNS- and CPF-CF-dependent termination activity. (A)-(H) The *DEF1*-lacZ reporter and the control reporters *CYC1*-lacZ (CPF-CF-NNS hybrid terminator) and *SNR13*-lacZ (NNS terminator) were transformed into the indicated strains above. Cultures were grown at permissive temperature (25°C or 30°C) and diluted prior to shifting to 30°C for 2 hr and 37°C (non-permissive) for 2 hr. β -galactosidase activity was measured following cell lysis and incubation with ONPG substrate, using absorption at OD₆₀₀ for cell density and OD₄₂₀ for β -gal production. Experiments were performed in biological triplicate, and errors bars show standard deviation.

six of the mutants and analyzed the ratio of attenuated:read-through transcripts via RT-PCR. The spot test and RT-PCR assays exhibited strong agreement with respect to attenuator activity. The WT *DEF1* attenuator resulted in no growth (-) on 0.6 mM copper and 78% usage of the attenuator (Figure 6B, C). The A-1G and T+6 mutants were most defective for attenuator recognition, resulting in strong copper-resistant growth (+++) and decreased attenuator usage (6–15%). The A+43G, A+16G, and A-5G mutants were of intermediate strength, and the T-31C mutant had the weakest effect of the mutants tested.

We arranged the mutants into four regions (I-IV) based on their clustering pattern and similarity to known 3'-end processing elements (Figure 6A) (Tian and Graber 2011). The most commonly used *DEF1* pA¹ site in our previously reported RNA-Seq data was A+43 (Graber *et al.* 2013). Our identification of A+43G as a read-through mutant validates our genetic approach and defines region IV as a putative cleavage site (CS). Region II of the *DEF1* attenuator contains a consensus match to the yeast efficiency element (EE), and 5/22 unique DNA point mutations targeted the TATATA sequence. The mutations within




Figure 4 Trans-acting mutants *ma14*, *ma15*, *glc7*, *ssu72*, *sen1*, and *ctk1* result in Pol II read-through (>twofold) of the *DEF1* attenuator, representing a contribution of both NNS- and CPF-CF-dependent termination activity. (A)-(F) The *DEF1-lacZ* reporter and the control reporters *CYC1-lacZ* (CPF-CF-NNS hybrid terminator) and *SNR13-lacZ* (NNS terminator) were transformed into the indicated strains above. Cultures were grown at permissive temperature (25°C or 30°C) and diluted prior to shifting to 30°C for 2 hr and 37°C (non-permissive) for 2 hr. β -galactosidase activity was measured following cell lysis and incubation with ONPG substrate, using absorption at OD₆₀₀ for cell density and OD₄₂₀ for β -gal production. Experiments were performed in biological triplicate, and errors bars show standard deviation. (G) Total RNA was collected from indicated strains containing *DEF1-lacZ* grown at 30°C, and attenuated and read-through mRNAs were detected via RT-PCR as done in Fig. 2C. Negative control reactions lacking reverse transcriptase (-RT) were used to ensure that amplification was RNA-dependent.

the proposed EE include A-1G, which was the strongest and most commonly identified read-through mutant. In yeast, the UAUUAU element serves as an RNA binding site for Hrp1 (Chen and Hyman 1998; Valentini *et al.* 1999). Based on its position between the EE and CS, region III of the *DEF1* attenuator is likely to contain a positioning element (PE). Region III contains a partial match (AATTTA) to the consensus PE (AATAAA), and the A+11G and A+16G mutants alter this proposed PE. The PE serves as an RNA-binding site for Rna15 (Gross and Moore 2001). The mutations we have identified in the EE

and PE are consistent with *hrp1-1* and *rna15-58* mutants being defective for *DEF1* attenuator recognition (Figure 3G, 4B).

HRP1 overexpression suppresses read-through defects of cis-acting *DEF1* attenuator mutants

Our identification of a pA site efficiency element (EE) and our characterization of the *hrp1-1* mutant suggested that Hrp1 may recognize the *DEF1* attenuator. To test whether Hrp1 binds to the *DEF1* EE or other regions of the attenuator, we overexpressed *HRP1* by

mutant strain	Pol II Termination Pathway	CYC1-lacZ mt/WT ratio	DEF1-lacZ mt/WT ratio	SNR13-lacZ mt/WT ratio	DEF1 looks like?
<i>paf1Δ</i>	NNS	1.3	4.3	9.2	SNR13
<i>nrd1-5</i>	NNS	1.3	1.2	53.5	CYC1
<i>nab3-11</i>	NNS	1.0	2.0	124.3	CYC1
<i>pcf11-2</i>	NNS	1.1	1.0	9.5	CYC1
<i>pcf11-9</i>	NNS	0.9	1.3	15.3	CYC1
<i>pcf11-13</i>	NNS	0.8	1.1	11.9	CYC1
<i>hrp1-1</i>	CPF-CF	2.6	5.5	1.8	CYC1
<i>cft2-5001</i>	CPF-CF	3.4	3.6	1.5	CYC1
<i>rna14-5</i>	NNS/CPF-CF	6.3	51.3	165.2	SNR13
<i>rna15-58</i>	NNS/CPF-CF	3.2	10.2	5.4	BOTH
<i>glc7-12</i>	NNS/CPF-CF	4.3	9.4	48.1	CYC1
<i>ssu72-2</i>	NNS/CPF-CF	2.9	7.2	4.1*	BOTH
<i>sen1-1</i>	NNS/CPF-CF	3.1	5.5	41.3*	CYC1
<i>ctk1Δ</i>	NNS/CPF-CF	2.3	4.7	27.1	CYC1

Terminator read-through: little/no effect intermediate strong

Figure 5 Summary of *trans*-acting mutant effects on *DEF1* attenuator indicates similar behavior to *CYC1* hybrid CPF-CF-NNS terminator. The relative level of terminator read-through in mutant/WT was calculated from lacZ assays in Figs. 3 and 4 and summarized in the table (little/no effect: \leq twofold (pink); intermediate: 2-10 fold (yellow); strong: $>$ 10-fold (green)). Pol II termination pathways for each factor were assigned based on sensitivity of mutants to a known NNS-dependent terminator (*SNR13*) or hybrid CPF-CF-NNS-dependent terminator (*CYC1*). The asterisk indicates cases in which the kinetic slope of lacZ activity from the WT *SNR13*-lacZ reporter was undetectable, and an approximate value was utilized that was comparable to other experiments (40 B-gal units).

transforming *cis*-acting mutant strains with a plasmid version of *HRP1* in addition to the chromosome. We chose a low-copy (CEN) *HRP1* plasmid since we have previously observed that high-copy *HRP1* expression from a 2 μ plasmid is toxic (data not shown). We predicted that *HRP1* overexpression would enhance attenuator recognition, reduce *CUP1* reporter expression, and therefore increase copper sensitivity. As expected, the *CUP1* reporter lacking a terminator (No Term.) was copper-resistant, the wild-type *DEF1-CUP1* reporter was copper-sensitive, and *cis*-acting attenuator mutants conferred various degrees of copper resistance in the presence of the pRS314 empty vector control (Figure 7A). The A-1G, T-6C, and T-4C mutants were more copper-sensitive with *HRP1* overexpression (pRS314-*HRP1*) than an empty vector control, consistent with improved binding of Hrp1 to the mutant EE of region II. Surprisingly, the T-31C mutant was also more copper-sensitive with *HRP1* overexpression, suggesting that Hrp1 binds region I and/or that region I influences Hrp1 binding at region II. The effect of *HRP1* overexpression was allele-specific, with little-to-no genetic interaction observed between pRS314-*HRP1* and the *DEF1* attenuator mutants T+6C, A+16G, or A+43G in regions III and IV.

To confirm that increased copper-sensitivity was due to enhanced Pol II termination, we analyzed the ratio of attenuated:read-through transcripts by RT-PCR. As expected for enhanced attenuator recognition, *HRP1* overexpression increased attenuated RNA \sim twofold, resulting in less read-through mRNA compared to the empty vector control (Figure 7B). Taken together, our genetic analysis of the *DEF1* attenuator indicates that region II contains the EE and region IV contains the CS (Figure 7C). Region III is likely to contain the PE, and the role of region I is unclear but its activity is influenced by Hrp1.

Attenuator mutant results in *DEF1* overexpression and exacerbates toxicity of constitutively active Def1

Thus far, our characterization of the *DEF1* attenuator utilized a reporter gene construct, with the terminator positioned within an *ACT1* intron and under transcriptional control of a constitutive *TDH3* promoter. To study the *DEF1* attenuator in a more natural context, we cloned the full-length *DEF1* gene (promoter, 5'-UTR, ORF, 3'-UTR) into a high-copy 2 μ plasmid (pRS426) and transformed it into a *def1Δ* strain, where the plasmid was the sole source of *DEF1*. We designed measured *DEF1* attenuator activity based on the level of mRNA accumulation, using primers that amplified read-through RT-PCR products extending beyond pA¹ (Figure 8A). We observed a \sim 1.6 increase in read-through mRNA in the A-1G attenuator mutant compared to wild-type, indicating that the natural *DEF1* attenuator functions to suppress transcription (Figure 8B).

Given that Def1 protein expression exhibits post-translational regulation (Wilson *et al.* 2013; David 2013), we sought to establish biological significance for our observed transcriptional regulation. To test the biological significance of the *DEF1* attenuator in isolation from post-translational regulation we created a *def1* C1590A allele, which introduces a mutation (C1590A) that results in a premature stop codon (TAC \rightarrow TAA) (Figure 8A). The truncated version of Def1 (pr-Def1) produced from *def1* C1590A mimics a UV-dependent processing event (Wilson *et al.* 2013). At elevated temperatures, the pr-Def1 protein becomes activated via nuclear localization, triggers Pol II ubiquitination and degradation, and is toxic to cells. We predicted that attenuator mutations would elevate levels of *DEF1* mRNA and pr-Def1 protein, resulting in greater cell toxicity.




Figure 6 Cis-acting mutations result in Pol II read-through of the *DEF1* attenuator. (A) Schematic of *DEF1* attenuator sequences (boxed) that were fused to the *CUP1* reporter gene and used in a genetic selection for copper-resistant mutations. The individual point mutations resulting in a copper-resistant phenotype are numbered relative to +1 *ATG* start codon. Mutations that are underlined and in bold were the top 3 most frequent. Mutations were organized into regions I – IV based on clustering and similarity to pA site consensus elements. (B) Copper-resistant growth of yeast strains containing *DEF1-CUP1* reporter plated on $-/+$ copper plates after serial dilution. Higher levels of copper-resistance indicate higher levels of attenuator read-through (low-red, medium-yellow, high-green). (C) RNA analysis of *DEF1-CUP1* expression using RT-PCR (similar primers to Fig. 2C) to detect attenuated vs. full-length mRNA. (Top panel) The % attenuated vs. full-length RNA was determined by adding signal intensities together for both bands and determining the relative ratio. (Bottom panel) Negative control for RT-PCR reaction in which reverse transcriptase enzyme was withheld from the reaction.

We analyzed the growth of yeast strains containing an attenuator mutation (*def1* A-1G) in the context of pr-Def1 expression (*def1* C1590A). At the permissive temperature of 30°, all strains grew at a similar rate and density (Figure 8C). As expected, the *def1* C1590A mutation resulted in heat-sensitivity at 37° and 39°, similar to the *def1Δ* strain with an empty vector (Wilson *et al.* 2013). Transformation of *def1Δ* with pRS426-*DEF1* restored growth to wild-type levels, consistent with full complementation by the plasmid-based *DEF1* allele. The *def1* A-1G attenuator mutant grew similarly to wild-type, presumably because transcriptional overexpression was masked by post-translational protein control. Interestingly, the *def1* A-1G/C1590A double

mutant exhibited a synthetic sick phenotype at the non-permissive temperature of 37°. To better quantify this genetic interaction, we determined the growth rate of strains in liquid culture. The pRS426-*DEF1* strain had a doubling time of 2.2 hr and 2.5 hr, at 30° and 39° respectively. In contrast, the *def1* C1590A mutant had a doubling time of 3.0 hr and 5.3 hr, consistent with a heat-sensitive defect. The *def1* A-1G/C1590A mutant had a doubling time of 3.0 hr and 7.9 hr, exhibiting a more severe growth defect at 39° than the *def1* C1590A single mutant.

To confirm that *def1* A-1G enhanced *def1* C1590A toxicity due to pr-Def1 overexpression, we quantified mRNA and protein levels. The *def1* A-1G mutation increased mRNA expression by 1.6 compared to




Figure 7 Hrp1 overexpression partially restores recognition of *cis*-acting *DEF1* attenuator mutants. (A) Reporter strains containing WT or mutant *DEF1-CUP1* reporters were transformed with an empty vector (pRS314) or a plasmid containing *HRP1* (pRS314-*HRP1*) and grown on -Leu/Trp copper plates to assess *CUP1* expression. (B) Total RNA was collected from indicated strains grown at 30°C, and attenuated and read-through mRNAs were detected via RT-PCR. (C) Summary of genetic interactions between hyperactive *HRP1* allele and *cis*-acting *DEF1* attenuator mutants. The copper-sensitivity of mutants indicated in red was partially suppressed by *HRP1* overexpression, consistent with improved attenuator recognition and Hrp1-binding at RNA regions I and II.

def1 C1590 alone (Figure 8D). This mRNA overexpression correlated with ~twofold more Def1 protein in *def1* A-1G/C1590A compared to *def1* C1590A (Figure 8E). Overall, these data are consistent with Pol II attenuation contributing to *DEF1* regulation in yeast and serving an important biological function.

DISCUSSION

Termination is one of the least understood aspects of Pol II transcription, and a further knowledge gap exists for regulation by premature termination (attenuation). In this study we conducted a thorough characterization of an attenuator in the *DEF1* DNA repair gene, which bears unique features unseen in previous attenuator studies. The *DEF1* attenuator relies on a hybrid of termination factors for efficient recognition, with a bias for the CPF-CF pathway vs. the NNS pathway. We have identified nine termination factors

and four RNA sequence elements that contribute to attenuator activity, including the Hrp1 RNA-binding protein, a putative pA site efficiency element, and the region around the pA site itself. Furthermore, we have shown that disruption of the attenuator is biologically significant, supporting a new role for transcription attenuation in regulating a DNA damage response gene.

The *DEF1* attenuator exhibits a unique hybrid attenuator biased toward CPF-CF termination

Our mutational analysis indicates that *DEF1* attenuator recognition involves both CPF-CF and NNS termination pathways, but it is more reliant on CPF-CF recognizing a traditional pA site (Figure 6A; regions II, III, and IV). The importance of *DEF1* attenuator region I is unclear, but it may influence Hrp1 binding to region II, perhaps by forming a secondary structure. The *DEF1* attenuator exhibits Sen1-dependence but little-to-no dependence on Nrd1 or Nab3 despite sequence




Figure 8 The *def1* A-1G attenuator mutant increases Def1 mRNA and protein and reduces cell viability when overexpressing pr-Def1. (A) Schematic of the *DEF1* gene (not to scale). The relevant pA sites, efficiency element (EE), mutations (A-1G, C1590A), and RT-PCR primers are indicated. (B) Yeast strains containing chromosomal (CHR) *DEF1* or *def1*Δ were transformed with empty vector (pRS426), WT *DEF1* (pRS426-*DEF1*), or mutant (pRS426-*DEF1*, A-1G) plasmids. Total RNA was isolated, and Def1 read-through mRNA was detected via RT-PCR (using blue primers indicated in (A)). The signal intensity of the *DEF1* mRNA bands was normalized to the 18S loading control and then the *def1*Δ pRS426-*DEF1* sample. RT: Reverse Transcriptase. (C) Yeast strains containing WT chromosomal *DEF1* or *def1*Δ were transformed with empty vector (pRS426), WT *DEF1* (pRS426-*DEF1*), attenuator mutant (pRS426-*DEF1*, A-1G), pr-Def1 mutant (pRS426-*DEF1*-C1590A), or double mutant plasmids. Strains were spotted on -Ura plates and growth was assessed after 1 week at the indicated temperatures. (D) Total RNA was collected from strains in (C) containing *def1* mutants C1590A and C1590A/A-1G in a *def1*Δ strain grown at 30°C. *DEF1* read-through mRNA was detected via RT-PCR and quantified as in (B). (E) Western blot of extracts from strains in (C) following growth at 30°C and a temperature shift to 39°C for 0, 1, or 2 hr. Def1 protein levels were normalized to the actin loading control, and signal from *def1* C1590A/A-1G was normalized to *def1* C1590A at consistent time points.

similarity to consensus Nrd1 (GUAA, GUAG) and Nab3 RNA-binding sites (UCUU) (Steinmetz and Brow 1998; Carroll *et al.* 2004; Creamer *et al.* 2011; Porrua *et al.* 2012). The very limited role for Nrd1 and Nab3 at the *DEF1* attenuator is a contrast to what has been observed for most

other attenuators, including *NRD1*, *IMD2*, *URA2*, *FKS2*, *CLN3*, *GPH1*, and *GLT1*, which exhibit strong dependence on Nrd1, Nab3, or both (Arigo *et al.* 2006; Jenks *et al.* 2008; Kuehner and Brow 2008; Thiebaut *et al.* 2008; Kim and Levin 2011; Darby *et al.* 2012; Chen *et al.* 2017;

Merran and Corden 2017). The lack of Nrd1/Nab3 involvement that we observe is consistent with *DEF1* expression levels not increasing upon Nrd1/Nab3 depletion and Nrd1/Nab3 failing to crosslink to *DEF1* during *in-vivo* crosslinking studies (Jamonnak *et al.* 2011; Merran and Corden 2017).

DEF1 attenuator recognition is dependent on CFI component Hrp1 rather than Nrd1 or Nab3, which is somewhat surprising given its promoter-proximal location. In the reporter system, the TSS to EE distance is 365 bp, but this distance is only ~100 bp in the natural *DEF1* context, and we observe a termination defect in both cases with the *def1* A-1G mutant. At individual genes, Hrp1 has been shown to crosslink to coding regions ~twofold better than the promoter, and in some cases Hrp1 shows the strongest occupancy at the pA site near the 3'-ends of genes (Komarnitsky *et al.* 2000; Mayer *et al.* 2010). However, Hrp1 is proposed to bind within its own 5'-UTR as a mean of autoregulation, suggesting that it can act near gene promoters (Steinmetz *et al.* 2006b; Kuehner and Brow 2008; Chen *et al.* 2017). Furthermore, genome-wide crosslinking indicated that the majority of Hrp1 is bound to promoter-proximal regions of mRNAs (Tuck and Tollervey 2013), and transcriptome analysis in an *hrp1* mutant revealed sn/snoRNA termination defects on approximately one-third of the sn/snoRNA genes (Chen *et al.* 2017). Overall, these data indicate that Hrp1 acts more generally as both a CPF-CF and an NNS termination factor.

Another unique feature of the *DEF1* attenuator is its seeming lack of dependence on Pcf11. None of the *pcf11* mutants altered *DEF1* attenuator recognition despite the termination defects they exhibited for the *SNR13* control terminator. At other gene targets, the *pcf11-2* mutation impairs mRNA cleavage, the *pcf11-13* mutation impairs Pol II CTD-binding, and *pcf11-9* impairs both cleavage and Pol II CTD-binding. Accordingly, these mutants disrupt CPF-CF, NNS, or both termination pathways (Amrani *et al.* 1997; Sadowski *et al.* 2003; Kim *et al.* 2006; Grzechnik *et al.* 2015). The termination defect of *pcf11-13* corresponds with failed release of Nrd1, reduced Ser2 CTD phosphorylation, and lack of Sen1 recruitment (Grzechnik *et al.* 2015). Seemingly the *DEF1* attenuator does not require this Pcf11 function to elicit termination, perhaps because the Nrd1 and Nab3 proteins are not required either.

We have confirmed that *DEF1* attenuator recognition requires Hrp1, Rna14, Rna15, Ssu72, Ctk1, Glc7, Cft2, Paf1, and Sen1. Like Hrp1, several of these proteins may contribute to *DEF1* pA¹ site recognition. Rna15 and Rna14 are also members of CFI, and while Rna15 recognizes pA site positioning elements, Rna14 is capable of bridging Rna15 and Hrp1 in a CFI complex (Gross and Moore 2001; Barnwal *et al.* 2012). Cft2 is a component of the core CPF that binds the *CYC1* pA site *in vitro* and crosslinks near pA sites *in vivo*, as well as interacting with the Pol II CTD (Dichtl and Keller 2001; Kyburz *et al.* 2003; Baejen *et al.* 2014).

In lieu of direct *DEF1* pA¹ site recognition, some proteins may promote recruitment of termination factors. Ctk1 is a kinase that phosphorylates Ser2 residues of the Pol II CTD, an event that can occur relatively early in transcription and perhaps lead to promoter-proximal termination via recruitment of Sen1 (Mayer *et al.* 2010; Chinchilla *et al.* 2012; Lenstra *et al.* 2013). Glc7 and Ssu72 are components of the APT sub-complex of CPF, also termed the phosphatase module, which are required for termination of both mRNA and noncoding RNA genes (Mayfield *et al.* 2016; Casañal *et al.* 2017). Removal of Ser5-P from the Pol II CTD occurs via Ssu72, which is present at both the 3'-end of genes as well as promoter regions (Singh and Hampsey 2007; Zhang *et al.* 2012). Defects in Ssu72 increase Ser5-P, which may disrupt Ser2-P accumulation and Sen1:Pol II association. Glc7 promotes the removal of Tyr1-P from the Pol II CTD, and failure to remove Tyr1-P prevents recruitment of termination factors, including Nrd1, Pcf11, and Rtt103

(Mayer *et al.* 2012; Schrieck *et al.*, 2014). Paf1 has been implicated in both CPF-CF and NNS termination, and *paf1* mutants exhibit altered histone modification and reduced Ser2-P levels (Sheldon *et al.* 2005; Nordick *et al.* 2008; Tomson *et al.* 2011; Terzi *et al.* 2011). There is also precedence for Paf1 helping to recruit CPF to Ser5-P CTD, consistent with Paf1:Pol II enrichment on transcripts containing CPF proteins Cft2 and Mpe1 (Nordick *et al.* 2008; Fischl *et al.* 2017).

Sen1 may contribute most directly to termination at the *DEF1* attenuator via ATP-dependent RNA translocation and destabilization of paused Pol II, perhaps via melting of the RNA:DNA active site hybrid (Kim *et al.* 1999; Porrua and Libri 2013; Martin-Tumasch and Brow 2015; Han *et al.* 2017; Leonaitė *et al.* 2017). Given the limited role of Nrd1 and Nab3 in *DEF1* attenuation, Sen1 recruitment could occur instead through CPF component Glc7 or direct interaction with Ser2-P CTD, or connecting to CFI protein Hrp1 via CFI-CPF cross-factor interactions (Preker *et al.* 1995; Ohnacker *et al.* 2000; Kyburz *et al.* 2003; Holbein *et al.* 2011; Ghazy *et al.* 2012; Chinchilla *et al.* 2012).

Regulation of *DEF1* by dual transcriptional and post-translational mechanisms

Our evidence suggests that Pol II transcriptional attenuation contributes to biologically meaningful *DEF1* regulation in addition to a previously described post-translational mechanism (Wilson *et al.* 2013). Constitutive expression of a truncated protein that mimics Def1 activation is lethal to cells (Wilson *et al.* 2013), consistent with tight control of Def1 expression being vital for cell survival. *DEF1* expression and function contributes to several biological processes, including rescue of stalled Pol II, nucleotide excision repair, telomere maintenance, translesion synthesis, and Pol II initiation (Woudstra *et al.* 2002; Chen *et al.* 2005; Daraba *et al.* 2014; Damodaren *et al.* 2017). In addition, Def1 contributes resistance to DNA damage stress, salt stress, and heat shock stress (Woudstra *et al.* 2002; Vanacloig-Pedros *et al.* 2015; Damodaren *et al.* 2017). Given that both Def1 depletion and overexpression can be lethal, cells likely evolved multiple regulatory mechanisms to maintain *DEF1* homeostasis. An advantage of Pol II attenuator read-through vs. upregulation of initiation is that it could provide a more rapid response to an environmental stressor, akin to release of paused Pol II (Adelman and Lis 2012).

There is limited evidence to explain the mechanism by which Pol II attenuator read-through occurs for stress response genes. In the case of *IMD2*, depletion of the guanine nucleotide pool causes Pol II to shift from an upstream guanine TSS to a downstream adenine TSS, bypassing an NNS-dependent attenuator and allowing synthesis of full-length mRNA (Steinmetz *et al.* 2006b; Jenks *et al.* 2008; Kuehner and Brow 2008). The Levin and Manley labs have described an alternative mechanism for regulation of the *FKS2* attenuator. Cell wall damage activates the MAP kinase Mpk1, which associates with Pol II and Paf1, phosphorylates Tyr1 of the Pol II CTD, restricts Nrd1 recruitment, and allows Pol II to bypass an attenuator and terminate at a downstream pA site (Kim and Levin 2011; Yurko *et al.* 2017). We have identified several characteristics that are unique to *DEF1* vs. *IMD2* and *FKS2* attenuators, perhaps reflecting a novel mechanism for Pol II recognition and read-through. *DEF1* will be a useful model to expand our understanding of the signaling mechanisms that modulate Pol II termination, particularly during stress adaptation to DNA damage. It will also be of interest to explore connections between yeast Pol II attenuator function and alternative polyadenylation (APA) in metazoans. Splicing plays a dominant role in usage of promoter-proximal intronic pA sites, but upstream exonic pA sites remain relatively unexplored (Luo *et al.* 2013; Tian and Manley 2017).

ACKNOWLEDGMENTS

We thank Emmanuel College, the National Science Foundation (NSF-ROA to J. Kuehner and C. Moore, MCB-1244043), the Tri-Beta Biological Honor Society, and the National Institutes of Health (National Research Mentoring Network IRACDA+) awards to Tufts for funding; the David Brow, Jesper Svejstrup, David Levin, and Charlie Boone labs for reagents; and the Moore and Kuehner labs for helpful discussions. The authors declare that there are no conflicts of interest.

LITERATURE CITED

- Adelman, K., and J. T. Lis, 2012 Promoter-proximal pausing of RNA polymerase II: emerging roles in metazoans. *Nat. Rev. Genet.* 13: 720–731. <https://doi.org/10.1038/nrg3293>
- Amrani, N., M. Minet, F. Wyers, M. E. Dufour, L. P. Aggerbeck *et al.*, 1997 PCF11 encodes a third protein component of yeast cleavage and polyadenylation factor I. *Mol. Cell. Biol.* 17: 1102–1109. <https://doi.org/10.1128/MCB.17.3.1102>
- Arigo, J. T., K. L. Carroll, J. M. Ames, and J. L. Corden, 2006 Regulation of yeast NRD1 expression by premature transcription termination. *Mol. Cell* 21: 641–651. <https://doi.org/10.1016/j.molcel.2006.02.005>
- Arndt, K. M., and D. Reines, 2015 Termination of Transcription of Short Noncoding RNAs by RNA Polymerase II. *Annu. Rev. Biochem.* 84: 381–404. <https://doi.org/10.1146/annurev-biochem-060614-034457>
- Baejen, C., P. Torkler, S. Gressel, K. Essig, J. Söding *et al.*, 2014 Transcriptome maps of mRNP biogenesis factors define pre-mRNA recognition. *Mol. Cell* 55: 745–757. <https://doi.org/10.1016/j.molcel.2014.08.005>
- Barnwal, R. P., S. D. Lee, C. Moore, and G. Varani, 2012 Structural and biochemical analysis of the assembly and function of the yeast pre-mRNA 3' end processing complex CF I. *Proc. Natl. Acad. Sci. USA* 109: 21342–21347. <https://doi.org/10.1073/pnas.1214102110>
- Bernstein, J., and E. A. Toth, 2012 Yeast nuclear RNA processing. *World J. Biol. Chem.* 3: 7–26. <https://doi.org/10.4331/wjbc.v3.i1.7>
- Bowman, E. A., and W. G. Kelly, 2014 RNA polymerase II transcription elongation and Pol II CTD Ser2 phosphorylation: A tail of two kinases. *Nucleus* 5: 224–236. <https://doi.org/10.4161/nucl.29347>
- Carroll, K. L., D. A. Pradhan, J. A. Granek, N. D. Clarke, and J. L. Corden, 2004 Identification of cis elements directing termination of yeast non-polyadenylated snoRNA transcripts. *Mol. Cell. Biol.* 24: 6241–6252. <https://doi.org/10.1128/MCB.24.14.6241-6252.2004>
- Casañal, A., A. Kumar, C. H. Hill, A. D. Easter, P. Emsley *et al.*, 2017 Architecture of eukaryotic mRNA 3'-end processing machinery. *Science* 358: 1056–1059. <https://doi.org/10.1126/science.aao6535>
- Chen, S., and L. E. Hyman, 1998 A specific RNA-protein interaction at yeast polyadenylation efficiency elements. *Nucleic Acids Res.* 26: 4965–4974. <https://doi.org/10.1093/nar/26.21.4965>
- Chen, X., K. Poorey, M. N. Carver, U. Müller, S. Bekiranov *et al.*, 2017 Transcriptomes of six mutants in the Sen1 pathway reveal combinatorial control of transcription termination across the *Saccharomyces cerevisiae* genome. *PLoS Genet.* 13: e1006863. <https://doi.org/10.1371/journal.pgen.1006863>
- Chen, Y.-B., C.-P. Yang, R.-X. Li, R. Zeng, and J.-Q. Zhou, 2005 Def1p is involved in telomere maintenance in budding yeast. *J. Biol. Chem.* 280: 24784–24791. <https://doi.org/10.1074/jbc.M413562200>
- Chinchilla, K., J. B. Rodríguez-Molina, D. Ursic, J. S. Finkel, A. Z. Ansari *et al.*, 2012 Interactions of Sen1, Nrd1, and Nab3 with multiple phosphorylated forms of the Rpb1 C-terminal domain in *Saccharomyces cerevisiae*. *Eukaryot. Cell* 11: 417–429. <https://doi.org/10.1128/EC.05320-11>
- Colin, J., D. Libri, and O. Porrua, 2011 Cryptic transcription and early termination in the control of gene expression. *Genet. Res. Int.* 2011: 653494. <https://doi.org/10.4061/2011/653494>
- Contreras, X., M. Benkirane, and R. Kiernan, 2013 Premature termination of transcription by RNAP II: the beginning of the end. *Transcription* 4: 72–76. <https://doi.org/10.4161/trns.24148>
- Creamer, T. J., M. M. Darby, N. Jamonnak, P. Schaughency, H. Hao *et al.*, 2011 Transcriptome-wide binding sites for components of the *Saccharomyces cerevisiae* non-poly(A) termination pathway: Nrd1, Nab3, and Sen1. *PLoS Genet.* 7: e1002329. <https://doi.org/10.1371/journal.pgen.1002329>
- Damodaren, N., T. Van Eeuwen, J. Zamel, E. Lin-Shiao, N. Kalisman *et al.*, 2017 Def1 interacts with TFIIF and modulates RNA polymerase II transcription. *Proc. Natl. Acad. Sci. USA* 114: 13230–13235. <https://doi.org/10.1073/pnas.1707955114>
- Daraba, A., V. K. Gali, M. Halmaj, L. Haracska, and I. Unk, 2014 Def1 Promotes the Degradation of Pol3 for Polymerase Exchange to Occur During DNA-Damage-Induced Mutagenesis in *Saccharomyces cerevisiae*. *PLoS Biol.* 12: e1001771. <https://doi.org/10.1371/journal.pbio.1001771>
- Darby, M. M., L. Serebreni, X. Pan, J. D. Boeke, and J. L. Corden, 2012 The *Saccharomyces cerevisiae* Nrd1-Nab3 transcription termination pathway acts in opposition to Ras signaling and mediates response to nutrient depletion. *Mol. Cell. Biol.* 32: 1762–1775. <https://doi.org/10.1128/MCB.00050-12>
- David, R., 2013 Transcription: Proteasome power to Def1. *Nat. Rev. Mol. Cell Biol.* 14: 612. <https://doi.org/10.1038/nrm3664>
- Dheur, S., L. T. A. Vo, F. Voisinet-Hakil, M. Minet, J.-M. Schmitter *et al.*, 2003 Ptl1p and Ref2p found in association with the mRNA 3' end formation complex direct snoRNA maturation. *EMBO J.* 22: 2831–2840. <https://doi.org/10.1093/emboj/cdg253>
- Dichtl, B., and W. Keller, 2001 Recognition of polyadenylation sites in yeast pre-mRNAs by cleavage and polyadenylation factor. *EMBO J.* 20: 3197–3209. <https://doi.org/10.1093/emboj/20.12.3197>
- Fatica, A., M. Morlando, and I. Bozzoni, 2000 Yeast snoRNA accumulation relies on a cleavage-dependent/polyadenylation-independent 3'-processing apparatus. *EMBO J.* 19: 6218–6229. <https://doi.org/10.1093/emboj/19.22.6218>
- Fischl, H., F. S. Howe, A. Furger, and J. Mellor, 2017 Paf1 Has Distinct Roles in Transcription Elongation and Differential Transcript Fate. *Mol. Cell* 65: 685–698.e8. <https://doi.org/10.1016/j.molcel.2017.01.006>
- Ghazal, G., J. Gagnon, P.-E. Jacques, J.-R. Landry, F. Robert *et al.*, 2009 Yeast RNase III triggers polyadenylation-independent transcription termination. *Mol. Cell* 36: 99–109. <https://doi.org/10.1016/j.molcel.2009.07.029>
- Ghazy, M. A., J. M. B. Gordon, S. D. Lee, B. N. Singh, A. Bohm *et al.*, 2012 The interaction of Pcf11 and Clp1 is needed for mRNA 3'-end formation and is modulated by amino acids in the ATP-binding site. *Nucleic Acids Res.* 40: 1214–1225. <https://doi.org/10.1093/nar/gkr801>
- Goffeau, A., B. G. Barrell, H. Bussey, R. W. Davis, B. Dujon *et al.*, 1996 Life with 6000 genes. *Science* 274: 546–563–7. <https://doi.org/10.1126/science.274.5287.546>
- Graber, J. H., F. I. Nazeer, P.-C. Yeh, J. N. Kuehner, S. Borikar *et al.*, 2013 DNA damage induces targeted, genome-wide variation of poly(A) sites in budding yeast. *Genome Res.* 23: 1690–1703. <https://doi.org/10.1101/gr.144964.112>
- Gross, S., and C. L. Moore, 2001 Rna15 interaction with the A-rich yeast polyadenylation signal is an essential step in mRNA 3'-end formation. *Mol. Cell. Biol.* 21: 8045–8055. <https://doi.org/10.1128/MCB.21.23.8045-8055.2001>
- Grosso, A. R., A. P. Leite, S. Carvalho, M. R. Matos, F. B. Martins *et al.*, 2015 Pervasive transcription read-through promotes aberrant expression of oncogenes and RNA chimeras in renal carcinoma. *eLife Sciences* 4: 43. <https://doi.org/10.7554/eLife.09214>
- Grzechnik, P., M. R. Gdula, and N. J. Proudfoot, 2015 Pcf11 orchestrates transcription termination pathways in yeast. *Genes Dev.* 29: 849–861. <https://doi.org/10.1101/gad.251470.114>
- Guarente, L., and T. Mason, 1983 Heme regulates transcription of the CYC1 gene of *S. cerevisiae* via an upstream activation site. *Cell* 32: 1279–1286. [https://doi.org/10.1016/0092-8674\(83\)90309-4](https://doi.org/10.1016/0092-8674(83)90309-4)
- Gudipati, R. K., T. Villa, J. Boulay, and D. Libri, 2008 Phosphorylation of the RNA polymerase II C-terminal domain dictates transcription termination choice. *Nat. Struct. Mol. Biol.* 15: 786–794. <https://doi.org/10.1038/nsmb.1460>

- Han, Z., D. Libri, and O. Porrua, 2017 Biochemical characterization of the helicase Sen1 provides new insights into the mechanisms of non-coding transcription termination. *Nucleic Acids Res.* 45: 1355–1370. <https://doi.org/10.1093/nar/gkw1230>
- Holbein, S., S. Scola, B. Loll, B. S. Dichtl, W. Hübner *et al.*, 2011 The P-loop domain of yeast Clp1 mediates interactions between CF IA and CPF factors in pre-mRNA 3' end formation. *PLoS One* 6: e29139. <https://doi.org/10.1371/journal.pone.0029139>
- Hyman, L. E., S. H. Seiler, J. Whoriskey, and C. L. Moore, 1991 Point mutations upstream of the yeast ADH2 poly(A) site significantly reduce the efficiency of 3'-end formation. *Mol. Cell. Biol.* 11: 2004–2012. <https://doi.org/10.1128/MCB.11.4.2004>
- Jamonnak, N., T. J. Creamer, M. M. Darby, P. Schaugency, S. J. Wheelan *et al.*, 2011 Yeast Nrd1, Nab3, and Sen1 transcriptome-wide binding maps suggest multiple roles in post-transcriptional RNA processing. *RNA* 17: 2011–2025. <https://doi.org/10.1261/rna.2840711>
- Jenks, M. H., T. W. O'Rourke, and D. Reines, 2008 Properties of an intergenic terminator and start site switch that regulate IMD2 transcription in yeast. *Mol. Cell. Biol.* 28: 3883–3893. <https://doi.org/10.1128/MCB.00380-08>
- Jensen, T. H., A. Jacquier, and D. Libri, 2013 Dealing with pervasive transcription. *Mol. Cell* 52: 473–484. <https://doi.org/10.1016/j.molcel.2013.10.032>
- Keogh, M.-C., J.-A. Kim, M. Downey, J. Fillingham, D. Chowdhury *et al.*, 2006 A phosphatase complex that dephosphorylates gammaH2AX regulates DNA damage checkpoint recovery. *Nature* 439: 497–501. <https://doi.org/10.1038/nature04384>
- Kim, H. D., J. Choe, and Y. S. Seo, 1999 The sen1(+) gene of *Schizosaccharomyces pombe*, a homologue of budding yeast SEN1, encodes an RNA and DNA helicase. *Biochemistry* 38: 14697–14710. <https://doi.org/10.1021/bi991470c>
- Kim, K.-Y., and D. E. Levin, 2011 Mpk1 MAPK association with the Paf1 complex blocks Sen1-mediated premature transcription termination. *Cell* 144: 745–756. <https://doi.org/10.1016/j.cell.2011.01.034>
- Kim, M., L. Vasiljeva, O. J. Rando, A. Zhelkovsky, C. Moore *et al.*, 2006 Distinct Pathways for snoRNA and mRNA Termination. *Mol. Cell* 24: 723–734. <https://doi.org/10.1016/j.molcel.2006.11.011>
- Komarnitsky, P., E. J. Cho, and S. Buratowski, 2000 Different phosphorylated forms of RNA polymerase II and associated mRNA processing factors during transcription. *Genes Dev.* 14: 2452–2460. <https://doi.org/10.1101/gad.824700>
- Kuehner, J. N., and D. A. Brow, 2008 Regulation of a eukaryotic gene by GTP-dependent start site selection and transcription attenuation. *Mol. Cell* 31: 201–211. <https://doi.org/10.1016/j.molcel.2008.05.018>
- Kuehner, J. N., E. L. Pearson, and C. Moore, 2011 Unravelling the means to an end: RNA polymerase II transcription termination. *Nat. Rev. Mol. Cell Biol.* 12: 283–294. <https://doi.org/10.1038/nrm3098>
- Kwapisz, M., M. Wery, D. Després, Y. Ghavi-Helm, J. Soutourina *et al.*, 2008 Mutations of RNA polymerase II activate key genes of the nucleoside triphosphate biosynthetic pathways. *EMBO J.* 27: 2411–2421. <https://doi.org/10.1038/emboj.2008.165>
- Kyburz, A., M. Sadowski, B. Dichtl, and W. Keller, 2003 The role of the yeast cleavage and polyadenylation factor subunit Ydh1p/Cft2p in pre-mRNA 3'-end formation. *Nucleic Acids Res.* 31: 3936–3945. <https://doi.org/10.1093/nar/gkg478>
- Lenstra, T. L., A. Tudek, S. Clauder, Z. Xu, S. T. Pachis *et al.*, 2013 The role of Ctk1 kinase in termination of small non-coding RNAs. *PLoS One* 8: e80495. <https://doi.org/10.1371/journal.pone.0080495>
- Leonaitė, B., Z. Han, J. Basquin, F. Bonneau, D. Libri *et al.*, 2017 Sen1 has unique structural features grafted on the architecture of the Upf1-like helicase family. *EMBO J.* 36: 1590–1604. <https://doi.org/10.15252/embj.201696174>
- Lesser, C. F., and C. Guthrie, 1993 Mutational analysis of pre-mRNA splicing in *Saccharomyces cerevisiae* using a sensitive new reporter gene, CUP1. *Genetics* 133: 851–863.
- Loya, T. J., and D. Reines, 2016 Recent advances in understanding transcription termination by RNA polymerase II. *F1000 Res.* 5: 1478. <https://doi.org/10.12688/f1000research.8455.1>
- Luo, W., Z. Ji, Z. Pan, B. You, M. Hoque *et al.*, 2013 The conserved intronic cleavage and polyadenylation site of CstF-77 gene imparts control of 3' end processing activity through feedback autoregulation and by U1 snRNP. *PLoS Genet.* 9: e1003613. <https://doi.org/10.1371/journal.pgen.1003613>
- Martin-Tumasz, S., and D. A. Brow, 2015 *Saccharomyces cerevisiae* Sen1 Helicase Domain Exhibits 5' to 3' Helicase Activity with a Preference for Translocation on DNA Rather than RNA. *J. Biol. Chem.* 290: 22880–22889. <https://doi.org/10.1074/jbc.M115.674002>
- Mayer, A., M. Heidemann, M. Lidschreiber, A. Schrieck, M. Sun *et al.*, 2012 CTD tyrosine phosphorylation impairs termination factor recruitment to RNA polymerase II. *Science* 336: 1723–1725. <https://doi.org/10.1126/science.1219651>
- Mayer, A., M. Lidschreiber, M. Siebert, K. Leike, J. Söding *et al.*, 2010 Uniform transitions of the general RNA polymerase II transcription complex. *Nat. Struct. Mol. Biol.* 17: 1272–1278. <https://doi.org/10.1038/nsmb.1903>
- Mayfield, J. E., N. T. Burkholder, and Y. J. Zhang, 2016 Dephosphorylating eukaryotic RNA polymerase II. *Biochim. Biophys. Acta* 1864: 372–387. <https://doi.org/10.1016/j.bbapap.2016.01.007>
- Merran, J., and J. L. Corden, 2017 Yeast RNA-binding protein Nab3 regulates genes involved in nitrogen metabolism. *Mol Cell Biol* MCB.00154–17. <https://doi.org/10.1128/MCB.00154-17>
- Mischo, H. E., and N. J. Proudfoot, 2013 Disengaging polymerase: terminating RNA polymerase II transcription in budding yeast. *Biochim. Biophys. Acta* 1829: 174–185. <https://doi.org/10.1016/j.bbagr.2012.10.003>
- Morlando, M., P. Greco, B. Dichtl, A. Fatica, W. Keller *et al.*, 2002 Functional analysis of yeast snoRNA and snRNA 3'-end formation mediated by uncoupling of cleavage and polyadenylation. *Mol. Cell Biol.* 22: 1379–1389. <https://doi.org/10.1128/MCB.22.5.1379-1389.2002>
- Muhlrad, D., R. Hunter, and R. Parker, 1992 A rapid method for localized mutagenesis of yeast genes. *Yeast* 8: 79–82. <https://doi.org/10.1002/yea.320080202>
- Nagalakshmi, U., Z. Wang, K. Waern, C. Shou, D. Raha *et al.*, 2008 The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science* 5881: 1344–1349. <https://doi.org/10.1126/science.1158441>
- Natarajan, M., G. M. Schiralli Lester, C. Lee, A. Missra, G. A. Wasserman *et al.*, 2013 Negative elongation factor (NELF) coordinates RNA polymerase II pausing, premature termination, and chromatin remodeling to regulate HIV transcription. *J. Biol. Chem.* 288: 25995–26003. <https://doi.org/10.1074/jbc.M113.496489>
- Naville, M., and D. Gautheret, 2010 Transcription attenuation in bacteria: theme and variations. *Brief. Funct. Genomics* 9: 178–189. <https://doi.org/10.1093/bfpg/elq008>
- Neil, H., C. Malabat, Y. d'Aubenton-Carafa, Z. Xu, L. M. Steinmetz *et al.*, 2009 Widespread bidirectional promoters are the major source of cryptic transcripts in yeast. *Nature* 457: 1038–1042. <https://doi.org/10.1038/nature07747>
- Nordick, K., M. G. Hoffman, J. L. Betz, and J. A. Jaehning, 2008 Direct interactions between the Paf1 complex and a cleavage and polyadenylation factor are revealed by dissociation of Paf1 from RNA polymerase II. *Eukaryot. Cell* 7: 1158–1167. <https://doi.org/10.1128/EC.00434-07>
- Ohnacker, M., S. M. Barabino, P. J. Preker, and W. Keller, 2000 The WD-repeat protein pfs2p bridges two essential factors within the yeast pre-mRNA 3'-end-processing complex. *EMBO J.* 19: 37–47. <https://doi.org/10.1093/emboj/19.1.37>
- Ozsolak, F., P. Kapranov, S. Foissac, S. W. Kim, E. Fishilevich *et al.*, 2010 Comprehensive Polyadenylation Site Maps in Yeast and Human Reveal Pervasive Alternative Polyadenylation. *Cell* 143: 1018–1029. <https://doi.org/10.1016/j.cell.2010.11.020>
- Pearson, E., and C. Moore, 2014 The evolutionarily conserved Pol II flap loop contributes to proper transcription termination on short yeast genes. *Cell Reports* 9: 821–828. <https://doi.org/10.1016/j.celrep.2014.10.007>
- Porrua, O., and D. Libri, 2013 A bacterial-like mechanism for transcription termination by the Sen1p helicase in budding yeast. *Nat. Struct. Mol. Biol.* 20: 884–891. <https://doi.org/10.1038/nsmb.2592>

- Porrua, O., and D. Libri, 2015 Transcription termination and the control of the transcriptome: why, where and how to stop. *Nat. Rev. Mol. Cell Biol.* 16: 190–202. <https://doi.org/10.1038/nrm3943>
- Porrua, O., F. Hobor, J. Boulay, K. Kubicek, Y. D'Aubenton-Carafa *et al.*, 2012 In vivo SELEX reveals novel sequence and structural determinants of Nrd1-Nab3-Sen1-dependent transcription termination. *EMBO J.* 31: 3935–3948. <https://doi.org/10.1038/emboj.2012.237>
- Preker, P. J., J. Lingner, L. Minvielle-Sebastia, and W. Keller, 1995 The FIP1 gene encodes a component of a yeast pre-mRNA polyadenylation factor that directly interacts with poly(A) polymerase. *Cell* 81: 379–389. [https://doi.org/10.1016/0092-8674\(95\)90391-7](https://doi.org/10.1016/0092-8674(95)90391-7)
- Rondon, A. G., H. E. Mischo, and N. J. Proudfoot, 2008 Terminating transcription in yeast: whether to be a 'nerd' or a 'rat'. *Nat. Struct. Mol. Biol.* 15: 775–776. <https://doi.org/10.1038/nsmb0808-775>
- Rondón, A. G., H. E. Mischo, J. Kawauchi, and N. J. Proudfoot, 2009 Fail-safe transcriptional termination for protein-coding genes in *S. cerevisiae*. *Mol. Cell* 36: 88–98. <https://doi.org/10.1016/j.molcel.2009.07.028>
- Rutkowski, A. J., F. Erhard, A. L'Hernault, T. Bonfert, M. Schilhabel *et al.*, 2015 Widespread disruption of host transcription termination in HSV-1 infection. *Nat. Commun.* 6: 7126. <https://doi.org/10.1038/ncomms8126>
- Sadowski, M., B. Dichtl, W. Hübner, and W. Keller, 2003 Independent functions of yeast Pcf11p in pre-mRNA 3' end processing and in transcription termination. *EMBO J.* 22: 2167–2177. <https://doi.org/10.1093/emboj/cdg200>
- Schrieck, A., A. D. Easter, S. Etzold, K. Wiederhold, M. Lidschreiber *et al.*, 2014 RNA polymerase II termination involves C-terminal-domain tyrosine dephosphorylation by CPF subunit Glc7. *Nat. Struct. Mol. Biol.* 21: 175–179. <https://doi.org/10.1038/nsmb.2753>
- Schmidt, K., and J. S. Butler, 2013 Nuclear RNA surveillance: role of TRAMP in controlling exosome specificity. *WIREs RNA* 4: 217–231. <https://doi.org/10.1002/wrna.1155>
- Sheldon, K. E., D. M. Mauger, and K. M. Arndt, 2005 A Requirement for the *Saccharomyces cerevisiae* Paf1 complex in snoRNA 3' end formation. *Mol. Cell* 20: 225–236. <https://doi.org/10.1016/j.molcel.2005.08.026>
- Singh, B. N., and M. Hampsey, 2007 A transcription-independent role for TFIIB in gene looping. *Mol. Cell* 27: 806–816. <https://doi.org/10.1016/j.molcel.2007.07.013>
- Steinmetz, E. J., and D. A. Brow, 1998 Control of pre-mRNA accumulation by the essential yeast protein Nrd1 requires high-affinity transcript binding and a domain implicated in RNA polymerase II association. *Proc. Natl. Acad. Sci. USA* 95: 6699–6704. <https://doi.org/10.1073/pnas.95.12.6699>
- Steinmetz, E. J., and D. A. Brow, 1996 Repression of gene expression by an exogenous sequence element acting in concert with a heterogeneous nuclear ribonucleoprotein-like protein, Nrd1, and the putative helicase Sen1. *Mol. Cell Biol.* 16: 6993–7003. <https://doi.org/10.1128/MCB.16.12.6993>
- Steinmetz, E. J., and D. A. Brow, 2003 Ssu72 protein mediates both poly(A)-coupled and poly(A)-independent termination of RNA polymerase II transcription. *Mol. Cell Biol.* 23: 6339–6349. <https://doi.org/10.1128/MCB.23.18.6339-6349.2003>
- Steinmetz, E. J., N. K. Conrad, D. A. Brow, and J. L. Corden, 2001 RNA-binding protein Nrd1 directs poly(A)-independent 3'-end formation of RNA polymerase II transcripts. *Nature* 413: 327–331. <https://doi.org/10.1038/35095090>
- Steinmetz, E. J., S. B. H. Ng, J. P. Cloute, and D. A. Brow, 2006a cis- and trans-Acting determinants of transcription termination by yeast RNA polymerase II. *Mol. Cell Biol.* 26: 2688–2696. <https://doi.org/10.1128/MCB.26.7.2688-2696.2006>
- Steinmetz, E. J., C. L. Warren, J. N. Kuehner, B. Panbehi, A. Z. Ansari *et al.*, 2006b Genome-wide distribution of yeast RNA polymerase II and its control by Sen1 helicase. *Mol. Cell* 24: 735–746. <https://doi.org/10.1016/j.molcel.2006.10.023>
- Terzi, N., L. S. Churchman, L. Vasiljeva, J. Weissman, and S. Buratowski, 2011 H3K4 trimethylation by Set1 promotes efficient termination by the Nrd1-Nab3-Sen1 pathway. *Mol. Cell Biol.* 31: 3569–3583. <https://doi.org/10.1128/MCB.05590-11>
- Thibodeau, S. A., R. Fang, and J. K. Joung, 2004 High-throughput beta-galactosidase assay for bacterial cell-based reporter systems. *Biotechniques* 36: 410–415.
- Thiebaut, M., J. Colin, H. Neil, A. Jacquier, B. Séraphin *et al.*, 2008 Futile cycle of transcription initiation and termination modulates the response to nucleotide shortage in *S. cerevisiae*. *Mol. Cell* 31: 671–682. <https://doi.org/10.1016/j.molcel.2008.08.010>
- Tian, B., and J. H. Graber, 2012 Signals for pre-mRNA cleavage and polyadenylation. *WIREs RNA* 3: 385–396. <https://doi.org/10.1002/wrna.116>
- Tian, B., and J. L. Manley, 2017 Alternative polyadenylation of mRNA precursors. *Nat. Rev. Mol. Cell Biol.* 18: 18–30. <https://doi.org/10.1038/nrm.2016.116>
- Tomson, B. N., C. P. Davis, M. H. Warner, and K. M. Arndt, 2011 Identification of a Role for Histone H2B Ubiquitylation in non-coding RNA 3'-end Formation Through Mutational Analysis of Rtf1 in *Saccharomyces cerevisiae*. *Genetics* 188: 273–289. <https://doi.org/10.1534/genetics.111.128645>
- Tuck, A. C., and D. Tollervey, 2013 A transcriptome-wide atlas of RNP composition reveals diverse classes of mRNAs and lncRNAs. *Cell* 154: 996–1009. <https://doi.org/10.1016/j.cell.2013.07.047>
- Tudek, A., O. Porrua, T. Kabzinski, M. Lidschreiber, K. Kubicek *et al.*, 2014 Molecular basis for coordinating transcription termination with noncoding RNA degradation. *Mol. Cell* 55: 467–481. <https://doi.org/10.1016/j.molcel.2014.05.031>
- Valentini, S. R., V. H. Weiss, and P. A. Silver, 1999 Arginine methylation and binding of Hrp1p to the efficiency element for mRNA 3'-end formation. *RNA* 5: 272–280. <https://doi.org/10.1017/S1355838299981633>
- Van Oss, S. B., C. E. Cucinotta, and K. M. Arndt, 2017 Emerging Insights into the Roles of the Paf1 Complex in Gene Regulation. *Trends Biochem. Sci.* 42: 788–798. <https://doi.org/10.1016/j.tibs.2017.08.003>
- Vanacloig-Pedros, E., C. Bets-Plasencia, A. Pascual-Ahuir, and M. Proft, 2015 Coordinated gene regulation in the initial phase of salt stress adaptation. *J. Biol. Chem.* 290: 10163–10175. <https://doi.org/10.1074/jbc.M115.637264>
- Webb, S., R. D. Hector, G. Kudla, and S. Granneman, 2014 PAR-CLIP data indicate that Nrd1-Nab3-dependent transcription termination regulates expression of hundreds of protein coding genes in yeast. *Genome Biol.* 15: R8. <https://doi.org/10.1186/gb-2014-15-1-r8>
- Wilson, M. D., M. Harreman, M. Taschner, J. Reid, J. Walker *et al.*, 2013 Proteasome-mediated processing of def1, a critical step in the cellular response to transcription stress. *Cell* 154: 983–995. <https://doi.org/10.1016/j.cell.2013.07.028>
- Woudstra, E. C. E., C. C. Gilbert, J. J. Fellows, L. L. Jansen, J. J. Brouwer *et al.*, 2002 A Rad26-Def1 complex coordinates repair and RNA pol II proteolysis in response to DNA damage. *Nature* 415: 929–933. <https://doi.org/10.1038/415929a>
- Yurko, N., X. Liu, T. Yamazaki, M. Hoque, B. Tian *et al.*, 2017 MPK1/SLT2 Links Multiple Stress Responses with Gene Expression in Budding Yeast by Phosphorylating Tyr1 of the RNAP II CTD. *Mol. Cell* 68: 913–925.e3. <https://doi.org/10.1016/j.molcel.2017.11.020>
- Zhang, D. W., A. L. Mosley, S. R. Ramisetty, J. B. Rodríguez-Molina, M. P. Washburn *et al.*, 2012 Ssu72 phosphatase-dependent erasure of phospho-Ser7 marks on the RNA polymerase II C-terminal domain is essential for viability and transcription termination. *J. Biol. Chem.* 287: 8541–8551. <https://doi.org/10.1074/jbc.M111.335687>
- Zhang, Z., and F. Dietrich, 2005 Mapping of transcription start sites in *Saccharomyces cerevisiae* using 5' SAGE. *Nucleic Acids Res.* 33: 2838–2851. <https://doi.org/10.1093/nar/gki583>

Communicating editor: J. Rine